

# Energy Levels of Light Nuclei

## $A = 5$

F. Ajzenberg-Selove

*University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

**Abstract:** An evaluation of  $A = 5\text{--}10$  was published in *Nuclear Physics A320* (1979), p. 1. This version of  $A = 5$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed in 1978)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 5$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides:  ${}^5\text{n}$ ,  ${}^5\text{H}$ ,  ${}^5\text{He}$ ,  ${}^5\text{Li}$ ,  ${}^5\text{Be}$

B. Tables of Recommended Level Energies:

**Table 5.1:** Energy levels of  ${}^5\text{He}$

**Table 5.3:** Energy levels of  ${}^5\text{Li}$

C. References

D. Figures:  ${}^5\text{He}$ ,  ${}^5\text{Li}$ , Isobar diagram

E. Erratum to the Publication: [PS](#) or [PDF](#)

**$^5\text{n}$**   
(Not illustrated)

$^5\text{n}$  has not been observed: see ([1972AG01](#)) and ([1977DE08](#)).

**$^5\text{H}$**   
(Fig. 3)

From the work of ([1967AD05](#)) on the  $^3\text{He}(^3\text{He}, \text{n})^5\text{Be}$  reaction (see  $^5\text{Be}$ ) it follows that  $^5\text{H}$  is unstable by more than 2.1 MeV to decay into  $^3\text{H} + 2\text{n}$  (using Coulomb corrections based on the 16.7 MeV states in  $^5\text{He}-^5\text{Li}$ ). A study of  $^9\text{Be}(\alpha, ^8\text{B})^5\text{H}$  at  $E_\alpha = 129$  MeV shows no evidence for sharp  $^5\text{H}$  states for several MeV above  $^3\text{H} + 2\text{n}$  ([1968MC02](#)). In  $^3\text{H}(\text{t}, \text{p})^5\text{H}$ , at  $E_\text{t} = 22.25$  MeV, a broad peak appears in the proton spectrum which may correspond to a  $^5\text{H}$  state at  $^3\text{H} + 2\text{n} + 1.8$  MeV, but which could also result from a four-body breakup ([1968YO06](#)). A recent study of the  $^7\text{Li}(^6\text{Li}, ^8\text{B})^5\text{H}$  reaction at  $E(^6\text{Li}) = 93.3$  MeV shows no evidence for the formation of a narrow  $^5\text{H}$  ground state: at  $\theta_{\text{lab}} = 14.7^\circ$  the yield near the predicted location of the  $^5\text{H}_{\text{g.s.}}$ , above the phase-space curve, is  $\approx 100$  nb/sr. MeV compared to an order of magnitude greater yield for the final state interaction in  $^4\text{H}$  ([1977WE03](#), [1977WE1B](#)). Attempts to observe  $^5\text{H}$  formation in  $^7\text{Li} + \pi^-$  have also been unsuccessful: see ([1974AJ01](#)). See also ([1975BE31](#); theor.).

**$^5\text{He}$**   
(Figs. 1 and 3)

GENERAL: (See also ([1974AJ01](#))).

*Model calculations:* ([1974JA30](#), [1974LE22](#), [1975DI04](#)).

*Special states:* ([1974GO13](#), [1974IR04](#), [1974JA30](#), [1974LE22](#), [1975DI04](#), [1976IR1B](#)).

*Special reactions:* ([1973FE1A](#), [1974FE1A](#), [1975FE1A](#), [1976VA29](#), [1978ME1C](#)).

*Reactions involving pions:* ([1973BA30](#), [1978FI1E](#)).

*Other topics:* ([1974GO13](#), [1974IR04](#), [1976BI1A](#), [1976IR1B](#), [1977SH1B](#), [1978GO1D](#)).

*Ground state of  $^5\text{He}$ :* ([1975BE31](#), [1977HI09](#)).

$$1. \ ^3\text{H}(\text{d}, \gamma)^5\text{He} \quad Q_m = 16.70$$

At low energies the reaction is dominated by a resonance at  $E_\text{d} = 107$  keV; the mirror reaction shows resonance at  $E_\text{d} = 430$  keV. ([1969BE56](#)) have measured the cross section for emission of 16.7 MeV  $\gamma$ -rays for  $E_\text{d} = 25$  to 100 keV: the ratio  $\sigma(\text{d}, \gamma)/\sigma(\text{d}, \text{n})$  is approximately constant at

Table 5.1: Energy levels of  ${}^5\text{He}$ <sup>a</sup>

$E_x$ (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	$0.60 \pm 0.02$ <sup>a</sup>	n, $\alpha$	1, 4, 6, 8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 20, 21, 22, 23, 26, 27, 29, 30
4 $\pm$ 1	$\frac{1}{2}^-; \frac{1}{2}$	4 $\pm$ 1	n, $\alpha$	4, 6, 10, 18, 22
16.76 $\pm$ 0.13	$\frac{3}{2}^+; \frac{1}{2}$	0.10 $\pm$ 0.05	$\gamma$ , n, d, t, $\alpha$	1, 2, 5, 6, 8, 9, 11, 13, 14, 15, 21, 22
19.8 $\pm$ 0.4 <sup>b</sup>	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	2.5 $\pm$ 0.5	n, d, t, $\alpha$	2, 3, 5, 8, 9, 10, 13, 14, 15, 19, 21, 22
24 – 25 <sup>b</sup>		broad		21, 22

<sup>a</sup> See Table 5.2 in (1966LA04) and reactions 6 and 20 here.

<sup>b</sup> See (1974AJ01), pp. 7 – 8.

$(2.1 \pm 0.6) \times 10^{-4}$ , leading to  $\Gamma_\gamma = 14 \pm 4$  eV, where  $\Gamma_n$  is taken as 66 keV. (1963BU07) have measured thick target yields from  $E_d = 150$  to 1300 keV. The derived cross sections are analyzed into resonant and direct-capture contributions: the cross section at resonance is here reported as 60  $\mu\text{b}$  [versus 1 mb reported by (1969BE56)]. At  $E_d = 1025 \pm 47$  keV, the differential cross section is  $0.44 \pm 0.12 \mu\text{b}/\text{sr}$  ( $90^\circ$ ) and the  $\gamma$  to n branching ratio is an order of magnitude smaller:  $2.3 \times 10^{-5}$ . The angular distribution of the  $\gamma$ -rays is forward peaked and the total cross section is estimated to be 4.8  $\mu\text{b}$  (1970KO09). The cross section for  $\gamma_0$  has been measured (at  $90^\circ$ ) for  $E_d = 2.0$  to 12.0 MeV and angular distributions were obtained at  $E_d = 4, 8, 10$  and 12 MeV for  $\gamma_0$  (1975BA1E; and C.A. Barnes, private communication). See also (1974AJ01).

- |  |                 |               |
|--|-----------------|---------------|
| 2. (a) ${}^3\text{H}(d, n){}^4\text{He}$ | $Q_m = 17.5894$ | $E_b = 16.70$ |
| (b) ${}^3\text{H}(d, 2n){}^3\text{He}$   | $Q_m = -2.9884$ |               |
| (c) ${}^3\text{H}(d, pn){}^3\text{H}$    | $Q_m = -2.2246$ |               |

Below  $E_d = 100$  keV, the cross section for reaction (a) follows the Gamow function,  $\sigma = (A/E)\exp(-44.40E^{-1/2})$  (1953JA1A, 1954AR02). A strong resonance,  $\sigma(\text{peak}) = 5.0$  b, appears at  $E_d = 107$  keV: see Table 5.2. From  $E_d = 10$  to 500 keV, the cross section is well fitted with the assumption of s-wave formation of a  $J^\pi = \frac{3}{2}^+$  state (1952AR30, 1952CO35, 1955KU03). Excitation curves and angular distributions for reaction (a) have been measured from  $E_d = 8$  keV to 21 MeV: see (1974AJ01) for earlier references and (1976DR1A, 1976DR1B; preliminary work; angular distributions at  $E_d = 7.00$  and 10.00 MeV, 13.36 and 16.50 MeV (partial) and at

Table 5.2: Resonance parameters for the  $\frac{3}{2}^+$  states observed in  ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$  and  ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ <sup>a</sup>

$E_r$ (keV)	$\Gamma_{\text{lab}}$ (keV)	$l_{\text{d}}$	$J^\pi$	$l_{\text{np}}$	$R$ (fm)	$E_\lambda$ (keV)	$\gamma_{\text{d}}^2$ (keV)	$\gamma_{\text{np}}^2$ (keV)	$\theta_{\text{d}}^2$ <sup>d</sup>	$\theta_{\text{np}}^2$ <sup>d</sup>	$E_x$ (MeV)
107 <sup>b</sup>	135	0	$\frac{3}{2}^+$	2	5.0	-464	$2000 \pm 500$	$50 \pm 10$	1.0	0.018	16.76
					7.0	-126	715	17	0.7	0.011	
450 <sup>c</sup>	$\approx 450$	0	$\frac{3}{2}^+$	2	5.0	-391	2930	42	1.4	0.013	16.66
					7.0	129	780	12	0.7	0.008	

<sup>a</sup> See also (1974KR07, 1976AH1A, 1977BO24, 1977DO01).

<sup>b</sup>  ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$ .

<sup>c</sup>  ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ .

<sup>d</sup> Units of  $3\hbar^2/2MR^2$ .

$E_t = 10.48$  and  $14.98$  MeV;  $0^\circ$  excitation function for  $E_d = 7$  to  $16.5$  MeV), (1975MA28; angular distributions at  $E_d = 8$  to  $16$  MeV, in  $1$  MeV steps;  $0^\circ$  excitation function for same interval) and (1977JA07;  $d\sigma/d\Omega$  at  $E_t = 20.00$  MeV; very accurate).


A study of reaction (a) with polarized deuterons at  $E_d = 0.2$  to  $1.0$  MeV indicates intervention of the s-wave,  $J^\pi = \frac{1}{2}^+$  channel, as well as possible p-waves above  $E_d = 0.3$  MeV (1965TR01, 1971GR32). At higher energies, the neutron polarization  $P_1(\theta_1)$  shows an angular distribution that peaks typically at  $\theta_1 = 30^\circ$  lab, then goes negative with increasing angle and has a minimum between  $90^\circ$  and  $130^\circ$ , depending upon energy. (1971MU04) have made an extensive study of  $P_1(30^\circ)$  for  $E_d = 5$  to  $15$  MeV and, with deuterium as target, for  $E_t = 4.5$  to  $19.5$  MeV (neutrons near  $135^\circ$  lab). The polarization increases monotonically from  $0.03$  at  $E_d = 3$  MeV to  $\approx 0.5$  at  $E_d = 6.5$  MeV and then with a lower slope to  $0.69$  at  $E_d = 13$  MeV. The change in the slope may be caused by excited states of  ${}^5\text{He}$  near  $20$  MeV. Comparison with the  ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$  mirror reaction at corresponding c.m. energies shows excellent agreement between the polarization values in the two reactions up to  $E_d = 6$  MeV, but then the proton polarization becomes  $\approx 15\%$  higher, converging back to the neutron values at  $E_d \approx 12 - 13$  MeV. This may be due to experimental factors (1971MU04). Using polarized deuterons (1971HI07) find that the average ratio of vector analyzing power of  ${}^3\text{He}(\text{d}, \text{p})$  to  ${}^3\text{H}(\text{d}, \text{n})$  is  $1.016 \pm 0.015$  at  $E_d = 6$  MeV and  $1.035 \pm 0.020$  at  $10$  MeV. The vector analyzing power of the two reactions agree to within  $\pm 0.025$  at all angles. This agreement between the mirror reactions is in apparent conflict with the result of (1971MU04). For earlier polarization measurements see Table 5.3 in (1974AJ01). Polarization studies are also reported by (1976SU06;  $E_d = 7$  MeV), (1976LI15;  $E_d = 3.5$  to  $12.8$  MeV), (1976OH02;  $E_t = 5.37$  to  $10.50$  MeV) and (1975SA1A; preliminary;  $E_d = 38$  MeV; looked at pol. of  $50$  MeV neutrons). See also (1977DR1B). The  $K_y'(0^\circ)$  determined for  $E_t = 5.37$  to  $10.50$  MeV show a minimum at  $E_t = 9.6$  MeV (1976OH02). See also (1978CA13;  $E_d = 0.7$  MeV).

Reaction (b) has been studied for  $E_d = 10.9$  to  $83$  MeV: see (1974AJ01). A discussion and comparison of attempts to measure  ${}^1\text{S}_0$  nn scattering length is presented by (1971GR45;  $E_d = 13.4$

MeV): based on kinematically complete measurements and using the Watson-Migdal treatment, they find  $a_{nn} = -16.0 \pm 1.0$  fm. Intervention of n- $^3$ He final state interactions is believed to be small in this work ([1971GR45](#)). At  $E_t = 22$  MeV no clear evidence is seen for sequential processes via excited states of  $^4$ He ([1971GR12](#)).

A reanalysis of the work of ([1963PO02](#)) on reaction (c) by ([1974SC04](#)) leads to the suggestion of a resonance at  $E_{c.m.} = 2.9 \pm 0.3$  MeV [ $E_x = 19.6$  MeV],  $\Gamma_{c.m.} = 1.9 \pm 0.2$  MeV, consistent with  $J^\pi = \frac{3}{2}^-$  [see, however, Table [5.1](#)]. n-p final-state interaction enhancement has been studied by ([1973SL03](#)) at  $E_d = 35$  MeV. They find that only when  $E_{np} \leq 1$  MeV is in agreement with the Watson-Migdal theory obtained, even if data far from dominant n-t final-state interaction and p-t quasi-free scattering are chosen. See also ([1974AL01](#)).

See also ([1974AD11](#)), ([1974GE1B](#), [1974WI1D](#), [1975CR1A](#), [1975ST1A](#), [1975ST1B](#), [1976BL1A](#), [1976BR1B](#), [1976JA1C](#), [1976MO1A](#), [1977BE1H](#), [1977KI1C](#), [1977MC1C](#); applied topics), ([1973LI1B](#), [1975KU1C](#), [1975SE07](#), [1975SO1C](#), [1976HA1C](#), [1976SE1B](#), [1976WA1C](#), [1977SE09](#); reviews), ([1975FO19](#); astrophysics) and ([1974KE1B](#), [1974ST14](#), [1975DO1B](#); theor.).



$$E_b = 16.70$$

The elastic scattering has been studied for  $E_d = 2.6$  to  $11.0$  MeV. The excitation curves show an interference at  $E_x \approx 19$  MeV and a broad ( $\Gamma > 1$  MeV) resonance corresponding to  $E_x = 20.0 \pm 0.5$  MeV, similar to that seen in  $^3\text{He}(\text{d}, \text{d})$  [see  $^5\text{Li}$ ]. Together with data from  $^3\text{H}(\text{d}, \text{n})^4\text{He}$ , this work favors an assignment  $D_{3/2}$  or  $D_{5/2}$  with a mixture of doublet and quartet components (channel spin  $\frac{1}{2}$  and  $\frac{3}{2}$ ) if only one state is involved ([1967TO02](#), [1968IV01](#)) [any appreciable doublet component would, however, be in conflict with results from  $^7\text{Li}(\text{p}, ^3\text{He})^5\text{He}$ ]. Measurements of differential cross section and analyzing power using polarized deuterons with  $E_{\bar{d}} = 3.2$  to  $12.3$  MeV show resonance-like behavior in the vector analyzing power near  $E_d = 5$  MeV. The anomaly appears in the odd Legendre coefficients and is interpreted in terms of a  $(\frac{1}{2}, \frac{3}{2})^-$  excited state of  $^5\text{He}$  with  $E_x \approx 19.6$  MeV. Broad structure in the differential cross section near  $6$  MeV, principally in the even Legendre coefficients, corresponds to an even-parity state  $^5\text{He}^*(20.0)$  ([1971KI02](#)). Tensor analyzing powers have been measured for  $E_d = 5.0$  to  $11.5$  MeV ([1973DE51](#)), at  $E_t = 10.5$  MeV ([1976OH02](#)) and at  $E_{\bar{d}} = 11.4$  and  $14.4$  MeV ([1974DO05](#); angular distributions of tensor and vector polarizations). See also ([1974ST14](#), [1975AB1C](#), [1975DO1B](#), [1976BA1E](#); theor.).



At  $E_t = 0.5$  MeV, the reaction appears to proceed via three channels: (i) direct breakup into  $^4\text{He} + 2\text{n}$ , the three-body breakup shape being modified by the n-n interaction; (ii) sequential decay via  $^5\text{He}(0)$ ; (iii) sequential decay via a broad excited state of  $^5\text{He}$ . The branching ratios at  $\theta = 90^\circ$  are  $0.7 : 0.2 : 0.1$ . The width of  $^5\text{He}(0)$  is estimated to be  $0.74 \pm 0.18$  MeV. Some evidence is

also shown for  ${}^5\text{He}^*$  at  $E_x \approx 2$  MeV,  $\Gamma \approx 2.4$  MeV ([1965WO03](#)). For reaction (b), see  ${}^6\text{He}$  and ([1974LA02](#)).

5. (a) ${}^3\text{He}(t, p){}^5\text{He}$	$Q_m = 11.20$
(b) ${}^3\text{He}(t, pn){}^4\text{He}$	$Q_m = 12.0959$

Some evidence is reported at  $E_t = 22.25$  MeV in reaction (a) for a broad state of  ${}^5\text{He}$  at  $E_x \approx 20$  MeV, in addition to a sharp peak corresponding to  ${}^5\text{He}^*(16.7)$  ([1968YO06](#)). For reaction (b) see  ${}^6\text{Li}$  and ([1975SC28](#), [1977DEYM](#)).



The coherent scattering length (thermal, bound) is  $3.0 \pm 0.1$  fm ([1973MU14](#)). The thermal scattering cross section is  $0.773 \pm 0.009$  b and the absorption cross section at 2200 m/sec  $\sigma_{n,\gamma}^0$  is  $< 0.05$  b ([1969RO16](#)). ([1973MU14](#)) adopt  $0.76 \pm 0.01$  b for  $\bar{\sigma}_s$ . Total cross sections for  $E_n = 4 \times 10^{-4}$  eV to 150.9 MeV and at 10 GeV/c are summarized in ([1974AJ01](#), [1976GAYV](#)). Angular distribution measurements are summarized in ([1970GA1A](#), [1974AJ01](#)). Polarization studies are displayed in Table 5.4 of ([1974AJ01](#)): recent work includes that of ([1976BO05](#):  $E_n = 1.5$  to 6.0 MeV; used an absolutely calibrated source of polarized neutrons), ([1976LI15](#):  $E_n = 20$  to 30 MeV;  $A_y(\theta)$  is the same for  ${}^4\text{He} + n$  and  ${}^4\text{He} + p$ ) and ([1978YO1A](#):  $E_n = 50$  MeV). See also ([1974TO03](#)).

The total cross section has a peak of 7.6 b ([1973GO38](#)) at  $E_n = 1.15 \pm 0.05$  MeV,  $E_{\text{c.m.}} = 0.92 \pm 0.04$  MeV, with a width of about 1.2 MeV ([1964ST25](#)). A second resonance is observed at  $E_n = 22.15 \pm 0.12$  MeV, corresponding to the 16.7 MeV  $J^\pi = \frac{3}{2}^+$  state ([1959BO54](#), [1964SH30](#)):  $\Gamma_{\text{c.m.}} = 100 \pm 50$  keV,  $\Gamma_n = \Gamma_d = 50 \pm 35$  keV ([1960HU08](#)). [[1966HO07](#)] find that the data are fitted best by  $\Gamma_n < \Gamma_d$  although  $\Gamma_n < \Gamma_d$  is not excluded]. Attempts to detect additional resonances in the total cross section have been unsuccessful: see ([1966LA04](#)).

The  $P_{3/2}$  phase shift shows strong resonance behavior near 1 MeV, while the  $P_{1/2}$  shift changes more slowly, indicating a broad  $P_{1/2}$  level at several MeV excitation ([1952DO30](#), [1966HO07](#), [1970AR1B](#), [1972ST01](#)). ([1966HO07](#)) have constructed a set of phase shifts for  $E_n = 0$  to 31 MeV,  $l = 0, 1, 2, 3$ , using largely p- $\alpha$  phase shifts. They have measured differential cross sections from 6 to 30 MeV, with special attention to the region near 22.15 MeV and have fitted the data with the assumed phase shifts. At the  $\frac{3}{2}^+$  state the best fit to all data is given by  $E_{\text{res}} = 17.669$  MeV  $\pm 10$  keV,  $\gamma_d^2 = 2.0$  MeV  $\pm 25\%$ ,  $\gamma_n^2 = 50$  keV  $\pm 25\%$  (see Table 5.2). The polarization has been calculated from the phase shifts and is presented as a contour plot ([1966HO07](#)). The work of ([1976LI15](#)) indicates some discrepancies with the results of ([1966HO07](#)) [below  $E_n = 22$  MeV] and with the measurements of ([1972BR10](#)).

A recent  $R$ -function analysis of the  ${}^4\text{He} + n$  data below 21 MeV (including the absolute neutron analyzing power measurement of ([1976BO05](#)) and the accurate cross section measurements of

(1973GO38)) has led to a set of phase shifts and analyzing powers which are based on the  ${}^4\text{He} + \text{n}$  data alone (rather than also including the  ${}^4\text{He} + \text{p}$  data). At  $r = 3.3$  fm the values obtained for the  $P_{1/2}$  and  $P_{3/2}$  resonances are, respectively,  $E_{\text{c.m.}} = 1.97$  and  $0.77$  MeV,  $\Gamma_{\text{c.m.}} = 5.22$  and  $0.64$  MeV (1977BO24). See also (1974KR07). For earlier analyses, see (1974AJ01).

See also (1974MA1C, 1977YO1B), (1975FI1B, 1976OH1B, 1976WA1B, 1976WA1C) and (1974BA34, 1974DO10, 1974GH01, 1974JA1F, 1974ST14, 1975BA76, 1975CA05, 1975LO1A, 1975TH12, 1976AH1A, 1976AL1E, 1976CH1B, 1976SP1A, 1977BA1N, 1977BE50, 1977CH1C, 1977LA1C, 1977LE1D, 1977PH01, 1977ST1D, 1977TH09, 1978AH01, 1978FO1G, 1978FR02, 1978KO13, 1978LE1H, 1978MA37; theor.).

$$7. {}^4\text{He}(\text{n}, \text{d}){}^3\text{H} \quad Q_{\text{m}} = -17.5894 \quad E_{\text{b}} = -0.89$$

See (1976BR1B), (1974AJ01) and (1976BA1E; theor.).

$$\begin{array}{lll} 8. (\text{a}) {}^4\text{He}(\text{d}, \text{p}){}^5\text{He} & Q_{\text{m}} = -3.12 \\ (\text{b}) {}^4\text{He}(\text{d}, \text{pn}){}^4\text{He} & Q_{\text{m}} = -2.2246 \end{array}$$

A typical proton spectrum consists of a peak corresponding to formation of the ground state of  ${}^5\text{He}$ , plus a lower continuum of protons ascribed to deuteron breakup (reaction (b)). Ground-state protons show pronounced azimuthal asymmetry when the reaction is induced by 8.5, 10 and 11 MeV vector polarized deuterons. A DWBA calculation is in only qualitative agreement (1967TR05, 1971KE16).

Reaction (b) has been studied for  $E = 9$  to  $165$  MeV: see (1974AJ01) and  $E_{\text{d}} = 6.8$  and  $7.8$  MeV (1978KA11),  $8.9$  MeV (1977SA21), at  $E_{\bar{\text{d}}} = 15$  MeV (1978NA08),  $E_{\alpha} = 21.9$  and  $23.7$  MeV (1974RA10),  $22.5$  MeV (1977PR06),  $27.2$  MeV (1974GR40) and  $165$  MeV (1974HO11). See also  ${}^6\text{Li}$ .

At  $E_{\text{d}} = 8.9$  MeV, the FSI (final state interaction) is very important (1977SA21). In the kinematically complete experiment of (1977PR06) at  $E_{\alpha} = 22.5$  MeV, the results are in quite good agreement with the predictions of the sequential decay model: the slight deviations are assumed to be due to the omission of Coulomb interaction effects and to the restriction to partial waves with  $l \leq 2$ . At  $E_{\alpha} = 70$  MeV, a kinematically complete experiment shows evidence for sequential decay, proceeding through excited states of  ${}^5\text{He}$ . Peaks in the coincident yield of protons and deuterons are ascribed to narrow states at  $E_{\text{x}} = 16.7 \pm 0.1$  MeV,  $\Gamma = 80 \pm 30$  keV, at  $E_{\text{x}} = 18.6 \pm 0.1$ ,  $18.8 \pm 0.1$  and  $19.2 \pm 0.1$  MeV, all with  $\Gamma = 180 \pm 60$  keV (1973TR04). The fine structure near 19 MeV is not confirmed in other experiments [see, however, reaction 12].

See also (1974HE21, 1974NA10, 1975HE09, 1976KO21, 1976LI1E, 1977BR25, 1978FI1E, 1978KO13),  ${}^5\text{Li}$  and  ${}^6\text{Li}$ .



At  $E_\alpha = 70$  MeV, there is indication of the population of  ${}^5\text{He}^*(0, 16.7)$  and of a structure with  $E_x = 18 - 20$  MeV ([1974LA1A](#), [1975AL1A](#); preliminary results).



At  $E_\gamma = 60$  MeV, the proton spectrum shows two prominent peaks attributed to  ${}^5\text{He}^*(0 + 4.0, 20 \pm 2)$  ([1973GA16](#), [1976MA34](#)). See also ([1973DE17](#)), ([1975MA1E](#)) and  ${}^6\text{Li}$ .



At  $E_p = 1180$  MeV, the exciation of  ${}^5\text{He}^*(0, 16.7)$  is reported: the latter state is formed with the ejection of an s-proton ([1972AN24](#), [1972AN27](#), [1972AN29](#)). See also ([1975ME27](#); theor.).



Angular distributions of ground state deuterons have been studied at  $E_n = 6.57$  and  $6.77$  MeV ([1977RO01](#)),  $14.4$  MeV ([1965VA05](#)) and  $56.3$  MeV ([1977BR17](#)). In the latter experiment, angular distributions have also been obtained to  ${}^5\text{He}^*(16.7, 18.5 \pm 0.5, 20.5 \pm 0.5)$ . The observation of the two highest states is not certain: if they exist their widths are less than the instrumental width,  $1.6$  MeV ([1977BR17](#)). For reactrion (b) see ([1967VA12](#)). See also ([1974AJ01](#)).



See ([1976HUZK](#)).



At  $E_p = 100$  MeV the population of  ${}^5\text{He}^*(0, 16.7)$  and possibly of a broad structure at  $E_x \approx 19$  MeV is observed: momentum distributions for  ${}^5\text{He}^*(0, 16.7)$  and angular correlation measurements are also reported. The main features of the data are reasonably well described by DWIA ([1974BH03](#)). For the earlier work see ([1974AJ01](#)). See also ([1975VO04](#), [1977RO1E](#)) and ([1974GH03](#), [1974PR10](#), [1975CH1C](#), [1976OH04](#); theor.).

15. (a) ${}^6\text{Li}(\text{d}, {}^3\text{He}) {}^5\text{He}$	$Q_m = 0.90$
(b) ${}^6\text{Li}(\text{d}, \text{n} {}^3\text{He}) {}^4\text{He}$	$Q_m = 1.795$

At  $E_d = 14.5$  MeV,  ${}^5\text{He}_{\text{g.s.}}$  is populated ([1955LE24](#)). At  $E_d = 80$  MeV, the population of  ${}^5\text{He}^*(16.7)$  as well as a broad state near  $E_x = 19$  MeV is reported ([1975DI1A](#), [1975DI1B](#); abstracts). For reaction (b) see ([1974AJ01](#)).

16. ${}^6\text{Li}(\text{t}, \alpha) {}^5\text{He}$	$Q_m = 15.22$
---	---------------

See ([1974AJ01](#)).

17. (a) ${}^6\text{Li}(\alpha, {}^5\text{Li}) {}^4\text{He} + \text{n}$	$Q_m = -5.66$
(b) ${}^6\text{Li}(\alpha, \alpha \text{p}) {}^5\text{He}$	$Q_m = -4.59$

See ([1975GR42](#)) for reaction (a) and ([1978CA1E](#)) for reaction (b).

18. ${}^7\text{Li}(\gamma, \text{d}) {}^5\text{He}$	$Q_m = -9.62$
---	---------------

See ([1975DE37](#)) in  ${}^7\text{Li}$ .

19. (a) ${}^7\text{Li}(\pi^+, 2\text{p}) {}^5\text{He}$	$Q_m = 128.51$
(b) ${}^7\text{Li}(\pi^-, 2\text{n}) {}^5\text{He}$	$Q_m = 126.94$

The two-proton spectrum shows broad structures attributed to the ground state of  ${}^5\text{He}$ , and to  $\text{p}^{-1}\text{s}^{-1}$  and  $\text{s}^{-2}$  states at  $\approx 20$  and ( $\approx 35$ ) MeV excitation ([1965CH12](#)). For reaction (b) see ([1977BA1M](#)).

20. (a) ${}^7\text{Li}(\text{n}, \text{t}) {}^5\text{He}$	$Q_m = -3.36$
(b) ${}^7\text{Li}(\text{n}, \text{tn}) {}^4\text{He}$	$Q_m = -2.467$

The angular distribution of the  $t_0$  group has been measured at  $E_t = 14.4$  MeV: it is fairly well reproduced by DWBA ([1970MI05](#)). Reaction (b) at  $E_n = 14.4$  MeV proceeds as a sequential process primarily involving  ${}^7\text{Li}^*(4.63)$  and  ${}^5\text{He}_{\text{g.s.}}$  ([1974AN02](#)). See also ([1974TU1A](#)) and ([1974AJ01](#)) for the earlier work.

21. (a) ${}^7\text{Li}(\text{p}, {}^3\text{He}){}^5\text{He}$	$Q_m = -4.12$
(b) ${}^7\text{Li}(\text{p}, \text{pd}){}^5\text{He}$	$Q_m = -9.618$

At  $E_p = 43.7$  MeV, angular distributions of the  ${}^3\text{He}$  groups to the ground state of  ${}^5\text{He}$  ( $\Gamma = 0.80 \pm 0.04$  MeV) and to levels at 16.7 and  $19.9 \pm 0.4$  MeV ( $\Gamma = 2.7$  MeV) have been determined. The angular distribution of the  ${}^5\text{He}$  ground state group indicates substantial mixing of  $L = 0$  and  $L = 2$  transfer. The distribution to  ${}^5\text{He}^*(16.7)$  is consistent with  $L = 1$ . Since no transitions are observed in the  ${}^7\text{Li}(\text{p}, \text{t}){}^5\text{Li}$  reaction to the analogue 20 MeV state in  ${}^5\text{Li}$  [see  ${}^5\text{Li}$ ], the transition is presumably  $S$ -forbidden and the states in  ${}^5\text{He}-{}^5\text{Li}$  near 20 MeV are  ${}^4D_{3/2}$  or  ${}^4D_{5/2}$  [compare  ${}^3\text{H}(\text{d}, \text{d})$ ] ([1966CE05](#)). Particle-particle coincidence data have been obtained at  $E_p = 43.7$  MeV. They suggest the existence of  ${}^5\text{He}^*(20.0)$  with  $\Gamma = 3.0 \pm 0.6$  MeV and of a broad state at  $\approx 25$  MeV. No  $T = \frac{3}{2}$  states decaying via  $T = 1$  states in  ${}^4\text{He}$  were observed ([1968MC02](#)). In reaction (b) at  $E_p = 100$  MeV  ${}^5\text{He}^*(0, 19.9)$  are strongly populated while  ${}^5\text{He}^*(16.7)$  is very weak. Positive parity is suggested for  ${}^5\text{He}^*(19.9)$  ([1975CH1B](#); preliminary).

22. (a) ${}^7\text{Li}(\text{d}, \alpha){}^5\text{He}$	$Q_m = 14.23$
(b) ${}^7\text{Li}(\text{d}, \text{n}){}^4\text{He}{}^4\text{He}$	$Q_m = 15.123$

At  $E_d = 24$  MeV, the  $\alpha$ -particle spectrum from reaction (a) shows structures corresponding to the ground and 16.7 MeV states and to states at  $E_x \approx 20.2$  and 23.8 MeV with  $\Gamma \approx 2$  MeV and  $\approx 1$  MeV respectively ([1972BA30](#)). The width of the ground state of  ${}^5\text{He}$  is  $615 \pm 30$  keV ([1973GA28](#)). Angular distributions are reported for  $E_d = 0.6$  to 1.25 MeV.


Spectra measured in reaction (b) suggest the involvement of the  $P_{1/2}$  state: the values suggested are inconsistent with each other because of the difficulty of evaluating the contribution of other reactions: see ([1974AJ01](#)) for the earlier values and see ([1976FO21](#)) for a discussion of some of the problems involved in these. ([1976FO21](#)) suggest a width for  ${}^5\text{He}_{\text{g.s.}} = 0.6$  MeV and  $E_x = 4.1 \pm 0.2$ ,  $\Gamma_{\text{c.m.}} = 4.4 \pm 0.2$  MeV for the  $P_{1/2}$  state. Reaction (b) proceeds mainly via excited states of  ${}^8\text{Be}$  and  ${}^5\text{He}_{\text{g.s.}}$ : see, e.g., ([1967VA11](#)). See also ([1973HU12](#), [1974DA28](#), [1974GR44](#), [1978AR10](#), [1978SP03](#)). Attempts to observe n- $\alpha$  rescattering effects following formation of  ${}^8\text{Be}^*(16.63, 16.91)$  have been unsuccessful: an upper limit of 1% of the yield from sequential decay is given by ([1968VA12](#)). See ([1974AJ01](#)) for more complete references to the older work. See also  ${}^8\text{Be}$  and  ${}^9\text{Be}$ .

23. ${}^7\text{Li}({}^3\text{He}, \text{d}{}^3\text{He}){}^5\text{He}$	$Q_m = -9.62$
--	---------------

See ([1976WA12](#)).

24. ${}^7\text{Li}(\alpha, {}^6\text{Li}){}^4\text{He} + \text{n}$	$Q_m = -7.251$
--	----------------

See (1975GR42).



See  ${}^9\text{Be}$ .



Studies of this reaction have been carried out at  $E_p = 26.0$  to  $57$  MeV [see (1974AJ01)] and at  $E_p = 100$  MeV (1977RO02). In the latter experiment good agreement is found with DWIA calculations: the average value of  $S_\alpha = 0.45 \pm 0.02$  (1977RO02). See also  ${}^9\text{Be}$ .



See (1975VO08).



See  ${}^9\text{Be}$ .



Reaction (a) has been studied at  $E_\alpha = 28$  to  $37.4$  MeV: see (1974AJ01). See also (1974GR42, 1977BR1E). Reaction (b) has been studied at  $E_\alpha = 65$  MeV: only the ground state of  ${}^5\text{He}$  is observed for  $E_x \lesssim 25$  MeV;  $S_\alpha = 0.53$  (1976WO11).



At  $E_d = 3.75$  MeV (1978GR07),  $10.4$  and  $12.0$  MeV (1971RE19), this reaction involves  ${}^5\text{He}_{g.s.}$  and states in  ${}^8\text{Be}$  and  ${}^9\text{Be}$  (1971RE19, 1978GR07) and  ${}^{12}\text{C}$  (1978GR07).



See  ${}^{12}\text{C}$  in (1975AJ02).

**$^5\text{Li}$**   
(Figs. 2 and 3)

GENERAL: (See also (1974AJ01).)

*Model calculations:* (1975KR1A).

*Special states:* (1974GO13, 1974IR04, 1976IR1B).

*Astrophysical questions:* (1974RA1C, 1978ME1C).

*Special reactions:* (1975BR1A, 1976VA29, 1978ME1C).

*Reactions involving pions:* (1973AR1B, 1974AM01).

*Applied topics:* (1975HU1A).

*Other topics:* (1974GO13, 1974IR04, 1976IR1B, 1978GO1D).

*Ground state of  ${}^5\text{Li}$ :* (1975BE31).

$$1. \ ^3\text{He}(\text{d}, \gamma){}^5\text{Li} \quad Q_m = 16.39$$

Excitation curves and angular distributions have been measured for  $E_{\text{d}} = 0.2$  to 5 MeV and  $E({}^3\text{He}) = 2$  to 26 MeV: see (1974AJ01).

A broad maximum in the cross section is observed at  $E_{\text{d}} = 0.45 \pm 0.04$  MeV [ ${}^5\text{Li}^*(16.66)$ ]:  $\sigma = 50 \pm 10 \mu\text{b}$ ,  $\Gamma_\gamma = 11 \pm 2$  eV (1954BU06:  $\gamma_0 + \gamma_1$ );  $\sigma_{\gamma_0} = 21 \pm 4 \mu\text{b}$ ,  $\Gamma_{\gamma_0} = 5 \pm 1$  eV (1968BU09). The radiation at resonance is isotropic, consistent with s-wave capture: see (1954BU06, 1968BU09, 1968KR03). Study of  $\gamma_0$  and  $\gamma_1$  yield  $\Gamma = 2.6 \pm 0.4$  MeV for the ground state width, and  $E_x = 7.5 \pm 1.0$  MeV,  $\Gamma = 6.6 \pm 1.2$  MeV for the  $\frac{1}{2}^-$  state. The ratio of  $\gamma_0$  to  $\gamma_1$  is  $1.00 \pm 0.2$  and  $1.9 \pm 0.4$  at  $E_{\text{d}} = 480$  and 1025 keV (1968BU09).

An excess in the cross section at higher bombarding energies is interpreted by (1972KI01) as being due to a state at  $E_x \approx 18$  MeV: even parity is deduced from the relative intensity of  $\gamma_0$  and  $\gamma_1$ . It is presumed to be the  $\frac{1}{2}^+$  state reported in reactions 3 and 7. A broad peak is also observed at  $E_x \approx 20.7$  MeV in the  $\gamma_0$  cross section. The cross section for  $\gamma_1$  is  $\approx 0$ . The observations are consistent with  $J^\pi = \frac{5}{2}^+$ : angular distributions appear to require at least one other state with significant strength near 19 MeV (1972KI01). (1970SC18) also report this state but find  $E_x = 19.7 \pm 0.2$  MeV [ $\Gamma = 5.0$  MeV]:  $(2J+1)\Gamma_{\gamma_0} = 1.32$  keV ( $\pm 50\%$ ), suggesting E1 (if only one state is involved).

$$2. \ ^3\text{He}(\text{d}, \text{n}){}^4\text{Li} \quad E_b = 16.39$$

Table 5.3: Energy levels of  ${}^5\text{Li}$  <sup>a</sup>

$E_x$ (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	$\approx 1.5$	p, $\alpha$	1, 5, 6, 7, 9, 13, 14, 15, 16, 17
5 – 10	$\frac{1}{2}^-; \frac{1}{2}$	$5 \pm 2$	p, $\alpha$	1, 5, 7, 13, 15, 16, 17
$16.66 \pm 0.07$	$\frac{3}{2}^+; \frac{1}{2}$	$\approx 0.3$	$\gamma, p, d, {}^3\text{He}, \alpha$	1, 3, 4, 7, 13, 15
$(18 \pm 1)$	$(\frac{1}{2}^+); \frac{1}{2}$	broad	$\gamma, p, d, {}^3\text{He}, \alpha$	1, 3, 7
$(20.0 \pm 0.5)$	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	$\approx 5$	$\gamma, p, d, {}^3\text{He}, \alpha$	1, 3, 4, 5, 7, 13, 15
a				

<sup>a</sup> See also discussion in reactions 3, 4, 7, 14 and 15.

This reaction has not been observed: see (1966LA04) and (1974POZN; abstract). See also (1978MC1C).

- |   |                 |               |
|---|-----------------|---------------|
| 3. (a) ${}^3\text{He}(d, p){}^4\text{He}$ | $Q_m = 18.3532$ | $E_b = 16.39$ |
| (b) ${}^3\text{He}(d, np){}^3\text{He}$   | $Q_m = -2.2246$ |               |
| (c) ${}^3\text{He}(d, 2p){}^3\text{H}$    | $Q_m = -1.4608$ |               |
| (d) ${}^3\text{He}(d, 2d){}^1\text{H}$    | $Q_m = -5.4936$ |               |

Below 100 keV the cross section follows the simple Gamow form:  $\sigma = (18.2 \times 10^3 / E) \exp(-91E^{-1/2})$  b ( $E$  in keV) (1953JA1A, 1954AR02). The zero-energy cross section factor  $S_0 = 6700 \text{ keV} \cdot \text{b}$  (1964PA1A). A pronounced resonance occurs at  $E_d = 430 \text{ keV}$ ,  $\Gamma \approx 450 \text{ keV}$ . The peak cross section is given as  $695 \pm 14 \text{ mb}$  (1952BO68, 1955KU03): see Table 5.2. Excitation functions for ground state protons have also been reported for  $E({}^3\text{He}) = 0.39$  to  $1.46 \text{ MeV}$  and  $18.7$  to  $44.1 \text{ MeV}$  and for  $E_d = 2.8$  to  $17.8 \text{ MeV}$ : see (1974AJ01). Angular distributions have been measured for  $E_d = 0.25$  to  $27 \text{ MeV}$  and  $E({}^3\text{He}) = 18.7$  to  $44.1 \text{ MeV}$  [see Table 5.6 in (1974AJ01)]. At  $E_d = 10.02 \text{ MeV}$  differential cross sections have been measured at four angles to better than 3% (1974JA15). Polarization measurements have been carried out at many energies: see Table 5.6 in (1974AJ01) and Table 5.4 here.

Below  $E_d = 1.1 \text{ MeV}$ , the polarization analyzing power of the reaction is minimally affected by p-wave contributions (less than a few percent). However, at  $E_d = 500 \text{ keV}$ , the analyzing power is reduced by  $(12 \pm 5)\%$  by intervention of the s-wave  $J^\pi = \frac{1}{2}^+$  channel (1966BR02, 1967MC01, 1971LE27). Contour maps,  $T_{ab}$  versus  $\theta, E_d$ , for the vector and tensor analyzing powers are presented for  $E_d = 0$  to  $12 \text{ MeV}$  by (1971GR47). Energy dependence of the angular distributions indicate resonance-like behavior at  $E_x = 16.6, 17.5, 20.0, 20.9$  and  $22.4 \text{ MeV}$  in  ${}^5\text{Li}$  (1971GR47). At  $E_d \approx 6 \text{ MeV}$  (1971KL02) find the angular distribution more complex than could be accounted for by a single  ${}^2D_{3/2}$  or  ${}^2D_{5/2}$  level at  $E_x \approx 20 \text{ MeV}$  [see  ${}^3\text{He}(d, d)$ ]. For

Table 5.4: Polarization measurements in  ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ <sup>a</sup>

$E_{\bar{\text{d}}} (\text{MeV})$	Refs.
0.344, 0.465, 0.727	(1974GA21)
0.34 – 11.60	(1976SC15)
0.75, 1.5 <sup>b</sup>	(1976BR1C) <sup>c</sup>
1 – 6	(1976DRZU, 1977DR1B) <sup>c</sup>
6.44 <sup>d</sup>	(1977ST06)
6.6 – 15.8	(1974TR02)
8.5 – 10.5 <sup>e</sup>	(1976GR10)
12 <sup>b,f</sup>	(1974BE67)
15 <sup>g</sup>	(1976ME13)
15 – 40	(1975RO1C, 1976RO1G) <sup>c</sup>
27 <sup>h</sup>	(1976OK1A) <sup>c</sup>
30, 35, 40 <sup>i</sup>	(1976RO1H) <sup>c</sup>

<sup>a</sup> See Table 5.6 in (1974AJ01) for a listing of the earlier work.

<sup>b</sup>  $E_{\text{d}}$ .

<sup>c</sup> Preliminary results.

<sup>d</sup> Measurement to better than 1% of  $iT_{11}$ .

<sup>e</sup> Maximum value of the tensor analyzing power ( $A_{yy}$ ) at 9.28 MeV is nearly 1.

<sup>f</sup>  ${}^3\vec{\text{He}}$  target.

<sup>g</sup> Reactions 3(b) and 3(c).

<sup>h</sup>  $E({}^3\vec{\text{He}})$ .

<sup>i</sup> Reaction 3(b).

$E_d$ (c.m.) = 7 to 18 MeV, differential cross sections fall monotonically with no indication of states of  ${}^5\text{Li}$  with  $23.4 < E_x < 34$ , with  $\Gamma > 0.8$  MeV and with appreciable widths for decay into  ${}^3\text{He} + \text{d}$  ([1972KI02](#)).

Table 5.5: Measurements of elastic scattering and polarization in  ${}^4\text{He}(p, p){}^4\text{He}$  <sup>a</sup>

Elastic scattering angular distribution and cross sections		Polarization	
$E_p$ (MeV)	Refs.	$E_{\vec{p}}$ (MeV)	Refs.
0.5 – 3.0	( <a href="#">1974KR07</a> )	3.6, 6.35, 12.2	( <a href="#">1974LI1C</a> , <a href="#">1976LIZV</a> )
11.16, 12.04, 13.60, 14.23	( <a href="#">1977DO01</a> )	3.74 – 8.94	( <a href="#">1976BR17</a> )
85.0	( <a href="#">1974VO05</a> )	6.36	( <a href="#">1977ME06</a> )
156	( <a href="#">1975CO27</a> )	11.93, 17.00	( <a href="#">1977HA06</a> )
185 – 500	( <a href="#">1978MC06</a> )	45.04, 52.34	( <a href="#">1977SA1E</a> )
200, 350, 500	( <a href="#">1977ST30</a> )	185 – 500	( <a href="#">1978MC06</a> )
350, 650, 1050, 1150	( <a href="#">1977AS01</a> )	560, 800, 1030,	( <a href="#">1977KL03</a> )
438, 648, 1036	( <a href="#">1976BE45</a> )	1270, 1730	
$[E_\alpha = 4.00, 5.08, 6.09 \text{ GeV}/c]$			
560, 800, 1030, 1270, 1730	( <a href="#">1977KL08</a> )		
580, 720	( <a href="#">1975VE09</a> )		
600	( <a href="#">1976FA09</a> )		
788	( <a href="#">1978FO33</a> )		
1000	( <a href="#">1977AL1E</a> )		
1050	( <a href="#">1974BA14</a> , <a href="#">1978BE30</a> )		
$E_\alpha = 7.0 \text{ GeV}/c$	( <a href="#">1977GE01</a> )		
$[E_p = 1050]$			
2680	( <a href="#">1975IG1B</a> , <a href="#">1977IG1A</a> , <a href="#">1977IG1B</a> , <a href="#">1977IG1C</a> )		
4.89 GeV	( <a href="#">1978NA1H</a> )		
50 – 500 GeV/c	( <a href="#">1978SM1C</a> )		

<sup>a</sup> See also Tables 5.5 in ([1966LA04](#)) and 5.7 in ([1974AJ01](#)) for a listing of the earlier work.

Reactions (b), (c) and (d) have been studied at  $E_d = 22.3$  and 35 MeV and at  $E({}^3\text{He}) = 30$ , 33.5 and 52.5 MeV by ([1977SL04](#)) and analyzed with a PWIA: Fourier transforms of the wave functions were obtained. At  $E({}^3\text{He}) = 35.9$  MeV, the spectra in reactions (b) and (c) are dominated by the nucleon-nucleon FSI: the results were fitted with a fully antisymmetrized PWBA and with DWBA ([1974WA06](#), [1975WA31](#)). At  $E_{\bar{d}} = 15$  MeV ([1976ME13](#)) have measured the vector and the two tensor analyzing powers for reactions (b) and (c). ([1974SC04](#), [1976SC26](#)) have studied the excitation function for reaction (c) for  $E_d = 2.2$  to 6 MeV in a kinematically complete experiment. They have extracted the p + t FSI going via  ${}^4\text{He}^*(20.1)$  [ $J^\pi = 0^+$ ] and suggest that the reaction goes primarily via a  $J^\pi = \frac{3}{2}^-$ ,  $T = \frac{1}{2}$  state of  ${}^5\text{Li}$  located  $0.8 \pm 0.2$  MeV above threshold [i.e.,  $E_x = 18.9 \pm 0.2$  MeV]. This suggests that the attraction of a  $p_{3/2}$  nucleon to  ${}^4\text{He}^*(0^+)$  is stronger

than is the attraction of such a nucleon to  ${}^4\text{He}_{\text{g.s.}}$ . See also (1975RU04) for reaction (c). See (1974AJ01) for the earlier work.

See also (1974TA1A, 1975GL08, 1976GO1B), (1974LA1C, 1974PR1B, 1975HE1C, 1975HU1A; applied topics), (1975SE07, 1975SE1D, 1975SE1E, 1976SE03, 1976SE1B, 1976SE1C, 1977SE09, 1977SE1C, 1978FI1E, 1978SL1B; reviews) and (1974AL01, 1974CH02, 1974DO10, 1974KE1B, 1975DO1B, 1975NE11, 1975YA12, 1977CH1D, 1977CO1B, 1978KO13, 1978SE01; theor.).



In the range  $E_{\text{d}} = 380$  to  $570$  keV, the scattering cross section is consistent with s-wave formation of the  $J^\pi = \frac{3}{2}^+$  state at  $16.66$  MeV (1954BR05). The excitation curves for  $E_{\text{d}} = 1.96$  to  $10.99$  MeV show a broad resonance ( $\Gamma > 1$  MeV) corresponding to  $E_x = 20.0 \pm 0.5$  MeV. From the behavior of the angular distributions an assignment of  ${}^2D_{3/2}$  or  $({}^2D, {}^4D)_{5/2}$  is favored, if only one state is involved (1967TO02). [There is some evidence that there is more than one  $D$ -wave state in this  $E_x$  region: see reaction 3]. See also (1976AL22:  $E_{\text{d}} = 1.01$  to  $2.95$  MeV). In the range  $E_{\text{d}}(\text{c.m.}) = 7$  to  $18$  MeV differential cross sections show only a monotonic variation with energy. There is no evidence for any other resonances from  $E_x = 23$  to  $34$  MeV (1972BA30, 1972KI02). Angular distributions have been measured at a number of energies from  $E({}^3\text{He}) = 18.7$  to  $44.1$  MeV (1972KI02) and at  $E({}^3\text{He}) = 10.00$  MeV (1974JA15,  $d\sigma/d\Omega$  to  $\pm 1\%$ ) and  $E_{\text{d}} = 39$  MeV (1975RU04).

Polarization measurements have been reported for  $E_{\text{d}} = 4.0$  to  $12$  MeV [see (1974AJ01)] and polarization analyzing powers have been studied at  $E_{\bar{\text{d}}} = 2.0$  to  $11.5$  MeV (1975JE1A; abstract),  $E_{\bar{\text{d}}} = 3.73$  to  $11.89$  MeV (1974OH05;  ${}^3\vec{\text{He}}$ ; also spin correlation parameters),  $E_{\text{d}} = 5.35$  to  $9.2$  MeV (1974LO06; polarization transfer coefficients),  $E_{\bar{\text{d}}} = 15$  to  $40$  MeV (1976RO1C, 1976ROZM; vector analyzing powers) and  $E({}^3\vec{\text{He}}) = 33$  MeV (1977KA10).

See also (1975TO1A) and (1973HA1F, 1974CH02, 1974CH1E, 1974LY03, 1975DO1B, 1975HA1E, 1975KA1D, 1975NE11, 1975TA1A, 1976DO1B, 1976LY1A, 1977KA1H, 1977SA1C; theor.).



Angular distributions have been measured at  $E({}^3\text{He}) = 2.0$  to  $5.5$  MeV for  $n_0$  ( $\Gamma_{\text{g.s.}} = 2.03 \pm 0.15$  MeV) and at  $3.5$  to  $5.5$  MeV (partial distributions) for  $n_1$  to an excited state at  $E_x = 10.21 \pm 0.28$  MeV,  $\Gamma = 1.49 \pm 0.61$  MeV (1975AB11). At  $E({}^3\text{He}) = 14$  to  $26$  MeV, the spectra show the  $n_0$  group and a broad resonance with  $E_x = 20.5 \pm 0.8$  MeV (1974CH15). Reaction (b) has been studied at  $E({}^3\text{He}) = 0.31$  to  $2.80$  to attempt to detect a possible resonant energy dependence of the spin-singlet ( $T = 1$ ) n-p FSI: a marked effect is not present (1975SC28). See also (1977DEYM) and  ${}^6\text{Li}$ .

6. (a) ${}^3\text{He}({}^3\text{He}, \text{p}){}^5\text{Li}$	$Q_m = 10.89$
(b) ${}^3\text{He}({}^3\text{He}, 2\text{p}){}^4\text{He}$	$Q_m = 12.8596$
(c) ${}^3\text{He}({}^3\text{He}, 3\text{p}){}^3\text{H}$	$Q_m = -6.9544$

The spectrum of protons shows a pronounced peak at  $E({}^3\text{He}) = 3$  to 18 MeV corresponding to  ${}^5\text{Li}_{\text{g.s.}}$  superposed on a continuum: see (1974AJ01). At  $E({}^3\text{He}) = 13.6$  MeV differential cross sections and polarizations have been measured for the  $p_0$  group (1976IR02). A deuteron cluster transfer process may be involved (1976AS05). At  $E({}^3\text{He}) = 43.7$  and 53.0 MeV, the spectra show a prominent peak at the high energy end whose angular distributions exhibit a pronounced diffraction pattern (1967SL01, 1968MO10). Searches for three-proton enhancement (reaction (c)) have been unsuccessful: see (1974AJ01). See also (1976HE07; theor.).

$$7. {}^4\text{He}(\text{p}, \text{p}){}^4\text{He} \quad E_b = -1.97$$

Differential cross sections and polarizations have been measured at many energies: see Tables 5.5 in (1966LA04), 5.7 in (1974AJ01) and 5.5 here.

Phase shifts below  $E_p = 18$  MeV have been determined by (1977DO01) based on all the available cross section and polarization measurements, using an  $R$ -matrix analysis program. The  $P_{3/2}$  phase shift shows a pronounced resonance corresponding to  ${}^5\text{Li}_{\text{g.s.}}$  while the  $P_{1/2}$  shift changes slowly over a range of several MeV, suggesting that the first excited state is very broad and located 5 – 10 MeV above the ground state. (1977DO01) find that the reduced widths of the  $P$ -wave resonance states are nearly the same. See also (1974KR07:  $E_x = 2.3 \pm 0.5$  MeV,  $\Gamma = 9.0 \pm 1.5$  MeV for the  $P_{1/2}$  state). The  $D_{5/2}$ ,  $D_{3/2}$ ,  $F_{7/2}$  and  $F_{5/2}$  phase shift become greater than  $1^\circ$  at  $E_p \approx 11, 13, 14$  and 16 MeV, respectively (1977DO01). A contour plot of  $A_y$  is shown in (1973AR1N): the results are not significantly different from those of (1977DO01). The work of (1977DO01) relies heavily on the very accurate elastic scattering cross sections which they measured (better than 1%) and on the polarization measurements of (1977HA06) [absolute precision of  $\pm 0.01$ ]. It should be noted that  $A_y = 0.984 \pm 0.005$  at  $E_p = 11.93$  MeV,  $\theta_{\text{c.m.}} = 128.28^\circ$  (1977HA06). For other phase-shift analyses, see (1976BR17) and (1974AJ01).

A resonance is observed at  $E_p = 23$  MeV, corresponding to the known  $\frac{3}{2}^+$  state at  $E_x = 16.7$  MeV (1968AL1B, 1968DA04). An anomaly in the polarization is also observed at this energy (1966WE03, 1968DA04, 1972BA24). A further broad feature in the polarization excitation function ( $\theta = 102^\circ$ ) is observed at  $E_p = 30$  MeV ( $E_x = 22$  MeV) (1972BA24). Cross sections for  $23 < E_p < 45$  MeV show no further evidence of excited states (1969BU10). An extensive phase-shift analysis, using complex phases with  $l \leq 4$  over the range  $E_p = 20$  to 40 MeV has been made by (1972PL02), using mainly the polarization and differential cross-section data of (1972BA24). The  $D_{3/2}$  level is fit with  $R$ -matrix formalism, including background interference, with the following parameters:  $E_x = 16.68$  MeV,  $\gamma_p^2 = 122$  keV,  $\gamma_d^2(l=0) = 1.58$  MeV (negative sign),  $\gamma_d^2(l=2) = 1.58$  MeV,  $\theta_p^2 = 0.014$ ,  $\theta_d^2 = 0.765$ .

The  $E_p = 30$  MeV structure is not reflected in anomalous behavior of any single phase shift. Strong absorption of even partial waves may indicate broad overlapping positive-parity levels,  $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$ , of d +  ${}^3\text{He}$  character near  $E_x = 22$  MeV, but there is no unambiguous identification of excited states other than the  $\frac{3}{2}^+$  state in the p- $\alpha$  results (1972PL02). In an analysis of the data of (1971PL07) for  $E_p = 25$  to 29 MeV, (1971RA27) report evidence for a  $\frac{5}{2}^+$  level at  $E_x = 20$  MeV [ $\Gamma_{\text{el}}/\Gamma_{\text{total}} = 0.15$ ,  $\Gamma_{\text{tot}} = 8$  MeV] and a  $\frac{1}{2}^+$  level at  $E_x = 18 \pm 1$  MeV.

A recent analysis by (1975PL01) of the phase shifts below 50 MeV suggests a small  $D$ -state admixture to the dominant  $S$ -state configuration of  ${}^4\text{He}_{\text{g.s.}}$ .

Total reaction cross sections have been measured at  $E_p = 18$  to 48 MeV (1974SO06, 1976SO01) and at 0.87 and 2.1 GeV (1975JA1A). At an equivalent  $E_p$  of 1.05 GeV, the elastic distribution is characterized by a shallow first minimum (1977GE01) and is said to be in excellent agreement with the predictions of multiple-diffraction theory (1977WA06). Alpha-proton bremsstrahlung is observed at  $E_p = 7.0$  to 12.0 MeV, at  $E_\alpha = 20$  MeV [see (1974AJ01)] and at  $E_p = 22$ , 40 and 45 MeV (1975AN18). Coherent single pion production has been studied at  $E_p = 18.6$  GeV/c (1977BR1C). See also (1973BE1A, 1978GL1B), (1975IG1A, 1975RO1B, 1975TO1A, 1976GR1D, 1977BR1E, 1977RI1B, 1978BR1E, 1978IG1B), (1973GU1A, 1973HA1F, 1974BA34, 1974BA38, 1974CH02, 1974CH1E, 1974CI02, 1974DO10, 1974HA43, 1974JA1F, 1974LY02, 1974SA1G, 1974UL1A, 1974WA02, 1975AH07, 1975BA05, 1975BA1G, 1975BA76, 1975BL04, 1975CA05, 1975CO1A, 1975DU1A, 1975GR1C, 1975GU17, 1975KA1D, 1975NA07, 1975RU01, 1975RU07, 1975TA1A, 1975WA16, 1976AH1A, 1976AL1E, 1976AR12, 1976AU04, 1976CH1B, 1976DO1C, 1976DU06, 1976LE22, 1976LE1E, 1976NA04, 1976RU04, 1976SP1A, 1977AL06, 1977AR01, 1977CH1C, 1977DY01, 1977JA12, 1977KA1E, 1977KA1F, 1977KA1H, 1977LA1C, 1977LE1E, 1977PH01, 1977TH07, 1977TH09, 1977WA1C, 1977WA1E, 1977YO04, 1978CL1C, 1978HE2B, 1978LA02, 1978LE23, 1978MA37, 1978ME05, 1978YE1A, 1978YO1B; theor.).

8. (a) ${}^4\text{He}(p, d){}^3\text{He}$	$Q_m = -18.3532$	$E_b = -1.97$
(b) ${}^4\text{He}(p, pn){}^3\text{He}$	$Q_m = -20.5778$	
(c) ${}^4\text{He}(p, 2p){}^3\text{H}$	$Q_m = -19.8140$	
(d) ${}^4\text{He}(p, pd){}^2\text{H}$	$Q_m = -23.8467$	

Angular distributions of  ${}^3\text{He}$  ions (reaction (a)) have been measured for  $E_p = 27.9$  to 155.4 MeV [see (1974AJ01)] and at  $E_p = 85$  MeV (1974VO05), 275 to 500 MeV (1978CA05: back angles), 770 MeV (1976CO1C, 1977BA19) and at  $E_\alpha = 3.98$  GeV/c (1977BE34). The 85 MeV data are in agreement with DWBA at forward angles but the back angle behavior probably results from the need to include two-nucleon pickup in the analysis (1974VO05). The differential cross section is strongly backward peaked at  $E_p = 275$  and 500 MeV and the differential cross section is similar in slope and magnitude to that for  ${}^3\text{He}(p, p)$ , suggesting that deuteron exchange may be the major component of the reaction mechanism (1978CA05). The excitation function shows no indication of resonance for  $E_p = 38.5$  to 44.6 MeV (1969BU10). Polarization measurements have been reported for  $E_p = 32$  to 63 MeV [see (1974AJ01)] and at 45.04 and 52.34 MeV (1977SA1E).

and 350, 400 and 500 MeV ([1978CA05](#)). At  $E_\alpha = 6.85 \text{ GeV}/c$ , the integrated cross section for  ${}^3\text{He}$  production (reactions (a) + (b)) is  $24.1 \pm 1.9 \text{ mb}$ , suggesting an important component of  ${}^3\text{He}$  in  ${}^4\text{He}$  ([1977BI05](#)).

Reaction (b) and (c) have also been reported at  $E_p = 46.8$  to 156 MeV [see ([1974AJ01](#))]. The (p, 2p) reaction (reaction (c)) has also been studied at 350 and 500 MeV ([1978EP1C](#), [1978KO1F](#)) and 600 MeV ([1969PE15](#)), while both that reaction and reaction (d) have been studied at  $E_p = 156 \text{ MeV}$  ([1975FR14](#)). See also ([1973KR1A](#), [1978GL1B](#)), ([1977TE1A](#), [1978IG1B](#)), ([1976CO1B](#); astrophysics) and ([1974BA34](#), [1974CH02](#), [1974CH29](#), [1974DO10](#), [1974HA36](#), [1974KA1D](#), [1974RO19](#), [1974YO01](#), [1975HA03](#), [1975KA1D](#), [1977CL1A](#), [1977KA1D](#), [1977KA1G](#), [1977RE1A](#), [1978BI1C](#), [1978RO06](#); theor.).

$$\begin{aligned} 9. \quad & (a) {}^4\text{He}(d, n){}^5\text{Li} & Q_m = -4.19 \\ & (b) {}^4\text{He}(d, np){}^4\text{He} & Q_m = -2.2246 \end{aligned}$$

Reaction (b) has been studied at  $E_d = 14.2 \text{ MeV}$  and at  $E_\alpha = 18.0$  to 42 MeV [see ([1974AJ01](#))] and at  $E_\alpha = 18 \text{ MeV}$  ([1978SA07](#)) and 21.9 and 23.7 MeV ([1974RA10](#)). The data show that at 42 MeV, direct breakup, with quasi-free  $\alpha$ -p scattering taking place and the n acting as a spectator, is at least as important a mechanism as the FSI in the  ${}^5\text{Li}_{g.s.}$  ([1968WA01](#)). See also ([1974HE21](#), [1976KO21](#), [1976LI1E](#), [1977BR25](#); theor.) and  ${}^6\text{Li}$ .

$$10. \quad {}^4\text{He}(t, 2np){}^4\text{He} \quad Q_m = -8.3820$$

See ([1974LA1A](#), [1975AL1A](#);  $E_\alpha = 70 \text{ MeV}$ ; abstracts).

$$11. \quad {}^4\text{He}({}^3\text{He}, np){}^5\text{Li} \quad Q_m = -9.68$$

See ([1973HA50](#)).

$$12. \quad {}^6\text{Li}(\gamma, n){}^5\text{Li} \quad Q_m = -5.66$$

See  ${}^6\text{Li}$ .

$$\begin{aligned} 13. \quad & (a) {}^6\text{Li}(p, d){}^5\text{Li} & Q_m = -3.44 \\ & (b) {}^6\text{Li}(p, pd){}^4\text{He} & Q_m = -1.4735 \\ & (c) {}^6\text{Li}(p, pn){}^5\text{Li} & Q_m = -5.66 \end{aligned}$$

Angular distributions have been measured at  $E_p = 18.6$  to  $156$  MeV [see ([1974AJ01](#))] and at  $E_p = 185$  MeV ([1974KA28](#), [1976FA03](#)). In the latter experiment the spectra are characterized by a broad, asymmetric peak corresponding to  ${}^5\text{Li}_{\text{g.s.}}$ , a narrow peak [ ${}^5\text{Li}^*(16.7)$ ] and a broad peak at  $E_x \approx 20$  MeV. DWBA analysis leads to  $C^2S = 0.64$  and  $0.57$  for  ${}^5\text{Li}^*(0, 16.7)$  ([1976FA03](#)). The first excited state of  ${}^5\text{Li}$  is reported to be populated by ([1969BA05](#);  $E_p = 156$  MeV). See also ([1978IG1A](#):  $0.65$  and  $0.8$  GeV).

Reaction (b) has been studied at  $E_p = 9$  and  $10$  MeV, at  $45$  MeV [see ([1974AJ01](#))] and  $40$ ,  $45$  and  $50$  MeV ([1974BE46](#)): the p- $\alpha$  FSI corresponding to  ${}^5\text{Li}_{\text{g.s.}}$  is observed. See also ([1975MI1A](#)). For reaction (c) see ([1977WA05](#)). See also ([1975ST1C](#), [1975VO04](#), [1978CH1G](#)) and  ${}^6\text{Li}$ .

14. (a) ${}^6\text{Li}(\text{d}, \text{t}){}^5\text{Li}$	$Q_m = 0.59$
(b) ${}^6\text{Li}(\text{d}, \text{pt}){}^4\text{He}$	$Q_m = 2.5592$

Angular distributions of the  $t_0$  group have been measured at  $E_d = 15$  and  $20$  MeV: see ([1974AJ01](#)). For reaction (b) see ([1977RO18](#):  $E_d = 0.47$  MeV), ([1974MI10](#), [1977MI13](#);  $E_d = 7.5$  to  $10.5$  MeV) and ([1975KO1A](#); theor.).

15. (a) ${}^6\text{Li}({}^3\text{He}, \alpha){}^5\text{Li}$	$Q_m = 14.91$
(b) ${}^6\text{Li}({}^3\text{He}, \text{p}\alpha){}^4\text{He}$	$Q_m = 16.880$

At  $E({}^3\text{He}) = 25.5$  MeV, the spectra show  ${}^5\text{Li}^*(0, 16.7)$  and two broad peaks at  $E_x \approx 19.8$  and  $22.7$  MeV with  $\Gamma_{\text{c.m.}} = 2$  and  $1$  MeV, respectively ([1972BA30](#)). In a kinematically complete experiment at  $E({}^3\text{He}) = 36$  MeV the population of  ${}^5\text{Li}^*(16.7, 20.2, 22.6)$  is reported ([1976RAZW](#)). The decay of the ground state has also been studied at  $E({}^3\text{He}) = 1.25$  MeV ([1974LI10](#)) and at  $1.47$  to  $1.75$  MeV ([1978GU15](#)).

Cylindrical asymmetry observed in the breakup of  ${}^5\text{Li}(0)$  is attributed to the short lifetime of the  ${}^5\text{Li}$  intermediate state and to the memory retained by the proton of its localization at the time of formation of  ${}^5\text{Li}$  ([1967RE03](#), [1968RE10](#)). The first excited state of  ${}^5\text{Li}$  also appears to be involved ([1972TH08](#), [1972TH1B](#)): its parameters are given as  $E_x = 3.2 \pm 0.2$  MeV,  $\Gamma = 1.5 \pm 0.5$  MeV ([1975GA14](#)). See also ([1975GL08](#), [1976STYX](#), [1977AR09](#)) and  ${}^8\text{Be}$ .

16. (a) ${}^7\text{Li}(\text{p}, \text{t}){}^5\text{Li}$	$Q_m = -4.43$
(b) ${}^7\text{Li}(\text{p}, \text{p}\alpha){}^3\text{H}$	$Q_m = -2.467$

At  $E_p = 43.7$  MeV, a triton group is observed to  ${}^5\text{Li}(0)$  ( $\Gamma = 1.55 \pm 0.15$  MeV): the angular distribution is consistent with a substantial mixing of  $L = 0$  and  $2$  transfer. There is some evidence also for a very broad excited state between  $E_x = 2$  and  $5$  MeV.  ${}^5\text{Li}^*(16.7, 20.0)$  were not observed.

The formation of  ${}^5\text{Li}^*(16.7)$  ( ${}^4S_{3/2}$ ) would be  $S$ -forbidden: the absence of  ${}^5\text{Li}^*(20.0)$  would indicate that this state(s) is also of quartet character [see reaction 20 in  ${}^5\text{He}$ ] ([1966CE05](#)). Weak, broad states at  $E_x = 22.0 \pm 0.5$  MeV and  $25.0 \pm 0.5$  MeV and possibly 34 MeV are reported by ([1968MC02](#)) in a coincidence experiment in which three- and four-particle breakup was analyzed. The  $t_0$  angular distribution has also been studied at  $E_p = 16.6$  and 30.3 MeV: see ([1974AJ01](#)).



At  $E({}^3\text{He}) = 2.45$  and 6.00 MeV the reaction proceeds in part via the first two states of  ${}^5\text{Li}$  ([1966WA16](#)).



Not observed: see ([1972MA62](#)).

**${}^5\text{Be}$**   
(Fig. 3)

The absence of any group structure in the neutron spectrum in the reaction  ${}^3\text{He}({}^3\text{He}, n){}^5\text{Be}$  at  $E({}^3\text{He}) = 18.0$  to 26.0 MeV indicates that  ${}^5\text{Be}(0)$  is at least 4.2 MeV unstable with respect to  ${}^3\text{He} + 2p$  [ $(M - A) > 33.7$  MeV]. With Coulomb corrections adjusted to match the 16.7 MeV states of  ${}^5\text{He}-{}^5\text{Li}$ , this observation places the first  $T = \frac{3}{2}$  level in these nuclei above  $E_x = 21.4$  MeV ([1967AD05](#)). See also ([1975BE31](#); theor.).

## References

(Closed 1978)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1952AR30 H.V. Argo, R.F. Taschek, H.M. Agnew, A. Hemmendinger and W.T. Leland, Phys. Rev. 87 (1952) 612
- 1952BO68 T.W. Bonner, J.P. Conner and A.B. Lillie, Phys. Rev. 88 (1952) 473
- 1952CO35 J.P. Conner, T.W. Bonner and J.R. Smith, Phys. Rev. 88 (1952) 468
- 1952DO30 D.C. Dodder and J.L. Gammel, Phys. Rev. 88 (1952) 520
- 1953JA1A Jarvis and Roaf, Proc. Roy. Soc. 218 (1953) 432
- 1954AR02 W.A. Arnold, J.A. Phillips, G.A. Sawyer, E.J. Stovall, Jr. and J.L. Tuck, Phys. Rev. 93 (1954) 483
- 1954BR05 R.J.S. Brown, K.F. Famularo, H.D. Holmgren, D. Rankin and T.F. Stratton, Phys. Rev. 96 (1954) 80
- 1954BU06 W.H. Burke, J.R. Risser and G.C. Phillips, Phys. Rev. 93 (1954) 188
- 1955KU03 W.E. Kunz, Phys. Rev. 97 (1955) 456
- 1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
- 1959BO54 T.W. Bonner, F.W. Prosser, Jr. and J. Slattery, Phys. Rev. 115 (1959) 398
- 1960HU08 D.J. Hughes, B.A. Magurno and M.K. Brussel, BNL 325, 2nd Edition, Suppl. Vol. 1 (1960)
- 1963BU07 W. Buss, H. Waffler and B. Ziegler, Phys. Lett. 4 (1963) 198
- 1963PO02 C.H. Poppe, C.H. Holbrow and R.R. Borchers, Phys. Rev. 129 (1963) 733
- 1964PA1A Parker, Bahcall and Fowler, Astrophys. J. 139 (1964) 602
- 1964SH30 R.E. Shamu and J.G. Jenkin, Phys. Rev. 135 (1964) B99
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Edition, Suppl. Vol. 1 (1964)
- 1965CH12 G. Charpak, G. Gregoire, L. Massonnet, J. Saudinos, J. Favier, M. Gusakow and M. Jean, Phys. Lett. 16 (1965) 54
- 1965TR01 W. Trachslin, H. Burgisser, P. Huber, G. Michel and H.R. Striebel, Helv. Phys. Acta 38 (1965) 523

- 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331
- 1965WO03 C. Wong, J.D. Anderson and J.W. McClure, Nucl. Phys. 71 (1965) 106
- 1966BR02 L. Brown, H.A. Christ and H. Rudin, Nucl. Phys. 79 (1966) 459
- 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, Phys. Rev. 152 (1966) 950
- 1966HO07 B. Hoop, Jr. and H.H. Barschall, Nucl. Phys. 83 (1966) 65
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966WA16 M.A. Waggoner, J.E. Etter, H.D. Holmgren and C. Moazed, Nucl. Phys. 88 (1966) 81
- 1966WE03 W.G. Weitkamp and W. Haeberli, Nucl. Phys. 83 (1966) 46
- 1967AD05 E.G. Adelberger, A.B. McDonald, T.A. Tombrello, F.S. Dietrich and A.V. Nero, Phys. Lett. B25 (1967) 595
- 1967MC01 L.C. McIntyre and W. Haeberli, Nucl. Phys. A91 (1967) 369
- 1967RE03 M.A. Reimann, P.W. Martin and E.W. Vogt, Phys. Rev. Lett. 18 (1967) 246
- 1967SL01 R.J. Slobodian, J.S.C. McKee, W.F. Tivol, D.J. Clark and T.A. Tombrello, Phys. Lett. B25 (1967) 19
- 1967TO02 T.A. Tombrello, R.J. Spiger and A.D. Bacher, Phys. Rev. 154 (1967) 935
- 1967TR05 A. Trier and W. Haeberli, Phys. Rev. Lett. 18 (1967) 915
- 1967VA11 V. Valkovic, W.R. Jackson, Y.S. Chen, S.T. Emerson and G.C. Phillips, Nucl. Phys. A96 (1967) 241
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
- 1968AL1B Allison and Smythe, Nucl. Phys. A121 (1968) 97
- 1968BU09 W. Buss, W. Del Bianco, H. Waffler and B. Ziegler, Nucl. Phys. A112 (1968) 47
- 1968DA04 P. Darriulat, D. Garreta, A. Tarrats and J. Testoni, Nucl. Phys. A108 (1968) 316
- 1968IV01 M. Ivanovich, P.G. Young and G.G. Ohlsen, Nucl. Phys. A110 (1968) 441
- 1968KR03 L. Kraus, M. Suffert and D. Magnac-Valette, Nucl. Phys. A109 (1968) 593
- 1968MC02 R.L. McGrath, J. Cerny and S.W. Cosper, Phys. Rev. 165 (1968) 1126
- 1968MO10 B.J. Morton, E.E. Gross, E.V. Hungerford, J.J. Malanify and A. Zucker, Phys. Rev. 169 (1968) 825
- 1968RE10 M.A. Reimann, P.W. Martin and E.W. Vogt, Can. J. Phys. 46 (1968) 2241
- 1968VA12 V. Valkovic, C. Joseph, A. Niiler and G.C. Phillips, Nucl. Phys. A116 (1968) 497
- 1968WA01 R.E. Warner and R.W. Bercaw, Nucl. Phys. A109 (1968) 205
- 1968YO06 P.G. Young, R.H. Stokes and G.G. Ohlsen, Phys. Rev. 173 (1968) 949

- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BE56 V.M. Bezotosnyi, V.A. Zhmailo, L.M. Surov and M.S. Shvetsov, Yad. Fiz. 10 (1969) 225; Sov. J. Nucl. Phys. 10 (1970) 127
- 1969BU10 S.N. Bunker, J.M. Cameron, M.P. Epstein, G. PaiImage, J.R. Richardson, J.G. Rogers, P. TomaImage and J.W. Verba, Nucl. Phys. A133 (1969) 537
- 1969PE15 C.F. Perdrisat, L.W. Swenson, P.C. Gugelot, E.T. Boschitz, W.K. Roberts, J.S. Vincent and J.R. Priest, Phys. Rev. 187 (1969) 1201
- 1969RO16 D.C. Rorer, B.M. Ecker and R.O. Akyuz, Nucl. Phys. A133 (1969) 410
- 1970AR1B R.A. Arndt and L.D Roper, Phys. Rev. C1 (1970) 903
- 1970GA1A Garber et al., BNL 400, 3rd Edition, Vol. 1 (1970)
- 1970KO09 A. Kosiara and H.B. Willard, Phys. Lett. B32 (1970) 99
- 1970MI05 D. Miljanic, M. Furic and V. Valkovic, Nucl. Phys. A148 (1970) 312
- 1970SC18 H. Schroder and W. Mausberg, Z. Phys. 235 (1970) 234
- 1971GR12 E.E. Gross, E.V. Hungerford and J.J. Malanify, Nucl. Phys. A164 (1971) 376
- 1971GR32 H. Grunder, R. Gleyvod, G. Lietz, G. Morgan, H. Rudin, F. Seiler and A. Stricker, Helv. Phys. Acta 44 (1971) 662
- 1971GR45 R. Grotzschel, B. Kuhn, K. Moller, J. Mosner and G. Schmidt, Nucl. Phys. A176 (1971) 261
- 1971GR47 W. Gruebler, V. Konig, A. Ruh, P.A. Schmelzbach, R.E. White and P. Marmier, Nucl. Phys. A176 (1971) 631
- 1971HI07 D. Hilscher, P.A. Quin and J.C. Davis, Nucl. Phys. A173 (1971) 216
- 1971KE16 L.G. Keller and W. Haeberli, Nucl. Phys. A172 (1971) 625
- 1971KI02 K. Kilian, H. Treiber, R. Strausz and D. Fick, Phys. Lett. B34 (1971) 283
- 1971KL02 W. Klinger, F. Dusch and R. Fleischmann, Nucl. Phys. A166 (1971) 253
- 1971LE27 C. Leemann, H. Burgisser, P. Huber, U. Rohrer, H. Paetz gen.Schieck and F. Seiler, Ann. Phys. 66 (1971) 810
- 1971MU04 G.S. Mutchler, W.B. Broste and J.E. Simmons, Phys. Rev. C3 (1971) 1031
- 1971PL07 D.J. Plummer, K. Ramavataram, T.A. Hodges, D.G. Montague, A. Zucker and N.K. Ganguly, Nucl. Phys. A174 (1971) 193
- 1971RA27 K. Ramavataram, D.J. Plummer, T.A. Hodges and D.G. Montague, Nucl. Phys. A174 (1971) 204
- 1971RE19 D. Rendic, N.D. Gabitzsch, V. Valkovic, W. von Witsch and G.C. Phillips, Nucl. Phys. A178 (1971) 49

- 1972AG01 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, G.P. Gulkanyan, V.M. Sidorov and V.A. Yarba, *Yad. Fiz.* 15 (1972) 18; *Sov. J. Nucl. Phys.* 15 (1972) 10
- 1972AN24 Y.N. Antufev, V.L. Agranovich, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, *Pisma Zh. Eksp. Teor. Fiz.* 16 (1972) 77; *JETP Lett.* 16 (1972) 52
- 1972AN27 Y.P. Antoufiev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, *Phys. Lett.* B42 (1972) 347
- 1972AN29 Y.P. Antufev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, *Pisma Zh. Eksp. Teor. Fiz.* 16 (1972) 339; *JETP Lett.* 16 (1972) 240
- 1972BA24 A.D. Bacher, G.R. Plattner, H.E. Conzett, D.J. Clark, H. Grunder and W.F. Tivol, *Phys. Rev. C5* (1972) 1147
- 1972BA30 M.P. Baker, J.M. Cameron, N.S. Chant and N.F. Mangelson, *Nucl. Phys. A184* (1972) 97
- 1972BR10 W.B. Broste, G.S. Mutchler, J.E. Simmons, R.A. Arndt and L.D. Roper, *Phys. Rev. C5* (1972) 761
- 1972KI01 H.T. King, W.E. Meyerhof and R.G. Hirko, *Nucl. Phys. A178* (1972) 337
- 1972KI02 T.R. King and R. Smythe, *Nucl. Phys. A183* (1972) 657
- 1972MA62 A.M. MacLeod and G.R. Milne, *J. Phys. (London) A5* (1972) 1252
- 1972PL02 G.R. Plattner, A.D. Bacher and H.E. Conzett, *Phys. Rev. C5* (1972) 1158
- 1972ST01 T. Stammbach and R.L. Walter, *Nucl. Phys. A180* (1972) 225
- 1972TH08 D.T. Thompson and G.E. Tripard, *Phys. Rev. C6* (1972) 452
- 1972TH1B Thompson, Thesis, Washington State Univ. (1972)
- 1973AR1B Arthur, Thesis LA 5230 T (1973)
- 1973AR1N Arndt, Long and Roper, *Nucl. Phys. A209* (1973) 429
- 1973BA30 Y.D. Bayukov, L.S. Vorobev, V.M. Kolybasov, G.A. Leksin, V.L. Stolin, V.B. Fedorov and V.D. Khovanskii, *Yad. Fiz.* 17 (1973) 916; *Sov. J. Nucl. Phys.* 17 (1974) 479
- 1973BE1A Berthot et al., Univ. Claude Bernard de Lyon, Rept. No. LYCEN 7338 (1973)
- 1973DE17 V.P. Denisov, L.A. Kulchitskii and I.Y. Chubukov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 107; *Bull. Acad. Sci. USSR Phys. Ser.* 37:1 (1973) 94
- 1973DE51 A.A. Debenham, V. Konig, W. Gruebler, P.A. Schmelzbach, R. Risler and D.O. Boerma, *Nucl. Phys. A216* (1973) 42
- 1973FE1A N. Feather and D.G. Vass, *Proc. Roy. Soc. Edinburgh* A71 (1973) 233
- 1973GA16 S.N. Gardiner, J.L. Matthews and R.O. Owens, *Phys. Lett.* B46 (1973) 186
- 1973GA28 H.P. Garnir, G. Robaye and J.M. Delbrouck-Habaru, *Bull. Soc. Roy. Sci. Liege (Belgium)* 42 (1973) 195
- 1973GO38 C.A. Goulding, P. Stoler and J.D. Seagrave, *Nucl. Phys. A215* (1973) 253

- 1973GU1A Guardiola, Nucl. Many-Body Problem, 1972, Vol. 2 (1973) 433
- 1973HA1F Hackenbroich, Nucl. Many-Body Problem, 1972, Vol. 1 (1973) 706
- 1973HA50 M.L. Halbert, A. van der Woude and N.M. O'Fallon, Phys. Rev. C8 (1973) 1621
- 1973HU12 J. Hudomalj and P. Tomas, Fizika (Yugoslavia) 5 (1973) 203
- 1973KR1A Krebs et al., Phys. in Canada 29 (1973) 22
- 1973LI1B Liskien and Paulsen, Nucl. Data Tables 11 (1973) 569
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973SL03 I. Slaus, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, J.M. Lambert and D.L. Shannon, Phys. Rev. C8 (1973) 444
- 1973TR04 P.A. Treado, J.M. Lambert, R.J. Kane, L.A. Beach, E.L. Petersen and R.B. Theus, Phys. Rev. C7 (1973) 1742
- 1974AD11 A. Adam and J. Cabe, Nucl. Instrum. Meth. 121 (1974) 339
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974AL01 R.G. Allas, L.A. Beach, R.O. Bondelid, E.M. Diener, E.L. Petersen, J.M. Lambert, P.A. Treado and I. Slaus, Phys. Rev. C9 (1974) 787
- 1974AM01 J. Amato, R.L. Burman, R. Macek, J. Oostens, W. Shlaer, E. Arthur, S. Sobottka and W.C. Lam, Phys. Rev. C9 (1974) 501
- 1974AN02 B. Antolkovic, Nucl. Phys. A219 (1974) 332
- 1974BA14 S.D. Baker, R. Beurtey, G. Bruge, A. Chaumeaux, J.-M. Durand, J.C. Faivre, J.M. Fontaine, D. Garreta, D. Legrand, J. Saudinos et al., Phys. Rev. Lett. 32 (1974) 839
- 1974BA34 A.G. Baryshnickov, L.D. Blokhintsev, A.N. Safronov and V.V. Turovtsev, Nucl. Phys. A224 (1974) 61
- 1974BA38 R.H. Bassel and M. Rosen, Phys. Rev. C10 (1974) 928
- 1974BE46 J.L. Beveridge, R. Corfu, J.P. Egger, F. Foroughi, J. Piffaretti, J. Rossel, J. Arvieux, J.L. Durand and C. Perrin, Helv. Phys. Acta 47 (1974) 211
- 1974BE67 R. Beckmann, U. Holm and K. Lorenzen, Z. Phys. 271 (1974) 89
- 1974BH03 R.K. Bhowmik, C.C. Chang, P.G. Roos and H.D. Holmgren, Nucl. Phys. A226 (1974) 365
- 1974CH02 F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Rev. C9 (1974) 56
- 1974CH15 C.C. Chang and E. Ventura, Phys. Rev. C9 (1974) 1671
- 1974CH1E F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Lett. B46 (1974) 301
- 1974CH29 F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Rev. C10 (1974) 406
- 1974CI02 C. Ciofi degli Atti and R. Guardiola, Phys. Lett. B51 (1974) 26

- 1974DA28 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares and R. Potenza, Nuovo Cim. A22 (1974) 313
- 1974DO05 T.R. Donoghue, G.G. Ohlsen, N. Jarmie and P.W. Keaton, Jr., Phys. Rev. C10 (1974) 571
- 1974DO10 E.I. Dolinsky and A.M. Mukhamedzhanov, Phys. Lett. B52 (1974) 17
- 1974FE1A Feather and Vass, Proc. Roy. Soc. Edinburgh A71 (1974) 293
- 1974GA21 R. Garrett and W.W. Lindstrom, Nucl. Phys. A224 (1974) 186
- 1974GE1B Geisler et al., Science 186 (1974) 361
- 1974GH01 A. Ghovanlou and D.R. Lehman, Phys. Rev. C9 (1974) 1730
- 1974GH03 A. Ghovanlou and F. Prats, Phys. Rev. C10 (1974) 1309
- 1974GO13 G.S. Gogosadze and T.I. Kopaleishvili, Yad. Fiz. 19 (1974) 539; Sov. J. Nucl. Phys. 19 (1974) 272
- 1974GR40 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2148; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 115
- 1974GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2543; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 69
- 1974GR44 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, V.M. Pugach, B.A. Rudenko, M.V. Sokolov, B.G. Struzhko et al., Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2547; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 72
- 1974HA36 R.D. Haracz and T.K. Lim, Phys. Rev. C10 (1974) 431
- 1974HA43 H.H. Hackenbroich, P. Heiss and Le-Chi-Niem, Phys. Lett. B51 (1974) 334
- 1974HE21 P. Heiss, H.H. Hackenbroich and K. Prescher, Phys. Lett. B52 (1974) 411
- 1974HO11 E. Hourany, H. Nakamura, F. Takeutchi and T. Yuasa, Nucl. Phys. A222 (1974) 537
- 1974IR04 J.M. Irvine, G.S. Mani and M. Vallieres, Czech. J. Phys. B24 (1974) 1269
- 1974JA15 N. Jarmie and J.H. Jett, Phys. Rev. C10 (1974) 54
- 1974JA1F Jackson, Prog. in Phys. 37 (1974) 55
- 1974JA30 V.K. Jain and B.B. Srivastava, Indian J. Pure Appl. Phys. 12 (1974) 117
- 1974KA1D Kanada, Kaneko and Nomoto, Prog. Theor. Phys. 52 (1974) 725
- 1974KA28 J. Kallne, B. Fagerstrom, O. Sundberg and G. Tibell, Phys. Lett. B52 (1974) 313
- 1974KE1B Keaton, Gammel and Ohlsen, Ann. Phys. 85 (1974) 152
- 1974KR07 L. Kraus and I. Linck, Nucl. Phys. A224 (1974) 45

- 1974LA02 R. Larose-Poutissou and H. Jeremie, Nucl. Phys. A218 (1974) 559
- 1974LA1A Lambert et al., Bull. Amer. Phys. Soc. 19 (1974) 506
- 1974LA1C Langley, Picrajx and Vook, J. Nucl. Materials 53 (1974) 257
- 1974LE22 C. Leclercq-Willain and M. Libert-Heinemann, Nucl. Phys. A229 (1974) 15
- 1974LI10 D.L. Livesey and C.J. Piluso, Can. J. Phys. 52 (1974) 1167
- 1974LI1C Lisowski, Walter, Jacobs and Clegg, Bull. Amer. Phys. Soc. 19 (1974) 1075
- 1974LO06 P.A. Lovoi, G.C. Salzman, G.G. Ohlsen, C.K. Mitchell and W. Gruebler, Phys. Rev. C9 (1974) 1336
- 1974LY02 G.I. Lykasov and A.V. Tarasov, Yad. Fiz. 20 (1974) 489; Sov. J. Nucl. Phys. 20 (1975) 263
- 1974LY03 E.B. Lyovshin, O.F. Nemets and A.M. Yasnogorodsky, Phys. Lett. B52 (1974) 392
- 1974MA1C Maayouf and Galloway, Nucl. Instrum. Meth. 118 (1974) 343
- 1974MI10 D. Miljanic, J. Hudomalj, G.S. Mutchler, E. Andrade and G.C. Phillips, Phys. Lett. B50 (1974) 330
- 1974NA10 H. Nakamura, Nucl. Phys. A223 (1974) 599
- 1974OH05 G.G. Ohlsen, R.A. Hardekopf, D.P. May, S.D. Baker and W.T. Armstrong, Nucl. Phys. A233 (1974) 1
- 1974POZN C.H. Poppe, J.C. Davis, J.D. Anderson, S.M. Grimes and C. Wong, Bull. Amer. Phys. Soc. 19 (1974) 1035
- 1974PR10 F. Prats, Nucl. Phys. A227 (1974) 469
- 1974PR1B Pronko, J. Nucl. Materials 53 (1974) 252
- 1974RA10 T. Rausch, H. Zell, D. Wallenwein and W. von Witsch, Nucl. Phys. A222 (1974) 429
- 1974RA1C Ramaty and Kozlovsky, Astrophys. J. 193 (1974) 729
- 1974RO19 P.G. Roos, Phys. Rev. C9 (1974) 2437
- 1974SA1G Saudinos and Wilkin, Ann. Rev. Nucl. Sci. 24 (1974) 341
- 1974SC04 H. Schroder, K.K. Kern and D. Fick, Phys. Lett. B48 (1974) 206
- 1974SO06 A.M. Sourkes, N.E. Davison, S.A. Elbakr, J.L. Horton, A. Houdayer, W.T.H. van Oers and R.F. Carlson, Phys. Lett. B51 (1974) 232
- 1974ST14 M. Stingl and A.S. Rinat, Phys. Rev. C10 (1974) 1253
- 1974TA1A Tanabe et al., Bull. Inst. Chem. Res. Kyoto U. 52 (1974) 284
- 1974TO03 W. Tornow, Z. Phys. 266 (1974) 357
- 1974TR02 T.A. Trainor, T.B. Clegg and P.W. Lisowski, Nucl. Phys. A220 (1974) 533
- 1974TU1A Turk, Antolkovic, Dolenc and Winterhalter, Proc. 1973 Gaussig Conf., ZFK-271 (1974) 115

- 1974UL1A Ullo and Feshbach, Ann. Phys. 82 (1974) 156
- 1974VO05 L.G. Votta, P.G. Roos, N.S. Chant and R. Woody, III, Phys. Rev. C10 (1974) 520
- 1974WA02 W.L. Wang and R.G. Lipes, Phys. Rev. C9 (1974) 814
- 1974WA06 R.E. Warner, S.A. Gottlieb, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Nucl. Phys. A221 (1974) 593
- 1974WI1D Wiemer, Atomkernenergie 23 (1974) 261
- 1974YO01 S.K. Young and E.F. Redish, Phys. Rev. C10 (1974) 498
- 1975AB11 U. Abbondanno, F. Demanins, C. Tuniz and G. Nardelli, Nuovo Cim. A29 (1975) 187
- 1975AB1C Abulaffio and Peres, Nucl. Cross Sect. and Tech., Pt. 3, Washington, D.C., 1975 (1975) 701
- 1975AH07 A.A.Z. Ahmad, S. Ali, N. Ferdous and M. Ahmed, Nuovo Cim. A30 (1975) 385
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975AL1A Allas et al., Few Body Problems, Quebec, 1974 (1975) 422
- 1975AN18 G.A. Anzelon, I. Slaus, S.Y. Tin, W.T.H. Van Oers, R.M. Eisberg, M. Makino, C.N. Waddell and M.B. Epstein, Nucl. Phys. A255 (1975) 250
- 1975BA05 S. Barshay, C.B. Dover and J.P. Vary, Phys. Rev. C11 (1975) 360
- 1975BA1E Batay-Csorba and Barnes, Bull. Amer. Phys. Soc. 20 (1975) 829
- 1975BA1G Baldin, AIP Conf. Proc. 26 (1975) 621
- 1975BA76 A.I. Baz, V.F. Demin and M.V. Zhukov, Fiz. Elem. Chastits At. Yadra 6 (1975) 515; Sov. J. Part. Nucl. 6 (1975) 207
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BL04 M. Bleszynski and T. Jaroszewicz, Phys. Lett. B56 (1975) 427
- 1975BR1A Brown et al., Bull. Amer. Phys. Soc. 20 (1975) 1157
- 1975CA05 G. Cattapan, G. Pisent and V. Vanzani, Nucl. Phys. A241 (1975) 204
- 1975CH1B Chang and Bhowmik, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 317
- 1975CH1C Chang, Bhowmik, Chant and Roos, ORO-4856-32 (1975)
- 1975CO1A Coker and Hoffmann, Bull. Amer. Phys. Soc. 20 (1975) 1166
- 1975CO27 V. Comparat, R. Frascaria, N. Fujiwara, N. Marty, M. Morlet, P.G. Roos and A. Willis, Phys. Rev. C12 (1975) 251
- 1975CR1A Cranberg, Bull. Amer. Phys. Soc. 20 (1975) 1182
- 1975DE37 V.P. Denisov and I.Y. Chubukov, Yad. Fiz. 22 (1975) 897; Sov. J. Nucl. Phys. 22 (1976) 466
- 1975DI04 H. Dirim, J.P. Elliott and J.A. Evans, Nucl. Phys. A244 (1975) 301

- 1975DI1A Didelez et al., Bull. Amer. Phys. Soc. 20 (1975) 597
- 1975DI1B Didelez et al., Jul-Conf-16 (1975) 90
- 1975DO1B Dodder, Bull. Amer. Phys. Soc. 20 (1975) 821
- 1975DU1A Dubnicka and Dumbrajs, Phys. Rept. 19 (1975) 141
- 1975FE1A Feather, Commun. Roy. Soc. Edinburgh (Phys. Sci.) 2 (1975) 19
- 1975FI1B Firk et al., Nucl. Cross Sections and Tech., Pt. II, Washington, D.C., 1975 (1975) 875
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, Ann. Rev. Astron. Astrophys. 13 (1975) 69
- 1975FR14 R. Frascaria, P.G. Roos, M. Morlet, N. Marty, A. Willis, V. Comparat and N. Fujiwara, Phys. Rev. C12 (1975) 243
- 1975GA14 R.M. Gagne, C.M. Fou and S. Ward, J. Phys. (Paris) 36 (1975) 759
- 1975GL08 Y.A. Glukhov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, Yad. Fiz. 22 (1975) 679; Sov. J. Nucl. Phys. 22 (1976) 351
- 1975GR1C Green, Few Body Problems, Quebec, 1974 (1975) 352
- 1975GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2117; Bull. Acad. Sci. USSR Phys. Ser. 39 (1975) 95
- 1975GU17 S.A. Gurvits, Y. Alexander and A.S. Rinat, Phys. Lett. B59 (1975) 22
- 1975HA03 R.D. Haracz and T.K. Lim, Phys. Rev. C11 (1975) 634
- 1975HA1E Hackenbroich, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 107
- 1975HE09 P. Heiss, Z. Phys. A272 (1975) 267
- 1975HE1C J. Henley and H.W. Meldner, Phys. Rev. C12 (1975) 407
- 1975HU1A Hufschmidt, Heintze, Moller and Kamke, Nucl. Instrum. Meth. 124 (1975) 573
- 1975IG1A Igo, AIP Conf. Proc. 26 (1975) 63
- 1975IG1B Igo et al., Bull. Amer. Phys. Soc. 20 (1975) 1166
- 1975JA1A Jaros et al., Bull. Amer. Phys. Soc. 20 (1975) 666
- 1975JE1A Jenny et al., Helv. Phys. Acta 48 (1975) 558
- 1975KA1D Kanada, Kaneko and Nomoto, Prog. Theor. Phys. 54 (1975) 1707
- 1975KO1A S. Kohmoto and Y. Sakamoto, Phys. Rev. Lett. 34 (1975) 550
- 1975KR1A Kramer, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 56
- 1975KU1C Kuhn, Few Body Problems, Quebec, 1974 (1975) 122
- 1975LO1A M.P. Locher, Nucl. Phys. A251 (1975) 493
- 1975MA1E Matthews, AIP Conf. Proc. 26 (1975) 437

- 1975MA28 E. Magiera, M. Bormann, W. Scobel and P. Heiss, Nucl. Phys. A246 (1975) 413
- 1975ME27 Y.P. Melnik and E.V. Inopin, Ukr. Fiz. Zh. 20 (1975) 2032
- 1975MI1A Miljanic, Zabel and Phillips, Few Body Problems, Quebec, 1974 (1975) 669
- 1975NA07 M.A. Nagarajan, W.L. Wang, D.J. Ernst and R.M. Thaler, Phys. Rev. C11 (1975) 1167
- 1975NE11 O.F. Nemets, Y.S. Stryuk and A.M. Yasnogorodskii, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2066; Bull. Acad. Sci. USSR Phys. Ser. 39:10 (1975) 49
- 1975PL01 G.R. Plattner, R.D. Viollier and K. Alder, Phys. Rev. Lett. 34 (1975) 830
- 1975RO1B Roos and Chant, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 242, 265
- 1975RO1C Roy et al., Bull. Amer. Phys. Soc. 20 (1975) 693
- 1975RU01 D.W. Rule and Y. Hahn, Phys. Rev. Lett. 34 (1975) 332
- 1975RU04 R.W. Rutkowski and E.E. Gross, Phys. Rev. C12 (1975) 362
- 1975RU07 D.W. Rule and Y. Hahn, Phys. Rev. C12 (1975) 1616
- 1975SA1A Sagle et al., Few Body Problems, Quebec, 1974 (1975) 621
- 1975SC28 G. Schrieder, H. Genz, A. Richter and W. von Witsch, Nucl. Phys. A247 (1975) 203
- 1975SE07 F. Seiler, Nucl. Phys. A244 (1975) 236
- 1975SE1D Seiler, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 191
- 1975SE1E Seiler, Few Body Problems, Quebec, 1974 (1975) 595
- 1975SO1C Soleilhac, CEA-N-1812 (1975)
- 1975ST1A Stengl, Vonach and Fabian, Nucl. Instrum. Meth. 126 (1975) 235
- 1975ST1B Stewart and Hale, USNDC-CTR-2, LA-5828-MS (1975)
- 1975ST1C Strobel, Jul-Conf-16 (1975) 89
- 1975TA1A Tang and Thompson, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 119
- 1975TH12 D.R. Thompson and Y.C. Tang, Phys. Rev. C12 (1975) 1432; Erratum Phys. Rev. C13 (1976) 2597
- 1975TO1A Tombrello, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 167
- 1975VE09 S.L. Verbeck, J.C. Fong, G. Igo, C.A. Whitten, Jr., D.L. Hendrie, Y. Terrien, V. Perez-Mendez and G.W. Hoffmann, Phys. Lett. B59 (1975) 339
- 1975VO04 W. Von Witsch, G.S. Mutchler and D. Miljanic, Nucl. Phys. A248 (1975) 485
- 1975VO08 W. von Oertzen and E.R. Flynn, Ann. Phys. 95 (1975) 326
- 1975WA16 S.J. Wallace, Phys. Rev. C12 (1975) 179
- 1975WA31 R.E. Warner, R.L. Ruyle, W.G. Davies, G.C. Ball, A.J. Ferguson and J.S. Forster, Nucl. Phys. A255 (1975) 95
- 1975YA12 A.M. Yasnogorodsky, Phys. Lett. B58 (1975) 139

- 1976AH1A M.U. Ahmed and P.E. Shanley, Phys. Rev. Lett. 36 (1976) 25
- 1976AL1E Ali, Proc. 7th Int. Conf. on Few Body Problems in Nucl. Part. Phys., Delhi, 1975 (1976) 438
- 1976AL22 M. Allab and H. Beaumevieille, Nuovo Cim. A36 (1976) 95
- 1976AR12 L.G. Arnold, B.C. Clark, R.L. Mercer, D.G. Ravenhall and A.M. Saperstein, Phys. Rev. C14 (1976) 1878
- 1976AS05 J. Asai and R.J. Slobodrian, Phys. Lett. B64 (1976) 257
- 1976AU04 J.P. Auger, J. Gillespie and R.J. Lombard, Nucl. Phys. A262 (1976) 372
- 1976BA1E Baz and Filippov, Proc. Dubna Conf. JINR-9920, Vol. II (1976) 268
- 1976BE45 J. Berger, J. Duflo, L. Goldzahl, F. Plouin, J. Oostens, M. Van Den Bossche, L. Vu Hai, G. Bizard, C. Le Brun, F.L. Fabbri et al., Phys. Rev. Lett. 37 (1976) 1195
- 1976BI1A A. Billoire and M. Chemtob, Phys. Lett. B64 (1976) 389
- 1976BL1A Block, Bull. Amer. Phys. Soc. 21 (1976) 1342
- 1976BO05 J.E. Bond and F.W.K. Firk, Nucl. Phys. A258 (1976) 189
- 1976BR17 M.E. Brandan, G.R. Plattner and W. Haeberli, Nucl. Phys. A263 (1976) 189
- 1976BR1B Brady et al., Lowell Conf., CONF-760715-P2 (1976) 1273
- 1976BR1C Brook et al., Polarization, Zurich, 1975 (1976) 546
- 1976CH1B Chuu, J. Natl. Chiao Tung Univ. (Taiwan) 1 (1976) 295
- 1976CO1B Colgate, Audouze and Fowler, OAP-458 (1976)
- 1976CO1C Couvert, Meson-Nucl. Phys., 1976 (1976) 310
- 1976DO1B Dodder, Polarization, Zurich, 1975 (1976) 167
- 1976DO1C Dodder et al., LA 6186 MS (1976)
- 1976DR1A Drosg, Lowell Conf., CONF-760715-P2 (1976) 1384
- 1976DR1B Drosg, Smith and Woods, LA 6262 MS (1976)
- 1976DRZU L.J. Dries, H.W. Clark, K.R. Crosthwaite, J.L. Regner and T.R. Donoghue, Bull. Amer. Phys. Soc. 21 (1976) 926, C2
- 1976DU06 O. Dumbrajs and M. Staszek, J. Phys. (London) G2 (1976) 795
- 1976FA03 B. Fagerstrom, J. Kallne, O. Sundberg and G. Tibell, Phys. Scr. 13 (1976) 101
- 1976FA09 J. Fain, J. Gardes, A. Lefort, L. Meritet, J.F. Pauty, G. Peynet, M. Querrou, F. Vazeille and B. Ille, Nucl. Phys. A262 (1976) 413
- 1976FO21 C.-M. Fou, Y.-C. Liu, C.-C. Hsu and S.-L. Huang, J. Phys. (London) G2 (1976) 847
- 1976GAYV D.I. Garber and R.R. Kinsey, BNL 325, 3rd Edition, Vol. 2 (1976)
- 1976GO1B Goddard, Rohrig, Knutson and Haeberli, Nucl. Instrum. Meth. 137 (1976) 451

- 1976GR10 W. Gruebler, P.A. Schmelzbach, V. Konig, B. Jenny, R. Risler, H.R. Burgi and J. Nurzynski, Nucl. Phys. A271 (1976) 29
- 1976GR1D Gruebler, Polarization, Zurich, 1975 (1976) 307
- 1976HA1C Hackenbroich, Polarization, Zurich, 1975 (1975) 133
- 1976HE07 P. Heiss and H.H. Hackenbroich, Phys. Lett. B61 (1976) 339
- 1976HUZK J. Hudomalj-Gabitsch, J. Clement, W. Dragoset, R. Felder, G.S. Mutchler, T.W. Williams, G.C. Phillips, E.V. Hungerford, M. Warneke, L. Pinsky et al., Bull. Amer. Phys. Soc. 21 (1976) 998, DE6
- 1976IR02 M. Irshad, J. Asai, S. Sen, R. Pigeon and R.J. Slobodian, Nucl. Phys. A265 (1976) 349
- 1976IR1B Irvine et al., Ann. Phys. 102 (1976) 129
- 1976JA1C Jarvis, J. Phys. (London) G2 (1976) 603
- 1976KO21 Y. Koike, Prog. Theor. Phys. 55 (1976) 2016
- 1976LE1E Levin and Strikman, Sov. J. Nucl. Phys. 23 (1976) 216
- 1976LE22 H. Lesniak, L. Lesniak and A. Tekou, Nucl. Phys. A267 (1976) 503
- 1976LI15 P.W. Lisowski, R.L. Walter, G.G. Ohlsen and R.A. Hardekopf, Phys. Rev. Lett. 37 (1976) 809
- 1976LI1E Lindner, Polarization, Zurich, 1975 (1976) 817
- 1976LIZV P.W. Lisowski, W. Tornow, R.C. Byrd, S.E. Skubic and R.L. Walter, Bull. Amer. Phys. Soc. 21 (1976) 534, BF8
- 1976LY1A Lyovshin, Nemets and Yasnogorodsky, Polarization, Zurich, 1975 (1976) 542
- 1976MA34 J.L. Matthews, D.J.S. Findlay, S.N. Gardiner and R.O. Owens, Nucl. Phys. A267 (1976) 51
- 1976ME13 H.O. Meyer, G.G. Ohlsen, R.A. Hardekopf, R.V. Poore and J.W. Sunier, Nucl. Phys. A265 (1976) 280
- 1976MO1A Motonaga et al., Polarization, Zurich, 1975 (1976) 837
- 1976NA04 N. Nakamura, Y. Takahashi and T. Kohmura, Phys. Lett. B60 (1976) 439
- 1976OH02 G.G. Ohlsen, R.A. Hardekopf, N. Jarmie, R.V. Poore, R.L. Walter and P.W. Lisowski, Phys. Rev. C14 (1976) 1688
- 1976OH04 G. Ohlen and T. Berggren, Nucl. Phys. A272 (1976) 21
- 1976OH1B Ohlsen, Polarization, Zurich, 1975 (1976) 315
- 1976OK1A Okumusoglu, Blyth and Dahme, Polarization, Zurich, 1975 (1976) 560
- 1976RAZW K. Ramavataram, R. Larue, R. Gagnon, W.G. Davies, G.C. Ball, A.J. Ferguson and R.W. Warner, Bull. Amer. Phys. Soc. 21 (1976) 774, BA5

- 1976RO1C Roy et al., Polarization, Zurich, 1975 (1976) 540
- 1976RO1G Roy et al., Polarization, Zurich, 1975 (1976) 548
- 1976RO1H Roy, Rad, Conzett and Seiler, Polarization, Zurich, 1975 (1976) 554
- 1976ROZM R. Roy, H.E. Conzett, R.M. Larimer, F.N. Rad and F. Seiler, Bull. Amer. Phys. Soc. 21 (1976) 774, BA1
- 1976RU04 D.W. Rule and Y. Hahn, Phys. Rev. C14 (1976) 1102
- 1976SC15 P.A. Schmelzbach, W. Gruebler, V. Konig, R. Risler, D.O. Boerma and B. Jenny, Nucl. Phys. A264 (1976) 45
- 1976SC26 H. Schroder, K.K. Kern, K. Schmidt and D. Fick, Nucl. Phys. A269 (1976) 74
- 1976SE03 F. Seiler, Phys. Lett. B61 (1976) 144
- 1976SE1B Seiler, Polarization, Zurich, 1975 (1976) 159
- 1976SE1C Seiler, Roy, Conzett and Rad, Polarization, Zurich, 1975 (1976) 550
- 1976SO01 A.M. Sourkes, A. Houdayer, W.T.H. van Oers, R.F. Carlson and R.E. Brown, Phys. Rev. C13 (1976) 451
- 1976SP1A D.A. Sparrow and W.J. Gerace, Nucl. Phys. A260 (1976) 474
- 1976STYX D.P. Stahel, R. Jahn, G.J. Wozniak and J. Cerny, Bull. Amer. Phys. Soc. 21 (1976) 1295, CC8
- 1976SU06 J.W. Sunier, R.V. Poore, R.A. Hardekopf, L. Morrison, G.C. Salzman and G.G. Ohlsen, Phys. Rev. C14 (1976) 8
- 1976VA29 G.V. Valskii, Yad. Fiz. 24 (1976) 270; Sov. J. Nucl. Phys. 24 (1976) 140
- 1976WA12 R.E. Warner, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Nucl. Phys. A269 (1976) 286
- 1976WA1B Walter, Polarization, Zurich, 1975 (1976) 377
- 1976WA1C Walter and Lisowski, Lowell Conf., CONF-760715-P2 (1976) 1061
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jolley, Phys. Rev. C14 (1976) 815
- 1977AL06 G.D. Alkhazov, Nucl. Phys. A280 (1977) 330
- 1977AL1E Alkhazov et al., Pisma Zh. Eksp. Teor. Fiz. 26 (1977) 110
- 1977AR01 L.G. Arnold, B.C. Clark and R.L. Mercer, Lett. Nuovo Cim. 18 (1977) 151
- 1977AR09 N. Arena, S. Cavallaro, M. Lattuada, R. Potenza, M.L. Sperduto, V. D'Amico, G. Fazio, S. Femino and S. Jannelli, Lett. Nuovo Cim. 19 (1977) 125
- 1977AS01 E. Aslanides, T. Bauer, R. Bertini, R. Beurtey, A. Boudard, F. Brochard, G. Bruge, A. Chaumeaux, H. Catz, J.M. Fontaine et al., Phys. Lett. B68 (1977) 221
- 1977BA19 T. Bauer, A. Boudard, H. Catz, A. Chaumeaux, P. Couvert, M. Garcon, J. Guyot, D. Legrand, J.C. Lugol, M. Matoba et al., Phys. Lett. B67 (1977) 265

- 1977BA1M Bassalleck et al., Proc., Zurich-Sin (1977) 35
- 1977BA1N D. Baye, J. Deenen and Y. Salmon, Nucl. Phys. A289 (1977) 511
- 1977BE1H Beynon and Constantine, J. Phys. (London) G3 (1977) 81
- 1977BE34 J. Berger, J. Duflo, L. Goldzahl, F. Plouin, G. Bizard, C. Le Brun, J. Oostens, M. Van Den Bossche, L. Vu Hai, F.L. Fabbri et al., Lett. Nuovo Cim. 19 (1977) 287
- 1977BE50 F. Becker and I. Strykowski, Nucl. Phys. A289 (1977) 425
- 1977BI05 G. Bizard, C. le Brun, J. Berger, J. Duflo, L. Goldzahl, F. Plouin, J. Oostens, M. van den Bossche, L. Vu Hai, F.L. Fabbri et al., Nucl. Phys. A285 (1977) 461
- 1977BO24 J.E. Bond and F.W.K. Firk, Nucl. Phys. A287 (1977) 317
- 1977BR17 F.P. Brady, N.S.P. King, B.E. Bonner, M.W. McNaughton, J.C. Wang and W.W. True, Phys. Rev. C16 (1977) 31
- 1977BR1C Bruton et al., Proc., Zurich-Sin (1977) 158
- 1977BR1E Bruge, High Energy Phys. Nucl. Phys. Struct., Zurich, 1977 (1977) 69
- 1977BR25 W. Bretfeld, W. Burgmer, H. Eichner, D. Gola, C. Heinrich, H.J. Helten, H. Kretzer, K. Prescher and W. Schnorrenberg, Z. Phys. A282 (1977) 365
- 1977CH1C Chuu and Han, J. Phys. (London) G3 (1977) 555
- 1977CH1D Chwieroth, Tang and Thompson, Fiz. Suppl. 9 (1977) 13
- 1977CL1A W.L. Clinton, K.P. Mandal, and J.A. McClure, Phys. Rev. Lett. 38 (1977) 143
- 1977CO1B H.E. Conzett and F. Seiler, Nucl. Phys. A290 (1977) 93
- 1977DE08 C. Detraz, Phys. Lett. B66 (1977) 333
- 1977DEYM H. de Jong, D. Kamke, J. Krug, K. Lorenzen and U. Weber, in Tokyo (1977) 18
- 1977DO01 D.C. Dodder, G.M. Hale, N. Jarmie, J.H. Jett, P.W. Keaton, Jr., R.A. Nisley and K. Witte, Phys. Rev. C15 (1977) 518
- 1977DR1B Dries et al., Bull. Amer. Phys. Soc. 22 (1977) 1016
- 1977DY01 R. Dymarz and A. Malecki, Phys. Lett. B66 (1977) 413
- 1977GE01 J.V. Geaga, M.M. Gazzaly, G.J. Igo, J.B. McClelland, M.A. Nasser, A.L. Sagle, H. Spinka, J.B. Carroll, V. Perez-Mendez and E.T.B. Whipple, Phys. Rev. Lett. 38 (1977) 1265
- 1977HA06 R.A. Hardekopf and G.G. Ohlsen, Phys. Rev. C15 (1977) 514
- 1977HI09 R.-P. Hirsekorn and L.J. Weigert, Nucl. Phys. A288 (1977) 439
- 1977IG1A Igo et al., Bull. Amer. Phys. Soc. 22 (1977) 562
- 1977IG1B Igo et al., in Tokyo (1977) 13, 15
- 1977IG1C Igo et al., Proc., Zurich-Sin. (1977) 219
- 1977JA07 N. Jarmie and J.H. Jett, Phys. Rev. C16 (1977) 15

- 1977JA12 N. Jarmie, Phys. Rev. C16 (1977) 2090
- 1977KA10 O. Karban, A.K. Basak, C.O. Blyth, W. Dahme, J.B.A. England, J.M. Nelson, N.T. Okumusoglu, S. Roman, G.G. Shute and R. Vlastou, J. Phys. (London) G3 (1977) 571
- 1977KA1D Kaneko and Kamada, Prog. Theor. Phys. 57 (1977) 1277
- 1977KA1E Kanada and Kaneko, in Tokyo (1977) 39
- 1977KA1F Kanada, Kaneko, Nagata and Nomoto, in Tokyo (1977) 40
- 1977KA1G Kanada and Kaneko, in Tokyo (1977) 471
- 1977KA1H Kanada and Kaneko, in Tokyo (1977) 472
- 1977KI1C Kim, Nucl. Instrum. Meth. 145 (1977) 9
- 1977KL03 R. Klem, G. Igo, R. Talaga, A. Wriegat, H. Courant, K. Einsweiler, T. Joyce, H. Kagan, Y. Makdisi, M. Marshak et al., Phys. Rev. Lett. 38 (1977) 1272
- 1977KL08 R. Klem, G. Igo, R. Talaga, A. Wriegat, H. Courant, K. Einsweiler, T. Joyce, H. Kagan, Y. Makdisi, M. Marshak et al., Phys. Lett. B70 (1977) 155
- 1977LA1C M. Lassaut and N. Vinh Mau, Phys. Lett. B70 (1977) 14
- 1977LE1D D.R. Lehman and B.F. Gibson, Phys. Rev. C16 (1977) 1275
- 1977LE1E Lesniak and Lesniak, Proc., Zurich-Sin (1977) 193
- 1977MC1C McNally, Bull. Amer. Phys. Soc. 22 (1977) 581
- 1977ME06 H.O. Meyer, T.P. Cleary and W. Haeberli, Nucl. Instrum. Meth. 140 (1977) 85
- 1977MI13 D. Miljanic, E. Andrade, J. Hudomalj-Gabitzsch, G.S. Mutchler and G.C. Phillips, Nucl. Phys. A290 (1977) 27
- 1977PH01 R.J. Philpott, Nucl. Phys. A289 (1977) 109
- 1977PR06 K. Prescher, W. Bretfeld, W. Burgmer, H. Eichner, H.J. Helten, H. Klein, H. Kretzer, H. Stehle and W.A. Wohlfarth, Nucl. Phys. A286 (1977) 142
- 1977RE1A Redish, AIP Conf. Proc. 37 (1977) 111
- 1977RI1B Rinat, High Energy Phys. Nucl. Struct., Zurich, 1977 (1977) 303
- 1977RO01 E. Rosario-Garcia and R.E. Benenson, Nucl. Phys. A275 (1977) 453
- 1977RO02 P.G. Roos, N.S. Chant, A.A. Cowley, D.A. Goldberg, H.D. Holmgren and R. Woody, III, Phys. Rev. C15 (1977) 69
- 1977RO18 J. Roth, W. Bretfeld, W. Burgmer, H. Eichner, D. Gola, C. Heinrich, H.J. Helten, H. Kretzer, K. Prescher and W. Schnorrenberg, Z. Phys. A283 (1977) 345
- 1977RO1E Roos, AIP Conf. Proc. 37 (1977) 32
- 1977SA1C Saito, Prog. Theor. Phys. Suppl. 62 (1977) 11
- 1977SA1E Saito et al., in Tokyo (1977) 12

- 1977SA21 K. Sagara, M. Hara, N. Takahashi, T. Motobayashi, F. Takeutchi, F. Soga and Y. Nogami, *J. Phys. Soc. Jpn.* 42 (1977) 732
- 1977SE09 F. Seiler, *Nucl. Phys.* A286 (1977) 1
- 1977SE1C F. Seiler and H.W. Roser, *Phys. Lett.* B69 (1977) 411
- 1977SH1B Shimodaya, in *Tokyo* (1977) 43
- 1977SL04 I. Slaus, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, J.M. Lambert, P.A. Treado and R.A. Moyle, *Nucl. Phys.* A286 (1977) 67
- 1977ST06 K. Stephenson, L.D. Knutson and W. Haeberli, *Nucl. Phys.* A277 (1977) 365
- 1977ST1D Stubeda and Tang, Proc. Int. Conf. Nucl. Structure, *Tokyo* (1977); *J. Phys. Soc. Jpn.* Suppl. 44 (1978) 236
- 1977ST30 A.W. Stetz, J.M. Cameron, D.A. Hutcheon, R.H. McCamis, C.A. Miller, G.A. Moss, G. Roy, J.G. Rogers, C.A. Goulding and W.T.H. van Oers, *Nucl. Phys.* A290 (1977) 285
- 1977TE1A Terrien, in *Tokyo* (1977) 843
- 1977TH07 D.R. Thompson, R.E. Brown, M. LeMere and Y.C. Tang, *Phys. Rev.* C16 (1977) 1
- 1977TH09 D.R. Thompson, M. LeMere and Y.C. Tang, *Nucl. Phys.* A286 (1977) 53
- 1977WA05 C.N. Waddell, E.M. Diener, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, A.G. Pieper, R.B. Theus, C.C. Chang and N.S. Chant, *Nucl. Phys.* A281 (1977) 418
- 1977WA06 S.J. Wallace and Y. Alexander, *Phys. Rev. Lett.* 38 (1977) 1269
- 1977WA1C Wallace, *Bull. Amer. Phys. Soc.* 22 (1977) 561
- 1977WA1E Wallace, *Bull. Amer. Phys. Soc.* 22 (1977) 1264
- 1977WE03 R.B. Weisenmiller, N.A. Jolley, D. Ashery, K.H. Wilcox, G.J. Wozniak, M.S. Zisman and J. Cerny, *Nucl. Phys.* A280 (1977) 217
- 1977WE1B R.B. Weisenmiller, Thesis, Univ. of California, Berkeley (1976); LBL-5077 (1977)
- 1977YO04 S.K. Young and C.W. Wong, *Phys. Rev.* C15 (1977) 2146
- 1977YO1B York et al., *Bull. Amer. Phys. Soc.* 22 (1977) 532
- 1978AH01 M.U. Ahmed and P.E. Shanley, *Phys. Rev.* C17 (1978) 1505
- 1978AR10 N. Arena, S. Cavallaro, R. Potenza, V. D'Amico, G. Fazio, S. Femino, G. Giardina, S. Jannelli and M. Lombardi, *Lett. Nuovo Cim.* 23 (1978) 1
- 1978BE30 J. Berger, J. Duflo, L. Goldzahl, J. Oostens, F. Plouin, F.L. Fabbri, P. Picozza, L. Satta, C. Le Brun and D. Legrand, *Phys. Rev. Lett.* 41 (1978) 152
- 1978BI1C Bizard and Tekou, Proc., Cluster, *Winnipeg* (1978) D1
- 1978BR1E Brown, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, *Winnipeg* (1978)

- 1978CA05 J.M. Cameron, L.G. Greeniaus, D.A. Hutcheon, R.H. McCamis, M.S. de Jong, B.T. Murdoch, W.T.H. van Oers, J.G. Rogers and A.W. Stetz, Phys. Lett. B74 (1978) 31
- 1978CA13 G. Calvi, S. Cavallaro, A.S. Figuera, R. Fonte and C. Milone, Lett. Nuovo Cim. 22 (1978) 536
- 1978CA1E Carey et al., Bull. Amer. Phys. Soc. 23 (1978) 594
- 1978CH1G Chant, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, Winnipeg (1978)
- 1978CL1C Clark, Arnold and Mercer, Bull. Amer. Phys. Soc. 23 (1978) 964
- 1978EP1C Epstein et al., Bull. Amer. Phys. Soc. 23 (1978) 625
- 1978FI1E Fick, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, Winnipeg (1978)
- 1978FO1G Fonseca, Revai and Matveenko, Proc., Cluster, Winnipeg (1978) B1
- 1978FO33 J. Fong, T.S. Bauer, G.J. Igo, G. Paulette, R. Ridge, R. Rolfe, J. Soukup, C.A. Whitten, Jr., G.W. Hoffmann, N. Hintz et al., Phys. Lett. B78 (1978) 205
- 1978FR02 H. Friedrich, Nucl. Phys. A294 (1978) 81
- 1978GL1B V.V. Glagolev, R.M. Lebedev, I.S. Saitov, V.N. Streletsov, L.I. Zuravleva, A.N. Gorbunov, K.U. Khayretdinov, G. Martinska, I. Patocka, M. Seman et al., Phys. Rev. C18 (1978) 1382
- 1978GO1D J. Gosset, J.I. Kapusta and G.D. Westfall, Phys. Rev. C18 (1978) 844
- 1978GR07 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.V. Ostashko, V.M. Pugach, V.A. Pilipchenko, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 147; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 126
- 1978GU15 O.P. Gupta, A. Bond and P.M.S. Lesser, Phys. Rev. C18 (1978) 1075
- 1978HE2B E.M. Henley, Nucl. Phys. A300 (1978) 273
- 1978IG1A Igo et al., Bull. Amer. Phys. Soc. 23 (1978) 47
- 1978IG1B Igo, Rev. Mod. Phys. 50 (1978) 523
- 1978KA11 T. Kambara, M. Takai, M. Nakamura and S. Kobayashi, J. Phys. Soc. Jpn. 44 (1978) 704
- 1978KO13 Y. Koike, Nucl. Phys. A301 (1978) 411
- 1978KO1F Koene et al., Proc., Cluster, Winnipeg (1978) D9
- 1978LA02 V. Layly and R. Schaeffer, Phys. Rev. C17 (1978) 1145
- 1978LE1H Le Mere and Tang, Proc., Cluster, Winnipeg (1978) A21
- 1978LE23 S.W-L. Leung and H.S. Sherif, Can. J. Phys. 56 (1978) 1116
- 1978MA37 Z. Majka, Phys. Lett. B76 (1978) 161

- 1978MC06 R.H. McCamis, J.M. Cameron, L.G. Greeniaus, D.A. Hutcheon, C.A. Miller, G.A. Moss, G. Roy, M.S. de Jong, B.T. Murdoch, W.T.H. van Oers et al., Nucl. Phys. A302 (1978) 388
- 1978MC1C McNally, Nucl. Fusion 18 (1978) 133
- 1978ME05 G. Mechtersheimer, G. Buche, U. Klein, W. Kluge, H. Matthay, D. Munchmeyer and A. Moline, Phys. Lett. B73 (1978) 115
- 1978ME1C A.Z. Mekjian, Phys. Rev. C17 (1978) 1051
- 1978NA08 H. Nakamura, H. Noya, S.E. Darden and S. Sen, Nucl. Phys. A305 (1978) 1
- 1978NA1H M.A. Nasser, M.M. Gazzaly, J V. Geaga, B. Hoistad, G. Igo, J.G. McClelland, A.L. Sagle, H. Spinka, J.B. Carroll, V. Perez-Mendez et al., Phys. Rev. C17 (1978) 1748
- 1978RO06 E. Rost, J.R. Shepard and D.A. Sparrow, Phys. Rev. C17 (1978) 1513
- 1978SA07 K. Sagara, T. Motobayashi, N. Takahashi, Y. Hashimoto, M. Hara, Y. Nogami, H. Nakamura and H. Noya, Nucl. Phys. A299 (1978) 77
- 1978SE01 F. Seiler, F.N. Rad, H.E. Conzett and R. Roy, Nucl. Phys. A296 (1978) 205
- 1978SL1B Slaus, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 57
- 1978SM1C Smirnov et al., Bull. Amer. Phys. Soc. 23 (1978) 51
- 1978SP03 C. Spitaleri, M. Lattuada, F. Riggi, N. Arena and D. Vinciguerra, Lett. Nuovo Cim. 21 (1978) 345
- 1978YE1A Yeh, Bull. Amer. Phys. Soc. 23 (1978) 553
- 1978YO1A York et al., Bull. Amer. Phys. Soc. 23 (1978) 602
- 1978YO1B S.K. Young and C.W. Wong, Phys. Lett. B77 (1978) 41

