

Energy Levels of Light Nuclei $A = 8$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A320* (1979), p. 1. This version of $A = 8$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed 1978)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 8$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^8\text{n}\$](#) , [\${}^8\text{He}\$](#) , [\${}^8\text{Li}\$](#) , [\${}^8\text{Be}\$](#) , [\${}^8\text{B}\$](#) , [\${}^8\text{C}\$](#)

B. Tables of Recommended Level Energies:

[Table 8.1](#): Energy levels of ${}^8\text{Li}$

[Table 8.3](#): Energy levels of ${}^8\text{Be}$

[Table 8.11](#): Energy levels of ${}^8\text{B}$

C. [References](#)

D. Figures: [\${}^8\text{Li}\$](#) , [\${}^8\text{Be}\$](#) , [\${}^8\text{B}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

⁸n
(Not illustrated)

⁸n has not been observed in the interaction of 0.7 and 400 GeV protons with uranium: the cross section is $< 2.3 \times 10^{-5} \mu\text{b}$ (1977TU02) at 0.7 GeV and $< 0.2 \mu\text{b}$ at 400 GeV (1977TU03). See also (1977DE08) and (1978SA1E).

⁸He
(Figs. 11 and 14)

GENERAL: (See also (1974AJ01).)

See (1973AL1B, 1973TO16, 1974IR04, 1974MA1E, 1975AB1D, 1975BE31, 1976BE1G, 1976IR1B, 1976VA29, 1978KO1H, 1978NA07).

Mass of ⁸He: The atomic mass excess of ⁸He is 31596 ± 7 keV (1977TR07). See also (1974CE05, 1975JA10, 1975KO18, 1978RO01). ⁸He is then stable with respect to decay into ⁶He + 2n by 2.141 MeV. See also (1974AJ01, 1976TR01) and (1975JE02; theor.). The IMME coefficients based on the latest masses of the $T = 2$ states in $A = 8$ are calculated by (1977TR07).

1. ⁸He(β^-)⁸Li $Q_m = 10.649$

The half-life is 122 ± 2 msec. The decay takes place 88% to ⁸Li*(0.98) [$\log ft = 4.19^\dagger$] [consistent with $J^\pi = 1^+$ for ⁸Li*(0.98)] and $(12 \pm 1)\%$ via ⁸Li states which decay by neutron emission (1965PO06). See also (1973AN11, 1976DE31).

2. ¹⁸O(α , ¹⁴O)⁸He $Q_m = -37.962$

At $E_\alpha = 58$ MeV, $\theta_{\text{lab}} = 8^\circ$, the differential cross for formation of ⁸He_{g.s.} is 40 ± 15 nb/sr (1975JA10).

3. ²⁶Mg(α , ²²Mg)⁸He $Q_m = -44.989$

This reaction has been studied at $E_\alpha = 80$ MeV (1966CE01) and 110.6 MeV (1974CE05).

[†] M.J. Martin, Nucl. Data Project, private communication.

${}^8\text{Li}$
(Figs. 11 and 14)

GENERAL: (See also (1974AJ01).)

Nuclear models: (1975KH1A, 1977ST24).

Special states: (1974IR04, 1976IR1B, 1978KH03).

Electromagnetic interactions: (1974KU06, 1976KU07).

Special reactions: (1973SI38, 1974BA70, 1974BA1N, 1974BO08, 1975FE1A, 1975ZE01, 1976BE67, 1976BO08, 1976BU16, 1977FE1B, 1977PR05, 1977ST1J, 1977YA1B, 1978DI04).

Muon and neutrino interactions: (1977BA1P).

Pion and kaon reactions (See also reaction 16.): (1968ZE1A, 1973AG05, 1973AG06, 1973CA1C, 1974BO1F, 1974ZI1A, 1975CO06, 1975KU1J, 1977JU1B, 1977MA1K, 1978MO01, 1978VE1B).

Astrophysical questions: (1977SC1D, 1977ST1J).

Other topics: (1974IR04, 1974SE1B, 1975KH1A, 1976IR1B, 1977TR07, 1978GA1C, 1978RO01).

Ground state properties: (1974KU06, 1974SHYR, 1975BE31, 1976FU06, 1977TR07, 1978RO01, 1978ZA1D).

$$J = 2 \text{ (1973NE10);}$$

$$\mu = +1.65335 \pm 0.00035 \text{ nm (1973HA12). See also (1974AJ01);}$$

$$Q = 24 \pm 2 \text{ mb (1975MI04);}$$

$$= 32 \pm 6 \text{ mb (1974AC03);}$$

$$= 32 \text{ mb (1977DU06).}$$

1. ${}^8\text{Li}(\beta^-){}^8\text{Be}$ $Q_m = 16.005$

The β -decay leads mainly to ${}^8\text{Be}^*(2.9)$: see ${}^8\text{Be}$. The adopted half-life [see (1974AJ01)] = 842 ± 6 msec, $\log ft = 5.61$ (B. Zimmerman, private communication). The distribution of recoil momenta indicates $J^\pi = 2^+$ for ${}^8\text{Li}_{\text{g.s.}}$. The coefficient for the angular correlation between β^- , $\bar{\nu}_e$ and α -particles, $b = -0.88 \pm 0.08$ (1965GR25), -1.01 ± 0.07 (1966EI02), in substantial agreement with $b = -1$, expected from axial vector coupling. (1975MI04) have looked at the symmetry of the β -decay in polarized ${}^8\text{Li}$ nuclei and have determined the quadrupole moment for ${}^8\text{Li}_{\text{g.s.}}$: see the “GENERAL” section here. See also reaction 28 in ${}^8\text{Be}$.

Table 8.1: Energy levels of ${}^8\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$2^+; 1$	$\tau_{1/2} = 842 \pm 6$ msec	β^+	1, 2, 3, 10, 11, 13, 14, 15, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 31
0.9808 ± 0.1	$1^+; 1$	$\tau_m = 12 \pm 4$ fsec	γ	2, 10, 12, 16, 17, 18, 19, 23, 25, 26, 27, 28, 29
2.255 ± 3	$3^+; 1$	$\Gamma = 33 \pm 6$ keV	γ, n	2, 3, 4, 17, 18, 19, 23, 26
3.21	$1; 1$	≈ 1000	n	5
5.4		≈ 650	n	4, 5
6.1 ± 100	$(3); 1$	≈ 1000	n	4, 5
6.53 ± 20	≥ 4	35 ± 15	n	2, 4, 18, 19, 23
7.1 ± 100		≈ 400	n	4
(9)		≈ 6000		17, 18
10.8222 ± 5.5	$0^+; 2$	< 12		21

 2. ${}^6\text{Li}(t, p){}^8\text{Li}$

$$Q_m = 0.801$$

Angular distributions have been obtained at $E_t = 23$ MeV for the proton groups to ${}^8\text{Li}^*(0, 0.98, 2.26, 6.54 \pm 0.03)$; $\Gamma_{\text{c.m.}}$ for ${}^8\text{Li}^*(2.26, 6.54) = 35 \pm 10$ and 35 ± 15 keV, respectively; J for the latter is ≥ 4 (1978AJ02).

 3. ${}^7\text{Li}(n, \gamma){}^8\text{Li}$

$$Q_m = 2.033$$

The cross section for capture radiation has been measured for $E_n = 40$ to 1000 keV: it decreases from $50 \mu\text{b}$ to $5 \mu\text{b}$ over the interval. The cross section shows the resonance corresponding to ${}^8\text{Li}^*(2.26)$: $\Gamma_\gamma = 0.07 \pm 0.03$ eV (1959IM04); $E_{\text{res}} = 254 \pm 3$ keV, $\Gamma_n = 31 \pm 7$ keV (1971AL09): see Table 8.2. For unpublished data see (1974AJ01). See also (1976GAYV).

 4. ${}^7\text{Li}(n, n){}^7\text{Li}$

$$E_b = 2.033$$

The thermal cross section is 1.07 ± 0.04 b (1960HU08); the coherent scattering length (thermal, bound) is -2.1 ± 0.1 fm (see (1973MU14)). See also (1974RO20). Total cross-section measurements have been reported for $E_n = 5$ eV to 30 MeV [see (1974AJ01, 1976GAYV)] and

Table 8.2: Resonance parameters for ${}^8\text{Li}^*(2.26)$ (1970ME1C) ^a

E_{res} (keV) ^b	261.2
E_x (MeV) ^c	2.261
Γ (keV)	35 ± 5 ^d
$\Gamma_n(E_r)$ (keV)	36.5
Γ_γ (eV) ^c	0.07 ± 0.03 ^e
γ_n^2 (keV)	594
θ^2	0.091
radius (fm)	3.30
σ_{max} ^b	12.0
J^π	3^+
l_n	1

^a Energies in lab system except for those labeled ^c.

^b $E_{\text{res}} = 254 \pm 3$ keV, $\Gamma_n = 31 \pm 7$ keV (1971AL09).

^c Energies in c.m. system.

^d (1960HU08).

^e (1959IM04).

by (1976KN1D, 1976LA1C; $E_n = 4$ to 7.5 MeV), (1974HY01; 14.1 MeV) and (1976MI1C; 14.6 MeV). For polarization measurements see (1966LA04, 1974AJ01). For angular distribution measurements see ${}^7\text{Li}$.

A pronounced resonance is observed at $E_n = 261$ keV with $J^\pi = 3^+$, formed by p-waves (Table 8.2) (1970ME1C). A good account of the polarization is given by the assumption of levels at $E_n = 0.25$ and 3.4 MeV, with $J^\pi = 3^+$ and 2^- , together with a broad $J^\pi = 3^-$ level at higher energy (1964LA19). Broad peaks are reported at $E_n = 4.6$ and 5.8 MeV (± 0.1 MeV) [${}^8\text{Li}^*(6.1, 7.1)$] with $\Gamma \approx 1.0$ and 0.4 MeV, respectively, and there is indication of a narrow peak at $E_n = 5.1$ MeV [${}^8\text{Li}^*(6.5)$] with $\Gamma \ll 80$ keV and of a weak, broad peak at $E_n = 3.7$ MeV (1971FO1P, 1971FO24) [and see (1974AJ01)]. See also (1977ST24, 1978ST20; theor.).

5. ${}^7\text{Li}(n, n'){}^7\text{Li}^*$

$$E_b = 2.033$$

The excitation function for 0.48 MeV γ -rays shows an abrupt rise from threshold (indicating s-wave formation and emission) and a broad maximum ($\Gamma \approx 1$ MeV) at $E_n = 1.35$ MeV. A good

fit is obtained with either $J^\pi = 1^-$ or 1^+ (2^+ not excluded), $\Gamma_{\text{lab}} = 1.14$ MeV (1955FR10). At higher energies a prominent peak is observed at $E_n = 3.8$ MeV ($\Gamma_{\text{lab}} = 0.75$ MeV) and there is some indication of a broad resonance ($\Gamma_{\text{lab}} = 1.30$ MeV) at $E_n = 5.0$ MeV. The two resonances are interpreted as being due to states with $J^\pi = 2^+$ or 3^+ [$^8\text{Li}^*(5.4)$] [see, however, (1978ST20)] and $J^\pi = 3^+$ or 3^- [$^8\text{Li}^*(6.4)$] [(1972PR03): Γ_λ , γ_λ^2 and θ_λ^2 are listed for these two states under various assumptions]. (1977BA62) predicts a 2^- state at $E_x = 2.74$ MeV and 1^- states at 3.52 and 5.4 MeV with $\Gamma \approx 1.8, 2.5$ and 3.8 MeV. The excitation function for 0.48 MeV γ -rays has been measured at 125° for $E_n = 0.4$ to 20 MeV: it shows the structures described above followed by a decrease in the cross section from $E_n = 6$ to 20 MeV with some evidence of structures at $E_n = 6.8$ and 8 MeV (1977DI11).

Cross-section measurements have also been reported for $E_n = 0.57$ to 4.0 MeV (1976SM08), 4.0 to 7.5 MeV (1977KN1B) and 7 to 13 MeV (1976BI1B; n_{0+1} , n_2). See also (1976MI1C, 1978LI1C) and the review in (1976GAYV). For applications see (1974ST1D).

$$6. \text{}^7\text{Li}(n, 2n)^6\text{Li} \qquad Q_m = -7.251 \qquad E_b = 2.033$$

See (1974KO35, 1976GAYV).

$$7. \text{}^7\text{Li}(n, p)^7\text{He} \qquad Q_m = -10.40 \qquad E_b = 2.033$$

See (1977RI07) and ^7He . See also (1976SL2A).

$$8. \text{}^7\text{Li}(n, d)^6\text{He} \qquad Q_m = -7.751 \qquad E_b = 2.033$$

See ^6He . See also (1976SL2A).

$$9. \text{}^7\text{Li}(n, t)^4\text{He} + n \qquad Q_m = -2.467 \qquad E_b = 2.033$$

The cross section rises to 450 mb at $E_n \approx 8$ MeV and thereafter decreases slowly to 300 mb at $E_n = 15$ MeV (1976GAYV). See also (1975VO1C, 1978QA01), (1974ST1D; applied) and (1976SL2A).

$$10. \text{}^7\text{Li}(d, p)^8\text{Li} \qquad Q_m = -0.192$$

Angular distributions of the p_0 and p_1 groups [$l_n = 1$] at $E_d = 12$ MeV have been analyzed by DWBA: $S_{\text{exp}} = 0.87$ and 0.48 respectively for ${}^8\text{Li}^*(0, 0.98)$ [$S_{\text{exp}}/S_{\text{theor}} = 0.84, 1.09$] (1967SC29). Angular distributions have also been measured at several energies in the range $E_d = 0.49 \rightarrow 3.44$ MeV (p_0) and 0.95 to 2.94 MeV (p_1) by (1975MC02). The first excited state decays by emitting a 980 ± 10 keV γ -ray. The transition is M1 (1962CH14). The lifetime of ${}^8\text{Li}^*(0.98)$ is 10.1 ± 4.5 fsec (1971TH02). See also (1975MI04, 1978HA1K), (1974AJ01, 1976HA1J) and (1974LO10, 1975KU27, 1976KU07; theor.).

11. ${}^7\text{Li}(t, d){}^8\text{Li}$ $Q_m = -4.225$

See ${}^{10}\text{Be}$ in (1974AJ01).

12. ${}^8\text{He}(\beta^-){}^8\text{Li}$ $Q_m = 10.649$

See ${}^8\text{He}$.

13. ${}^9\text{Be}(\gamma, p){}^8\text{Li}$ $Q_m = -16.888$

See ${}^9\text{Be}$.

14. ${}^9\text{Be}(e, ep){}^8\text{Li}$ $Q_m = -16.888$

At $E_e = 801$ MeV two groups are observed at $Q = -28.7 \pm 0.1$ and -16.9 ± 0.1 MeV corresponding to the ejection of $1s$ and $1p$ protons (1974GO35). See also ${}^9\text{Be}$ and (1977BE1Q, 1977MO1F).

15. ${}^9\text{Be}(n, d){}^8\text{Li}$ $Q_m = -14.663$

See ${}^{10}\text{Be}$.

16. ${}^9\text{Be}(\pi^+, \pi^+p){}^8\text{Li}$ $Q_m = -16.888$

The population of ${}^8\text{Li}^*(0.98)$ and its subsequent γ -decay are reported by (1978MO01).

$$17. \text{}^9\text{Be}(p, 2p)\text{}^8\text{Li} \quad Q_m = -16.888$$

The summed proton spectrum at $E_p = 156$ MeV shows peaks corresponding to ${}^8\text{Li}(0)$ and ${}^8\text{Li}^*(0.98 + 2.26)$ [unresolved]. In addition s-states [$J^\pi = 1^-, 2^-$] are suggested at $E_x = 9$ and 16 MeV, with $\Gamma_{\text{c.m.}} \approx 6$ and 8 MeV: the latter may actually be due to continuum protons (1967RO06). See also ${}^9\text{Be}$.

$$18. \text{}^9\text{Be}(d, \text{}^3\text{He})\text{}^8\text{Li} \quad Q_m = -11.394$$

Angular distributions have been reported for the ${}^3\text{He}$ ions to ${}^8\text{Li}^*(0, 0.98, 2.26, 6.53)$ at $E_d = 27.97$ MeV (1977OO01; $C^2S(\text{abs.}) = 1.63, 0.61, 0.48, 0.092$) and 52 MeV (1975SC41; $S_{p_{3/2}} = 1.50, 0.63, 0.50$ for the first three states, with the ground state value normalized to the predicted theoretical value). The distributions to ${}^8\text{Li}^*(6.53)$ [$\Gamma < 100$ keV (1974DI1A, 1975DI1B)] are featureless (1975DI1B, 1975SC41, 1977OO01). See reaction 32 [${}^9\text{Be}(d, t)$] in ${}^8\text{Be}$ (1977OO01). At $E_d = 80$ MeV, (1974DI1A, 1975DI1B; abstracts) report a broad peak at $E_x \approx 9$ MeV corresponding to s-shell pickup. See also (1975RO1E).

$$19. \text{}^9\text{Be}(t, \alpha)\text{}^8\text{Li} \quad Q_m = 2.926$$

At $E_t = 12.98$ MeV, α -particle groups are observed to ${}^8\text{Li}^*(0, 0.98, 2.26)$ and to a state at $E_x = 6.530 \pm 0.020$ MeV with $\Gamma_{\text{c.m.}} < 40$ keV (1965WA12). For angular distributions to these four states, see (1968AJ01): the distribution of the α -particles to ${}^8\text{Li}^*(6.53)$ is rather featureless and does not involve a forward maximum, suggesting $l > 1$. A large l -transfer is consistent with the narrow width of this unbound state (1968AJ01). See also (1974AJ01) for unpublished measurements and earlier work.

The mean lifetime, $\tau_m = 14 \pm 5$ fsec for ${}^8\text{Li}^*(0.98)$: $E_x = 980.80 \pm 0.10$ keV (1972CO09). See also ${}^{12}\text{B}$ in (1975AJ02).

$$20. \text{}^9\text{Be}(\text{}^7\text{Li}, \text{}^8\text{Be})\text{}^8\text{Li} \quad Q_m = 0.368$$

See (1974KE1C) and (1974AJ01).

$$21. \text{}^{10}\text{Be}(p, \text{}^3\text{He})\text{}^8\text{Li} \quad Q_m = -15.982$$

At $E_p = 45$ MeV ^3He ions are observed to a state of $E_x = 10.8222 \pm 0.0055$ MeV ($\Gamma_{\text{c.m.}} < 12$ keV): the angular distributions for the transition to this state, and to its analog ($^8\text{Be}^*(27.49)$), measured in the mirror reaction [$^{10}\text{Be}(p, t)^8\text{Be}$] are very similar. They are both consistent with $L = 0$ using a DWBA (LZR) analysis (1975RO01). The particle decay of the $T = 2$ state has been studied by (1975ADZV; abstract).

$$22. \ ^{10}\text{Be}(d, \alpha)^8\text{Li} \quad Q_m = 2.372$$

See ^{12}B in (1975AJ02).

$$23. \ ^{10}\text{B}(^6\text{Li}, ^8\text{B})^8\text{Li} \quad Q_m = -17.730$$

At $E(^6\text{Li}) = 80$ MeV, $^8\text{Li}^*(0, 0.98, 2.26, 6.53)$ are populated (1976WE09, 1977WE1B).

$$24. \ ^{11}\text{B}(\gamma, ^3\text{He})^8\text{Li} \quad Q_m = -27.210$$

See (1963NE07).

$$25. \ ^{11}\text{B}(n, \alpha)^8\text{Li} \quad Q_m = -6.633$$

Angular distributions for the α_0 and α_1 groups have been measured at $E_n = 14.1$ MeV (1973BO26). See also ^{12}B in (1975AJ02) and (1977AN1C).

$$26. \ ^{12}\text{C}(\alpha, ^8\text{B})^8\text{Li} \quad Q_m = -41.444$$

At $E_\alpha = 129$ MeV, ^8B groups are observed to $^8\text{Li}^*(0, 0.98, 2.26)$ (1968MC02).

$$27. \ ^{13}\text{C}(^3\text{He}, ^8\text{B})^8\text{Li} \quad Q_m = -25.812$$

See (1976BE1L).

$$28. \ ^{13}\text{C}(^7\text{Li}, ^{12}\text{C})^8\text{Li} \quad Q_m = -2.914$$

At $E(^7\text{Li}) = 34$ MeV angular distributions have been measured for the transitions to $^8\text{Li}_{\text{g.s.}} + ^{12}\text{C}_{\text{g.s.}}$, $^8\text{Li}_{0.98}^* + ^{12}\text{C}_{\text{g.s.}}$, $^8\text{Li}_{\text{g.s.}} + ^{12}\text{C}_{4.4}^*$ and $^8\text{Li}_{0.98}^* + ^{12}\text{C}_{4.4}^*$ (1973SC26).

29. $^{14}\text{N}(^3\text{He}, ^9\text{C})^8\text{Li}$ $Q_{\text{m}} = -32.064$

At $E(^3\text{He}) = 76$ MeV $^8\text{Li}^*(0, 0.98)$ are populated (1976RO04).

30. $^{18}\text{O}(^7\text{Li}, ^{17}\text{O})^8\text{Li}$ $Q_{\text{m}} = -6.012$

See (1974AJ01).

31. $^{24}\text{Mg}(^3\text{He}, ^{19}\text{Na})^8\text{Li}$ $Q_{\text{m}} = -32.876$

See (1975BE38).

⁸Be
(Figs. 12 and 14)

GENERAL: (See also (1974AJ01).)

Shell model: (1973AR1C, 1974KA11, 1975GO07, 1975SC1K, 1976AR07, 1976JI1A, 1976ST04).

Collective, rotational and deformed models: (1974BO25, 1974LE04, 1975AR28, 1975KH1A).

Cluster and α -particle models: (1970YU02, 1973AB1A, 1973AR1C, 1974CH01, 1974DR05, 1974KA11, 1975AB1E, 1975SC1K, 1976BA1N, 1976ST04, 1977AR08, 1977BE50, 1977FU1D, 1977FU1F, 1977HE10, 1977HE1C, 1977KA1K, 1977KI10, 1977SE1D, 1977SM1A, 1977ZA04, 1978AR1M, 1978ST10, 1978TO09).

Special levels: (1973DO1B, 1974HA1G, 1974IR04, 1974KA11, 1974KU06, 1974MC04, 1974NE1B, 1975SH01, 1976IR1B, 1976JI1A, 1977BA62, 1977FU1F, 1977KA1K, 1978AR1M, 1978BA31, 1978BA66, 1978SE1C, 1978SH04, 1978ST10).

Electromagnetic transitions: (1974HA1C, 1974KU06, 1976KU07, 1977HE1C, 1977KU20, 1978TO09).

Special reactions: (1974GA1E, 1974KO16, 1975FE1A, 1975UN1A, 1976AR02, 1976VA29, 1976WO11, 1977GA1F, 1978ZO1A).

Pion and kaon reactions: (1973AR1B, 1973CA1C, 1974AM01, 1974BO1F, 1974ZI1A, 1977BA51).

Other topics: (1973DO1B, 1973KR1B, 1973PE05, 1974BE1L, 1974BEYR, 1974CH01, 1974DZ03, 1974IR04, 1974KU06, 1974MC04, 1975AR10, 1975AR28, 1975ER09, 1975GO07, 1975KH1A, 1975KO1C, 1975SH01, 1976AR07, 1976IR1B, 1976KH04, 1976KU07, 1976SA16, 1977AR08, 1977BE1U, 1977HE10, 1977TR07, 1978BA31, 1978JE1B, 1978KA1G, 1978MC04, 1978RO01, 1978RO17, 1978SH04).

Ground state of ⁸Be: (1973AB1A, 1973DO1B, 1973FA1C, 1973PA20, 1973PE05, 1974DR05, 1974DZ03, 1975BE31, 1976BE1G, 1976IR1B, 1976OV1B, 1978RO17).

1. ${}^8\text{Be} \rightarrow {}^4\text{He}{}^4\text{He}$ $Q_m = 0.09188$

$\Gamma_{\text{c.m.}}$ for the ${}^8\text{Be}_{\text{g.s.}} = 6.8 \pm 1.7$ eV (1968BE02). See also (1974FE1B, 1976FE1A) and (1974LI16; theor.).

2. ${}^4\text{He}(\alpha, \gamma){}^8\text{Be}$ $Q_m = -0.09188$

The yield of γ_1 has been measured for $E_\alpha = 32$ to 36 MeV: see (1975NA12, GA77F, 1977PA26). An angular correlation measurement at the resonances corresponding to ${}^8\text{Be}^*(16.6 + 16.9) [2^+; T = 0 + 1]$ gives $\delta = 0.19 \pm 0.03$ (1978BO30); $\Gamma_\gamma(\text{M1}) = 6.1 \pm 0.53$ eV (1978BO30), 7.4 ± 1.0 eV (1977PA26), 5.50 ± 0.60 eV (1977LO1D, 1977LO1E). On the basis of these results there is no evidence for violation of CVC or for the existence of second-class currents: see (1978BO30). (1977PA26) find $E_x = 3.18 \pm 0.05$ MeV for the energy of the first excited state (see also Table 8.4 in (1974AJ01)). The yield of γ_0 for $E_\alpha = 33$ to 38 MeV is twenty times lower than that for γ_1 , consistent with E2 decay (1977GAZZ). See also (1975NA1C) and (1977GA1E).

Table 8.3: Energy Levels of ${}^8\text{Be}$ ^a

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$0^+; 0$	6.8 ± 1.7 eV	α	1, 2, 4, 11, 12, 13, 14, 15, 21, 22, 23, 24, 25, 27, 30, 31, 32, 33, 34, 36, 38, 39, 40, 41, 42, 43, 45, 47, 48, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 64, 66, 68, 69
2.94 ± 30	$2^+; 0$	1560 ± 30	α	2, 4, 12, 13, 14, 15, 21, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 37, 38, 39, 40, 41, 42, 43, 45, 47, 48, 50, 51, 52, 53, 54, 57, 61
11.4 ± 300	$4^+; 0$	≈ 3500 ^b	α	4, 13, 21, 24, 31, 33, 40, 43, 45, 52, 53, 54
16.626 ± 3	$2^+; 0 + 1$	108.1 ± 0.5	γ, α	2, 4, 12, 14, 15, 21, 23, 24, 29, 31, 32, 33, 38, 40, 45, 47, 48, 52, 53
16.922 ± 3	$2^+; 0 + 1$	74.0 ± 0.4	γ, α	2, 4, 12, 14, 15, 21, 23, 24, 31, 32, 33, 38, 40, 45, 47, 48, 52, 53
17.641 ± 1.5	$1^+; 1$	10.7 ± 0.5	γ, p	5, 12, 15, 17, 21, 23, 31, 32, 33, 40, 45, 47, 53
18.150 ± 4	$1^+; 0$	138 ± 6	γ, p	12, 15, 17, 21, 31, 32, 33, 40, 45

Table 8.3: Energy Levels of ${}^8\text{Be}$ ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
18.91	2^-	48 ± 20	γ, n, p	12, 15, 16, 17, 21, 31, 45
19.069 ± 10	$3^+; (1)$	270 ± 20	γ, p	12, 15, 17, 21, 32, 45
19.24 ± 25	$3^+; (0)$	230 ± 30	n, p	16, 17, 21, 31, 32, 33, 40
19.4	1^-	≈ 650	n, p	12, 16, 17
19.86 ± 50	$4^+; 0$	700 ± 100	p, α	4, 12, 14, 20, 24, 32, 48
20.1	$2^+; 0$	≈ 1100	n, p, α	3, 4, 16, 20, 24, 32, 48
20.2	$0^+; 0$	< 1000	α	4
20.9	4^-	1600 ± 200	p	17
21.5	$3^{(+)}$	1000	n, p	16, 31
22.0 ^c	$1^-; 1$	≈ 4000	γ, p	15
22.05 ± 100		270 ± 70		33
22.2	$2^+; 0$	≈ 800	n, p, d, α	3, 4, 10, 17, 20
22.63 ± 100		100 ± 50		14, 33
22.98 ± 100		230 ± 50		31, 33
24.0 ^c	$(1, 2)^-; 1$	≈ 7000	γ, p	15
25.2	$2^+; 0$		p, d, α	4, 10, 20, 33
25.5	$4^+; 0$	broad	d, α	4, 10
27.494 ± 2 ^d	$0^+; 2$	5.5 ± 2.0	γ, p, d, α	3, 5, 7, 8, 10, 35
(28.6)		broad	γ, p	15

^a See also Table 8.7.

^b I am greatly indebted to Prof. F.C. Barker for enlightening discussions concerning the width of ${}^8\text{Be}^*(11.4)$.

^c Giant resonance: see reaction 15.

^d See Table 8.4.

3. (a) ${}^4\text{He}(\alpha, n){}^7\text{Be}$ $Q_m = -18.992$ $E_b = -0.09188$
 (b) ${}^4\text{He}(\alpha, p){}^7\text{Li}$ $Q_m = -17.347$
 (c) ${}^4\text{He}(\alpha, d){}^6\text{Li}$ $Q_m = -22.3733$

Table 8.4: Parameters of the first $T = 2$ state in ${}^8\text{Be}$

E_x (MeV)	27.4922 ± 0.0026	(1975RO01)
	27.4958 ± 0.0024	(1976NO07)
Γ (keV)	10 ± 3	(1969BL14)
	14.7 ± 4.0	(1976HI04)
	12 ± 3	(1975RO01)
	5.5 ± 2.0	(1976NO07)
$\Gamma_{\gamma_5}^b$	23 ± 4 eV	(1976NO07)
Γ_{p_0}/Γ	$0.012 < \Gamma_{p_0}/\Gamma < 0.023^a$	(1976HI04)
Γ_{p_2}/Γ	> 0.005	(1976HI04)
Γ_{d_0}/Γ	> 0.0014	(1976HI04)
Γ_{d_1}/Γ	> 0.0018	(1976HI04)
Γ_{α_0}/Γ	< 0.003	(1976HI04)
c		

^a Lower limit from ${}^4\text{He}(\alpha, p_0)$; upper limit from ${}^7\text{Li}(p, p_0)$ (1976HI04).

^b $\Gamma_{\gamma_1} < 0.01$ eV (1977LO1F; prelim.).

^c $\Gamma_d\Gamma_\gamma/\Gamma < 2.5$ eV (1975GE1F; prelim.).

The cross sections for formation of ${}^7\text{Li}^*(0, 0.48)$ [$E_\alpha = 39$ to 49.5 MeV] and ${}^7\text{Be}^*(0, 0.43)$ [39.4 to 47.4 MeV] both show structures at $E_\alpha \approx 40.0$ and ≈ 44.5 MeV: they are due predominantly to the 2^+ states ${}^8\text{Be}^*(20.1, 22.2)$. The cross sections for (a) and (b) are of interest in accounting for the abundance of ${}^7\text{Li}$ in the galaxy (1977KI12). Cross sections for p_{0+1} are also reported at $E_\alpha = 60.2, 92.4$ and 140.0 MeV (1975KI14). See also (1977RA1B). The cross section for reaction (c) has been measured at three energies in the range $E_\alpha = 46.7$ to 49.5 MeV (1977KI12). See also (1977KI05). (1976HI04) have measured the excitation functions for p_0, p_2, d_0, d_1 for $E_\alpha = 54.96$ to 55.54 MeV to study the decay of the first $T = 2$ state in ${}^8\text{Be}$: see Table 8.4. See also (1975MA1G), (1977AU1B, 1977KO1J; astrophys.) and ${}^6\text{Li}, {}^7\text{Li}$ and ${}^7\text{Be}$.

4. ${}^4\text{He}(\alpha, \alpha){}^4\text{He}$

$$E_b = -0.09188$$

The $\alpha\alpha$ scattering reveals the ground state as a resonance with $Q_0 = 92.12 \pm 0.05$ keV, $\Gamma_{c.m.} = 6.8 \pm 1.7$ eV, [$\tau = (0.97 \pm 0.24) \times 10^{-16}$ sec] (1968BE02). For $E_\alpha = 30$ to 70 MeV the $l = 0$ phase shift shows resonant behavior at $E_\alpha = 40.7$ MeV, corresponding to a 0^+ state at $E_x = 20.2$ MeV,

Table 8.5: Some ${}^8\text{Be}$ states with $16.6 < E_x < 23.0$ MeV ^a

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Reaction	Refs.
16.627 ± 5	113 ± 3	${}^7\text{Li}({}^3\text{He}, \text{d})$	(1967MA12) ^b
	90 ± 5	${}^{10}\text{B}(\text{d}, \alpha)$	(1971NO04)
16.623 ± 3	107.7 ± 0.5	${}^4\text{He}(\alpha, \alpha)$	(1978HI04) ^c
16.630 ± 3	108.5 ± 0.5	${}^4\text{He}(\alpha, \alpha)$	(1978HI04) ^d
16.626 ± 3	108.1 ± 0.5	best value	
16.901 ± 5	77 ± 3	${}^7\text{Li}({}^3\text{He}, \text{d})$	(1967MA12) ^b
	70 ± 5	${}^{10}\text{B}(\text{d}, \alpha)$	(1971NO04)
16.925 ± 3	74.4 ± 0.4	${}^4\text{He}(\alpha, \alpha)$	(1978HI04) ^c
16.918 ± 3	73.6 ± 0.4	${}^4\text{He}(\alpha, \alpha)$	(1978HI04) ^d
16.922 ± 3	74.0 ± 0.4	best value	
17.641 ± 1.5	10.7 ± 0.5	${}^7\text{Li}(\text{p}, \gamma)$	Table 8.6 ^a
18.156 ± 5	147	${}^7\text{Li}(\text{p}, \gamma)$	Table 8.6
18.150 ± 5	138 ± 6	${}^{10}\text{B}(\text{d}, \alpha)$	(1970CA12)
18.144 ± 5		${}^9\text{Be}(\text{d}, \text{t})$	(1977OO01)
18.150 ± 4	138 ± 6	best value	
19.06 ± 20	270 ± 20	${}^7\text{Li}(\text{p}, \gamma)$	Table 8.6
19.071 ± 10	270 ± 30	${}^9\text{Be}(\text{d}, \text{t})$	(1977OO01)
19.069 ± 10	270 ± 20	best value	^h
19.21	208 ± 30	${}^9\text{Be}(\text{p}, \text{d})$	(1967KU10)
19.22 ± 30	265 ± 30	${}^9\text{Be}({}^3\text{He}, \alpha)$	(1976AJ01) ^g
19.26 ± 30	220 ± 30	${}^9\text{Be}(\text{d}, \text{t})$	(1977OO01)
19.24 ± 25 ^e	230 ± 30	best value	
19.86 ± 50	700 ± 100	${}^9\text{Be}(\text{d}, \text{t})$	(1977OO01)

Table 8.5: Some ${}^8\text{Be}$ states with $16.6 < E_x < 23.0$ MeV ^a (continued)

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Reaction	Refs.
22.05 ± 100 ^f	270 ± 70	${}^9\text{Be}({}^3\text{He}, \alpha)$	(1976AJ01)
22.63 ± 100	100 ± 50	${}^9\text{Be}({}^3\text{He}, \alpha)$	(1976AJ01)
22.98 ± 100 ^f	230 ± 50	${}^9\text{Be}({}^3\text{He}, \alpha)$	(1976AJ01)

^a See also Table 8.5 in (1974AJ01) and Tables 8.6, 8.8, 8.9 here.

^b See also (1970CA12, 1971PI06, 1977OO01, 1978HI04).

^c R -matrix theory.

^d Complex eigenvalue theory.

^e See also (1969SU02): 19.16 ± 0.07 MeV.

^f The states at 22.0 ± 0.15 and 22.9 ± 0.15 reported by (1969SU02) in the ${}^9\text{Be}(p, d)$ reaction have $\Gamma_{\text{c.m.}} \geq 1$ MeV.

^g And private communication.

^h See, however, (1978BA66).

$\Gamma < 1$ MeV, $\Gamma_\alpha/\Gamma < 0.5$. No evidence for other 0^+ states is seen above $E_\alpha = 43$ MeV (1972BA83): see also (1974AJ01).

The d -wave phase shift becomes appreciable for $E_\alpha > 2.5$ MeV and passes through resonance at $E_\alpha = 6$ MeV ($E_x = 3.18$ MeV, $\Gamma = 1.5$ MeV, $J^\pi = 2^+$) (1963TO02): see Table 8.4 in (1974AJ01). Five 2^+ levels are observed from $l = 2$ phase shifts measured from $E_\alpha = 30$ to 70 MeV: ${}^8\text{Be}^*(16.6, 16.9)$ with $\Gamma_\alpha = \Gamma$ [see (1978HI04) and Table 8.5], and states with $E_x = 20.2, 22.2$ and 25.2 MeV. The latter has a small Γ_α (1972BA83).

The $l = 4$ shift rises from $E_\alpha \approx 11$ MeV and indicates a broad 4^+ level at $E_x = 11.4 \pm 0.3$ MeV (1959BR71), 11.7 ± 0.4 MeV [$\Gamma = 4.0 \pm 0.4$ MeV] (1974CH45). A rapid rise of δ_4 at $E_\alpha = 40$ MeV corresponds to a 4^+ state at 19.9 MeV with $\Gamma_\alpha/\Gamma \approx 0.96$; $\Gamma < 1$ MeV and therefore $\Gamma_\alpha < 1$ MeV, which is $< 5\%$ of the Wigner limit. A broad 4^+ state is also observed near $E_\alpha = 51.3$ MeV ($E_x = 25.5$ MeV) (1972BA83, 1978LE13).

Over the range $E_\alpha = 30$ to 70 MeV a gradual increase in δ_6 is observed (1972BA83). Some indications of a 6^+ state at $E_x \approx 28$ MeV and of an 8^+ state at ≈ 57 MeV have been reported by (1965DA1A) with $\Gamma_{\text{c.m.}} \approx 20$ and ≈ 73 MeV, respectively. A resonance is not observed at the first $T = 2$ state, ${}^8\text{Be}^*(27.49)$: see Table 8.4 (1976HI04). The elastic scattering has also been studied at $E_\alpha = 160$ MeV (1978RO1G), at 650 and 850 MeV (1976FO03), 850 and 1370 MeV (1977TE1A) and at 4.30 and 5.05 GeV/ c (1977BE1R, 1977BE1T). The total cross section has been measured at 0.87 and 2.1 GeV/ A (1975JA1A).

The bremsstrahlung cross section has been measured for $E_\alpha = 9.35$ to 18.7 MeV: see (1974AJ01). See also (1975GR1C; theor.).

See also (1975IG1A, 1975TO1A, 1976SL02, 1977KI05, 1978BR1E, 1978CH1G, 1978FI1E) and (1973CO1C, 1973FA1C, 1973HA1F, 1973TA1B, 1974AT1A, 1974BA1P, 1974FR1B, 1974FR06, 1974LU05, 1974RA16, 1974TH04, 1974WA02, 1974WA1D, 1975AN1F, 1975BA76, 1975BE1M, 1975BE16, 1975CL1C, 1975DU09, 1975GR05, 1975GR41, 1975GR1C, 1975HE07, 1975HO1E, 1975KU09, 1975MA27, 1975TA1D, 1976AL1E, 1976BA1E, 1976BR32, 1976CO18, 1976CU07, 1976GO12, 1976HA18, 1976HA1L, 1976HO1A, 1976JA1F, 1976LE1J, 1976LI1E, 1976ME1C, 1976MI17, 1976MO31, 1976RA1E, 1977AL1G, 1977AN1B, 1977BA17, 1977BE50, 1977BU01, 1977CA06, 1977CH1C, 1977DA20, 1977FL13, 1977FR12, 1977FR18, 1977FU1E, 1977HO1D, 1977HO1E, 1977HU1A, 1977IK1A, 1977KO1K, 1977LA1D, 1977MA1J, 1977SA1C, 1977ST1G, 1977TH09, 1977ZA04, 1977ZI01, 1978AR1M, 1978FO1G, 1978FR1F, 1978FR1H, 1978LE13, 1978MA20, 1978NO1B, 1978SA24, 1978SA2B, 1978ST10, 1978TO09, 1978VA03, 1978YA1A; theor.).

5. ${}^6\text{Li}(d, \gamma){}^8\text{Be}$ $Q_m = 22.2814$

The yield of γ -rays to ${}^8\text{Be}^*(17.64) [1^+; T = 1]$ has been measured for $E_d = 6.85$ to 7.10 MeV: a resonance is observed at $E_d = 6962.8 \pm 3.0$ keV [$E_x = 27495.8 \pm 2.4$ keV, $\Gamma_{\text{c.m.}} = 5.5 \pm 2.0$ keV]; $\Gamma_\gamma = 23 \pm 4$ eV [1.14 ± 0.20 W.u.] for this M1 transition from the first $0^+; T = 2$ state in ${}^8\text{Be}$, in good agreement with the intermediate coupling model (1976NO07). The γ_1 transition from ${}^8\text{Be}^*(27.49)$ to ${}^8\text{Be}^*(2.9)$ is not observed: assuming $\Gamma_{d_0}/\Gamma = 0.78$, $\Gamma_\gamma^{\Delta T=2}/\Gamma < 2 \times 10^{-6}$. If $\Gamma_{\text{c.m.}} = 5.5$ keV, $\Gamma_\gamma^{\Delta T=2} < 0.01$ eV [$< 1.5 \times 10^{-3}$ W.u.] for this E2, $\Delta T = 2$ transition (1977LO1F).

6. (a) ${}^6\text{Li}(d, n){}^7\text{Be}$ $Q_m = 3.382$ $E_b = 22.2814$
 (b) ${}^6\text{Li}(d, n){}^4\text{He} + {}^3\text{He}$ $Q_m = 1.7955$

The yield curve has been measured for $E_d = 0.06$ to 5.5 MeV and 12 to 17 MeV: see (1974AJ01) and (1977SZ05: 0.10 to 0.18 MeV; activation cross section), (1977EL09: 0.20 to 0.90 MeV; cross sections for n_0 and n_1) and (1974MCZS, 1975MC02: 0.5 to 3.0 MeV; $n_1\gamma$). Polarization measurements have been reported at $E_d = 0.27$ to 0.60 MeV (1966MI06: n_1) and $E_d = 0.6, 0.8$ and 1.0 MeV (1977GL05; n_1) and at $E_d = 2.5$ to 3.7 MeV (1970TH08; n_0, n_1). See also ${}^7\text{Be}$.

Comparisons of the populations of ${}^7\text{Be}^*(0, 0.43)$ and of ${}^7\text{Li}^*(0, 0.48)$ have been made at many energies, up to $E_d = 7.2$ MeV. The n/p ratios are closely equal for analog states, as expected from charge symmetry: see (1974AJ01) and (1977EL09). See also (1973GL1A, 1974AU1B) and (1976MO23, 1977EL11, 1977MC1C; applications).

7. (a) ${}^6\text{Li}(d, p){}^7\text{Li}$ $Q_m = 5.026$ $E_b = 22.2814$
 (b) ${}^6\text{Li}(d, p){}^4\text{He} + {}^3\text{H}$ $Q_m = 2.5592$

Excitation functions have been measured for $E_d = 30$ keV to 5.4 MeV: see (1974AJ01). Recent cross-section measurements are reported at $E_d = 0.1$ to 1.0 MeV (1977EL09; p_0, p_1) and 0.5 to 3.4 MeV (1974MCZS, 1975MC02; $p_0, p_1, p_1\gamma$). There is no resonance at $E_d = 0.4$ MeV: see (1976MO23). An anomaly is observed in the p_1/p_0 intensity ratio at $E_d = 6.945$ MeV, corresponding to the first $0^+; T = 2$ state: $E_x = 27.483 \pm 0.010$ MeV, $\Gamma = 10 \pm 3$ keV, $\Gamma_{p_0} \ll \Gamma_{p_1}, \Gamma_{p_0} < \Gamma_d$ (1969BL14). See also (1975GE1F).

Polarization measurements for p_0 and p_1 have been reported at $E_d = 0.6$ and 0.96 MeV (1977GL05) and $E_d = 2.1$ to 10.9 MeV (1968DU09, 1968FI07, 1970FI07). The latter report pronounced differences in the angular distributions of the vector analyzing power of the two $l_n = 1$ transitions to ${}^7\text{Li}^*(0, 0.48)$. See also ${}^7\text{Li}$, (1974AU1B, 1978VA1D), (1974AJ01, 1975SE07, 1976MO23) and (1977EL11; applications).

8. ${}^6\text{Li}(d, d){}^6\text{Li}$

$$E_b = 22.2814$$

The yield of elastically scattered deuterons has been measured for $E_d = 2$ to 7.14 MeV [see (1974AJ01)]: no resonances are observed, except, possibly, for a very weak anomaly at $E_d = 6.96$ MeV (1975GE1F; abstract). See also ${}^6\text{Li}$ and (1974CH58; theor.).

9. (a) ${}^6\text{Li}(d, t){}^5\text{Li}$

$$Q_m = 0.59$$

$$E_b = 22.2814$$

(b) ${}^6\text{Li}(d, {}^3\text{He}){}^5\text{He}$

$$Q_m = 0.90$$

The cross section for tritium production rises rapidly to 190 mb at 1 MeV, then more slowly to 290 mb near 4 MeV. There is evidence of deviation from isotropy near 0.4 MeV (1955MA20). See also ${}^5\text{Li}$. For reaction (b) see ${}^5\text{He}$.

10. (a) ${}^6\text{Li}(d, \alpha){}^4\text{He}$

$$Q_m = 22.3733$$

$$E_b = 22.2814$$

(b) ${}^6\text{Li}(d, \alpha p){}^3\text{H}$

$$Q_m = 2.5592$$

(c) ${}^6\text{Li}(d, \alpha n){}^3\text{He}$

$$Q_m = 1.7954$$

(d) ${}^6\text{Li}(d, 2d){}^4\text{He}$

$$Q_m = -1.4735$$

Cross sections and angular distributions (reaction (a)) have been measured at $E_d = 0.03$ to 12 MeV [see (1974AJ01)] and at $E_d = 0.118$ to 0.975 MeV (1977EL09: very accurate total reaction cross sections), 0.50 to 3.4 MeV (1974MCZS, 1975MC02; σ_R), 1.0 to 11.5 MeV (1977RI09; $\sigma(\theta)$: see below) and 13.6 MeV (1978FU03; ang. distrib. to ${}^4\text{He}^*(20.1, 21.1, 25.5)$) and $E({}^6\text{Li}) = 10$ to 31 MeV (1979WA02: 30° yield of α_0 and α^* to ${}^4\text{He}^*(20.08) [0^+]$). A critical analysis of the low energy data has led to a calculation of the reaction rate parameter for thermonuclear reactions for plasma temperatures of 2 keV to 1 MeV (1978CL07). Polarization measurements are reported at

$E_d = 0.40$ to 11.8 MeV and at $E(^6\text{Li}) = 0.6$ MeV [see (1974AJ01)] and at $E_{\bar{d}} = 1.5$ to 11.5 MeV (1977RI09: vector and tensor analyzing powers). At $E_{\bar{d}} = 5, 6, 8$ and 9 MeV, $A_{yy} = 1$ is observed at certain angles: see (1978SE01).

Pronounced variations are observed in the cross sections and in the analyzing powers (1977RI09). Maxima are seen at $E_d = 0.8$ MeV, $\Gamma_{\text{lab}} \approx 0.8$ MeV and $E_d = 3.75$ MeV, $\Gamma_{\text{lab}} \approx 1.4$ MeV (1963ME09, 1964PA06). The 4 MeV peak is also observed in the tensor component coefficients with $L = 0, 4$ and 8 and in the vector component coefficients: two overlapping resonances are suggested. At higher energies all coefficients show a fairly smooth behavior which suggests that only broad resonances can exist (1977RI09). The results of this and other experiments are in good agreement with the results of (1972BA83) [$^4\text{He}(\alpha, \alpha)$: reaction 4], that is with two 2^+ states at $E_x = 22.2$ and 25.2 MeV and a 4^+ state at 25.5 MeV (1977RI10). A strong resonance is seen in the α^* channel [to $^4\text{He}^*(20.08)$, $J^\pi = 0^+$] presumably due to $^8\text{Be}^*(25.2, 25.5)$. In addition the ratio of the α^*/α differential cross sections at 30° show a broad peak centered at $E_x \approx 26.5$ MeV (which may be due to interference effects) and suggest a resonance-like anomaly at $E_x \approx 28$ MeV (1979WA02). See also the discussion in (1974AJ01). At $E_d = 6.945$ MeV, the α_0 yield shows an anomaly corresponding to $^8\text{Be}^*(27.49)$, the 0^+ ; $T = 2$ analog of $^8\text{He}_{\text{g.s.}}$ (1969BL14) [(1975GE1F); prelim.] find $E_d = 6962 \pm 7$ keV, $\Gamma \approx 10$ keV.]

For reaction (b) see (1977RO18: $E_d = 0.465$; evidence for QSI) and (1977MI13: $E_d = 7.5, 9, 10, 10.5$ MeV; see $^5\text{Li}, ^7\text{Li}$). For reaction (c) see (1974AJ01). For reaction (d) see (1977MI13, 1978FU03). See also ^6Li .

See also (1975GL08, 1975WI25, 1976UL1C, 1978VA1D), (1975SE07, 1977SE1C, 1977SE09, 1978FI1E), (1976MO23, 1977EL11, 1978MO1J; applications) and (1974LI1E; theor.).

- | | |
|---|-----------------|
| 11. (a) $^6\text{Li}(t, n)^8\text{Be}$ | $Q_m = 16.0240$ |
| (b) $^6\text{Li}(t, n)^4\text{He}^4\text{He}$ | $Q_m = 16.1159$ |

See (1966LA04, 1974AJ01).

- | | |
|---|-----------------|
| 12. (a) $^6\text{Li}(^3\text{He}, p)^8\text{Be}$ | $Q_m = 16.7878$ |
| (b) $^6\text{Li}(^3\text{He}, p)^4\text{He}^4\text{He}$ | $Q_m = 16.8797$ |

Proton groups are observed to $^8\text{Be}^*(0, 2.9, 16.63, 16.92, 17.64)$: see Tables 8.4 and 8.5 in (1974AJ01). The excitation of $^8\text{Be}^*(18.15, 19.0, 19.4, 19.9)$ is also reported by (1971GL07). Angular distributions have been measured at $E(^3\text{He}) = 1.4$ to 17 MeV [see (1974AJ01)] and at $3, 4.5$ and 6 MeV (1976GO1F; p_0, p_1) and 14 MeV (1977IR01; p_0, p_1).

Reaction (b) proceeds via $^8\text{Be}^*(16.63, 16.92)$: $\Gamma = 117 \pm 10$ and 85 ± 10 keV, respectively. Interference effects are reported (1969VI05): see also Table 8.5. See also (1976GO1F, 1977AR09), (1977AS04; theor.) and ^9B .

13. (a) ${}^6\text{Li}(\alpha, d){}^8\text{Be}$ $Q_m = -1.5654$
 (b) ${}^6\text{Li}(\alpha, 2\alpha){}^2\text{H}$ $Q_m = -1.4735$

Deuteron groups have been observed to ${}^8\text{Be}^*(0, 2.9, 11.3 \pm 0.4)$. Angular distributions have been measured at $E_\alpha = 20$ to 48 MeV [see (1974AJ01)] and at $E_\alpha = 15.8$ to 25.1 MeV (1974KO24; d_0, d_1), 17.3, 23.3, 25.1 MeV (1974LE14; d_0, d_1), 20 and 24 MeV (1974GR21; d_0) and 29.4 MeV (1974MA49; d_0, d_1). Large differential cross sections at back angles are analyzed by (1974GR21) using both direct and exchange amplitudes.

A study of reaction (b) shows that the peak due to ${}^8\text{Be}^*(2.9)$ is best fitted by using $\Gamma = 1.2 \pm 0.3$ MeV (1969BA18): see also Table 8.4 in (1974AJ01). At $E_\alpha = 42$ MeV the α - α FSI is dominated by ${}^8\text{Be}^*(0, 2.9)$ (1977BO22). See also (1974MA49, 1976LE1K).

14. (a) ${}^6\text{Li}({}^6\text{Li}, \alpha){}^8\text{Be}$ $Q_m = 20.808$
 (b) ${}^6\text{Li}({}^6\text{Li}, \alpha){}^4\text{He}{}^4\text{He}$ $Q_m = 20.900$

This reaction proceeds via ${}^8\text{Be}^*(0, 2.9, 16.6, 16.9, 22.5)$, and there is indication also that the direct three-body break-up (reaction (b)) is possible (1971GA21, 1972GA32: $E_{\max}({}^6\text{Li}) = 13.0$ MeV). The involvement of a state at $E_x = 19.9$ MeV ($\Gamma = 1.3$ MeV) is suggested by (1966MA40). Good agreement with the shapes of the peaks corresponding to ${}^8\text{Be}^*(16.6, 16.9)$ is obtained by using a simple two-level formula with interference, corrected for the effect of final state Coulomb interaction, assuming $\Gamma(16.6) = 90$ keV and $\Gamma(16.9) = 70$ keV: see also Table 8.5 (1971NO04). The ratio of the intensities of the groups corresponding to ${}^8\text{Be}^*(16.6, 16.9)$ remains constant for $E({}^6\text{Li}) = 4.3$ to 5.5 MeV: $I(16.6)/I(16.9) = 1.22 \pm 0.08$ (1966MA40, 1966KI09). Partial angular distributions for the α_0 group have been measured at fourteen energies for $E({}^6\text{Li}) = 4$ to 24 MeV (1970FR06). See also (1975NO1C; theor.) and (1974AJ01).

15. ${}^7\text{Li}(p, \gamma){}^8\text{Be}$ $Q_m = 17.255$

Cross sections and angular distributions have been reported from $E_p = 30$ keV to 18 MeV. Gamma rays are observed to the ground (γ_0) and to the broad, 2^+ , excited state at 2.9 MeV (γ_1) and to ${}^8\text{Be}^*(16.6, 16.9)$ (γ_3, γ_4). Resonances for both γ_0 and γ_1 occur at $E_p = 0.44$ and 1.03 MeV, and for γ_1 alone at 2, 4.9, 6.0, 7.3, and possibly at 3.1 and 11.1 MeV. In addition broad resonances are reported at $E_p \approx 5$ MeV (γ_0), $\Gamma \approx 4 - 5$ MeV, and at $E_p \approx 7.3$ MeV (γ_1), $\Gamma \approx 8$ MeV: see Table 8.6. The $E_p \approx 5$ MeV resonance ($E_x \approx 22$ MeV) represents the giant dipole resonance based on ${}^8\text{Be}(0)$ while the γ_1 resonance, ≈ 2.2 MeV higher, is based on ${}^8\text{Be}^*(2.9)$. The γ_0 and γ_1 giant resonance peaks each contain about 10% of the dipole sum strength (1976FI1C). The main trend between $E_p = 8$ and 17.5 MeV is a decreasing cross section (1976FI1C). (1977UL02) have studied the analyzing power for $E_{\bar{p}} = 0.38$ to 0.96 MeV: the data indicate some possibility of a 1^- state at 17.70 MeV.

Table 8.6: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, \gamma){}^8\text{Be}$

E_{res} (keV)	Γ_{lab} (keV)	${}^8\text{Be}^*$ (MeV)	l_p	J^π	Res. ^c	Refs.
441.4 ± 0.5 ^a	12.2 ± 0.5	17.641	1	1^+	$\gamma_0, \gamma_1, \gamma_3, \gamma_4$	A
1030 ± 5	168	18.156	1	1^+	$\gamma_0, \gamma_1, \gamma_3, \gamma_4$	A, (1976FI1C) ^d
1890	150 ± 50	18.91		(2^-)	γ_3, γ_4	(1969SW01)
2060 ± 20	310 ± 20	19.06		$J = 1, 2, 3, \pi = (-)$ ^b	γ_1	A, (1976FI1C)
(3100)		(20.0)			γ_1	(1976FI1C)
4900		21.5			γ_1	(1976FI1C)
5000	≈ 4500	21.6	0	$1^-; T = 1$	γ_0	A, (1976FI1C)
6000		22.5			γ_1	(1976FI1C)
7500	≈ 8000	23.8	(0)	$(1^-, 2^-); T = 1$	γ_1	A, (1976FI1C)
(11100)		(27.0)			γ_1	(1976FI1C)
13000	broad	28.6				(1967FE04)

A: see references listed for this resonance in Table 8.6 of (1974AJ01).

^a See (1959AJ76).

^b (1964SC19). See, however, reaction 17.

^c $\gamma_0, \gamma_1, \gamma_3, \gamma_4$ represent transitions to ${}^8\text{Be}^*(0, 2.9, 16.6, 16.9)$, respectively.

^d See also (1970FI1B).

At the $E_p = 0.44$ MeV resonance ($E_x = 17.64$ MeV) the radiation is nearly isotropic consistent with p-wave formation, $J^\pi = 1^+$, with channel spin ratio $\sigma(J_c = 2)/\sigma(J_c = 1) = 3.2 \pm 0.5$ (1961ME10). Radiative widths for the γ_0 and γ_1 decay are displayed in Table 8.7. A careful study of the α -breakup of ${}^8\text{Be}^*(16.63, 16.92)$ [both $J^\pi = 2^+$] for $E_p = 0.44$ to 2.45 MeV shows that the non-resonant part of the cross section for production of ${}^8\text{Be}^*(16.63)$ is accounted for by an extranuclear direct-capture process. Resonances for production of ${}^8\text{Be}^*(16.63, 16.92)$ are observed at $E_p = 0.44, 1.03$ and 1.89 MeV [${}^8\text{Be}^*(17.64, 18.15, 18.9)$]. The results are consistent with the hypothesis of nearly maximal isospin mixing for ${}^8\text{Be}^*(16.63, 16.92)$: decay to these states is not observed from the 3^+ states at $E_x = 19$ MeV, but rather from the 2^- state at 18.9 MeV excitation (1969SW01). (1968PA09) find squared $T = 1$ components of 40% and 60% in ${}^8\text{Be}^*(16.6, 16.9)$ and of 95% and 5% in ${}^8\text{Be}^*(17.6, 18.2)$. See also (1974SO1D, 1978GO1F, 1978ZI1A; applied work), (1974VA1C; astrophys.) and (1977BA62, 1978BA66; theor.).

16. ${}^7\text{Li}(p, n){}^7\text{Be}$

$$Q_m = -1.644$$

$$E_b = 17.255$$

Measurements of cross sections have been reported for $E_p = 1.9$ to 52 MeV: see (1974AJ01) and the review article by (1975LI1E). Recent measurements have been carried out at $E_p = 1.9$ to

Table 8.7: Electromagnetic transitions in $^8\text{Be}^a$

Transition	Γ_γ (eV)	$ M ^2$ (W.u.)	Refs.
17.6 \rightarrow 0	16.7	0.15	see (1974AJ01)
17.6 \rightarrow 2.9	8.15 ± 0.07 (M1) ^b 0.15 ± 0.07 (E2)	0.12	(1961ME10) (1961ME10)
17.6 \rightarrow 16.6	0.032 ± 0.003 ^c	1.48 ± 0.15 (M1)	(1969SW01)
17.6 \rightarrow 16.9	0.0013 ± 0.0003	0.15 ± 0.04 (M1)	(1969SW01)
18.15 \rightarrow 0	3.0		(1976FI1C)
18.15 \rightarrow 2.9	3.8		(1976FI1C)
18.15 \rightarrow 16.6	0.077 ± 0.019	1.04 ± 0.26 (M1)	(1969SW01)
18.15 \rightarrow 16.9	0.062 ± 0.007	1.51 ± 0.17 (M1)	(1969SW01)
18.9 \rightarrow 16.6	0.168	0.053 (E1)	(1969SW01)
18.9 \rightarrow 16.9	0.099	0.045 (E1)	(1969SW01)
19.07 \rightarrow 2.9	10.5		(1976FI1C)

^a See also Table 8.7 in (1974AJ01) and reaction 2 here.

^b (1967CO19) report $\delta(\text{E2/M1}) = 0.21 \pm 0.04$, averaged over the energy of the final state.

^c Nearly pure M1: $\delta(\text{E2/M1}) = -0.014 \pm 0.013$ (1969SW02).

3.8 MeV (1974BU16: n_0 ; $d\sigma/d\Omega(0^\circ)$), 4.2 to 25.9 MeV (1976PO06: n_0, n_1 ; 3.5°), and for neutron production at 0° at 14.3, 19.4 and 29.6 MeV (1975MC18), 29.4, 39.2 and 50.6 MeV (1976RO10), and 24.7 to 44.7 MeV (1977SC37). Polarization measurements have been reported at $E_p = 2.05$ to 5.5 MeV and 30 and 50 MeV: see (1974AJ01). See also (1976DOZW, 1978LE10). For angular distributions see ^7Be .

The yield of ground state neutrons (n_0) rises steeply from threshold and shows pronounced resonances at $E_p = 2.25$ and 4.9 MeV (1963BO06). The yield of n_1 also rises steeply from threshold (1964BU08) and exhibits a broad maximum near $E_p = 3.2$ MeV (1961BE05, 1972PR03) and a broad dip at $E_p \approx 5.5$ MeV, also observed in the p_1 yield (1972PR03).

Multi-channel scattering length approximation analysis of the 2^- partial wave near the n_0 threshold indicates that the 2^- state at $E_x = 18.9$ MeV is virtual relative to the threshold and that its width $\Gamma = 50 \pm 20$ keV (1974AR10). The ratio of the cross section for $^7\text{Li}(p, \gamma)^8\text{Be}^*(18.9) \xrightarrow{\gamma} ^8\text{Be}^*(16.6 + 16.9)$ [obtained by (1969SW01)] to the thermal neutron capture cross section $^7\text{Be}(n, \gamma)^8\text{Be}^*(18.9) \xrightarrow{\gamma} ^8\text{Be}^*(16.6 + 16.9)$ [obtained by (1973BA1J)], provides a rough estimate of the isospin impurity of $^8\text{Be}^*(18.9)$: $\sigma_{p,\gamma}/\sigma_{n,\gamma} \approx 1.5 \times 10^{-5}$ and therefore the $T = 1$ isospin impurity is $< 4\%$ in intensity (1974AR10). See, however, (1977BA62; theor.) who estimates $\approx 10\%$ isospin mixing.

The structure at $E_p = 2.25$ MeV is ascribed to a 3^+ , $T = (1)$, $l = 1$ resonance with $\Gamma_n \approx \Gamma_p$ and $\gamma_n^2/\gamma_p^2 = 3$ to 10: see (1966LA04). At higher energies the broad peak in the n_0 yield at $E_p = 4.9$ MeV can be fitted by $J^\pi = 3^{(+)}$ with $\Gamma = 1.1$ MeV, $\gamma_n^2 \approx \gamma_p^2$ (1963BO06). The behavior of the n_1 cross section can be fitted by assuming a 1^- state at $E_x = 19.5$ MeV and a $J = 0, 1, 2$, positive-parity state at 19.9 MeV [presumably the 20.1–20.2 MeV states reported in reaction 4]. In addition the broad dip at $E_p \approx 5.5$ MeV may be accounted for by the interference of two 2^+ states (1972PR03). See Table 8.8.

The ratio of the cross sections of the (p, n_1) reaction to ${}^7\text{Be}^*(0.43)$ to that for the (p, p_1) reaction to the analog state ${}^7\text{Li}^*(0.48)$ has been measured for $E_p = 2.4$ to 10 MeV and 23 to 52 MeV [see (1974AJ01)] and at $E_p = 3$ to 25.9 MeV (1976PO06): at the lower energies the ratio varies very strongly with energy. [An average value for the ratio over the range 5 to 26 MeV is ≈ 0.3 .] See also (1973AT01, 1974AR05, 1976CY1A), (1974LE1E, 1974LO1B, 1975SO1C, 1976RI1B, 1976SL02), (1976BR42, 1976CAZG, 1977LO10, 1977ME1C; applications) and (1974CO1B, 1976AR1F, 1977ST18, 1978BA66, 1978DE37; theor.).

17. (a) ${}^7\text{Li}(p, p){}^7\text{Li}$

$$E_b = 17.255$$

(b) ${}^7\text{Li}(p, p'){}^7\text{Li}^*$

Absolute differential cross sections for elastic scattering have been reported for $E_p = 0.4$ to 12 MeV and at 14.5, 20.0 and 31.5 MeV. The yields of inelastically scattered protons (to ${}^7\text{Li}^*(0.48)$) and of 0.48 MeV γ -rays have been measured for $E_p = 0.8$ to 12 MeV: see (1974AJ01). (1976SOZW; prelim.) reports total cross-section measurements at $E_p = 25, 30, 35, 40, 45$ and 48 MeV. Polarization measurements have been reported at $E_p = 0.67$ to 10.6 MeV and at 14.5, 49.8, 152 and 155 MeV: see (1974AJ01). The polarization has also been studied by (1974DE45; p_0) for $E_p = 1.8$ to 3.0 MeV. For angular distribution see ${}^7\text{Li}$.

Anomalies in the elastic scattering appear at $E_p = 0.44, 1.03, 1.88, 2.1, 2.5, 4.2$ and 5.6 MeV. Resonances at $E_p = 1.03, 3$ and 5.5 MeV and an anomaly at $E_p = 1.88$ MeV appear in the inelastic channel. A phase-shift analysis and a review of the existing cross-section data by (1973BR13) show that the 0.44 and 1.03 MeV resonances are due to 1^+ states which are a mixture of 5P_1 and 3P_1 with a mixing parameter of $+25^\circ$; that the 2^- state at the neutron threshold ($E_p = 1.88$ MeV) has a width of about 50 keV [see also reaction 16]; and that the $E_p = 2.05$ MeV resonance corresponds to a 3^+ state. The anomalous behavior of the 5P_3 phase around $E_p = 2.2$ MeV appears to result from the coupling of the two 3^+ states [resonances at $E_p = 2.05$ and 2.25 MeV]. The 3S_1 phase begins to turn positive after 2.2 MeV suggesting a 1^- state at $E_p = 2.5$ MeV (1973BR13): see Table 8.9, (1972PR03) and (1977BA62, 1978BA66; theor.). The polarization data of (1974DE45) show structures at $E_p = 1.9$ and 2.3 MeV. See also (1974AR10, 1976MA64).

An attempt has been made to observe the $T = 2$ state [${}^8\text{Be}^*(27.47)$] in the p_0, p_1 and p_2 yields. None of these shows the effect of the $T = 2$ state. Table 8.4 displays the upper limit for Γ_{p_0}/Γ (1976HI04). See (1973GO41, 1978FR12), (1976IK01) and (1975MA1H, 1977ST18, 1978DE37; theor.).

$$18. \text{}^7\text{Li}(p, d)\text{}^6\text{Li} \qquad Q_m = -5.026 \qquad E_b = 17.255$$

The excitation function for d_0 measured for $E_p = 11.64$ to 11.76 MeV does not show any effect from the $T = 2$ state [${}^8\text{Be}^*(27.47)$] (1976HI04). See also ${}^6\text{Li}$.

$$19. \text{(a) } {}^7\text{Li}(p, t)\text{}^5\text{Li} \qquad Q_m = -4.43 \qquad E_b = 17.255$$

$$\text{(b) } {}^7\text{Li}(p, {}^3\text{He})\text{}^5\text{He} \qquad Q_m = -4.12$$

See ${}^5\text{Li}$ and ${}^5\text{He}$.

$$20. \text{}^7\text{Li}(p, \alpha)\text{}^4\text{He} \qquad Q_m = 17.347 \qquad E_b = 17.255$$

The cross section follows the expression $E^{-1}e^{-B/\sqrt{E}}$, with $B = 91.5 \pm 4.5$ keV $^{1/2}$, in the range $E_p = 23$ to 50 keV. The cross section in that interval rises from 0.013 to 2.4 μb (1967FI05). Taking into account ${}^8\text{Be}$ $J^\pi = 2^+$ levels at 16.7 , 16.9 and 20.6 MeV, (1972BA41) has made an R -matrix fit to the revised data of (1971SP05) for $E_p = 131$ to 561 keV and has obtained a quadratic energy dependence for the S -factor: $S = 0.065[1 + 1.82E - 2.51E^2]$ MeV \cdot b, over the energy range $E_p = 0$ to 600 keV.

Excitation functions and angular distributions have been measured at many energies in the range $E_p = 23$ keV to 45.2 MeV: see (1966LA04, 1974AJ01). Recent measurements include those of (1976SAZR: 1.6 to 12.0 MeV) and (1976HI04: 11.64 to 11.76 MeV). Polarization measurements have been carried out for $E_p = 0.8$ to 10.6 MeV: see (1974AJ01). In the range $E_p = 3$ to 10 MeV the asymmetry has one broad peak in the angular distribution at all energies except near 5 MeV; the peak value is 0.98 ± 0.04 at 6 MeV and is essentially 1.0 for $E_p = 8.5$ to 10 MeV (1968PL01, 1969KI04) [see Fig. 12 in (1969KI04) and Fig. 6 (1968PL01) for contour maps of the asymmetry].

Broad resonances are reported to occur at $E_p = 3.0$ MeV [$\Gamma \approx 1$ MeV] and at ≈ 5.7 MeV [≈ 1 MeV]. Structures are also reported at $E_p = 6.8$ MeV and at $E_p = 9.0$ MeV: see (1974AJ01) and (1976SAZR). The 9.0 MeV resonance is also reflected in the behavior of the A_2 coefficient (1968PL01). The experimental data on yields and on polarization have been analyzed by (1971KU10): the data appear to require including two 0^+ states [at $E_x \approx 19.7$ and 21.8 MeV] with very small α -particle widths, and four 2^+ states [at $E_x \approx 15.9$, 20.1 , 22.2 and 25 MeV]. See however reaction 4 and (1972BA83). A 4^+ state near 20 MeV was also introduced in the calculation but its contribution was negligible. The observed discrepancies are said to be probably due to the assumption of pure $T = 0$ for these states (1971KU10). At $E_p = 11.64$ to 11.76 MeV the excitation function does not show any effect due to the $T = 2$ state at $E_x = 27.47$ MeV (1976HI04).

For a study of spallation see (1976KO03) and (1978FI1E). See also (1974VA33), (1978PR1B; applied) and (1974LO1B, 1975CI1A; theor.).

Table 8.8: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, n){}^7\text{Be}$

E_p (MeV)	Γ_{lab} (keV)	${}^8\text{Be}^*$ (MeV)	J^π	Refs.
1.88	55 ± 20	18.90	2^-	(1974AR10)
2.25	220	19.22	3^+	(1957NE22, 1961BE05)
2.6 ^a	≈ 750	19.5	1^-	(1972PR03)
3.0	≈ 1250	19.9	(2^+)	(1972PR03)
4.9	1100	21.5	$3^{(+)}$	(1959GI47, 1963BO06)
5.5	broad	22.1	^b	(1972PR03)

^a $\gamma_{n_1}^2$ and $\gamma_{p_1}^2 \approx 1\%$ of the Wigner limit (1972PR03).

^b The broad dip in the n_1 yield at the same energy as the broad bump in the p_1 yield may be due to interference of two 2^+ states (1972PR03).

 Table 8.9: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, p_0){}^7\text{Li}$ and ${}^7\text{Li}(p, p_1){}^7\text{Li}^*$

E_p (MeV)	Γ_{lab} (keV)	${}^8\text{Be}^*$ (MeV)	J^π	$\Gamma_{p'}$ (keV)	Refs.
0.441	12.2 ^b	17.641	1^+		A, (1973BR13)
1.030 ± 0.005	168	18.156	1^+	≈ 6	A, (1973BR13)
1.88 ^a	55 ± 20	18.90	2^-		(1973BR13, 1974AR10, 1974DE45)
2.05	≈ 400	19.05	3^+ ^g	small	A, (1973BR13)
2.25		19.22	3^+ ^g	small	A, (1973BR13, 1974DE45)
2.5 ^c	≈ 750	19.4	1^-	res	(1972PR03, 1973BR13)
^d					
4.2 ± 0.2	1800 ± 200	20.9 ^e	4^-	(res)	(1965GL03)
5.6	broad	22.2	^f	res	(1965GL03, 1972PR03)

A: See references listed in (1974AJ01).

^a (p, n) threshold: see reaction 16.

^b $\theta_p^2 = 0.064$.

^c See also Table 8.8, $\gamma_{n_1}^2$ and $\gamma_{p_1}^2 \approx 1\%$ of the Wigner limit (1972PR03).

^d A 2^+ state at $E_x \approx 20$ MeV appears to be necessary to account for the cross sections: see Table 8.3 and reaction 4 (1972PR03).

^e Reduced width is 70% of the Wigner limit (1965GL03).

^f May be due to two 2^+ states (1972PR03). See also reaction 16.

^g See also (1978BA66; theor.).

21. (a) ${}^7\text{Li}(d, n){}^8\text{Be}$ $Q_m = 15.031$
 (b) ${}^7\text{Li}(d, n){}^4\text{He}{}^4\text{He}$ $Q_m = 15.123$

The population of ${}^8\text{Be}^*(0, 2.9, 16.6, 16.9, 17.6, 18.2, 18.9, 19.1, 19.2)$ has been reported in reaction (a): for the parameters of ${}^8\text{Be}^*(2.9)$ see Table 8.4 in (1974AJ01). Angular distributions of n_0 and n_1 have been reported at $E_d = 0.7$ to 3.0 MeV [see (1974AJ01)] and at $E_d = 15.25$ MeV (1975AZ02; also n to ${}^8\text{Be}^*(16.6 + 16.9)$; spectroscopic factors). The angular distributions of the neutrons to ${}^8\text{Be}^*(16.6, 17.6, 18.2)$ are fit by $l_p = 1$: see (1974AJ01). See also (1978BA66; theor.).

Reaction (b) appears to proceed primarily via ${}^8\text{Be}^*(2.9, 16.6, 16.9)$ and ${}^5\text{He}_{g.s.}$: see (1974AJ01). However, ${}^8\text{Be}^*(11.4)$ may also be involved [$E_x = 11.4 \pm 0.05$ MeV, $\Gamma_{c.m.} = 2.8 \pm 0.2$ MeV (1969HO11), 3.4 ± 0.1 MeV (1976FO21)] as may state(s) at $E_x \approx 20$ MeV (1978AR10). Attempts to observe n - α rescattering (“proximity scattering”) proceeding via ${}^8\text{Be}^*(16.6, 16.9)$ have been unsuccessful: see (1974AJ01). See, however, (1975GR43). See also (1973HU12, 1974DA28, 1974GR44, 1975DA15, 1977LO10), (1976EA1A, 1976NE1B; applications) and ${}^5\text{He}$ and ${}^9\text{Be}$.

22. ${}^7\text{Li}(t, 2n){}^8\text{Be}$ $Q_m = 8.774$

See (1975JE04).

23. (a) ${}^7\text{Li}({}^3\text{He}, d){}^8\text{Be}$ $Q_m = 11.762$
 (b) ${}^7\text{Li}({}^3\text{He}, pn){}^8\text{Be}$ $Q_m = 9.537$

Deuteron groups are observed to ${}^8\text{Be}^*(0, 2.9, 16.6, 16.9, 17.6)$. For the parameters of ${}^8\text{Be}^*(2.9)$ see Table 8.4 in (1974AJ01). The $J^\pi = 1^+$ mixed isospin states have been studied by (1967MA12): see Table 8.5. Angular distributions have been measured at a number of energies in the range $E({}^3\text{He}) = 0.9$ to 24.3 MeV [see (1974AJ01)] and at 1.2 to 2.5 MeV (1975BO56, 1977BO29). For reaction (b) see (1976DA24). See also (1977SL1B), (1974AL07), (1975KU27, 1975TR02, 1976KU07, 1977BA62; theor.) and ${}^9\text{Be}$.

24. (a) ${}^7\text{Li}(\alpha, t){}^8\text{Be}$ $Q_m = -2.559$
 (b) ${}^7\text{Li}(\alpha, \alpha t){}^4\text{He}$ $Q_m = -2.467$

Angular distributions have been measured up to $E_\alpha = 50$ MeV [see (1966LA04, 1974AJ01)] and at 23.2 and 25.0 MeV (1974DM01; t_0 , and t_1 at higher energy) and 29.4 MeV (1974MA49; t_0 , t_1). The ground state of ${}^8\text{Be}$ does decay isotropically in the cm system: $J^\pi = 0^+$ (1970LA14). At $E_\alpha = 10$ MeV an anomaly (“ghost”) is observed at $E_x \approx 0.5$ MeV: see (1971BE52) and reaction 31. Sequential decay (reaction (b)) is reported at $E_\alpha = 50$ MeV via ${}^8\text{Be}^*(0, 2.9, 11.4, 16.6, 16.9, 19.9)$ (1970LA14). See also (1974MA49). See also (1976LE1K) and (1977BE51; theor.).

25. ${}^7\text{Li}({}^7\text{Li}, {}^6\text{He}){}^8\text{Be}$ $Q_m = 7.281$

At $E({}^7\text{Li}) = 1.4, 1.7$ and 1.8 MeV, the angular distributions of ${}^6\text{He}$ ions leaving ${}^8\text{Be}$ in its ground and 2.9 MeV states are essentially isotropic (1968ST12).

26. (a) ${}^7\text{Be}(n, p){}^7\text{Li}$ $Q_m = 1.6443$ $E_b = 18.900$
 (b) ${}^7\text{Be}(n, \alpha){}^4\text{He}$ $Q_m = 18.992$
 (c) ${}^7\text{Be}(n, \gamma\alpha){}^4\text{He}$ $Q_m = 18.992$

At thermal energies, the (n, p) cross section is $(4.8 \pm 0.9) \times 10^4$ b (1973MU14), the (n, α) cross section is ≤ 0.1 mb and the $(n, \gamma\alpha)$ cross section is 155 mb (1963BA34). These values, and comparison of the (p, n) cross section with that of reaction (a), support the $J^\pi = \frac{3}{2}^-$ assignment for ${}^7\text{Be}(0)$ (1957NE22, 1963BA34). See also (1974AJ01) and (1977BA62; theor.).

27. ${}^7\text{Be}(d, p){}^8\text{Be}$ $Q_m = 16.675$

${}^8\text{Be}^*(0, 2.9)$ is populated in this reaction at $E_d = 0.8$ to 1.7 MeV. See also Table 8.4 in (1974AJ01).

28. ${}^8\text{Li}(\beta^-){}^8\text{Be}$ $Q_m = 16.005$

${}^8\text{Li}$ decays mainly to the broad 2.9 MeV, 2^+ level of ${}^8\text{Be}$, which decays into two α particles. Both the β -spectrum and the resulting α -spectrum have been extensively studied. There appears to be an increasing excess of α -particles with E_α which may reflect transitions into the tail of the $J^\pi = 2^+$ level at $E_x = 16.63$ MeV. See (1955AJ61, 1966LA04) for earlier references. See also ${}^8\text{B}(\beta^+)$.

Studies of the distribution of recoil momenta and neutrino recoil correlation indicate that the decay is overwhelmingly GT, axial vector [see reaction 1 in ${}^8\text{Li}$] and that the ground state of ${}^8\text{Li}$ has $J^\pi = 2^+$: see (1966LA04).

Angular correlations have been measured for the decays of ${}^8\text{Li}$ and ${}^8\text{B}$ as a test of the conserved vector current theory of β -decay. The values of the coefficients are displayed in Table 8.10 (1962NO02, 1963GR11, 1966EI02, 1975TR06). The experimental value of δ [$\delta \equiv B({}^8\text{Li}) - B({}^8\text{B})$] is $(5.4 \pm 0.4) W_\beta$, consistent with CVC theory (1966EI02). There is no evidence for second class currents in this decay: see, e.g., (1975TR06). See also (1973AN11, 1974TR01, 1978FO1H, 1978HA1K, 1978MC1E), (1973BR1C, 1975CA1F, 1976CA1E, 1977GA1E, 1978BO30) and (1973KU1D, 1973TA30, 1974HO1D, 1974KU06, 1975KR14, 1975WI1E, 1976KU1B, 1977KU1E, 1977KU20, 1977OK1A, 1977SA23, 1977TE1B, 1977WI02, 1978KH03; theor.).

Table 8.10: $\alpha - \beta$ angular correlation coefficients in ${}^8\text{Li}$, ${}^8\text{B}$ ^a

Nuclide	A/W_β	B/W_β	W_β (MeV)	δ/W_β ^b	Refs.
${}^8\text{Li}$	$(-8.7 \pm 0.7) \times 10^{-3}$	$(+3.2 \pm 0.6) \times 10^{-3}$	11	$(7.0 \pm 1.2) \times 10^{-3}$	(1962NO02)
${}^8\text{Li}$	$(-8.3 \pm 1.1) \times 10^{-3}$	$(+3.7 \pm 1.0) \times 10^{-3}$	7.5		(1963GR11)
${}^8\text{Li}$	$(-9.7 \pm 0.7) \times 10^{-3}$	$(+3.1 \pm 0.3) \times 10^{-3}$	6.6	$(5.4 \pm 0.4) \times 10^{-3}$	(1966EI02)
${}^8\text{Li}$	$(-8.09 \pm 0.13) \times 10^{-3}$	$(+3.25 \pm 0.58) \times 10^{-3}$	6.9		(1975TR06)
${}^8\text{Li}$	$(-9.34 \pm 0.14) \times 10^{-3}$	$(+3.18 \pm 0.62) \times 10^{-3}$	10.9		(1975TR06)
${}^8\text{Li}$	$(-9.78 \pm 0.26) \times 10^{-3}$	$(+4.29 \pm 1.16) \times 10^{-3}$	12.9		(1975TR06)
${}^8\text{B}$	$(-8.7 \pm 0.9) \times 10^{-3}$	$(-3.9 \pm 1.0) \times 10^{-3}$	11		(1962NO02)
${}^8\text{B}$	$(-11.1 \pm 1.3) \times 10^{-3}$	$(-2.3 \pm 0.3) \times 10^{-3}$	7.0		(1966EI02)
${}^8\text{B}$	$(-7.09 \pm 0.18) \times 10^{-3}$	$(-4.75 \pm 0.55) \times 10^{-3}$	6.9		(1975TR06)
${}^8\text{B}$	$(-8.66 \pm 0.16) \times 10^{-3}$	$(-4.66 \pm 0.54) \times 10^{-3}$	10.9		(1975TR06)
${}^8\text{B}$	$(-9.34 \pm 0.22) \times 10^{-3}$	$(-4.96 \pm 0.82) \times 10^{-3}$	12.9		(1975TR06)

^a $W(\theta) = 1 + A \cos\theta + B \cos^2\theta$.

^b $\delta \equiv B({}^8\text{Li}) - B({}^8\text{B})$.

29. ${}^8\text{B}(\beta^+){}^8\text{Be}$

$$Q_m = 17.980$$

The decay proceeds mainly to ${}^8\text{Be}^*(2.9)$ [see Table 8.4 in (1974AJ01) for its parameters]. Detailed study of the high energy portion of the α -spectrum reveals a maximum near $E_\alpha = 8.3$ MeV, corresponding to transitions to ${}^8\text{Be}^*(16.63)$, for which parameters $E_x = 16.67$ MeV, $\Gamma = 150$ to 190 keV or $E_x = 16.62$ MeV, $\Gamma = 95$ keV are derived. $\log ft = 2.9$ (1964MA35), 3.33 (1969BA43). The $\log ft$ value is consistent with the identification $J^\pi = 2^+$, $T = 1$ for ${}^8\text{Be}^*(16.63)$ (1964MA35) [actually this state is of mixed $T = 0$ and 1 isospin]. The energy distribution of α -particles has also been measured by (1969CL10). Analysis of this data and of data from α - α scattering in a three level R -matrix formalism indicate a 2^+ state of ${}^8\text{Be}$ at $E_x = 12.0_{+3.5}^{-3.0}$ MeV and of $\Gamma = 14_{+4}^{-3}$ MeV ($a_2 = 6.0 \pm 0.5$ fm) (1969CL10).

For angular correlation measurements see reaction 28 and Table 8.10 (1962NO02, 1966EI02, 1975TR06). See also (1974TR01). There is no evidence for second-class currents in this decay: see, e.g., (1975TR06). See also (1973BR1C, 1975CA1F, 1976CA1E, 1977GA1E, 1978BO30) and (1974HO1D, 1974KU06, 1975KR14, 1975WI1E, 1976KU1B, 1976KU07, 1977KU1E, 1977OK1A, 1977TE1B, 1977WI02, 1978KH03; theor.).

30. (a) ${}^9\text{Be}(\gamma, n){}^8\text{Be}$

$$Q_m = -1.6652$$

(b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$

$$Q_m = -1.6652$$

(c) ${}^9\text{Be}(p, pn){}^8\text{Be}$

$$Q_m = -1.6652$$

(d) ${}^9\text{Be}(t, tn){}^8\text{Be}$

$$Q_m = -1.6652$$

$$(e) \text{}^9\text{Be}(\alpha, \alpha n)\text{}^8\text{Be} \quad Q_m = -1.6652$$

Neutron groups to ${}^8\text{Be}^*(0, 2.94)$ have been studied for $E_\gamma = 18$ to 26 MeV (1977BU02). See also (1974AJ01) and ${}^9\text{Be}$. Reaction (b) appears to proceed largely via excited states of ${}^9\text{Be}$ with subsequent decay mainly to ${}^8\text{Be}^*(2.9)$: see (1966LA04, 1974AJ01), ${}^9\text{Be}$ and ${}^{10}\text{Be}$. Reaction (c) has been studied at $E_p = 45$ and 47 MeV: the reaction primarily populates ${}^8\text{Be}^*(0, 2.9)$ (1977WA05). See also (1977JE01, 1978CH07), ${}^9\text{Be}$ and ${}^9\text{B}$. For reactions (d) and (e) see (1974AJ01) and ${}^9\text{Be}$. For reaction (e) see (1974GR42).

$$31. (a) \text{}^9\text{Be}(p, d)\text{}^8\text{Be} \quad Q_m = 0.5594$$

$$(b) \text{}^9\text{Be}(p, d)\text{}^4\text{He}\text{}^4\text{He} \quad Q_m = 0.6513$$

Angular distributions of deuteron groups have been reported at $E_p = 0.11$ to 185 MeV [see (1974AJ01)] and at $E_p = 4.6$ MeV (1974YA1C; d_0), 17.7 MeV (1977GU14; d_0, d_1) and $E_p = 15$ MeV (1976DA15; d_0, d_1). In the latter work S for ${}^8\text{Be}^*(0, 2.9)$ are 0.80 and 1.05 (see also reaction 32) (1976DA15). For other work on spectroscopic factors see (1974AJ01).

Besides ${}^8\text{Be}^*(2.9)$ [see Table 8.4 in (1974AJ01)] ${}^8\text{Be}^*(11.4, 16.6, 16.9, 17.6, 18.2, 18.9, 19.2, (20.0), 21.5, 22.0, 23.0, (24.5), 26.0)$ are populated. See also Table 8.5. (1969SU02) report $E_x = 11.3 \pm 0.3$ MeV for the 4^+ state. At $E_p = 46$ and 100 MeV the angular distributions to ${}^8\text{Be}^*(16.9, 17.6, 18.2, 19.1)$ are consistent with $l = 1$ (1967VE01, 1968LE01, 1971SC26).

An anomaly has been reported in deuteron spectra between the d_0 and d_1 groups. At $E_p = 39.9$ MeV it is seen at $E_x \approx 0.6$ MeV. An R -matrix analysis using a three-level approximation for both the 0^+ and the 2^+ contributions shows that the anomaly is due to a “ghost” of the 0^+ state which produces the ground state group (1976BA67). See (1974AJ01) for the earlier work. Reaction (b) at $E_p = 9$ MeV is dominated by strong FSI through ${}^8\text{Be}^*(0, 2.9)$ and ${}^6\text{Li}^*(2.19)$ with little or no yield from a direct three-body decay (1971EM01). See also (1974AD1B, 1974LO1B), (1978PR1A; applied) (1976BO15, 1978BA66; theor.) and ${}^{10}\text{B}$.

$$32. (a) \text{}^9\text{Be}(d, t)\text{}^8\text{Be} \quad Q_m = 4.5921$$

$$(b) \text{}^9\text{Be}(d, t)\text{}^4\text{He}\text{}^4\text{He} \quad Q_m = 4.6840$$

Angular distributions have been measured for $E_d = 0.3$ to 20 MeV [see (1966LA04, 1974AJ01)] and at 0.15 to 2.5 MeV (1974CH1L; t_0), 0.6 to 2.2 MeV (1974FR02; t_0), 0.9 to 2.5 MeV (1974BO42, 1975BO1C; t_0), 0.9 to 3.1 MeV (1975ZW01; t_0), 12.35 and 14.06 MeV (1978TA04; t_0, t_1), 13.6 MeV (1973ZA06; t_0), 28.0 MeV (1977OO01: see below) and at $E_d = 15$ MeV (1976DA15; t_0, t_1 ; $S = 0.37, 0.55$: see also reaction 31).

At $E_d = 27.97$ MeV angular distributions of triton groups to ${}^8\text{Be}^*(16.6, 16.9, 17.6, 18.2, 19.1, 19.2, 19.8)$ have been analyzed using DWUCK: absolute C^2S are 0.074, 1.56, 0.22, 0.17, 0.41,

0.48, 0.40, respectively. See also Table 8.5. An isospin amplitude impurity of 0.21 ± 0.03 is found for ${}^8\text{Be}^*(17.6, 18.2)$ (1977OO01). See also ${}^9\text{Be}(d, {}^3\text{He})$ [reaction 18] in ${}^8\text{Li}$ and (1978BA66; theor.).

A kinematically complete study of reaction (b) at $E_d = 26.3$ MeV indicates the involvement of ${}^8\text{Be}^*(0, 2.9, 11.4, 16.9, 19.9 + 20.1)$ (1973SO08). See also (1974SO1C, 1975DO04, 1975ZA06, 1976HE1F), (1974AJ01) and ${}^{11}\text{B}$ in (1980AJ01).

33. (a) ${}^9\text{Be}({}^3\text{He}, \alpha){}^8\text{Be}$	$Q_m = 18.9126$
(b) ${}^9\text{Be}({}^3\text{He}, \alpha){}^4\text{He}{}^4\text{He}$	$Q_m = 19.0045$

Angular distributions have been measured in the range $E({}^3\text{He}) = 3.0$ to 26.7 MeV [see (1974AJ01)] and at $E({}^3\text{He}) = 1.75, 2.12, 2.43$ and 2.95 MeV (1974SA1K; α_0), 4, 5 and 7 MeV (1975BI14: to ${}^8\text{Be}^*(17.6, 18.2)$), 4 to 7 MeV (1978BI15; α_0), 9 and 10 MeV (1975RO09; α_0) and at $E({}^3\text{He}) = 33.3$ MeV (1976KA23: to ${}^8\text{Be}^*(16.9, 17.6, 19.2)$) [$S = 1.74, 0.72, 1.17$, assuming mixed isospin for ${}^8\text{Be}^*(16.9)$]. The parameters of observed states are displayed in Table 8.4 of (1974AJ01) and Table 8.5 here. In a recent experiment (1974CA32) suggest $E_x = 2.80 \pm 0.05$ and 11.8 ± 0.5 MeV for the first two excited states of ${}^8\text{Be}$. In addition to three relatively sharp states reported in the range $22 < E_x < 23$ MeV [see Table 8.5] (1976AJ01) suggest the possibility of a broad state at $E_x \approx 25$ MeV. See also (1978BA66; theor.).

Reaction (b) has been studied at $E({}^3\text{He}) = 3.0$ and 4.0 MeV (1966SU04, 1968MO05, 1972TA04, 1975KA05) and at 2.9 to 10 MeV (1975RO09). See also (1976AR11). The reaction proceeds via ${}^8\text{Be}^*(0, 2.9, 11.4, 16.6, 16.9, 19.9, 22.5)$ (1972TA04). See, however, (1975RO09): $\Gamma_{\text{c.m.}}$ for ${}^8\text{Be}^*(2.9)$ is reported to be 0.94 ± 0.04 MeV. $J^\pi = 2^+$ is indicated for ${}^8\text{Be}^*(16.6, 16.9)$ (1966SU04, 1968MO05). See also (1976STYX), (1976RO1L) and ${}^{12}\text{C}$ in (1980AJ01).

34. (a) ${}^9\text{Be}({}^6\text{Li}, {}^7\text{Li}){}^8\text{Be}$	$Q_m = 5.585$
(b) ${}^9\text{Be}({}^7\text{Li}, {}^8\text{Li}){}^8\text{Be}$	$Q_m = 0.368$
(c) ${}^9\text{Be}({}^{12}\text{C}, {}^{13}\text{C}){}^8\text{Be}$	$Q_m = 3.2812$
(d) ${}^9\text{Be}({}^{13}\text{C}, {}^{14}\text{C}){}^8\text{Be}$	$Q_m = 6.5114$
(e) ${}^9\text{Be}({}^{16}\text{O}, {}^{17}\text{O}){}^8\text{Be}$	$Q_m = 2.479$
(f) ${}^9\text{Be}({}^{18}\text{O}, {}^{19}\text{O}){}^8\text{Be}$	$Q_m = 2.292$
(g) ${}^9\text{Be}({}^{19}\text{F}, {}^{20}\text{F}){}^8\text{Be}$	$Q_m = 4.9360$

At $E({}^6\text{Li}) = 5.5$ and 6.5 MeV angular distributions involving ${}^8\text{Be}_{\text{g.s.}}$ and ${}^7\text{Li}^*(0, 0.48)$ have been measured (1974VO06). For reaction (b) see (1974KE1C). For reaction (c) see (1977UH1A, 1978CH02). For reaction (d) see (1975SE03, 1975SE04). Reaction (e) has been studied at $E({}^{18}\text{O}) = 16$ and 20 MeV (1971KN05). See also (1974AJ01) for the earlier work.

35. $^{10}\text{Be}(p, t)^8\text{Be}$ $Q_m = 0.0049$

At $E_p = 34$ MeV tritons are observed to the first $T = 2$ state, $E_x = 27.4922 \pm 0.0026$ MeV, $\Gamma_{\text{c.m.}} = 12 \pm 3$ keV: the angular distributions for the transitions to this state and to $^8\text{Li}^*(10.82)$, reached in the $(p, ^3\text{He})$ reaction, are very similar. They are both consistent with $L = 0$ using a DWBA (LZR) analysis (1975RO01). The particle decay of the $T = 2$ state has been studied by (1975ADZV; abstract).

36. $^{10}\text{B}(\gamma, d)^8\text{Be}$ $Q_m = -6.0259$

See ^{10}B .

37. $^{10}\text{B}(\pi^-, 2n)^8\text{Be}$ $Q_m = 130.534$

Using stopped pions, ^8Be states at ≈ 3 and ≈ 19 MeV are populated. The 19 MeV structure may be due to a superposition of three peaks at 17, 19 and 22 MeV (1977BA51).

38. (a) $^{10}\text{B}(n, t)^8\text{Be}$ $Q_m = 0.2314$
 (b) $^{10}\text{B}(n, t)^4\text{He}^4\text{He}$ $Q_m = 0.3233$

Angular distributions have been measured at $E_n = 14.4$ MeV (1964VA14; t_0, t_1). See also (1974TU1A). For reaction (b) see (1967VA12). See also ^{11}B in (1980AJ01).

39. $^{10}\text{B}(p, ^3\text{He})^8\text{Be}$ $Q_m = -0.5323$

At $E_p = 39.4$ MeV angular distribution measurements have been carried out for the ^3He groups to $^8\text{Be}^*(0, 2.9, 16.6, 16.9)$: the ratio $d\sigma(16.63)/d\sigma(16.91)$ has a mean value of 0.65 ± 0.05 for $\theta = 15^\circ$ to 30° , suggesting possibly a preferential excitation of the $T = 1$ components of these two states. The ratio of the differential cross sections $d\sigma(p, t)$ [to $^8\text{B}_{\text{g.s.}}$] to $d\sigma(p, ^3\text{He})$ [to $^8\text{Be}^*(16.63, 16.92)$] (15° to 30°) seems to also suggest this (1971SQ01).

40. (a) $^{10}\text{B}(d, \alpha)^8\text{Be}$ $Q_m = 17.8208$
 (b) $^{10}\text{B}(d, \alpha)^4\text{He}^4\text{He}$ $Q_m = 17.9127$

Angular distributions have been reported at $E_d = 0.5$ to 7.5 MeV [see (1974AJ01)] and at $E_d = 0.8$ to 9.0 MeV (1978BU04), 1.83 MeV (1974LA29; α_0, α_1) and 5 MeV (1975RO09; α_0). At $E_d = 7.5$ MeV the total cross section for formation of ${}^8\text{Be}^*(16.63)$, $\sigma_t(16.63)$, is about $1.15\sigma_t(16.92)$, consistent with the mixed isospin character of these two states. $\sigma_t(18.15)$ is $\approx 0.85\sigma_t(16.92)$, but the other nearby 1^+ state ${}^8\text{Be}^*(17.64)$ has $\sigma_t(17.64) \approx 0.07\sigma_t(16.92)$, consistent with the nearly pure $T = 1$ nature of ${}^8\text{Be}^*(17.64)$ (1966BR08). These four states [${}^8\text{Be}^*(16.63, 16.92, 17.64, 18.15)$] have been studied for $E_d = 4.0$ to 12.0 MeV. Interference between the 2^+ states [${}^8\text{Be}^*(16.63, 16.92)$] varies as a function of energy. The cross-section ratios for formation of ${}^8\text{Be}^*(17.64, 18.15)$ vary in a way consistent with a change in the population of the $T = 1$ part of the wave function over the energy range: at the higher energies, there is little isospin violation. At higher E_x only the 3^+ state at $E_x = 19.2$ MeV is observed, the neighboring 3^+ state at $E_x = 19.07$ MeV is not seen (1970CA12). There is some question as to whether a two-level fit can be made for the α groups to ${}^8\text{Be}^*(16.63, 16.92)$. ((1970CA12) and W.D. Callender, private communication) are dubious about this, feeling that other 2^+ states have to be brought into the calculation. Based on a two-level fit they find the following average values: $\Gamma_{16.6} = 113$ keV, $\Gamma_{16.9} = 80$ keV, $\Delta Q = 302$ keV. However, (1971NO04) state that the two-level fit is appropriate of the spectra are properly corrected for effects of final state Coulomb interactions: $\Gamma_{16.6} = 90 \pm 5$ keV, $\Gamma_{16.9} = 70 \pm 5$ keV, $\Delta Q = 290 \pm 7$ keV: see (1978HI04) and Table 8.5. See also Table 8.4 in (1974AJ01) and (1977BA62, 1978BA66; theor.).

Reaction (b) [$E_d < 5$ MeV] takes place mainly by a sequential process involving ${}^8\text{Be}^*(0, 2.9, 11.4, 16.6, 16.9)$: see (1975RO09, 1975VA04) and the earlier work in (1974AJ01). See also (1970HU1B, 1976GR22, 1977LA13, 1977NO10), (1974AD1B), and ${}^{12}\text{C}$ in (1980AJ01).

$$41. {}^{10}\text{B}({}^3\text{He}, p\alpha){}^8\text{Be} \quad Q_m = 12.3273$$

At $E({}^3\text{He}) = 2.45$ and 6.00 MeV this reaction proceeds primarily by sequential decay via ${}^8\text{Be}^*(0, 2.9)$ and via ${}^5\text{Li}$, ${}^9\text{B}$ and ${}^{12}\text{C}$ states (1966WA16).

$$42. {}^{10}\text{B}(\alpha, {}^6\text{Li}){}^8\text{Be} \quad Q_m = -4.552$$

Angular distributions have been obtained at $E_\alpha = 46$ MeV for the transitions to ${}^8\text{Be}^*(0, 2.9)$ (1970ZE03). See also ${}^6\text{Li}$ (1974WO1C, 1976WO11).

$$43. {}^{10}\text{B}({}^7\text{Li}, {}^9\text{Be}){}^8\text{Be} \quad Q_m = 10.670$$

Angular distributions have been obtained at $E({}^7\text{Li}) = 24$ MeV to ${}^8\text{Be}^*(0, 2.9, 11.4)$ (1977KO27).

44. (a) $^{11}\text{B}(\gamma, t)^8\text{Be}$ $Q_m = -11.2238$
 (b) $^{11}\text{B}(\gamma, t)^4\text{He}^4\text{He}$ $Q_m = -11.1319$

See ^{11}B in (1980AJ01).

45. (a) $^{11}\text{B}(p, \alpha)^8\text{Be}$ $Q_m = 8.5902$
 (b) $^{11}\text{B}(p, \alpha)^4\text{He}^4\text{He}$ $Q_m = 8.6821$

Angular distributions have been measured at $E_p = 0.78$ to 45 MeV [see (1974AJ01)] and at $E_p = 5.08$ to 38 MeV (1975BU1B, 1978BU1D; α_0) and 7.3 MeV (1974KA1J; α_0, α_1). Observed parameters for $^8\text{Be}^*(2.9)$ are shown in Table 8.4 of (1974AJ01). (1977FU09) find that the width of this state varies with E_p . $^8\text{Be}_{\text{g.s.}}$ decays isotropically, as expected from its 0^+ character (1976KO18).

Reaction (b) has been studied for $E_p = 0.15$ to 10.5 MeV. The reaction proceeds predominantly by sequential two-body decay via $^8\text{Be}^*(0, 2.9)$: see (1974AJ01), ^{12}C in (1975AJ02, 1980AJ01) and (1974KA1J, 1975VA04, 1976GR22, 1977LA13, 1977OH1A). See also (1972TH1C, 1977GR10) and (1978BI1E; applied).

46. $^{11}\text{B}(d, n\alpha)^8\text{Be}$ $Q_m = 6.3656$

See ^9Be and (1978GR07).

47. $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$ $Q_m = 4.5702$

At $E(^3\text{He}) = 25.6$ MeV angular distributions have been obtained for the ^6Li ions to $^8\text{Be}^*(0, 16.6, 16.9, 17.6)$. In the range $E(^3\text{He}) = 25.2$ to 26.3 MeV, the group to $^8\text{Be}^*(18.2)$ [$J^\pi = 1^+$; $T = 0$] is not observed: its intensity is < 0.15 of the intensity to $^8\text{Be}^*(17.6)$ [$J^\pi = 1^+$; $T = 1$] (1974DE25). For the earlier work see (1974AJ01), (1972YO02) and ^6Li .

48. $^{11}\text{B}(\alpha, ^7\text{Li})^8\text{Be}$ $Q_m = -8.757$

Angular distributions have been reported at $E_\alpha = 28.4$ and 29.0 MeV for $^8\text{Be}_{\text{g.s.}} + ^7\text{Li}_{\text{g.s.}}$, $^8\text{Be}_{\text{g.s.}} + ^7\text{Li}_{0.48}^*$ and $^8\text{Be}_{2.9}^* + ^7\text{Li}$ (29 MeV only) by (1968KA24) and at 42 MeV for $^8\text{Be}_{\text{g.s.}} + ^7\text{Li}_{\text{g.s.}}$ and $^8\text{Be}_{\text{g.s.}} + ^7\text{Li}_{0.48}^*$ by (1968MI05). At $E_\alpha = 65$ MeV $^8\text{Be}^*(16.6 + 16.9, 20.0)$ are apparently also excited (1973WO06). See also (1974WO1C, 1974WO1D), (1973SC1B) and ^7Li .

49. (a) $^{12}\text{C}(\gamma, \alpha)^8\text{Be}$ $Q_m = -7.3667$
 (b) $^{12}\text{C}(e, e\alpha)^8\text{Be}$ $Q_m = -7.3667$

See ^{12}C in (1975AJ02, 1980AJ01).

50. $^{12}\text{C}(n, n\alpha)^8\text{Be}$ $Q_m = -7.3667$

This reaction proceeds via $^8\text{Be}^*(0, 2.9)$ at $E_n = 13$ to 18 MeV and via states in ^5He , ^9Be and ^{12}C (1966MO05, 1975AN01). See also (1975AN1G, 1976AN16) and ^{12}C in (1975AJ02, 1980AJ01).

51. (a) $^{12}\text{C}(p, p\alpha)^8\text{Be}$ $Q_m = -7.3667$
 (b) $^{12}\text{C}(p, d^3\text{He})^8\text{Be}$ $Q_m = -25.720$

Reaction (a) has been studied for $13 < E_p < 160$ MeV. At low energies it involves $^8\text{Be}_{g.s.}$; at higher energies $^8\text{Be}^*(0, 2.9)$: see (1974AJ01) and (1969EP01, 1977IO1A, 1977RO02). See also ^{12}C in (1975AJ02, 1980AJ01), (1978CH1G, 1978DE1J) and (1977CH02; theor.). For reaction (b) see (1977CO07; 100 MeV; $^8\text{Be}(0)$) [$S_\alpha = 0.56 \pm 0.12$].

52. $^{12}\text{C}(d, ^6\text{Li})^8\text{Be}$ $Q_m = -5.893$

Angular distributions have been obtained at $E_d = 19.5, 28$ and 51.8 MeV [see (1974AJ01)] and at $E_d = 12.7$ and 13.2 MeV (1975GO36; $^8\text{Be}_{g.s.}$), 13.6 and 14.6 MeV (1974GA30; $^8\text{Be}_{g.s.}$), at $E_d = 16$ MeV (1976JA1G; $^8\text{Be}_{g.s.}$), at $E_d = 35$ MeV (1975BE01; $^8\text{Be}_{g.s.}$) and at 54.3 MeV (1977TA1D: to $^8\text{Be}^*(0, 2.9, 11.4, 16.6 + 16.9)$). See also (1973FO1A) and (1974DO03, 1976NA05, 1978TA1F; theor.).

53. $^{12}\text{C}(^3\text{He}, ^7\text{Be})^8\text{Be}$ $Q_m = -5.781$

Angular distributions have been measured involving $^8\text{Be}_{g.s.} + ^7\text{Be}_{g.s.}$ and $^8\text{Be}_{g.s.} + ^7\text{Be}^*(0.43)$ at $E(^3\text{He}) = 25.5, 27.0$ and 29.0 MeV (1976PI10) and at 28 MeV (1970DE12). In addition at $E(^3\text{He}) = 70$ MeV the distributions involving $^8\text{Be}^*(2.9, 11.4, 16.6)$ and $^7\text{Be}_{0+1}$, $^8\text{Be}^*(16.9)$ and $^7\text{Be}^*(0.43)$, and $^8\text{Be}^*(17.6)$ and $^7\text{Be}^*(0, 0.43)$ have been studied by (1976ST11). See also (1975AU01), (1974DO03; theor.) and ^7Be .

54. (a) $^{12}\text{C}(\alpha, 2\alpha)^8\text{Be}$ $Q_m = -7.3667$
 (b) $^{12}\text{C}(\alpha, ^8\text{Be})^8\text{Be}$ $Q_m = -7.4586$

This reaction has been studied up to $E_\alpha = 104$ MeV: see (1974AJ01). Angular correlations involving $^8\text{Be}^*(0, 2.9)$ have been studied at $E_\alpha = 90$ MeV: $S_\alpha(\text{PWIA}) = 2.9 \pm 0.4$ and 2.8 ± 0.3 , respectively; $S_\alpha(\text{DWIA})$ for $^8\text{Be}_{\text{g.s.}} = 2.4 \pm 0.4$ (1976SH02).

Angular distributions at $E_\alpha = 65$ MeV (reaction (b)) lead to $S_\alpha = 0.55$ and 0.75 (DWBA) for $^8\text{Be}^*(0, 2.9)$. $^8\text{Be}^*(11.4)$ was also observed (1974WO1C, 1976WO11). See also (1977KA1J).

See also ^{12}C in (1975AJ02, 1980AJ01), ^{16}O in (1977AJ02), (1973RU09, 1974RU06, 1975CO1C, 1977MC07, 1977YA1A) and (1976ME20; theor.).

55. (a) $^{12}\text{C}(^{11}\text{B}, ^{15}\text{N})^8\text{Be}$ $Q_m = 3.6246$
 (b) $^{12}\text{C}(^{12}\text{C}, ^{16}\text{O})^8\text{Be}$ $Q_m = -0.2047$

At $E(^{12}\text{C}) \approx 37$ MeV angular distributions (reaction (b)) are reported involving $^8\text{Be}_{\text{g.s.}} + ^{16}\text{O}_{\text{g.s.}}$ (1975EB03, 1976EB01). See also ^{16}O in (1977AJ02), ^{24}Mg in (1978EN06), (1974AJ01) and (1974HO30, 1976AR02, 1977WA04). For reaction (a) see (1978FR20) and ^{15}N in (1981AJ01).

56. $^{12}\text{C}(^{16}\text{O}, ^{20}\text{Ne})^8\text{Be}$ $Q_m = -2.6358$

See (1974ME09) and ^{20}Ne in (1978AJ03).

57. $^{13}\text{C}(\text{p}, ^6\text{Li})^8\text{Be}$ $Q_m = -8.615$

Angular distributions have been measured at $E_p = 45$ MeV for the transitions to $^8\text{Be}^*(0, 2.9)$ (1971BR07). See also ^6Li .

58. $^{13}\text{C}(\text{d}, ^7\text{Li})^8\text{Be}$ $Q_m = -3.589$

See ^7Li .

59. $^{13}\text{C}(^3\text{He}, ^8\text{Be})^8\text{Be}$ $Q_m = 8.1728$

Angular distributions have been obtained at $E(^3\text{He}) = 3.3, 5.0$ and 5.8 MeV for the transition to $^8\text{Be}_{\text{g.s.}} + ^8\text{Be}_{\text{g.s.}}$ (1968JA07). See also ^{12}C in (1975AJ02), ^{16}O in (1977AJ02), (1974BA42) and (1973OG1A).

60. (a) $^{14}\text{N}(\text{n}, ^7\text{Li})^4\text{He}^4\text{He}$ $Q_{\text{m}} = -8.823$
(b) $^{14}\text{N}(\text{n}, \text{t})^4\text{He}^4\text{He}^4\text{He}$ $Q_{\text{m}} = -11.2899$

See (1976TU04, 1977TU1D). See also ^7Li .

61. $^{14}\text{N}(\text{d}, 2\alpha)^8\text{Be}$ $Q_{\text{m}} = 6.2076$

See (1972FA07).

62. $^{14}\text{N}(\alpha, ^{10}\text{B})^8\text{Be}$ $Q_{\text{m}} = -11.7051$

See (1974WO1C, 1976WO11).

63. $^{14}\text{N}(^{10}\text{B}, 6\alpha)$ $Q_{\text{m}} = 0.3655$

See (1974AJ01).

64. $^{15}\text{N}(\alpha, ^{11}\text{B})^8\text{Be}$ $Q_{\text{m}} = -11.0832$

See (1974WO1C, 1974WO1D, 1976WO11).

65. $^{16}\text{O}(\text{p}, \text{p})^4\text{He}^4\text{He}^4\text{He}$ $Q_{\text{m}} = -14.4368$

See ^{16}O in (1977AJ02).

66. $^{16}\text{O}(\alpha, ^{12}\text{C})^8\text{Be}$ $Q_{\text{m}} = -7.2538$

See (1974WO1C, 1974WO1D, 1976WO11), (1973SC1B) and ^{12}C in (1980AJ01).

$$67. \text{}^{16}\text{O}({}^6\text{Li}, 2\alpha){}^{14}\text{N} \quad Q_{\text{m}} = 1.6370$$

See (1975MI1A).

$$68. \text{}^{16}\text{O}({}^{16}\text{O}, {}^{24}\text{Mg}){}^8\text{Be} \quad Q_{\text{m}} = -0.4852$$

See (1976AR02), ^{20}Ne in (1978AJ03) and ^{24}Mg in (1978EN06).

$$69. \text{}^{19}\text{F}(\text{p}, \text{}^{12}\text{C}){}^8\text{Be} \quad Q_{\text{m}} = 0.8599$$

See ^{20}Ne in (1978AJ03).

⁸B
(Figs. 13 and 14)

GENERAL: (See also (1974AJ01).)

Nuclear models: (1975KH1A).

Special states: (1974IR04, 1976IR1B, 1978KH03).

Electromagnetic interactions: (1974KU06, 1976KU07).

Special reactions: (1974BA70, 1976BE1K, 1976WE09, 1977WE1B, 1977YA1B).

Pion and kaon reactions: (1973CA1C, 1977JU1B).

Astrophysical questions: (1973BA1H, 1973DE1D, 1973TR1C, 1974DA18, 1976BA1J, 1977BA1V, 1977SC1D).

Other topics: (1974IR04, 1975KH1A, 1976IR1B, 1977TR07, 1978RO01).

Ground state properties: (1974SHYR, 1975BE31, 1976FU06).

$$\mu = 1.0355 \pm 0.003 \text{ nm (1973MIYZ)}.$$

1. ${}^8\text{B}(\beta^+){}^8\text{Be}$ $Q_m = 17.980$

The β^+ decay leads mainly to ${}^8\text{Be}^*(2.9)$. The mean of half-lives listed in (1974AJ01) is 770 ± 3 msec; $\log ft = 5.64$ (1966BA1A). There is also a branch to a ${}^8\text{Be}$ state at ≈ 16.6 MeV; $\log ft = 2.9$ (1964MA35), 3.33 (1969BA43). See reaction 29 in ${}^8\text{Be}$. See (1973TR1B, 1974BA1Q, 1974CL1E, 1974UL1B, 1974VA1C, 1975UL1A, 1977HA1M) for astrophysical considerations.

2. ${}^6\text{Li}({}^3\text{He}, n){}^8\text{B}$ $Q_m = -1.975$

Angular distributions for the n_0 group have been reported at $E({}^3\text{He}) = 4.8$ to 5.7 MeV. The appearance of a forward peak indicates an $L = 0$ transfer and hence a knockout mechanism ($L = 0$ is forbidden for simple diproton stripping) (1967VA24). The first excited state is at 767 ± 12 (1965FA03), 783 ± 10 keV (1965MC06: $\Gamma = 40 \pm 10$ keV). See also (1966LA04) and ${}^9\text{B}$.

Table 8.11: Energy levels of ^8B

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$2^+; 1$	$\tau_{1/2} = 770 \pm 3$ msec	β^+	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11
0.778 ± 7		$\Gamma = 40 \pm 10$	γ, p	2, 6, 8
2.32 ± 30	$3^+; 1$	350 ± 40		6, 8
10.619 ± 9	$0^+; 2$	< 60		8

3. $^7\text{Be}(p, \gamma)^8\text{B}$ $Q_m = 0.137$

Absolute cross sections have been measured for $E_p = 0.165$ to 10.0 MeV [see (1974AJ01)] and at $E_p = 0.36$ MeV (1977WI05). A resonance at $E_p = 724$ keV [$^9\text{B}^*(0.77)$] with $\Gamma_{\text{lab}} \approx 46$ keV [$\sigma_{\text{peak}} = 2.20 \pm 0.22$ μb , $\Gamma_\gamma = 50 \pm 25$ meV] is reported by (1966PA16, 1972KA1B).

The zero-energy cross-section factor $S(0) = 0.032 \pm 0.003$ keV \cdot b, $dS/dE = -3 \times 10^{-5}$ b ((1972KA1B) and P.D. Parker, private communication). The change in the S -value is due to a remeasurement of the $^7\text{Li}(d, p)$ cross section: see ^9Be . See also (1974AJ01, 1977WI05). The relevance of this reaction to astrophysics is discussed by (1971BA1A, 1973BA1H, 1974VA1C, 1974HO11). See also (1974AJ01, 1974LO1B).

4. $^9\text{Be}(p, 2n)^8\text{B}$ $Q_m = -20.428$

See (1974DA18) and ^{10}B .

5. $^{10}\text{B}(\pi^+, d)^8\text{B}$ $Q_m = 115.563$

The ground state transition has been studied at $E_{\pi^+} = 50$ MeV (1977AM1B).

6. $^{10}\text{B}(p, t)^8\text{B}$ $Q_m = -18.531$

At $E_p = 49.5$ MeV angular distributions have been measured for the tritons to $^8\text{B}^*(0, 2.32)$ (1970SQ01): $L = 2$ and $L = 0 + 2$, leading to $J^\pi = 2^+$ and 3^+ , respectively. See also $^{10}\text{B}(p, ^3\text{He})^8\text{B}$. The energy of the excited state is 2.29 ± 0.05 MeV (1970SQ01), 2.34 ± 0.04 MeV (1968BR23): $\Gamma_{\text{lab}} = 390 \pm 40$ keV (1967MC14). $^8\text{B}^*(0.78)$ is also observed.

7. $^{10}\text{B}(^6\text{Li}, ^8\text{Li})^8\text{B}$ $Q_m = -17.730$

See (1976WE09, 1977WE1B) and ^8Li .

8. $^{11}\text{B}(^3\text{He}, ^6\text{He})^8\text{B}$ $Q_m = -16.917$

At $E(^3\text{He}) = 50$ MeV, ^6He groups are observed to the first three states of ^8B (1967MC14). At $E(^3\text{He}) = 72$ MeV the first $T = 2$ state is observed at $E_x = 10.619 \pm 0.009$ MeV, $\Gamma < 60$ keV: $d\sigma/d\Omega(\text{lab}) = 190$ nb/sr at $\theta_{\text{lab}} = 9^\circ$. No other states are observed within 2.4 MeV of this state (1975RO01).

9. $^{12}\text{C}(^3\text{He}, ^7\text{Li})^8\text{B}$ $Q_m = -22.899$

This reaction has been studied at $E(^3\text{He}) = 40.7$ MeV (1971DE37). See also (1975PA11).

10. $^{12}\text{C}(\alpha, ^8\text{Li})^8\text{B}$ $Q_m = -41.444$

See (1968MC02).

11. $^{13}\text{C}(^3\text{He}, ^8\text{Li})^8\text{B}$ $Q_m = -25.812$

See (1976BE1L).

^8C (Fig. 14)

Mass of ^8C : The atomic mass excess of ^8C is 35096 ± 26 keV, $\Gamma_{\text{c.m.}} = 230 \pm 50$ keV: see (1977TR07). See also (1974AJ01, 1974RO17, 1976TR01, 1978RO01). ^8C is stable with respect to $^7\text{B} + \text{p}$ ($Q = -0.13$ MeV) and unstable with respect to $^6\text{Be} + 2\text{p}$ ($Q = 2.143$), $^5\text{Li} + 3\text{p}$ ($Q = 1.55$), $^4\text{He} + 4\text{p}$ ($Q = 3.514$). At $E(^3\text{He}) = 76$ MeV the differential cross section for formation of $^8\text{C}_{\text{g.s.}}$ in the $^{14}\text{N}(^3\text{He}, ^9\text{Li})$ reaction is ≈ 5 nb/sr at $\theta_{\text{lab}} = 10^\circ$ (1976RO04). The $^{12}\text{C}(\alpha, ^8\text{He})^8\text{C}$ reaction has been studied at $E_\alpha = 156$ MeV: $d\sigma/d\Omega \approx 20$ nb/sr at $\theta_{\text{lab}} = 20^\circ$ (1974RO17). See also (1976RO1M) and (1974IR04, 1975BE31, 1976IR1B; theor.).

References

(Closed 1978)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1955AJ61 F. Ajzenberg and T. Lauritsen, *Rev. Mod. Phys.* 27 (1955) 77
- 1955FR10 J.M. Freeman, A.M. Lanem and B. Rose, *Phil. Mag.* 46 (1955) 17
- 1955MA20 R.L. Macklin and H.E. Banta, *Phys. Rev.* 97 (1955) 753
- 1957NE22 H.W. Newson, R.M. Williamson, K.W. Jones, J.H. Gibbons and H. Marshak, *Phys. Rev.* 108 (1957) 1294
- 1959AJ76 F. Ajzenberg and T. Lauritsen, *Nucl. Phys.* 11 (1959) 1
- 1959BR71 D.J. Bredin, W.E. Burcham, D. Evans, W.M. Gibson, M.S.C. McKee, D.J. Prowse, J. Rotblat and J.N. Snyder, *Proc. Roy. Soc.* 251 (1959) 143
- 1959GI47 J.H. Gibbons and R.L. Macklin, *Phys. Rev.* 114 (1959) 571
- 1959IM04 W.L. Imhof, R.G. Johnson, F.J. Vaughn and M. Walt, *Phys. Rev.* 114 (1959) 1037
- 1960HU08 D.J. Hughes, B.A. Magurno and M.K. Brussel, *BNL 325, 2nd Edition, Suppl. Vol. 1* (1960)
- 1961BE05 P.R. Bevington, W.W. Rolland and H.W. Lewis, *Phys. Rev.* 121 (1961) 871
- 1961ME10 V. Meyer, H. Muller, H.H. Staub and R. Zurmuhle, *Nucl. Phys.* 27 (1961) 284
- 1962CH14 L.F. Chase, Jr., R.G. Johnson, F.J. Vaughn and E.K. Warburton, *Phys. Rev.* 127 (1962) 859
- 1962NO02 M.E. Nordberg, Jr., F.B. Morinigo and C.A. Barnes, *Phys. Rev.* 125 (1962) 321
- 1963BA34 P. Bassi, B. Ferretti, G. Venturini, G.C. Bertolini, F. Cappellani, V. Mandl, G.B. Restelli and A. Rota, *Nuovo Cim.* 28 (1963) 1049
- 1963BO06 R.R. Borchers and C.H. Poppe, *Phys. Rev.* 129 (1963) 2679
- 1963GR11 W. Gruhle, K.H. Lauterjung, B. Schimmer, U. Schimmer and U. Schmidt-Rohr, *Nucl. Phys.* 42 (1963) 321
- 1963ME09 V. Meyer, W. Pfeifer and H.H. Staub, *Helv. Phys. Acta* 36 (1963) 465
- 1963NE07 B.M.K. Nefkens, *Phys. Rev. Lett.* 10 (1963) 243
- 1963TO02 T.A. Tombrello and L.S. Senhouse, *Phys. Rev.* 129 (1963) 2252
- 1964BU08 S.G. Buccino, C.E. Hollandsworth and P.R. Bevington, *Nucl. Phys.* 53 (1964) 375

1964LA19 R.O. Lane, A.J. Elwyn and A. Langsdorf, Jr., Phys. Rev. 136 (1964) B1710
1964MA35 E. Matt, H. Pfander, H. Rieseberg and V. Soergel, Phys. Lett. 9 (1964) 174
1964PA06 P. Paul and K.P. Lieb, Nucl. Phys. 53 (1964) 465
1964SC19 D.J. Schlueter, R.W. Krone and F.W. Prosser, Jr., Nucl. Phys. 58 (1964) 254
1964VA14 V. Valkovic, Nucl. Phys. 54 (1964) 465
1965DA1A Darriulat, Igo, Pugh and Holmgren, Phys. Rev. 137 (1965) B315
1965FA03 U. Fasoli, D. Toniolo and G. Zago, Nuovo Cim. 37 (1965) 349
1965GL03 R. Gleyvod, N.P. Heydenburg and I.M. Naqib, Nucl. Phys. 63 (1965) 650
1965GR25 W. Gruhle, K.H. Lauterjung, B. Schimmer and U. Schmidt-Rohr, Nucl. Phys. 74 (1965) 129
1965MC06 W.R. McMurray, P. Van Der Merwe and I.J. Van Heerden, Phys. Lett. 18 (1965) 319
1965PO06 A.M. Poskanzer, R.A. Esterlund and R. McPherson, Phys. Rev. Lett. 15 (1966) 1030
1965WA12 J.W. Watson, F. Ajzenberg-Selove and R. Middleton, Phys. Lett. 18 (1965) 302
1966BA1A Bahcall, Nucl. Phys. 75 (1966) 10
1966BR08 C.P. Browne and J.R. Erskine, Phys. Rev. 143 (1966) 683
1966CE01 J. Cerny, S.W. Cosper, G.W. Butler, R.H. Pehl, F.S. Goulding, D.A. Landis and C. Detraz, Phys. Rev. Lett. 16 (1966) 469
1966EI02 H. Eichner, K.H. Lauterjung, H. Meinhardt, B. Schimmer and U. Schmidt-Rohr, Z. Naturforsch. A21 (1966) 908
1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932
1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
1966MA40 M.D. Mancusi and E. Norbeck, Phys. Rev. 151 (1966) 830
1966MI06 G. Michel, R.E. Benenson, H. Burgisser, P. Huber, W.A. Schier and H.R. Striebel, Helv. Phys. Acta 39 (1966) 267
1966MO05 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. 75 (1966) 113
1966PA16 P.D. Parker, Phys. Rev. 150 (1966) 851
1966SU04 D.J. Sullivan and P.B. Treacy, Nucl. Phys. 78 (1966) 225
1966WA16 M.A. Waggoner, J.E. Etter, H.D. Holmgren and C. Moazed, Nucl. Phys. 88 (1966) 81
1967CO19 P.M. Cockburn and I.S. Grant, Phys. Lett. B25 (1967) 111
1967FE04 L. Feldman, B.B. Baliga and M. Nessin, Phys. Rev. 157 (1967) 921
1967FI05 O. Fiedler and P. Kunze, Nucl. Phys. A96 (1967) 513
1967KU10 L.A. Kull, Phys. Rev. 163 (1967) 1066

- 1967MA12 J.B. Marion, P.H. Nettles, C.L. Cocke and G.J. Stephenson, Jr., Phys. Rev. 157 (1967) 847
- 1967MC14 R.L. McGrath, J. Cerny and E. Norbeck, Phys. Rev. Lett. 19 (1967) 1442
- 1967RO06 J.C. Roynette, M. Arditì, J.C. Jacmart, F. Mazloum, M. Riou and C. Ruhla, Nucl. Phys. A95 (1967) 545
- 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
- 1967VA24 P. van der Merwe, W.R. McMurray and I.J. van Heerden, Nucl. Phys. A103 (1967) 474
- 1967VE01 J.W. Verba, H. Willmes, R.F. Carlson, I. Slaus, J.R. Richardson and E.L. Petersen, Phys. Rev. 153 (1967) 1127
- 1968AJ01 F. Ajzenberg-Selove, R.D. Wardaski and R. Middleton, Nucl. Phys. A116 (1968) 481
- 1968BE02 J. Benn, E.B. Dally, H.H. Muller, R.E. Pixley, H.H. Staub and H. Winkler, Nucl. Phys. A106 (1968) 296
- 1968BR23 H. Brunnader, J.C. Hardy and J. Cerny, Phys. Rev. 174 (1968) 1247
- 1968DU09 W. Durr, G. Clausnitzer, D. Fick, R. Fleischmann, H.M. Hofmann and K. Kilian, Nucl. Phys. A122 (1968) 153
- 1968FI07 D. Fick, G. Clausnitzer, W. Durr, R. Fleischmann, H.M. Hoffmann and K. Kilian, Phys. Lett. B27 (1968) 366
- 1968JA07 K.S. Jayaraman and H.D. Holmgren, Phys. Rev. 172 (1968) 1015
- 1968KA24 S. Kakigi, N. Fujiwara, K. Fukunaga, D.-C. Nguyen, S. Yamashita and T. Yanabu, J. Phys. Soc. Jpn. 25 (1968) 1214
- 1968LE01 J.K.P. Lee, S.K. Mark, P.M. Portner and R.B. Moore, Nucl. Phys. A106 (1968) 357
- 1968MC02 R.L. McGrath, J. Cerny and S.W. Cospèr, Phys. Rev. 165 (1968) 1126
- 1968MI05 P.F. Mizera and J.B. Gerhart, Phys. Rev. 170 (1968) 839
- 1968MO05 C. Moazed and H.D. Holmgren, Phys. Rev. 166 (1968) 977
- 1968PA09 P. Paul, D. Kohler and K.A. Snover, Phys. Rev. 173 (1968) 919
- 1968PL01 G.R. Plattner, T.B. Clegg and L.G. Keller, Nucl. Phys. A111 (1968) 481
- 1968ST12 R.A. Stryk and J.M. Blair, Phys. Rev. 169 (1968) 767
- 1968ZE1A Zelenskaya, El-Nager and Yarba, Sov. J. Nucl. Phys. 7 (1968) 322
- 1969BA18 K. Bahr, T. Becker, O.M. Bilaniuk and R. Jahr, Phys. Rev. 178 (1969) 1706
- 1969BA43 F.C. Barker, Aust. J. Phys. 22 (1969) 293
- 1969BL14 J.L. Black, W.J. Caelli, D.L. Livesey and R.B. Watson, Phys. Lett. B30 (1969) 100

- 1969CL10 G.J. Clark, P.B. Treacy and S.N. Tucker, *Aust. J. Phys.* 22 (1969) 663
- 1969EP01 M. Epstein, H.D. Holmgren, M. Jain, H.G. Pugh, P.G. Roos, N.S. Wall, C.D. Goodman and C.A. Ludemann, *Phys. Rev.* 178 (1969) 1698
- 1969HO11 G. Hofmann and D. Kamke, *Z. Phys.* 224 (1969) 446
- 1969KI04 K. Kilian, G. Clausnitzer, W. Durr, D. Fick, R. Fleischmann and H.M. Hofmann, *Nucl. Phys.* A126 (1969) 529
- 1969SU02 O. Sundberg and J. Kallne, *Ark. Fys.* 39 (1969) 323
- 1969SW01 W.E. Sweeney, Jr. and J.B. Marion, *Phys. Rev.* 182 (1969) 1007
- 1969SW02 W.E. Sweeney, Jr., *Phys. Rev.* 182 (1969) 1022
- 1969VI05 B. Vignon, E. Ligeon and J.P. Longequeue, *J. Phys. (Paris)* 30 (1969) 913
- 1970CA12 W.D. Callender and C.P. Browne, *Phys. Rev.* C2 (1970) 1
- 1970DE12 C. Detraz, H.H. Duhm and H. Hafner, *Nucl. Phys.* A147 (1970) 488
- 1970FI07 D. Fick, R. Kankowsky, K. Kilian and E. Salzborn, *Phys. Rev. Lett.* 24 (1970) 1503
- 1970FI1B Fisher, Thesis, Stanford Univ. (1970)
- 1970FR06 B. Frois, L. Marquez, J.L. Quebert, J.N. Scheurer, J.P. Laugier, G. Gruber, E. Heinicke and K. Meier-Ewert, *Nucl. Phys.* A153 (1970) 277
- 1970HU1B Humphrey, Helms and McGimsey, Conf-700322, USAEC (1970) 133
- 1970LA14 J.M. Lambert, P.A. Treado, L.A. Beach, R.B. Theus and E.L. Petersen, *Nucl. Phys.* A152 (1970) 516
- 1970ME1C Meadows and Whalen, *Nucl. Sci. Energ.* 41 (1970) 351
- 1970SQ01 G.T.A. Squier, A.R. Johnston, E.W. Spiers, S.A. Harbison and N.M. Stewart, *Nucl. Phys.* A141 (1970) 158
- 1970TH08 R.S. Thomason, G. Spalek and R.L. Walter, *Nucl. Phys.* A155 (1970) 659
- 1970YU02 T. Yukawa and S. Yoshida, *Phys. Let.* B33 (1970) 334
- 1970ZE03 B. Zeidman, H.T. Fortune and A. Richter, *Phys. Rev.* C2 (1970) 1612
- 1971AL09 B.J. Allen and R.L. Macklin, *Phys. Rev.* C3 (1971) 1737
- 1971BA1A Barnes, *Advances in Nucl. Phys.* 4 (1971) 133
- 1971BE52 E.H. Berkowitz, G.L. Marolt, A.A. Rollefson and C.P. Browne, *Phys. Rev.* C4 (1971) 1564
- 1971BR07 K.H. Bray, M. Jain, K.S. Jayaraman, G. LoBianco, W.T.H. Van Oers and Y.I. Wu, *Nucl. Phys.* A163 (1971) 649
- 1971DE37 C. Detraz, C.D. Zafiratos, C.E. Moss and C.S. Zaidins, *Nucl. Phys.* A177 (1971) 258
- 1971EM01 S.T. Emerson, V. Valkovic, W.R. Jackson, C. Joseph, A. Niiler, W.D. Simpson and G.C. Phillips, *Nucl. Phys.* A169 (1971) 317

- 1971FO1P D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 604
- 1971FO24 D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 576
- 1971GA21 L.L. Gadeken and E. Norbeck, Phys. Rev. Lett. 27 (1971) 952
- 1971GL07 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, Yad. Fiz. 13 (1971) 277; Sov. J. Nucl. Phys. 13 (1971) 154
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971KU10 N. Kumar and F.C. Barker, Nucl. Phys. A167 (1971) 434
- 1971NO04 E. Norbeck, L.L. Gadeken and F.D. Ingram, Phys. Rev. C3 (1971) 2073
- 1971PI06 C.J. Piluso, R.H. Spear, K.W. Carter, D.C. Kean and F.C. Barker, Aust. J. Phys. 24 (1971) 459
- 1971SC26 J.L. Schoonover, T.Y. Li and S.K. Mark, Nucl. Phys. A176 (1971) 567
- 1971SP05 H. Spinka, T. Tombrello and H. Winkler, Nucl. Phys. A164 (1971) 1; Erratum Nucl. Phys. A196 (1972) 634
- 1971SQ01 G.T.A. Squier, A.R. Johnston, E.W. Spiers, S.A. Harbison and N.M. Stewart, Nucl. Phys. A160 (1971) 602
- 1971TH02 M.J. Throop, D.H. Youngblood and G.C. Morrison, Phys. Rev. C3 (1971) 536
- 1972BA41 F.C. Barker, Astrophys. J. 173 (1972) 477
- 1972BA83 A.D. Bacher, F.G. Resmini, H.E. Conzett, R. de Swiniarski, H. Meiner and J. Ernst, Phys. Rev. Lett. 29 (1972) 1331
- 1972CO09 G. Costa, F.A. Beck and D. Magnac-Valette, Nucl. Phys. A181 (1972) 174
- 1972FA07 M.A. Fawzi, Z. Phys. 250 (1972) 120
- 1972GA32 L.L. Gadeken and E. Norbeck, Phys. Rev. C6 (1972) 1172
- 1972KA1B Kavanagh, Cosmolgy, Fusion and Other Matters; Eds., F. Reines and G. Gamow, Memorial Vol. (1972) 169
- 1972PR03 G. Presser and R. Bass, Nucl. Phys. A182 (1972) 321
- 1972TA04 M.C. Taylor, V. Valkovic and G.C. Phillips, Nucl. Phys. A182 (1972) 558
- 1972TH1C Thompson, Thesis, Yale Univ. (1972)
- 1972YO02 F.C. Young and A.R. Knudson, Nucl. Phys. A184 (1972) 563
- 1973AB1A Abe, Hiura and Tanaka, Prog. Theor. Phys. 49 (1973) 800
- 1973AG05 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, V.I. Genchev, N.F. Golovanova, G.R. Gulkanyan, V.M. Sidorov, K.M. Chernev and V.A. Yarba, Yad. Fiz. 18 (1973) 256; Sov. J. Nucl. Phys. 18 (1974) 131

- 1973AG06 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, V.I. Genchev, N.F. Golovanova, G.R. Gulkanyan, V.M. Sidorov, K.M. Chernev and V.A. Yarba, *Yad. Fiz.* 18 (1973) 264; *Sov. J. Nucl. Phys.* 18 (1974) 135
- 1973AL1B Aleshin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 1775
- 1973AN11 V.N. Andreev, V.G. Nedopekin and V.I. Rogov, *Yad. Fiz.* 17 (1973) 897; *Sov. J. Nucl. Phys.* 17 (1974) 469
- 1973AR1B Arthur, Thesis LA 5230 T (1973)
- 1973AR1C Arima, in Munich, 2 (1973) 184
- 1973AT01 M.A. Atta and M.C. Scott, *J. Nucl. Energy* 27 (1973) 875
- 1973BA1H Bahcall, in Munich, 2 (1973) 682
- 1973BA1J Balestra et al., in Munich, 1 (1973) 645
- 1973BO26 M. Bormann, D. Kaack, V. Schroder, W. Scobel and L. Wilde, *Z. Phys.* 258 (1973) 285
- 1973BR13 L. Brown, E. Steiner, L.G. Arnold and R.G. Seyler, *Nucl. Phys.* A206 (1973) 353
- 1973BR1C Bromley, in Munich, 2 (1973) 22
- 1973CA1C Catala et al., *An. Fis.* 69 (1973) 267
- 1973CO1C Coelho, *Rev. Bras. Fis. (Brazil)* 3 (1973) 463
- 1973DE1D Demarque, Mengel and Sweigart, *Nature* 246 (1973) 33
- 1973DO1B Dolgintseva and Fomenko, *Vastn. Leningr. Univ. Fiz. Khim.* 4 (1973) 43
- 1973FA1C Fang-Landau and Locher, *Nucl. Phys.* B66 (1973) 210
- 1973FO1A Fortune, Heavy Ion Lecture Series, Kansas State Univ. (1973) 1
- 1973GL1A Glor and Rudin, *Helv. Phys. Acta* 46 (1973) 441
- 1973GO41 Y.P. Gorin, S.P. Denisov, S.V. Donskov, R.N. Krasnokutskii, A.I. Petrukhin, Y.D. Prokoshkin and D.A. Stoyanova, *Yad. Fiz.* 18 (1973) 336; *Sov. J. Nucl. Phys.* 18 (1974) 173
- 1973HA12 R.C. Haskell and L. Madansky, *Phys. Rev.* C7 (1973) 1277
- 1973HA1F Hackenbroich, *Nucl. Many-Body Problem, 1972, Vol. 1* (1973) 706
- 1973HU12 J. Hudomalj and P. Tomas, *Fizika* 5 (1973) 203
- 1973KR1B Kravtsov and Skachkov, *At. Data Nucl. Data Tables* 12 (1973) 143
- 1973KU1D Kubodera, Delorme and Rho, *Nucl. Phys.* B66 (1973) 253
- 1973MIYZ T. Minamisono, Y. Nojiri, A. Mizobuchi and K. Sugimoto, *J. Phys. Soc. Jpn. Suppl.* 34 (1973) 156, 324
- 1973MU14 S.F. Mughabghab and D.I. Garber, *BNL 325, 3rd Edition, Vol. 1* (1973)
- 1973NE10 R. Neugart, *Z. Phys.* 261 (1973) 237

- 1973OG1A Ogloblin, Sov. J. Part. Nucl. 3 (1973) 467
- 1973PA20 K.H. Passler, P.G. Zint and U. Mosel, Phys. Lett. B47 (1973) 419
- 1973PE05 A.K. Petrauskas, G.P. Kamuntavicius and K.I. Jankauskas, Yad. Fiz. 17 (1973) 519; Sov. J. Nucl. Phys. 17 (1974) 266
- 1973RU09 N.Y. Rutkevich, A.P. Klyucharev, M.A. Kuzmichev and A.R. Linetskii, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1697; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 114
- 1973SC1B Scott, in Munich, 2 (1973) 216
- 1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, Nucl. Phys. A212 (1973) 573
- 1973SI38 J. Sieminska and K. Soltynski, Acta Phys. Pol. B4 (1973) 639
- 1973SO08 M.A.A. Sonnemans, J.C. Waal and R. Van Dantzig, Phys. Rev. Lett. 31 (1973) 1359
- 1973TA1B Tanabe, Tohsaki and Tamagaki, Prog. Theor. Phys. 50 (1973) 1774
- 1973TA30 K. Takahashi, M. Yamada and T. Kondoh, At. Data Nucl. Data Tables 12 (1973) 101
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973TR1B Truran, Cosmochemistry; Ed., Cameron (1973) 23
- 1973TR1C Truran, Space Sci. Rev. 15 (1973) 23
- 1973WO06 G.J. Wozniak, N.A. Jelley and J. Cerny, Phys. Rev. Lett. 31 (1973) 607
- 1973ZA06 N.I. Zaika, Y.V. Kibkalo, A.V. Mokhnach, O.F. Nemets, V.S. Semenov, V.P. Tokarev, E.D. Chesnokov and P.L. Shmarin, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1073; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 140
- 1974AC03 H. Ackermann, D. Dubbers, M. Grupp, P. Heitjans and H.-J. Stockmann, Phys. Lett. B52 (1974) 54
- 1974AD1B Adelberger, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 641
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974AL07 T.K. Alexander, G.J. Costa, J.S. Forster, O. Hausser, A.B. McDonald, A. Olin and W. Witthuhn, Phys. Rev. C9 (1974) 1748
- 1974AM01 J. Amato, R.L. Burman, R. Macek, J. Oostens, W. Schlaer, E. Arthur, S. Sobottka and W.C. Lam, Phys. Rev. C9 (1974) 501
- 1974AR05 U.R. Arifkhanov, M. Gulyamov, B.I. Islamov, M. Kayumov and E. Ergashov, Yad. Fiz. 19 (1974) 25; Sov. J. Nucl. Phys. 19 (1974) 12
- 1974AR10 L.G. Arnold, R.G. Seyler, L. Brown, T.I. Bonner and E. Steiner, Phys. Rev. Lett. 32 (1974) 895
- 1974AT1A Atkinson, Kok, De Roo and Johnson, Nucl. Phys. B77 (1974) 109
- 1974AU1B Austin, Herschberger, Riess and Simon, Bull. Amer. Phys. Soc. 19 (1974) 451

- 1974BA1N Bayukov et al., JETP Lett. 19 (1974) 399
- 1974BA1P Baye and Heenen, Nucl. Phys. A233 (1974) 304
- 1974BA1Q J.N. Bahcall, S.B. Treiman and A. Zee, Phys. Lett. B52 (1974) 275
- 1974BA42 D.P. Balamuth, R.W. Zurmuhle and S.L. Tabor, Phys. Rev. C10 (1974) 975
- 1974BA70 G. Baroni, S. Di Liberto, S. Petrera and G. Romano, Nucl. Phys. A231 (1974) 521
- 1974BE1L Beck, Borysowitz, Brink and Mihailovic, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 245
- 1974BEYR R. Beck, D.M. Brink, M.V. Mihailovic, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 1 (1974) 49
- 1974BO08 V.I. Bogatin, V.K. Bondarev, V.F. Litvin, O.V. Lozhkin, N.A. Perfilov, Y.P. Yakovlev and V.P. Bochyn, Yad. Fiz. 19 (1974) 32; Sov. J. Nucl. Phys. 19 (1974) 16
- 1974BO1F Bohm et al., Nucl. Phys. B74 (1974) 237
- 1974BO25 M. Bouten and M.-C. Bouten, Lett. Nuovo Cim. 10 (1974) 161
- 1974BO42 I. Bondouk, F. Asfour and F. Machali, Rev. Roum. Phys. 19 (1974) 551
- 1974BU16 C.A. Burke, M.T. Lunnnon and H.W. Lefevre, Phys. Rev. C10 (1974) 1299
- 1974CA32 J. Catala, A. Garcia and V. Martinez, An. Fis. 70 (1974) 186
- 1974CE05 J. Cerny, N.A. Jelley, D.L. Hendrie, C.F. Maguire, J. Mahoney, D.K. Scott and R.B. Weisenmiller, Phys. Rev. C10 (1974) 2654
- 1974CH01 F.C. Chang, Phys. Rev. C9 (1974) 1
- 1974CH1L Chiang et al., Acta Phys. Sin. (China) 23 (1974) 362
- 1974CH45 W.S. Chien and R.E. Brown, Phys. Rev. C10 (1974) 1767
- 1974CH58 I.P. Chernov, B.I. Kuznetsov, V.V. Kozyr and V.A. Matusevich, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2530; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 58
- 1974CL1E Clayton, Nature 249 (1974) 131
- 1974CO1B H.E. Conzeit, Phys. Lett. B51 (1974) 445
- 1974DA18 N.E. Davison, M.J. Canty, D.A. Dohan and A. McDonald, Phys. Rev. C10 (1974) 50
- 1974DA28 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares and R. Potenza, Nuovo Cim. A22 (1974) 313
- 1974DE25 C. Detraz, F. Pougheon, M. Bernas, M. Langevin, P. Roussel and J. Vernotte, Nucl. Phys. A228 (1974) 39
- 1974DE45 A.S. Deineko, Y.G. Mashkarov, R.P. Slabospitskii, I.I. Zalyubovskii, B.S. Bezvershuk and I.Y. Malakhov, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 1694; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 115
- 1974DI1A Didelez et al., Bull. Amer. Phys. Soc. 19 (1974) 1022

- 1974DM01 T.A. Dmitrieva, V.M. Lebedev, V.G. Markman, A.V. Spasskii and I.B. Teplov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2567; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 91
- 1974DO03 E.I. Dolinskii, V.V. Turovtsev and R. Yarmukhamedov, *Yad. Fiz.* 19 (1974) 1003; *Sov. J. Nucl. Phys.* 19 (1975) 514
- 1974DR05 R.M. Dreizler, H. Galbraith and L. Lin, *Nucl. Phys.* A234 (1974) 253
- 1974DZ03 R.I. Dzhibuti and H.M. Sallam, *Yad. Fiz.* 19 (1974) 75; *Sov. J. Nucl. Phys.* 19 (1974) 40
- 1974FE1B Feather, *Proc. Roy. Soc. Edinburgh A*71 (1974) 468
- 1974FR02 E. Friedland, H.W. Alberts and J.C. van Staden, *Z. Phys.* 267 (1974) 97
- 1974FR06 H. Friedrich, *Nucl. Phys.* A224 (1974) 537; *Erratum Nucl. Phys.* A239 (1975) 531
- 1974FR1B H. Friedrich, H. Husken and A. Weiguny, *Nucl. Phys.* A220 (1974) 125
- 1974GA1E Galstyan and Zverev, *Sov. J. Nucl. Phys.* 19 (1974) 601
- 1974GA30 A.S. Gass, O.Y. Goryunov, V.N. Dobrikov, M.G. Makowska-Rzeszutko, O.F. Nemets and A.T. Rudchik, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 873; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 184
- 1974GO35 V.A. Goldshtein, V.B. Shostak, N.G. Afanasev, V.G. Vlasenko, E.L. Kuplennikov and V.I. Startsev, *Pisma Zh. Eksp. Teor. Fiz.* 19 (1974) 695; *JETP Lett.* 19 (1974) 358
- 1974GR21 M.B. Greenfield, M.F. Werby and R.J. Philpott, *Phys. Rev.* C10 (1974) 564
- 1974GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2543; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 69
- 1974GR44 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, V.M. Pugach, B.A. Rudenko, M.V. Sokolov, B.G. Struzhko et al., *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2547; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 72
- 1974HA1C Hanna, *Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2* (1974) 249
- 1974HA1G Hanna, *Nukleonika (Poland)* 19 (1974) 655
- 1974HO11 E. Hourany, H. Nakamura, F. Takeuchi and T. Yuasa, *Nucl. Phys.* A222 (1974) 537
- 1974HO1D Holstein, *Rev. Mod. Phys.* 46 (1974) 789
- 1974HO30 H. Ho, W. Dunnweber, D. Dehnhard, K. Mudersbach and J.P. Wurm, *Nucl. Phys.* A233 (1974) 361
- 1974HY01 M. Hyakutake, M. Sonoda, A. Katase, Y. Wakuta, M. Matoba, H. Tawara and I. Fujita, *J. Nucl. Sci. Tech. (Japan)* 11 (1974) 407
- 1974IR04 J.M. Irvine, G.S. Mani and M. Vallieres, *Czech. J. Phys.* B24 (1974) 1269

- 1974KA11 S.G. Kadenskii, A.A. Martynov and Y.I. Kharitonov, *Yad. Fiz.* 19 (1974) 529; *Sov. J. Nucl. Phys.* 19 (1974) 267
- 1974KA1J Kakigi et al., *Bull. Inst. Chem. Res. Kyoto Univ.* 52 (1974) 218
- 1974KE1C Kemper, Moore, Puigh and White, *Bull. Amer. Phys. Soc.* 19 (1974) 503
- 1974KO16 R.L. Kozub, D. Logan, J.M. Miller and A.M. Zebelman, *Phys. Rev. C* 10 (1974) 1246
- 1974KO24 S.D. Kostyuchenkov, V.M. Lebedev, A.V. Spasskii and I.B. Teplov, *Vestn. Mosk. Univ. Fiz. Astron.* 15 (1974) 424
- 1974KO35 E. Kondaiyah, *J. Phys. (London)* 7A (1974) 1457
- 1974KU06 N. Kumar, *Nucl. Phys.* A225 (1974) 221
- 1974LA29 M. Lattuada and R. Potenza, *Lett. Nuovo Cim.* 11 (1974) 662
- 1974LE04 L.B. Leinson, *Yad. Fiz.* 19 (1974) 283; *Sov. J. Nucl. Phys.* 19 (1974) 139
- 1974LE14 V.M. Lebedev, A.V. Spassky, I.B. Teplov and N.S. Zelenskaya, *Nucl. Phys.* A223 (1974) 351
- 1974LE1E Leung, Rept. TPB-ND-7, Phys. Tech. Bund., Braunschweig, Germany (1974)
- 1974LI16 L. Lin, R.M. Dreizler and H. Galbraith, *Lett. Nuovo Cim.* 11 (1974) 627
- 1974LI1E A. Lindner and H. Ebinghaus, *Nucl. Phys.* A230 (1974) 487
- 1974LO10 J.M. Lombaard and E. Friedland, *Z. Phys.* 268 (1974) 413
- 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325
- 1974LU05 A. Lumbroso, *Phys. Rev. C* 10 (1974) 1271
- 1974MA1E Maguin, *Nuovo Cim.* A19 (1974) 638
- 1974MA49 S. Matsuki, S. Yamashita, N. Fujiwara, K. Fukunaga, D.C. Nguyen and T. Yanabu, *Bull. Inst. Chem. Res. Kyoto Univ.* 52 (1974) 202
- 1974MC04 R.J. McCarthy and G.E. Walker, *Phys. Rev. C* 9 (1974) 809
- 1974MCZS C.R. McClenahan, Thesis, Northwestern Univ. (1974); ANL-8088 (1974)
- 1974ME09 A. Menchaca-Rocha, *Nucl. Instrum. Meth.* 114 (1974) 425
- 1974NE1B Negele, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 618
- 1974RA16 M. Rahman, D. Husain and S. Ali, *Phys. Rev. C* 10 (1974) 1
- 1974RO17 R.G.H. Robertson, S. Martin, W.R. Falk, D. Ingham and A. Djalois, *Phys. Rev. Lett.* 32 (1974) 1207
- 1974RO20 P. Roubeau, A. Abragam, G.L. Bacchella, H. Glattli, A. Malinovski, P. Meriel, J. Piesvaux and M. Pinot, *Phys. Rev. Lett.* 33 (1974) 102

- 1974RU06 N.Y. Rutkevich, A.P. Klyucharev, G.A. Zachepilo, M.A. Kuzmichev and A.R. Linetskii, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 1737; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 150
- 1974SA1K Sakamoto, Ogino, Tomita and Inagaki, *Bull. Inst. Chem. Res. Kyoto Univ.* 52 (1974) 146
- 1974SE1B Serov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 84
- 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)
- 1974SO1C Sonnemans, Thesis, IKO, Amsterdam (1974)
- 1974SO1D Sorriaux, Rubio and Boudet, *Nucl. Instrum. Meth. Phys. Res.* 118 (1974) 213
- 1974ST1D Steiner and Tobias, *Nucl. Fusion* 14 (1974) 153
- 1974TH04 D.R. Thompson, Y.C. Tang and F.S. Chwieroth, *Phys. Rev. C* 10 (1974) 987
- 1974TR01 R.E. Tribble and G.T. Garvey, *Phys. Rev. Lett.* 32 (1974) 314
- 1974TU1A Turk, Antolkovic, Dolenc and Winterhalter, *Proc. of 1973 Gaussig Conf., ZFK-271* (1974) 115
- 1974UL1B Ulrich, *Astrophys. J.* 188 (1974) 369
- 1974VA1C Vasilev, Kocharov and Levkovskii, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 1827
- 1974VA33 M. Varnagy, J. Csikai, J. Szabo and S. Szegedi, J. Banhalmi, *Nucl. Instrum. Meth.* 119 (1974) 451
- 1974VO06 P.L. Von Behren, E. Norbeck and G.L. Payne, *Phys. Rev. C* 10 (1974) 550
- 1974WA02 W.L. Wang and R.G. Lipes, *Phys. Rev. C* 9 (1974) 814
- 1974WA1D W.L. Wang, *Phys. Lett.* B52 (1974) 143
- 1974WO1C Wozniak, LBL-2999 (1974)
- 1974WO1D Wozniak, Jelley and Cerny, *Nucl. Instrum. Meth.* 120 (1974) 29
- 1974YA1C Yasue et al., *Bull. Inst. Chem. Res. Kyoto Univ.* 52 (1974) 177
- 1974ZI1A D. Zieminska and R.H. Dalitz, *Nucl. Phys.* B74 (1974) 248
- 1975AB1D Abdul-Magd, *Prog. Theor. Phys.* 53 (1975) 997
- 1975AB1E Abe, *Clustering Phenom. in Nucl., II, ORO-4856-26* (1975) 500
- 1975ADZV E.G. Adelberger, S.J. Freedman, A.V. Nero, A.B. McDonald, R.G.H. Robertson and D.R. Goosman, *Bull. Amer. Phys. Soc.* 20 (1975) 596, DE 3
- 1975AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A248 (1975) 1
- 1975AN01 B. Antolkovic and Z. Dolenc, *Nucl. Phys.* A237 (1975) 235
- 1975AN1F Ando, Ikeda and Suzuki, *Clustering Phenom. in Nucl., II, ORO-4856-26* (1975) 227
- 1975AN1G Antolkovic and Turk, *Fiz. Suppl.* 7 (1975) 55

- 1975AR10 F. Arickx, P. Van Leuven and M. Bouten, *Z. Phys.* 273 (1975) 205
- 1975AR28 F. Arickx, P. Van Leuven and M. Bouten, *Nucl. Phys.* A252 (1975) 416
- 1975AU01 G. Audi, C. Detraz, M. Langevin and F. Pougheon, *Nucl. Phys.* A237 (1975) 300
- 1975AZ02 S.A. Azimov, U.R. Arifkhanov, M. Gulyamov, B.I. Islamov, T. Iskhakov, A.M. Mukhamedzhanov, E. Ergashov, Y.I. Faizullaev and R. Yarmukhamedov, *Yad. Fiz.* 22 (1975) 225; *Sov. J. Nucl. Phys.* 22 (1976) 115
- 1975BA76 A.I. Baz, V.F. Demin and M.V. Zhukov, *Fiz. Elem. Chastits At. Yad. (USSR)* 6 (1975) 515; *Sov. J. Part. Nucl.* 6 (1975) 207
- 1975BE01 F.D. Becchetti, L.T. Chua, J. Janecke and A.M. Vander Molen, *Phys. Rev. Lett.* 34 (1975) 225
- 1975BE16 R. Beck, J. Borysowicz, D.M. Brink and M.V. Mihailovic, *Nucl. Phys.* A244 (1975) 58
- 1975BE1M R. Beck, J. Borysowicz, D.M. Brink and M.V. Mihailovic, *Nucl. Phys.* A244 (1975) 45
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, *Nucl. Phys.* A249 (1975) 1
- 1975BE38 W. Benenson, A. Guichard, E. Kashy, D. Mueller, H. Nann and L.W. Robinson, *Phys. Lett.* B58 (1975) 46
- 1975BI14 B. Bilwes, R. Bilwes, J.L. Ferrero, A. Garcia and V. Martinez, *J. Phys. (Paris)* 36 (1975) 1055
- 1975BO1C Bondouk, Asfour and Machali, *Ann. Phys.* 32 (1975) 255
- 1975BO56 I. Bondouk, F. Asfour, F. Machali and Z. Saleh, *Rev. Roum. Phys.* 20 (1975) 1111
- 1975BU1B Buck et al., *Clustering Phenom. in Nucl., II*, ORO-4856-26 (1975) 429
- 1975CA1F Calaprice, *Proc. Int. Symp. on Interaction Studies in Nucl., Germany, 1975* (1975) 83
- 1975CI1A Ciric et al., *Fiz. Suppl.* 7 (1975) 62
- 1975CL1C Clement, Kanellopoulos and Wildermuth, *Acta Phys. Austriaca* 42 (1975) 29
- 1975CO06 B. Coupat, P.Y. Bertin, D.B. Isabelle, P. Vernin, A. Gerard, J. Miller, J. Morgenstern, J. Picard and B. Saghai, *Phys. Lett.* B55 (1975) 286
- 1975CO1C Cowley and Heymann, *Clustering Phenom. in Nucl., II*, ORO-4856-26 (1975) 323
- 1975DA15 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares and R. Potenza, *Lett. Nuovo Cim.* 13 (1975) 489
- 1975DI1B Didelez et al., *Jul-Conf-16* (1975) 90
- 1975DO04 L.R. Dodd, M.A.A. Sonnemans and R. van Dantzig, *Lett. Nuovo Cim.* 12 (1975) 597
- 1975DU09 S. Dubnicka and O. Dumbrajs, *Phys. Lett.* B57 (1975) 327
- 1975EB03 K.A. Eberhard, E. Mathiak, J. Stettmeier, W. Trombik, A. Weidinger, L.N. Wustefeld and K.G. Bernhardt, *Phys. Lett.* B56 (1975) 445

- 1975ER09 K.M. Erokhin, N.V. Orlova, Y.F. Smirnov and K.V. Shitikova, *Yad. Fiz.* 22 (1975) 1102; *Sov. J. Nucl. Phys.* 22 (1975) 574
- 1975FE1A Feather, *Commun. Roy. Soc. Edinburgh (Phys. Sci.)* 2 (1975) 19
- 1975GE1F Geesman, Noe, Paul and Suffert, *Bull. Amer. Phys. Soc.* 20 (1975) 597
- 1975GL08 Y.A. Glukhov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, *Yad. Fiz.* 22 (1975) 679; *Sov. J. Nucl. Phys.* 22 (1976) 351
- 1975GO07 P. Goldhammer, *Phys. Rev.* C11 (1975) 1422
- 1975GO36 O.Y. Goryunov, V.N. Dobrikov, O.F. Nemets, E.P. Reuk and A.T. Rudchik, *Ukr. Fiz. Zh.* 20 (1975) 1775
- 1975GR05 A.M. Green and J. Niskanen, *Nucl. Phys.* A240 (1975) 263
- 1975GR1C Green, *Few Body Problems, Quebec, 1974* (1975) 352
- 1975GR41 K.A. Gridnev and A.A. Ogloblin, *Fiz. Elem. Chastits At. Yad. (USSR)* 6 (1975) 393; *Sov. J. Part. Nucl.* 6 (1976) 158
- 1975GR43 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, V.M. Pugach, M.V. Sokolov, B.G. Struzhko, B.A. Rudenko et al., *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 1717; *Bull. Acad. Sci. USSR Phys. Ser.* 39 (1975) 127
- 1975HE07 A.C. Heemskerk, L.P. Kok and M. De Roo, *Nucl. Phys.* A244 (1975) 15
- 1975HO1E Horiuchi, *Prog. Theor. Phys.* 53 (1975) 447
- 1975IG1A Igo, *AIP Conf. Proc.* 26 (1975) 63
- 1975JA10 J. Janecke, F.D. Becchetti, L.T. Chua and A.M. VanderMolen, *Phys. Rev.* C11 (1975) 2114
- 1975JA1A Jaros et al., *Bull. Amer. Phys. Soc.* 20 (1975) 666
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, *Phys. Rev.* C11 (1975) 2049
- 1975JE04 H. Jeremie and J. Baltar, *Can. J. Phys.* 53 (1975) 2639
- 1975KA05 J. Kasagi, T. Nakagawa, N. Sekine, T. Tohei and H. Ueno, *Nucl. Phys.* A239 (1975) 233
- 1975KH1A Khadkikar, Kamble and Kulkarni, *Proc. Int. Conf. Nucl. Self-Consistent Fields, Italy, 1975* (1975) 85
- 1975KI14 King et al., *Phys. Rev. Lett.* 35 (1975) 988
- 1975KO18 R. Kouzes and W.H. Moore, *Phys. Rev.* C12 (1975) 1511; *Erratum Phys. Rev.* C13 (1976) 890
- 1975KO1C Kohler, *Phys. Rept.* C18 (1975) 217
- 1975KR14 S. Krol and S. Szpikowski, *Acta Phys. Pol.* B6 (1975) 409
- 1975KU09 V.I. Kukulkin, V.G. Neudatchin and Y.F. Smirnov, *Nucl. Phys.* A245 (1975) 429

- 1975KU1J V. Kumar, G.P. Sharma and A.P. Sharma, Phys. Lett. B59 (1975) 95
- 1975KU27 N. Kumar and S.K. Soni, Indian J. Pure Appl. Phys. 13 (1975) 826
- 1975LI1E Liskien and Paulsen, At. Data Nucl. Data Tables 15 (1975) 57
- 1975MA1G Mathews et al., Bull. Amer. Phys. Soc. 20 (1975) 710
- 1975MA1H Mashkarov, Chegoryan and Zalyubovskii, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 560
- 1975MA27 V.V. Malyarov, I.V. Poplavskii and M.N. Popushoi, Yad. Fiz. 21 (1975) 987; Sov. J. Nucl. Phys. 21 (1975) 506
- 1975MC02 C.R. McClenahan and R.E. Segel, Phys. Rev. C11 (1975) 370
- 1975MC18 M.W. McNaughton, N.S.P. King, F.P. Brady, J.L. Romero and T.S. Subramanian, Nucl. Instrum. Meth. Phys. Res. 130 (1975) 555
- 1975MI04 T. Minamisono, J.W. Hugg, D.G. Mavis, T.K. Saylor, S.M. Lazarus, H.F. Glavish and S.S. Hanna, Phys. Rev. Lett. 34 (1975) 1465
- 1975MI1A Miljanic, Zabel and Phillips, Few Body Problems, Quebec, 1974 (1975) 669
- 1975NA12 A.M. Nathan, G.T. Garvey, P. Paul and E.K. Warburton, Phys. Rev. Lett. 35 (1975) 1137; Erratum Phys. Rev. Lett. 49 (1982) 1056
- 1975NA1C Nathan, Thesis, Princeton Univ. (1975)
- 1975NO1C Norbeck, Fox and Payne, Bull. Amer. Phys. Soc. 20 (1975) 1174
- 1975PA11 G.L. Payne and R.R. Carlson, Phys. Rev. C12 (1975) 1260
- 1975RO01 R.G.H. Robertson, W.S. Chien and D.R. Goosman, Phys. Rev. Lett. 34 (1975) 33
- 1975RO09 R. Roy, J. Birchall, B. Frois, U. Von Moellendorff, C.R. Lamontagne and R.J. Slobodrian, Nucl. Phys. A245 (1975) 87
- 1975RO1E Rogge et al., Jul-Conf-16 (1975) 91
- 1975SC1K Schenzle, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 85
- 1975SC41 U. Schwinn, G. Mairle, G.J. Wagner and C. Ramer, Z. Phys. A275 (1975) 241
- 1975SE03 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 151
- 1975SE04 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 159
- 1975SE07 F. Seiler, Nucl. Phys. A244 (1975) 236
- 1975SH01 P.E. Shanley, Phys. Rev. Lett. 34 (1975) 218
- 1975SO1C Soleilhac, CEA-N-1812 (1975)
- 1975TA1D Tanabe, Tohsaki and Tamagaki, Prog. Theor. Phys. 53 (1975) 677
- 1975TO1A Tombrello, Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 167

- 1975TR02 P.B. Treacy, Nucl. Phys. A240 (1975) 413
- 1975TR06 R.E. Tribble and G.T. Garvey, Phys. Rev. C12 (1975) 967
- 1975UL1A Ulrich, Science 190 (1975) 619
- 1975UN1A Unternahrer, Lang, Buta and Muller, Helv. Phys. Acta 48 (1975) 558
- 1975VA04 V. Valkovic, D. Miljanic, R.B. Liebert and G.C. Phillips, Nucl. Phys. A239 (1975) 260
- 1975VO1C Von Behren, Oliver, Risser and Phillips, Bull. Amer. Phys. Soc. 20 (1975) 560
- 1975WI1E Wilkinson, Proc. Int. Symp. on Interaction Studies in Nucl., Germany, 1975 (1975) 147
- 1975WI25 F. Wittchow, J. Ulbricht, U. Holm, K.D. Stahl and H. Ebinghaus, Phys. Lett. B59 (1975) 29
- 1975ZA06 N.I. Zaika and P.L. Shmarin, Ukr. Fiz. Zh. 20 (1975) 238
- 1975ZE01 A.M. Zebelman, A.M. Poskanzer, J.D. Bowman, R.G. Sextro and V.E. Viola, Jr., Phys. Rev. C11 (1975) 1280
- 1975ZW01 Zwiegliniski, Saganek, Sledzinska and Wilhelmi, Nucl. Phys. A250 (1975) 93
- 1976AJ01 F. Ajzenberg-Selove, C.F. Maguire, D.L. Hendrie, D.K. Scott and J. Mahoney, Phys. Rev. C13 (1976) 46
- 1976AL1E Ali, Proc. 7th Int. Conf. on Few Body Problems in Nucl. Part. Phys., Delhi, 1975 (1976) 438
- 1976AN16 B. Antolkovic, Fizika 8 (1976) 163
- 1976AR02 J.L. Artz, M.B. Greenfield and N.R. Fletcher, Phys. Rev. C13 (1976) 156
- 1976AR07 F. Arickx, Nucl. Phys. A268 (1976) 347
- 1976AR11 N. Arena, D. Vinciguerra, F. Riggi and C. Spitaleri, Lett. Nuovo Cim. 17 (1976) 231
- 1976AR1F Arnould, Polarization, Zurich, 1975 (1976) 503
- 1976BA1E Baz and Filippov, Proc. Dubna Conf. JINR-9920, Vol. II (1976) 268
- 1976BA1J Bahcall and Davis, Science 191 (1976) 264
- 1976BA1N Baz, Goldberg, Gridnev and Semenov, Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 96
- 1976BA67 F.C. Barker, G.M. Crawley, P.S. Miller and W.F. Steele, Aust. J. Phys. 29 (1976) 245
- 1976BE1G Berlovich, Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 756
- 1976BE1K Bertini, Spallation Nucl. Reactions and Their Applications; Eds., Shen and Merker (1976) 27
- 1976BE1L Benenson, Kashy, Mueller and Nann, Cargese Conf., CERN 76-13 (1976) 235

- 1976BE67 B.N. Belyaev, V.D. Domkin, Y.G. Korobulin and L.M. Krizhanskii, *Yad. Fiz.* 24 (1976) 1177; *Sov. J. Nucl. Phys.* 24 (1976) 617
- 1976BI1B Bilpuch et al., Lowell Conf., CONF-760715-P2 (1976) 1309
- 1976BO08 V.I. Bogatin, V.F. Litvin, O.V. Lozhkin, N.A. Perfilov and Y.P. Yakovlev, *Nucl. Phys.* A260 (1976) 446
- 1976BO15 I. Borbely, *Nucl. Phys.* A262 (1976) 244
- 1976BR32 R.E. Brown and Y.C. Tang, *Phys. Rev.* C14 (1976) 1675
- 1976BR42 J.D. Brandenberger, F.D. Snyder, J.D. Dawson, T.W. Burrows and F.D. McDaniel, *Nucl. Instrum. Meth.* 138 (1976) 321
- 1976BU16 M. Buenerd, C.K. Gelbke, B.G. Harvey, D.L. Hendrie, J. Mahoney, A. Menchaca-Rocha, C. Olmer and D.K. Scott, *Phys. Rev. Lett.* 37 (1976) 1191
- 1976CA1E Calaprice, *Bull. Amer. Phys. Soc.* 21 (1976) 507
- 1976CAZG J. Campbell and M.C. Scott, *Bull. Amer. Phys. Soc.* 21 (1976) 1337, RT67
- 1976CO18 W.R. Coker and T. Tamura, *Phys. Lett.* B62 (1976) 374
- 1976CU07 R.Y. Cusson, R. Hilko and D. Kolb, *Nucl. Phys.* A270 (1976) 437
- 1976CY1A Cybulska, Douglas, Farrell-Pessoa and Wylie, Lowell Conf., CONF-76015-P2 (1976) 1388
- 1976DA15 S.E. Darden, G. Murillo and S. Sen, *Nucl. Phys.* A266 (1976) 29
- 1976DA24 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares, N. Arena, G. Calvi and R. Potenza, *Lett. Nuovo Cim.* 15 (1976) 563
- 1976DE31 G.C. Deka, K. Pathak and I. Laskar, *Can. J. Phys.* 54 (1976) 1905
- 1976DOZW Sr.M.A. Doyle, H.W. Clark, K.R. Crosthwaite, L.J. Dries, J.L. Regner and T.R. Donoghue, *Bull. Amer. Phys. Soc.* 21 (1976) 636, HG1
- 1976EA1A Earwacker and Walker, Lowell Conf., CONF-760715-P2 (1976)
- 1976EB01 K.A. Eberhard and K.G. Bernhardt, *Phys. Rev.* C13 (1976) 440
- 1976FE1A Feather, *Commun. Roy. Soc. Edinburgh (Phys. Sci.)* 7 (1976) 73
- 1976FI1C Filippov, Ovcharenko, Maksimenko and Steshenko, *Prob. of Vibrational Nucl.*, Zagreb, 1974 (1975) 333
- 1976FO03 J.C. Fong, M.M. Gazzaly, G. Igo, A.D. Liberman, R.J. Ridge, S.L. Verbeck, C.A. Whitten, Jr., V. Perez-Mendez and W.R. Coker, *Nucl. Phys.* A262 (1976) 365
- 1976FO21 C.-M. Fou, Y.-C. Liu, C.-C. Hsu and S.-L. Huang, *J. Phys. (London)* G2 (1976) 847
- 1976FU06 G.H. Fuller, *J. Phys. Chem. Ref. Data* 5 (1976) 835
- 1976GAYV D.I. Garber and R.R. Kinsey, *BNL 325, 3rd Edition, Vol. 2* (1976)
- 1976GO12 G.-H. Goritz and U. Mosel, *Z. Phys.* A277 (1976) 243

- 1976GO1F Gould and Boyce, Nucl. Sci. Eng. 60 (1976) 477
- 1976GR22 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, M.V. Sokolov, B.G. Struzhko and B.A. Rudenko, Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 203; Bull. Acad. Sci. USSR Phys. Ser. 40 (1976) 172
- 1976HA18 H.H. Hackenbroich, T.H. Seligman, W. Zahn and D. Fick, Phys. Lett. B62 (1976) 121
- 1976HA1J Hanna, Polarization, Zurich, 1975 (1976) 407
- 1976HA1L Han, J. Natl. Chiao Tung Univ. (Taiwan) 1 (1976) 311
- 1976HE1F Hermans et al., Polarization, Zurich, 1975 (1976) 813
- 1976HI04 F. Hinterberger, P.V. Rossen, E. Lebsanft, P. Luttkicken, J. Bisping, R. Jahn and B. Schuller, Nucl. Phys. A259 (1976) 385
- 1976HO1A Horiuchi, Invited Talk at VII Int. Conf. on Few Body Problems, Delhi, 1975-1976 (1976)
- 1976IK01 P.G. Ikossi, W.J. Thompson, T.B. Clegg, W.W. Jacobs and E.J. Ludwig, Phys. Rev. Lett. 36 (1976) 1357
- 1976IR1B Irvine et al., Ann. Phys. 102 (1976) 129
- 1976JA1F H.R. Jaqaman and A.Z. Mekjian, Nucl. Phys. A259 (1976) 157
- 1976JA1G Jacobs, Clegg, Ikossi and Ludwig, Polarization, Zurich, 1975 (1976) 697
- 1976JI1A Jing-Shang and Yi-Zhong, Acta Phys. Sin. 25 (1976) 293
- 1976KA23 O. Karban, A.K. Basak, J.B.A. England, G.C. Morrison, J.M. Nelson, S. Roman and G.G. Shute, Nucl. Phys. A269 (1976) 312
- 1976KH04 S.B. Khadkikar and V.B. Kamble, Phys. Lett. B64 (1976) 131
- 1976KN1D Knox, White and Lane, Lowell Conf., CONF-760715-P2 (1976) 1307
- 1976KO03 V.I. Komarov, G.E. Kosarev, E.S. Kuzmin, A.G. Molokanov, G.P. Reshetnikov, O.V. Savchenko and S. Tesch, Nucl. Phys. A256 (1976) 362
- 1976KO18 R.P. Kolalis, V.S. Sadkovskii, G.A. Feofilov, A.E. Denisov, L.I. Vinogradov and I.S. Altarev, Yad. Fiz. 24 (1976) 246; Sov. J. Nucl. Phys. 24 (1976) 127
- 1976KU07 N. Kumar and O. Prakash, Z. Phys. A278 (1976) 105
- 1976KU1B Kubodera, Suppl. Prog. Theor. Phys. 60 (1976) 29
- 1976LA1C Lane, White and Knox, Lowell Conf., CONF-760715-P2 (1976) 1042
- 1976LE1J Letourneux and Onsi, Bull. Amer. Phys. Soc. 21 (1976) 11
- 1976LE1K Lebedev et al., Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 98
- 1976LI1E Lindner, Polarization, Zurich, 1975 (1976) 817

- 1976MA64 Y.G. Mashkarov, A.S. Deineko and R.P. Slabospitskii, *Izv. Akad. Nauk SSSR Ser. Fiz.* 40 (1976) 2189; *Bull. Acad. Sci. USSR Phys. Ser.* 40 (1976) 140
- 1976ME1C Meboniya and Surmava, *Sov. J. Nucl. Phys.* 24 (1976) 253
- 1976ME20 D.V. Meboniya and I.G. Surmava, *Yad. Fiz.* 24 (1976) 308; *Sov. J. Nucl. Phys.* 24 (1976) 161
- 1976MI17 Y. Mito and M. Kamimura, *Prog. Theor. Phys.* 56 (1976) 583
- 1976MI1C Miljanic, Pecar, Rendic and Tomas, *Lowell Conf.*, CONF-760715-P2 (1976) 1343
- 1976MO23 J.E. Monahan, A.J. Elwyn and F.J.D. Serduke, *Nucl. Phys.* A269 (1976) 61
- 1976MO31 P. Moller and J.R. Nix, *Nucl. Phys.* A272 (1976) 502
- 1976NA05 P. Nagel and R.D. Koshel, *Phys. Rev.* C13 (1976) 907
- 1976NE1B Nelson, Purser, von Behren and Newson, *Lowell Conf.*, CONF-760715-P2 (1976) 1475
- 1976NO07 J.W. Noe, D.F. Geesaman, P. Paul and M. Suffert, *Phys. Lett.* B65 (1976) 125
- 1976OV1B Ovcharenko, Pavlenko and Fillippov, *Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct.*, Vol. 1, JINR D-9682 (1976) 24
- 1976PI10 D.J. Pisano and P.D. Parker, *Phys. Rev.* C14 (1976) 475
- 1976PO06 C.H. Poppe, J.D. Anderson, J.C. Davis, S.M. Grimes and C. Wong, *Phys. Rev.* C14 (1976) 438
- 1976RA1E Raichev, Roussev and Slavov, *Doklady Bolgarskoi Akad. Nauk* 29 (1976) 37
- 1976RI1B A.I.M. Ritchie, *J. Phys.* D9 (1976) 15
- 1976RO04 R.G.H. Robertson, W. Benenson, E. Kashy and D. Mueller, *Phys. Rev.* C13 (1976) 1018
- 1976RO10 J.L. Romero, F.P. Brady and J.A. Jungerman, *Nucl. Instrum. Meth.* 134 (1976) 537
- 1976RO1L Roman, *Polarization, Zurich, 1975* (1976) 255
- 1976RO1M Robertson, *At. Masses Fund. Constants; Eds.*, Sanders and Wapstra (1976) 147
- 1976SA16 R. Sartor, *Nucl. Phys.* A267 (1976) 29
- 1976SAZR H. Sandhu, L. Gonczi and B. Sundqvist, *Bull. Amer. Phys. Soc.* 21 (1976) 535, BF16
- 1976SH02 J.D. Sherman, D.L. Hendrie and M.S. Zisman, *Phys. Rev.* C13 (1976) 20
- 1976SL02 R.J. Slobodrian, *Phys. Lett.* B63 (1976) 5
- 1976SL2A I. Slaus, *Lowell Conf.*, CONF-760715-P1 (1976) 272
- 1976SM08 D.L. Smith, *Nucl. Sci. Eng.* 61 (1976) 540
- 1976SOZW A.M. Sourkes, D.K. Hasell, M.S. De Jong, T. Nasr, R.F. Carlson, A.J. Cox and D.J. Margaziotis, *Bull. Amer. Phys. Soc.* 21 (1976) 987, CD10
- 1976ST04 A.I. Steshenko, *Yad. Fiz.* 23 (1976) 44; *Sov. J. Nucl. Phys.* 23 (1976) 23

1976ST11 W.F. Steele, P.A. Smith, J.E. Finck and G.M. Crawley, Nucl. Phys. A266 (1976) 424

1976STYX D.P. Stahel, R. Jahn, G.J. Wozniak and J. Cerny, Bull. Amer. Phys. Soc. 21 (1976) 1295, CC8

1976TR01 R.E. Tribble, R.A. Kenefick and R.L. Spross, Phys. Rev. C13 (1976) 50

1976TU04 M. Turk, B. Antolkovic and D. Winterhalter, Nucl. Phys. A270 (1976) 381

1976UL1C Ulbright et al., Polarization, Zurich, 1975 (1976) 875

1976VA29 G.V. Valskii, Yad. Fiz. 24 (1976) 270; Sov. J. Nucl. Phys. 24 (1976) 140

1976WE09 R.B. Weisenmiller, N.A. Jelley, K.H. Wilcox, G.J. Wozniak and J. Cerny, Phys. Rev. C13 (1976) 1330

1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, Phys. Rev. C14 (1976) 815

1977AJ02 F. Ajzenberg-Selove, Nucl. Phys. A281 (1977) 1

1977AL1G Ali and Ferdous, in Tokyo (1977) 514

1977AM1B Amann et al., Bull. Amer. Phys. Soc. 22 (1977) 1007

1977AN1B Ando, Ikeda and Tohsaki, in Tokyo (1977) 587

1977AN1C Antolkovic and Rudar, Fiz. Suppl. 9 (1977) 50

1977AR08 F. Arickx, Nucl. Phys. A284 (1977) 264

1977AR09 N. Arena, S. Cavallaro, M. Lattuada, R. Potenza, M.L. Sperduto, V. D'Amico, G. Fazio, S. Femino and S. Jannelli, Lett. Nuovo Cim. 19 (1977) 125

1977AS04 J. Asai, C. Rioux, M. Irshad, R. Pigeon and R.J. Slobodrian, Phys. Rev. C16 (1977) 1762

1977AU1B Austin, MSUCL-254 (1977)

1977BA17 A.I. Baz, V.Z. Goldberg, K.A. Gridnev, V.M. Semjonov and E.F. Hefter, Z. Phys. A280 (1977) 171

1977BA1P Batusov and Eramzhyan, Fiz. Elem. Chastits At. Yad. (USSR) 8 (1977) 229

1977BA1V Bahcall, AIP Conf. Proc. 37 (1977) 53

1977BA51 B. Bassalleck, W.-D. Klotz, F. Takeutchi, H. Ullrich and M. Furic, Phys. Rev. C16 (1977) 1526

1977BA62 F.C. Barker, Aust. J. Phys. 30 (1977) 113

1977BE1Q Bernheim et al., Proc., Zurich-Sin (1977) 267

1977BE1R Berger et al., in Tokyo (1977) 761

1977BE1T Berger et al., Proc., Zurich-Sin (1977) 160

1977BE1U Bedjidian et al., Proc., Zurich-Sin (1977) 308

1977BE50 F. Becker and I. Strykowski, Nucl. Phys. A289 (1977) 425

- 1977BE51 F. Becker and I. Strykowski, Nucl. Phys. A289 (1977) 446
- 1977BO22 E. Bovet, F. Foroughi, C. Nussbaum and G. Pauletta, Lett. Nuovo Cim. 19 (1977) 270
- 1977BO29 I.I. Bondouk and S. Saad, Atomkernergie 29 (1977) 270
- 1977BU01 B. Buck, H. Friedrich and C. Wheatley, Nucl. Phys. A275 (1977) 246
- 1977BU02 A. Buchnea, R.G. Johnson and K.G. McNeill, Can. J. Phys. 55 (1977) 364
- 1977CA06 L.F. Canto and D.M. Brink, Nucl. Phys. A279 (1977) 85
- 1977CH02 N.S. Chant and P.G. Roos, Phys. Rev. C15 (1977) 57
- 1977CH1C Chuu and Han, J. Phys. (London) G3 (1977) 555
- 1977CO07 A.A. Cowley, P.G. Roos, N.S. Chant, R. Woody, III, H.D. Holmgren and D.A. Goldberg, Phys. Rev. C15 (1977) 1650
- 1977DA20 N.Z. Darwisch, K.A. Gridnev, E.F. Hefter and V.M. Semjonov, Nuovo Cim. A42 (1977) 303
- 1977DE08 C. Detraz, Phys. Lett. B66 (1977) 333
- 1977DI11 J.K. Dickens, G.L. Morgan, G.T. Chapman, T.A. Love, E. Newman and F.G. Perey, Nucl. Sci. Eng. 62 (1977) 515
- 1977DU06 D. Dubbers, K. Dorr, H. Ackermann, F. Fujara, H. Grupp, M. Grupp, P. Heitjans, A. Korblein and H.-J. Stockmann, Z. Phys. A282 (1977) 243
- 1977EL09 A.J. Elwyn, R.E. Holland, C.N. Davids, L. Meyer-Schutzmeister, J.E. Monahan, F.P. Mooring and W. Ray, Jr., Phys. Rev. C16 (1977) 1744
- 1977EL11 A.J. Elwyn, J.E. Monahan and F.J.D. Serduke, Nucl. Sci. Eng. 63 (1977) 343
- 1977FE1B Feshbach, High Energy Phys. Nucl. Struct., Zurich, 1977 (1977) 291
- 1977FL13 T. Fliessbach and P. Manakos, J. Phys. (London) G3 (1977) 643
- 1977FR12 V. Franco and A. Tekou, Phys. Rev. C16 (1977) 658
- 1977FR18 H. Friedrich and L.F. Canto, Nucl. Phys. A291 (1977) 249
- 1977FU09 K. Fukunaga, T. Ohsawa, S. Kakigi, S. Tanaka and N. Fujiwara, J. Phys. Soc. Jpn. 42 (1977) 725
- 1977FU1D Furutani, Horiuchi and Tamagaki, in Tokyo (1977) 145
- 1977FU1E Fukushima and Kamimura, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 225
- 1977FU1F Furutani, Horiuchi and Tamagaki, Fiz. Suppl. 9 (1977) 29
- 1977GA1E Garvey, AIP Conf. Proc. 37 (1977) 104
- 1977GA1F Galstyan, Yad. Fiz. 26 (1977) 225
- 1977GAZZ G.T. Garvey, T. Bowles and A. Nathan, Bull. Amer. Phys. Soc. 22 (1977) 27

- 1977GL05 M. Glor, H.P. Nagele, G. Morgan, R. Neff, H. Rudin and F. Seiler, Nucl. Phys. A286 (1977) 31
- 1977GR10 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.A. Pilipchenko, V.M. Pugach, B.A. Rudenko, M.V. Sokolov, B.G. Struzhko and A.M. Shur, Izv. Akad. Nauk SSSR Ser. Fiz. 41 (1977) 147; Bull. Acad. Sci. USSR Phys. Ser. 41 (1977) 121
- 1977GU14 M.G. Gulyamov, B.S. Mazitov, G.A. Radyuk, E.I. Rasulov and R. Yarmukhamedov, Izv. Akad. Nauk SSSR Ser. Fiz. 41 (1977) 2214; Bull. Acad. Sci. USSR Phys. Ser. 41 (1977) 189
- 1977HA1M W.C. Haxton and T.W. Donnelly, Phys. Lett. B66 (1977) 123
- 1977HE10 K.T. Hecht, Nucl. Phys. A283 (1977) 223
- 1977HE1C Hecht and Braunschweig, Bull. Amer. Phys. Soc. 22 (1977) 1012
- 1977HO1D Hocquenghem and Lumbroso, in Tokyo (1977) 145
- 1977HO1E Horiuchi, Invited Talk at Int. Symp. Nucl. Collisions and Their Microscopic Description, Bled, Yugoslavia (1977)
- 1977HU1A H. Husken, W. Wedekind and A. Weiguny, Nucl. Phys. A286 (1977) 163
- 1977IK1A Ikeda and Tamagaki, Prog. Theor. Phys. Suppl. 62 (1977) 1
- 1977IO1A Iori et al., in Tokyo (1977) 519
- 1977IR01 M. Irshad, C. Rioux, J. Asai, R. Pigeon and R.J. Slobodrian, Nucl. Phys. A286 (1977) 483
- 1977JE01 H. Jeremie and G. Kennedy, Can. J. Phys. 55 (1977) 261
- 1977JU1B Juric and Drndarevic, Nucl. Instrum. Meth. Phys. Res. 147 (1977) 251
- 1977KA1J Kato et al., in Tokyo (1977) 569
- 1977KA1K Kahn et al., J. Phys. (London) G3 (1977) L209
- 1977KI05 C.H. King, S.M. Austin, W.S. Chien and H.H. Rossner, Phys. Lett. B67 (1977) 389
- 1977KI10 V.S. Kinchakov, Yad. Fiz. 26 (1977) 261; Sov. J. Nucl. Phys. 26 (1977) 136
- 1977KI12 C.H. King, S.M. Austin, H.H. Rossner and W.S. Chien, Phys. Rev. C16 (1977) 1712
- 1977KN1B Knox, White and Lane, Bull. Amer. Phys. Soc. 22 (1977) 631
- 1977KO1J Kozlovsky and Ramaty, Astrophys. Lett. 19 (1977) 19
- 1977KO1K Kolesnikov, Krokhin, Kopylov and Chernov, Izv. Vuz Fiz. 6 (1977) 75
- 1977KO27 W. Kohler, G. Gruber, A. Steinhauser and K. Bethge, Nucl. Phys. A290 (1977) 233
- 1977KU1E K. Kubodera, J. Delorme and M. Rho, Phys. Rev. Lett. 38 (1977) 321
- 1977KU20 K. Kubodera and A. Arima, Prog. Theor. Phys. 57 (1977) 1599
- 1977LA13 M. Lattuada, F. Riggi, S. Barbarino, C. Spitaleri and D. Vinciguerra, Lett. Nuovo Cim. 20 (1977) 193

- 1977LA1D Lassila and Lary, Proc., Zurich-Sin (1977) 180
- 1977LO10 M.A. Lone, C.B. Bigham, J.S. Fraser, H.R. Schneider, T.K. Alexander, A.J. Ferguson and A.B. McDonald, Nucl. Instrum. Meth. 143 (1977) 331
- 1977LO1D Long and Bromley, Bull. Amer. Phys. Soc. 22 (1977) 541
- 1977LO1E Long and Bromley, in Tokyo (1977) 782
- 1977LO1F Long, Shay, Peschel and Bromley, in Tokyo (1977) 144
- 1977MA1J Malecki, Gluski and Picchi, Proc., Zurich-Sin (1977) 178
- 1977MA1K Mandal et al., Z. Naturf. A32 (1977) 17
- 1977MC07 R. McKeown and G.T. Harvey, Phys. Rev. C16 (1977) 482
- 1977MC1C McNally, Bull. Amer. Phys. Soc. 22 (1977) 581
- 1977ME1C Meadows, ANL/NDM-25 (1977)
- 1977MI13 D. Miljanic, E. Andrade, J. Hudomalj-Gabitzsch, G.S. Mutchler and G.C. Phillips, Nucl. Phys. A290 (1977) 27
- 1977MO1F Mougey, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 420
- 1977NO10 U. Nocken and J. Krug, Nucl. Phys. A292 (1977) 43
- 1977OH1A Ohsawa et al., in Tokyo (1977) 165
- 1977OK1A Oka and Kubodera, Proc., Zurich-Sin (1977) 329
- 1977OO01 M.A. Oothoudt and G.T. Garvey, Nucl. Phys. A284 (1977) 41
- 1977PA26 P. Paul, M. Suffert and P. Gorodetzky, Phys. Lett. B71 (1977) 71
- 1977PR05 P.B. Price, J. Stevenson and K. Frankel, Phys. Rev. Lett. 39 (1977) 177
- 1977RA1B Raisbeck, Yiou and Quechon, Proc., Zurich-Sin (1977) 182
- 1977RI07 P.J. Riley, C.W. Bjork, C.R. Newsom, M.L. Evans, G. Glass, J.C. Hiebert, M. Jain, R.A. Kenefick, L.C. Northcliffe, B.E. Bonner et al., Phys. Lett. B68 (1977) 217
- 1977RI09 R. Risler, W. Gruebler, A.A. Debenham, V. Konig, P.A. Schmelzbach and D.O. Boerma, Nucl. Phys. A286 (1977) 115
- 1977RI10 R. Risler, W. Gruebler, V. Konig and P.A. Schmelzbach, Nucl. Phys. A286 (1977) 131
- 1977RO02 P.G. Roos, N.S. Chant, A.A. Cowley, D.A. Goldberg, H.D. Holmgren and R. Woody, III, Phys. Rev. C15 (1977) 69
- 1977RO18 J. Roth, W. Bretfeld, W. Burgmer, H. Eichner, D. Gola, C. Heinrich, H.J. Helten, H. Kretzer, K. Prescher and W. Schnorrenberg, Z. Phys. A283 (1977) 345
- 1977SA1C Saito, Prog. Theor. Phys. Suppl. 62 (1977) 11
- 1977SA23 E.L. Saw and C.T. Yap, Nuovo Cim. A39 (1977) 643
- 1977SC1D Schramm and Wagoner, Ann. Rev. Nucl. Sci. 27 (1977) 37

- 1977SC37 S.D. Schery, L.E. Young, R.R. Doering, S.M. Austin and R.K. Bhowmik, Nucl. Instrum. Meth. 147 (1977) 399
- 1977SE09 F. Seiler, Nucl. Phys. A286 (1977) 1
- 1977SE1C F. Seiler and H.W. Roser, Phys. Lett. B69 (1977) 411
- 1977SE1D Seya et al., in Tokyo (1977) 149
- 1977SL1B Slobodrian, Pigeon, Roy and Irshad, in Tokyo (1977) 573
- 1977SM1A Yu.F. Smirnov and Yu.M. Tchuvil'sky, Phys. Rev. C15 (1977) 84
- 1977ST18 H. Stowe and W. Zahn, Nucl. Phys. A286 (1977) 89
- 1977ST1G Steiner, High Energy Phys. Nucl. Struct., Zurich, 1977 (1977) 261
- 1977ST1J J. Stevenson, P.B. Price and K. Frankel, Phys. Rev. Lett. 38 (1977) 1125
- 1977ST24 H. Stowe and W. Zahn, Nucl. Phys. A289 (1977) 317
- 1977SZ05 J. Szabo, Z.T. Body, S. Szegedi and M. Varnagy, Nucl. Phys. A289 (1977) 526
- 1977TA1D Takimoto et al., in Tokyo (1977) 567
- 1977TE1A Terrien, in Tokyo (1977) 843
- 1977TE1B Telegdi, High Energy Phys. Nucl. Struct., Zurich, 1977 (1977) 367
- 1977TH09 D.R. Thompson, M. LeMere and Y.C. Tang, Nucl. Phys. A286 (1977) 53
- 1977TR07 R.E. Tribble, J.D. Cossairt, D.P. May and R.A. Kenefick, Phys. Rev. C16 (1977) 1835
- 1977TU02 A. Turkevich, J.R. Cadieux, J. Warren, T. Economou, J. La Rosa and H.R. Heydegger, Phys. Rev. Lett. 38 (1977) 1129
- 1977TU03 A. Turkevich, J.R. Cadieux, J. Warren, T. Economou and J. LaRosa, Phys. Lett. B72 (1977) 163
- 1977TU1D Turk and Antolkovic, Fiz. Suppl. 9 (1977) 49
- 1977UH1A Uhlhorn, Lange and Gonsoir, in Tokyo (1977) 610
- 1977UL02 J. Ulbricht, W. Arnold, H. Berg, E. Huttel, H.H. Krause and G. Clausnitzer, Nucl. Phys. A287 (1977) 220
- 1977WA04 R. Wada, J. Schimizu and K. Takimoto, Phys. Rev. Lett. 38 (1977) 1341
- 1977WA05 C.N. Waddell, E.M. Diener, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, A.G. Pieper, R.B. Theus, C.C. Chang and N.S. Chant, Nucl. Phys. A281 (1977) 418
- 1977WE1B R.B. Weisenmiller, Thesis, Univ. of California, Berkeley (1976); LBL-5077 (1977)
- 1977WI02 D.H. Wilkinson, Phys. Lett. B66 (1977) 105
- 1977WI05 C. Wiezorek, H. Krawinkel, R. Santo and L. Wallek, Z. Phys. A282 (1977) 121
- 1977YA1A Yasue et al., in Tokyo (1977) 520
- 1977YA1B Yakovlev, Fiz. Elem. Chastits At. Yad. (USSR) 8 (1977) 255

- 1977ZA04 W. Zahn, J. Phys. (London) G3 (1977) L209
- 1977ZI01 P.G. Zint, Z. Phys. A281 (1977) 373
- 1978AJ02 F. Ajzenberg-Selove, E.R. Flynn and O. Hansen, Phys. Rev. C17 (1978) 1283.
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- 1978AR10 N. Arena, S. Cavallaro, R. Potenza, V. D'Amico, G. Fazio, S. Femino, G. Giardina, S. Jannelli and M. Lombardi, Lett. Nuovo Cim. 23 (1978) 1
- 1978AR1M Arnould, Proc., Cluster, Winnipeg (1978) B8
- 1978BA31 F.C. Barker, Aust. J. Phys. 31 (1978) 27
- 1978BA66 F.C. Barker and N. Ferdous, Aust. J. Phys. 31 (1978) 239
- 1978BI15 B. Bilwes, R. Bilwes, J.L. Ferrero and A. Garcia, J. Phys. (Paris) 39 (1978) 805
- 1978BI1E Biersack, Fink, Henkelmann and Muller, Nucl. Instrum. Meth. 149 (1978) 93
- 1978BO30 T.J. Bowles and G.T. Garvey, Phys. Rev. C18 (1978) 1447; Erratum Phys. Rev. C26 (1982) 2336
- 1978BR1E Brown, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, Winnipeg (1978)
- 1978BU04 W. Buck, T. Rohwer, G. Staudt, A. Ziemke and F. Vogler, Nucl. Phys. A297 (1978) 231
- 1978BU1D Buck et al., Proc., Cluster, Winnipeg (1978) F8
- 1978CH02 H.C. Cheung, M.D. High and B. Cujec, Nucl. Phys. A296 (1978) 333
- 1978CH07 J.-C. Chou, C.-M. Fou, C.-S. Lin, P.-S. Song, M. Wen and Y.-C. Liu, J. Phys. Soc. Jpn. 44 (1978) 1
- 1978CH1G Chant, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, Winnipeg (1978)
- 1978CL07 R.G. Clark, H. Horn, P.S. Ray and E.W. Titterton, Phys. Rev. C18 (1978) 1127
- 1978DE1J Devaux et al., Proc., Cluster, Winnipeg (1978) D4
- 1978DE37 A.S. Deineko, Y.G. Mashkarov and V.E. Storizhko, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 144; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 123
- 1978DI04 S. Di Liberto, F. Meddi, G. Romano, G. Rosa and C. Sgarbi, Nucl. Phys. A296 (1978) 519
- 1978EN06 P.M. Endt and C. van der Leun, Nucl. Phys. A310 (1978) 96
- 1978FI1E Fick, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. Nucl. Reactions, Winnipeg (1978)
- 1978FO1G Fonseca, Revai and Matveenko, Proc., Cluster, Winnipeg (1978) B1
- 1978FO1H Fou, Bull. Amer. Phys. Soc. 23 (1978) 500

- 1978FR12 S. Frankel, W. Frati, M. Gazzaly, G.W. Hoffmann, O. Van Dyck and R.M. Woloshyn, Phys. Rev. Lett. 41 (1978) 148
- 1978FR1F V. Franco and W.T. Nutt, Phys. Rev. C17 (1978) 1347
- 1978FR1H V. Franco and G.K. Varma, Phys. Rev. C18 (1978) 349
- 1978FR20 A.D. Frawley, J.F. Mateja, A. Roy and N.R. Fletcher, Phys. Rev. Lett. 41 (1978) 846
- 1978FU03 K. Fukunaga, T. Ohsawa, N. Fujiwara, S. Tanaka, A. Okihana and T. Yanabu, J. Phys. Soc. Jpn. 44 (1978) 1413
- 1978GA1C Gal, Soper and Dalitz, Ann. Phys. 113 (1978) 79
- 1978GO1F J. Golden, R.A. Mahaffey, J.A. Pasour, F.C. Young and C.A. Kapetanacos, Rev. Sci. Instrum. 49 (1978) 1384
- 1978GR07 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.V. Ostashko, V.M. Pugach, V.A. Pilipchenko, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 147; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 126
- 1978HA1K Hall, Clark, Freedman and Hanna, Bull. Amer. Phys. Soc. 23 (1978) 953
- 1978HI04 F. Hinterberger, P.D. Eversheim, P. von Rossen, B. Schuller, R. Schonhagen, M. Theene, R. Gorgen, T. Braml and H.J. Hartmann, Nucl. Phys. A299 (1978) 397
- 1978JE1B A.S. Jensen and S.E. Koonin, Phys. Lett. B73 (1978) 243
- 1978KA1G Kato and Bando, Proc., Cluster, Winnipeg (1978) B14
- 1978KH03 A.M. Khan, Prog. Theor. Phys. 60 (1978) 220
- 1978KO1H Kohno, Nagata and Tamagaki, Prog. Theor. Phys. 59 (1978) 1230
- 1978LE10 P. Leleux, M. Bosman, P. Lipnik, P. Macq, J.P. Meulders, C. Pirart and G. Valenduc, Nucl. Instrum. Meth. 152 (1978) 495
- 1978LE13 M. LeMere and Y.C. Tang, Phys. Rev. C18 (1978) 1114
- 1978LI1C Lisowski et al., Bull. Amer. Phys. Soc. 23 (1978) 635
- 1978MA20 A. Malecki and L. Satta, Lett. Nuovo Cim. 21 (1978) 457
- 1978MC04 A.B. McDonald and E.G. Adelberger, Phys. Rev. Lett. 40 (1978) 1692
- 1978MC1E McKeown and Garvey, Bull. Amer. Phys. Soc. 23 (1978) 500
- 1978MO01 C.L. Morris, R.L. Boudrie, J.J. Kraushaar, R.J. Peterson, R.A. Ristinen, G.R. Smith, J.E. Bolger, W.J. Braithwaite, C.F. Moore and L.E. Smith, Phys. Rev. C17 (1978) 227
- 1978MO1J Moller, Hufschmidt and Pfeiffer, Nucl. Instrum. Meth. Phys. Soc. 149 (1978) 73
- 1978NA07 R. Nayak and L. Satpathy, Nucl. Phys. A304 (1978) 64
- 1978NO1B Y. Nogami and C.S. Warke, Phys. Rev. C17 (1978) 1905
- 1978PR1A Pronko, Okamoto and Wiedersich, Nucl. Instrum. Meth. Phys. Res. 149 (1978) 77

1978PR1B Pretorius and Peisach, Nucl. Instrum. Meth. 149 (1978) 69
 1978QA01 S.M. Qaim and R. Wolffe, Nucl. Phys. A295 (1978) 150
 1978RO01 R.G.H. Robertson, E. Kashy, W. Benenson and A. Ledebuhr, Phys. Rev. C17 (1978) 4
 1978RO17 D. Robson, Nucl. Phys. A308 (1978) 381
 1978RO1G Roos et al., Bull. Amer. Phys. Soc. 23 (1978) 97
 1978SA1E Satpathy and Nayak, J. Phys. (London) G4 (1978) L161
 1978SA24 K.R. Sandhya Devi and M.R. Strayer, Phys. Lett. B77 (1978) 135
 1978SA2B K.R. Sandhya Devi and M.R. Strayer, J. Phys. (London) G4 (1978) L97
 1978SE01 F. Seiler, F.N. Rad, H.E. Conzett and R. Roy, Nucl. Phys. A296 (1978) 205
 1978SE1C Seylar and Arnould, Bull. Amer. Phys. Soc. 23 (1978) 928
 1978SH04 S. Shlomo and G.J. Wagner, Z. Phys. A285 (1978) 283
 1978ST10 H. Stowe and W. Zahn, Z. Phys. A286 (1978) 173
 1978ST20 H. Stowe and W. Zahn, J. Phys. (London) G4 (1978) 1423
 1978TA04 S. Tanaka, J. Phys. Soc. Jpn. 44 (1978) 1405
 1978TA1F Takimoto et al., Proc., Cluster, Winnipeg (1978) F16
 1978TO09 T. Tomoda and K. Kubodera, Phys. Lett. B76 (1978) 259
 1978VA03 G.K. Varma, Nucl. Phys. A294 (1978) 465
 1978VA1D Varnagy, Szabo and Szegedi, Nucl. Instrum. Meth. Phys. Res. 154 (1978) 557
 1978VE1B Verma, Indian J. Pure Appl. Phys. 16 (1978) 446
 1978YA1A Yamamoto, Bando and Nagata, Prog. Theor. Phys. 59 (1978) 817
 1978ZA1D P.C. Zalm, J.F.A. van Hienen and P.W.M. Glaudemans, Z. Phys. A287 (1978) 255
 1978ZI1A Ziegler et al., Nucl. Instrum. Meth. Phys. Res. 149 (1978) 19
 1978ZO1A Zolnowski et al., Bull. Amer. Phys. Soc. 23 (1978) 942
 1979WA02 R.E. Warner, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Phys. Rev. C19 (1979) 293
 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, Nucl. Phys. A336 (1980) 1
 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1
 GA77F Unknown Source

