

Energy Levels of Light Nuclei

$A = 14$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 13\text{--}15$ was published in *Nuclear Physics A152* (1970), p. 1. This version of $A = 14$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1969)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 14$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [¹⁴B](#), [¹⁴C](#), [¹⁴N](#), [¹⁴O](#)

B. Tables of Recommended Level Energies:

[Table 14.1](#): Energy levels of ¹⁴C

[Table 14.7](#): Energy levels of ¹⁴N

[Table 14.26](#): Energy levels of ¹⁴O

C. [References](#)

D. Figures: [¹⁴C](#), [¹⁴N](#), [¹⁴O](#), [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{14}B

^{14}B has been identified in the 5.3 GeV proton bombardment of uranium ([1966PO09](#)) and in the 3 GeV proton bombardment of gold ([1968TH04](#)). ^{14}B is particle stable ([1966PO09](#)). See also ([1966GA25](#), [1969AR13](#)).

^{14}C (Fig. 5)

GENERAL:

See ([1954JA1A](#), [1956EL1B](#), [1957VI1A](#), [1958BA1A](#), [1959OT1A](#), [1959SK1A](#), [1960TA1C](#), [1960WA12](#), [1961BA1F](#), [1961FR1C](#), [1962TA1E](#), [1963BL1B](#), [1963NA04](#), [1963SO04](#), [1963VL1A](#), [1964LI1B](#), [1964LO1B](#), [1965BA1X](#), [1965KO1D](#), [1965WA1J](#), [1965ZA1B](#), [1966BA42](#), [1966BO1R](#), [1966GU08](#), [1966MI1G](#), [1966ZA03](#), [1967GR1D](#), [1967HA10](#), [1967IN1B](#), [1967KO1N](#), [1967KO1F](#), [1968EI1C](#), [1968FA1B](#), [1968FR03](#), [1968NE1C](#), [1968RO1C](#), [1969AR13](#), [1969AT1A](#), [1969FR1E](#), [1969SH1A](#), [1969SO08](#), [1969SO1E](#)).

$$1. \ ^{14}\text{C}(\beta^-)^{14}\text{N} \quad Q_m = 0.156$$

Recent values are 5745 ± 50 y ([1961MA1L](#), [1961MA32](#), [1964HU09](#)), 5780 ± 65 y ([1961WA16](#)), 5680 ± 40 y ([1962OL04](#)), 5660 ± 30 y ([1968BE47](#)), leading to a weighted mean of 5692 ± 21 y. See also ([1962GO33](#)). Using Q_m , $\log ft = 9.03$ ([1966BA1A](#)), 9.02 ([1969KA1B](#)). The spectrum does not deviate from the allowed shape down to 3 keV ([1954MO84](#)). The β -decay rate of ^{14}C and the radiative width of $^{14}\text{N}^*(2.31)$ cannot simultaneously be explained by a conventional central force plus spin-orbit interaction together with configuration mixing: the nuclear force must include a tensor part ([1968RO1C](#)). For other discussions of the long lifetime of ^{14}C , see ([1959AJ76](#)) and ([1959JA1B](#), [1962AL1F](#), [1962VA1F](#), [1962WE1B](#), [1964KU1F](#), [1966ZA03](#), [1967BL24](#), [1968FR03](#), [1969DE16](#), [1969EL1B](#), [1969FR1E](#)).

2. (a) $^7\text{Li}(^7\text{Li}, 2n)^{12}\text{C}$	$Q_m = 13.672$	$E_b = 26.795$
(b) $^7\text{Li}(^7\text{Li}, n)^{13}\text{C}$	$Q_m = 18.619$	
(c) $^7\text{Li}(^7\text{Li}, p)^{13}\text{B}$	$Q_m = 5.964$	
(d) $^7\text{Li}(^7\text{Li}, d)^{12}\text{B}$	$Q_m = 3.309$	
(e) $^7\text{Li}(^7\text{Li}, t)^{11}\text{B}$	$Q_m = 6.197$	
(f) $^7\text{Li}(^7\text{Li}, \alpha)^{10}\text{Be}$	$Q_m = 14.783$	
(g) $^7\text{Li}(^7\text{Li}, ^8\text{Be})^6\text{He}$	$Q_m = 7.272$	
(h) $^7\text{Li}(^7\text{Li}, ^7\text{Li})^7\text{Li}$		

Table 14.1: Energy levels of ^{14}C

E_x in ^{14}C (MeV \pm keV)	$J^\pi; T$	τ or Γ (keV)	Decay	Reactions
g.s.	$0^+; 1$	$\tau_{1/2} = 5692 \pm 21$ y	β^-	1, 3, 4, 6, 7, 8, 9, 10, 11, 14, 15, 17, 18, 19, 20, 21, 22, 23, 27, 28, 29, 30, 31
6.0932 ± 1.3	1^-	$\tau_m < 0.3$ psec	γ	3, 4, 7, 9, 11, 14, 18, 20
6.5892 ± 1.4	0^+	> 0.6 psec	γ	4, 7, 9, 14, 18
6.7281 ± 1.4	3^-	97 ± 15 psec	γ	4, 7, 9, 14, 17, 18, 20
6.9012 ± 1.5	0^-	< 0.3 psec	γ	4, 9, 14, 17
7.0117 ± 5.2	2^+	< 0.12 psec	γ	4, 7, 9, 18
7.3414 ± 3.4	2^-	< 0.22 psec	γ	4, 7, 9, 14, 18
8.318 ± 5	$(1, 2)^+$	$\Gamma = 12$	n	9, 12, 14, 18
9.801 ± 8	(1)	19	n	12, 14, 18
10.433 ± 10	(2)	16	n	12, 14, 18
10.453 ± 10	≥ 1	9	n	12, 14, 18
10.74 \pm 20		< 15		7, 9
(11.35 \pm 50)				7
(11.66 \pm 50)				7
11.9 ± 300		950 ± 300		12, 14
12.601 ± 20		110 ± 20		14
12.854 ± 20				14
12.958 ± 20				14

These reactions have been studied with $E(^7\text{Li})$ to 6.5 MeV: see ([1957NO14](#), [1957NO17](#), [1962BE24](#), [1963CA09](#), [1963HU02](#), [1964DZ1A](#), [1966PI02](#), [1969CA1A](#)). For $E(^7\text{Li}) = 2.3$ to 5.8 MeV, the cross section for emission of α_0 , α_1 , α_{2+3+4} (to the first five states of ^{10}Be) increases monotonically with energy ([1964DZ1A](#), [1969CA1A](#)). The measured cross sections for reactions (c), (d), (e) and (f) indicate that the reactions probably occur by a mixture of compound nucleus and direct reaction mechanisms. The Coulomb field appears to have a marked effect on the cross sections ([1969CA1A](#): see also ([1963HU02](#))).



See ([1967SE08](#)) and ${}^{15}\text{N}$.

Table 14.2: ^{14}C levels from $^{12}\text{C}(\text{t}, \text{p})^{14}\text{C}$

E_x^a (MeV \pm keV)	(1964MI05) ^b L	(1960JA17) ^c L
0	0	0
6.090 ± 10	(2)	1
6.582 ± 10	1	1
6.725 ± 10	(2)	3
6.893 ± 10	e	e
7.009 ± 10	(2)	0
7.335 ± 10	e	e
8.32	2 ^f	
10.74 ± 20^d		

^a The excitation energies of the first six excited states have been determined by (1960JA17); the seventh comes from the $^{13}\text{C}(\text{n}, \text{n})^{13}\text{C}$ work of (1961CO05); the eighth has been determined by (1964MI05).

^b $E_{\text{t}} = 11$ MeV; except ground state $E_{\text{t}} = 8$ to 13 MeV.

^c $E_{\text{t}} = 5.5$ MeV.

^d $\Gamma < 15$ keV. No states are observed between this level and the broad state at 11.9 MeV (1964MI05).

^e Weak group.

^f (1964MI05) suggest $J^\pi = 2^+$; see, however, (1961CO05).

$$4. \ ^9\text{Be}(\text{d}, \text{d})^{14}\text{C} \quad Q_m = 10.102$$

At $E(^7\text{Li}) = 3.2$ MeV, the ground state deuteron group and the deuteron groups corresponding to the known levels with $E_x < 9$ MeV have been observed (1964CA05)[†]. At $E(^7\text{Li}) = 5.1$ MeV, the γ -decay of the six bound excited states has been studied by (1966CA07); see Table 14.3. Measurements at $E(^7\text{Li}) = 5.7$ MeV give $\tau_m < 0.32$ and < 0.12 psec, respectively, for $^{14}\text{C}^*(6.09, 7.01)$. $E_\gamma = 6.0945 \pm 0.0032$, 6.7281 ± 0.0014 and 7.0117 ± 0.0052 MeV for the ground state transitions for these two levels and $^{14}\text{C}^*(6.73)$ (1969TH01).

$$5. (\text{a}) \ ^{11}\text{B}(\text{t}, \text{n})^{13}\text{C} \quad Q_m = 12.422 \quad E_b = 20.598$$

[†] Angular distributions of the deuterons to $^{14}\text{C}^*(0, 6.09, 6.59 + 6.73, 6.90 + 7.01, 7.34, 8.32)$ have been measured at $E(^7\text{Li}) = 5.6, 5.8, 6.0$ and 6.2 MeV (1969SN02).

Table 14.3: Branching ratios of γ -rays in ^{14}C

E_i (MeV)	J_i^π	E_f (MeV)	Branch (%)			
			A	B	C	D
6.09	1^-	0			100	
6.59	0^+	0	1.0 ± 0.4 ^c			
		6.09	99.0 ± 0.4			
6.73	3^-	0	93 ± 2	97.3 ± 1	91 ± 5	
		6.09	7 ± 2	2.7 ± 1	9 ± 3	
6.90	0^-	6.09	100 ^a			
7.01	2^+	0		98.6 ± 0.7	100_{-5}^{+0}	
		6.09		1.4 ± 0.7	< 5	
7.34	2^-	0	18 ± 4	14 ± 4	13 ± 3	13 ± 5
		6.09	47 ± 4	52 ± 5 ^b	60 ± 5	29 ± 13
		6.73	35 ± 7	34 ± 4 ^b	27 ± 5	58 ± 7

A: (1966AL10): $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$.

B: (1968BE30): $^{12}\text{C}(\text{t}, \text{p})^{14}\text{C}$.

C: (1966CA07): $^9\text{Be}(^7\text{Li}, \text{d})^{14}\text{C}$.

D: (1965LA09): $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$ [see also (1966AL10)].

^a (1958WA02).

^b $\delta(M2/E1) = -0.04 \pm 0.09$ and $+0.07 \pm 0.30$, respectively (1968BE30).

^c Internal pairs.



Resonant structure has been observed in the yield of neutrons at $E_t = 1.200, 1.340, 1.567, 1.650, 1.700, 1.800, 1.940, 2.055, 2.245$ and 2.315 MeV corresponding to $E_x = 21.540, 21.650, 21.828, 21.893, 21.933, 22.011, 22.121, 22.211, 22.360$ and 22.415 MeV (1965VA13; natural B target). For reaction (b) see (1967SI1F).



The angular distributions of ground state protons have been measured for $E_\alpha = 2.5$ to 5.0 MeV (1963MA28). See also (1959AJ76) and ^{15}N .



At $E(\text{Li}) = 5$ MeV, α -particle groups are observed to the known states of ^{14}C with $E_x < 10$ MeV except $^{14}\text{C}^*(6.90)$, and to the (unresolved) 10.4 MeV states. There is some indication also of ^{14}C states at (10.71), 11.35, 11.66, (14.15), (14.73) and (15.07) MeV ($\pm \approx 50$ keV), in addition to the 12 MeV states. The wide state at $E_x = 11.9$ MeV is not observed. Angular distributions have been obtained for the α -particles to the ground state of ^{14}C and to the states at $E_x = 6.09$ and 8.32 MeV (1966MC05). See also (1963MI02, 1963MO1B).



See (1963HO1E).




Observed proton groups are displayed in Table 14.2. Angular distributions have been observed at $E_t = 5.5$ MeV (1960JA17) and $E_t = 8$ to 13 MeV (1964MI05). Aside from the ground state and groups corresponding to $^{14}\text{C}^*(6.59, 8.32)$, the stripping patterns are inconclusive (1964MI05). Particle- γ correlations have been studied by (1968BE30): see Table 14.3. The lifetime of $^{14}\text{C}^*(6.73)$ is 97 ± 15 psec (1968AL12). The $(6.73 \rightarrow 0)$ E3 transition is enhanced by 3.3 ± 0.8 W.u. (1968AL12).

See also (1960MU07, 1962GU01, 1962KU09, 1967KE1J, 1969ET01), (1964AB1B, 1965GL07, 1965SH1E, 1966GL1C, 1966SH1F, 1967TI1B, 1969SO08; theor.), (1959AJ76) and ^{15}N .



See (1968HU1H, 1968SC1H, 1969BR1D, 1969SU1E).



The thermal capture cross section is 0.9 ± 0.2 mb (1964ST25), 1.0 ± 0.2 mb (1963MO1C). Two γ -rays are observed with $E_\gamma = 8.174 \pm 0.002$ and 6.093 ± 0.002 MeV [$E_x = 6.094 \pm 0.002$ MeV], with intensities of 87 ± 5 and $13 \pm 1\%$. Intensities of transitions via other ^{14}C states are $< 2\%$ (1967TH05). See also (1968FO1A).

Table 14.4: Resonances in $^{13}\text{C}(\text{n}, \text{n})^{13}\text{C}$ ([1961CO05](#))

E_{res} (MeV \pm keV)	E_x (MeV \pm keV)	Γ_{lab} (keV)	σ^{a} (b)	J^π
0.153 ± 5	8.318 ± 5	13	7	1^+
1.751 ± 8	9.801 ± 8	20	[≈ 1.3]	1
2.432 ± 10	10.433 ± 10	17	[≈ 1.3]	2
2.454 ± 10	10.453 ± 10	10	[≈ 0.7]	≥ 1
3.8 ^b	11.7			

^a Corrected peak cross section, above background.

^b Broad resonance structure.

12. $^{13}\text{C}(\text{n}, \text{n})^{13}\text{C}$

$$E_b = 8.176$$

The coherent scattering length (thermal, bound) is 6.0 fm ([1961WI1A](#), [1969BA1P](#)). The total cross section has been measured for $E_n = 0.11$ to 9 MeV and 16 to 23 MeV. The observed resonances are listed in Table 14.4. For the 153 keV resonance, the shape excludes $l = 0$; θ^2 for $l = 2$ would be 2.3, thus $l = 1$. The peak cross section is too large for $J = 0$, but lower than expected for $J = 1$: it is concluded that $J^\pi = 1^+$. The peak cross section for the $E_n = 1.75$ MeV level is in excellent agreement with $J = 1$. Formation by $l = 0$ is excluded by the shape, but $l = 1, 2$ remain as possibilities. Peak cross sections for the states at $E_n = 2.43$ and 2.45 MeV indicate $J = 2$ and $J \geq 1$, respectively ([1961CO05](#)). See also ([1969HO1Y](#)).

13. $^{13}\text{C}(\text{n}, \alpha)^{10}\text{Be}$

$$Q_m = -3.836$$

$$E_b = 8.176$$

See ([1947HU03](#), [1954SA68](#), [1964GA1A](#)).

14. $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

$$Q_m = 5.952$$

The weighted mean of reported Q -values is 5.948 ± 0.003 MeV: see ([1965RY01](#)).

Proton groups reported by ([1954SP01](#), [1955MC75](#), [1961JA23](#)) are displayed in Table 14.5. See also ([1961TE02](#)). Angular distributions have been analyzed by PWBA and DWBA and have led to J^π assignments and to determinations of θ^2 : see Table 14.5 ([1955MC75](#), [1958WA02](#), [1963LI09](#), [1966GL01](#), [1967SC29](#)). See also ([1963DE19](#), [1965LA09](#)) and ^{15}N .

Observed γ -radiation assigned to ^{14}C is exhibited in Table 14.6 ([1955BE62](#), [1955MA36](#), [1958RA13](#), [1958WA02](#), [1965LA09](#), [1966AL10](#)) and in Table 14.3. The internally formed positron distribution shows that the decay of the 6.09 MeV level is E1 and hence that it has $J^\pi = 1^-$; it is presumably

Table 14.5: Proton groups from $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

(1954SP01)	(1955MC75)			(1958WA02, 1959WA04)		(1966GL01)		(1960MA32)
E_x (MeV \pm keV)	E_x (MeV \pm keV)	l_n ^c	J^π	J^π	θ_n^2 ^g	θ_n^2 ^{g,h}	θ_n^2 ^{g,i}	θ_n^2 ^{g,k}
0	0	1	$0^+, 1^+, 2^+$	0^+	0.10	0.067	0.031	0.063
6.091 ± 10 ^a	6.09	0	$0^-, 1^-$	1^-	0.40 ^j	0.20	0.13	0.20
	6.589 ± 20	$1, 2, 3$ ^f	$(1^-, 2, 3^-)$		$\lesssim 0.01$	0.008		0.006
6.723 ± 10 ^b	6.72	2	$1^-, 2^-, 3^-$	$3^-(2^-)$	0.11	0.065	0.052	0.07
6.894 ± 10 ^b	6.89	$0, 1$ ^f	$0, 1, 2^+$	0^-	0.39	0.24		
	7.346 ± 20	2	$1^-, 2^-, 3^-$	$2^-, 3^-$	0.11	0.072	0.061	0.06
	8.321 ± 20					0.009		0.0015
	9.800 ± 20							
	10.433 ± 20							
	10.505 ± 20							
	11.9 ± 300 ^d							
	12.601 ± 20 ^e							
	12.854 ± 20							
	12.958 ± 20							

^a 6.112 ± 0.012 MeV (1961JA23).^b The spacing of these two levels is 171 ± 3 keV (1954SP01).^c See also (1959AJ76).^d $\Gamma_{\text{lab}} = 1.10 \pm 0.30$ MeV.^e $\Gamma_{\text{lab}} = 0.130 \pm 0.020$ MeV.^f See footnotes 18 and 31 in (1958WA02).^g From PWBA analysis, assuming $J^\pi = 0^+, 1^-, 3^-, 0^-, 2^-$.^h $E_d = 12$ MeV.ⁱ $E_d = 8$ MeV.^j $E_d = 4.7$ and 6 MeV, $\theta^2 = 0.95 \pm 0.24$, DWBA analysis (1963LI09).^k $E_d = 14.8$ MeV.^l For $^{14}\text{C}^*(7.01)$.

Table 14.6: Gamma rays from $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

Transition	E_γ (MeV \pm keV)				
	(1955MA36)	(1955BE62)	(1958RA13)	(1958WA02)	(1966AL10)
6.09 \rightarrow g.s.	6.090 \pm 25 ^b	6.11 \pm 30 ^b	6.090 \pm 20 ^b	6.09	
6.73 \rightarrow g.s.	6.730 \pm 40 ^c	6.720 \pm 30 ^c	6.738 \pm 25 ^c	6.72	
7.34 \rightarrow g.s. ^a		(7.30 \pm 50)	7.323 \pm 25 ^b	7.35	
6.59 \rightarrow 6.09					0.4958 \pm 0.4
6.90 \rightarrow 6.09	0.811 \pm 3 ^d			0.813 \pm 8 ^d	0.8087 \pm 1.0
7.34 \rightarrow 6.73				0.621 \pm 7	
7.34 \rightarrow 6.09					1.248 \pm 3 ^e

^a $\Gamma_{7.34}/\Gamma_{6.72} \leq 0.22 \pm 0.07$, $\Gamma_{1.25}/\Gamma_{6.72} = 0.5 \pm 0.2$ ([1965LA09](#)).

^b Corrected for Doppler shift: see ([1958WA02](#)).

^c No Doppler shift: $\tau > 0.3$ psec ([1958WA02](#)).

^d A Doppler shift of 0.5–1.0% applies ([1958WA02](#)).

^e See also ([1965LA09](#)).

the analog of the 8.06 MeV level in ^{14}N ([1952TH24](#), [1958CH1A](#), [1958GO81](#), [1959CH28](#), [1959WA04](#), [1966WA1C](#)). The p- γ correlation is isotropic, consistent with $l = 0$ formation ([1963LI09](#), [1965LA09](#)). The 6.59 MeV state is observed in internal pairs but not in external pair formation. The transition to the ground state is E0, therefore $J^\pi = 0^+$; $\tau_m > 0.6$ psec ([1963AL21](#), [1964WA05](#), [1966AL10](#)). The mean lifetime of the 6.73 MeV state is greater than 2 psec ([1966AL10](#)) [see also reaction 9] and study of the internal pairs suggests that the ground state transition is E3 ([1964WA05](#), [1966WA1C](#)). The stripping pattern is rather clearly $l = 2$; (p- γ) angular correlations are consistent with $J^\pi = 3^-$: $J = 1$ and 2 are ruled out ([1965LA09](#)): see also ([1966AL10](#)).

The 0.81 MeV cascade transition from the 6.90 MeV state shows a Doppler shift ($\tau_m < 0.3$ psec) and hence is predominantly dipole. The angular correlation of 6.1 and 0.8 MeV γ -rays is consistent with $J = 0$ and excludes $J = 1$ or 2. $J = 0$ is also suggested by the absence of the direct ground state decay of the 6.90 MeV level ([1958WA02](#)). The plane polarization of the 0.81 MeV cascade has been measured in coincidence with the 6.09 MeV ground state transition: the parity of the 6.90 MeV state is negative ([1966RI02](#)).

Protons leading to the 7.34 MeV state exhibit a clear $l = 2$ pattern: therefore $J^\pi = 1^-$, 2^- or 3^- . The level decays via cascades through the 6.09 (1^-) and 6.73 (3^-) states: see Table 14.3 ([1958RA13](#), [1958WA02](#), [1965LA09](#), [1966AL10](#)). The strength of the cascade (7.34 – 6.09) compared to the ground state transition argues against $J^\pi = 1^-$ and 3^- for the 7.34 MeV state ([1958WA02](#), [1966AL10](#)); the angular correlation data is consistent with $J^\pi = 2^-$ and excludes $J = 3$ ([1965LA09](#)). Comparison of reduced widths and calculations of level shifts suggests the following associations of ^{14}C and ^{14}N levels: 6.09 – 8.06, 6.59 – 8.62, 6.73 – 8.91, 6.90 – 8.80,

(7.01 – 9.17), 7.34 – 9.51 and (8.32 – 10.43) ([1960WA12](#)). See also ([1959KU1C](#)).




At $E_t = 5.5$ MeV, the ground state deuterons have been observed ([1961BA10](#)).



Not reported.



See ([1967PO13](#), [1969BR1D](#)).



At $E_d = 14.9$ MeV, inelastic deuteron groups have been seen to the states at 6.09, 6.58, 6.72, 7.01 ± 0.02 , 7.34, 8.32, 9.80 and 10.5 MeV ([1959AR1A](#)). An angular distribution of elastically scattered deuterons has been obtained at $E_d = 3.4$ MeV by ([1967NE06](#)). See also ([1960WA12](#)). For reaction (a), see ([1969CU1D](#)).



See ([1968BA1E](#), [1968CE1C](#), [1969DA1P](#)).



The weighted mean of five Q -value determinations is 626 ± 1 keV ([1957VA11](#)). ([1965IS1A](#)) report 621 ± 6 keV. ([1969NY1A](#)) report γ -rays with $E_\gamma = 6.082 \pm 0.010$ (Doppler corrected) and 6.732 ± 0.005 MeV. τ_m of $^{14}\text{C}^*(6.09) \leq 0.3$ psec ([1969NY1A](#)). See also ([1959GA14](#), [1959HA13](#), [1963MO04](#), [1964MO1D](#), [1967AN08](#), [1969DI1B](#)), ([1959AJ76](#)) and ^{15}N .



See ([1968ST1U](#)).



See ([1964KO10](#)) and ^{15}N .



The angular distributions of ground state neutrons have been determined at $E_n = 14.1$ and 14.8 MeV ([1967FE06](#)). See also ([1956FR1A](#)).



Not reported.



Not reported.



Not reported.



See ([1961TA10](#)).



See ^{18}O and ([1965BE1N](#)).



See ([1963DR1B](#), [1964BL1C](#), [DE65C](#), [1966DA1C](#), [1966DE09](#)).



See ([1968GO1Q](#)) and reaction 10.



See ([1967DE03](#)).



Not observed: see ([1962LA15](#)).

¹⁴N
(Figs. 6 and 8)

GENERAL:

Model calculations: (1957HU1C, 1959BA1F, 1959BR1E, 1959OT1A, 1959SK1A, 1960PA08, 1960TA1C, 1960WA12, 1961BA1F, 1961BA1E, 1961FR1C, 1961TR1B, 1962IN1C, 1962TA1E, 1962WE1C, 1963KU1B, 1963NA04, 1963SE19, 1963TR02, 1963WA15, 1964AM1D, 1964BR1L, 1964FE02, 1964LO1B, 1964MA1G, 1964NE1E, 1964ST1B, 1964UL1A, 1965CO25, 1965GL09, 1966BO1R, 1966HA18, 1966HA31, 1966HE1E, 1966MAZY, 1966MI1G, 1966WI1E, 1967CO32, 1967EV1C, 1967KU1E, 1967LI06, 1967PA05, 1967SO1A, 1968CO13, 1968DE13, 1968EI1C, 1968GO01, 1968HO1H, 1968KU1E, 1968NO1C, 1968RA10, 1968SO1B, 1968ZH05, 1969UL03, 1969VA1C).

General calculations and reviews: (1960PA08, 1962MA1P, 1963BL1B, 1963VL1A, 1964LI1B, 1964MC1C, 1964TH03, 1964YO1B, 1965BE1B, 1965GI1B, 1965KO1D, 1965MA1N, 1966DA1E, 1966MAZY, 1966WI1E, 1966ZA03, 1967BI06, 1968FR03, 1968GA03, 1968HI1H, 1968LA1J, 1968RI1H, 1968RO1C, 1969AT1A, 1969FR1E, 1969WA1F).

Electromagnetic transitions: (1959FA1C, 1959WA16, 1959WA04, 1960WA12, 1962MO1A, 1962UN1A, 1963KU03, 1964HE1F, 1965BI10, 1966HA31, 1966WA1E, 1967KU1E, 1967PO1J, 1967WA1C, 1968RO1C, 1969VA1C).

Meson interactions: (1967BA2H, 1967BA78, 1967BU1D, 1967FO1A, 1967KO1D, 1967MI1B, 1968BA2G, 1968BA1M, 1968BA48, 1968CH1F, 1968GO1T, 1968KO1C, 1968NO1A, 1968RI1H, 1968TA1C, 1968WI1B, 1968ZU1A, 1969CH1C, 1969MO1E, 1969SA1L, 1969WU1A).

Spallation: (1965ZH1A, 1967AU1B).

Ground state: $J = 1; \mu = +0.40361 \text{ nm}$ (1964LI14).

$$Q = 16 \pm 7 \text{ mb} \quad (1960LI10).$$

See also (1959KA07, 1960PA08, 1961BR13, 1962BA1K, 1962HE1A, 1966MA1P, 1967SH05, 1967SH14, 1968PE16, 1968RO1C, 1969AL04, 1969FU11, 1969VA1C, 1969WU1A).

1. (a) ${}^9\text{Be}({}^6\text{Li}, \text{n}){}^{14}\text{N}$	$Q_m = 14.504$
(b) ${}^9\text{Be}({}^7\text{Li}, 2\text{n}){}^{14}\text{N}$	$Q_m = 7.251$

Gamma rays due to the $3.95 \rightarrow 2.31$, $4.91 \rightarrow 0$, $6.21 \rightarrow 2.31$ and $6.44 \rightarrow 0$ transitions have been observed in these reactions and in reactions 6 and 19: $E_\gamma = 1631.3 \pm 1.3$, 4913.8 ± 3.0 , 3883.0 ± 1.9 and $6443.7 \pm 1.8 \text{ keV}$, respectively. $\tau_m = 0.62 \pm 0.08 \text{ psec}$ for ${}^{14}\text{N}^*(6.44)$ [see also Table 14.12] (1969TH01). For reaction (b), see also (1957NO17).

Table 14.7: Energy levels of ^{14}N ^a

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
0	$1^+; 0$	stable	γ	1, 6, 7, 8, 9, 10, 11, 16, 17, 18, 19, 20, 21, 22, 23, 24, 30, 31, 32, 33, 34, 35, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 56, 57, 58, 59, 60, 61, 62, 63
2.31281 ± 0.06	$0^+; 1$	$\tau_m = 0.085 \pm 0.010$	γ	1, 6, 9, 10, 16, 17, 19, 24, 30, 31, 32, 36, 37, 39, 40, 41, 42, 43, 45, 46, 51, 52, 54, 57, 58
3.9447 ± 1.2	$1^+; 0$	0.0045 ± 0.0004	γ	1, 6, 7, 9, 10, 16, 17, 18, 19, 20, 22, 24, 30, 31, 32, 36, 37, 40, 41, 42, 43, 45, 46, 51, 52, 54, 57, 58
4.9134 ± 2.4	$(0, 1)^-; 0$	< 0.050	γ	1, 6, 7, 9, 17, 18, 19, 22, 24, 30, 31, 32, 37, 40, 41, 42, 43, 45, 46, 52, 54, 58
5.10587 ± 0.18	$2^-; 0$	12.4 ± 1.4	γ	6, 7, 16, 17, 18, 19, 22, 24, 30, 31, 32, 37, 40, 41, 42, 43, 45, 46, 52, 54, 58
5.691 ± 3	$1^-; 0$	< 0.036	γ	6, 7, 17, 18, 19, 24, 30, 31, 32, 33, 37, 41, 42, 43, 45, 46, 52, 54, 58
5.833 ± 2	$3^-; 0$	18 ± 2	γ	6, 7, 16, 17, 18, 19, 20, 24, 30, 31, 32, 33, 37, 40, 41, 42, 43, 45, 46, 52, 54, 58
6.1976 ± 2.0	$1^+; 0$		γ	1, 6, 7, 17, 18, 19, 24, 30, 31, 37, 42, 43, 45, 46, 52, 54, 58
6.4436 ± 1.6	$3^+; 0$	0.62 ± 0.06	γ	1, 6, 7, 17, 18, 20, 24, 30, 31, 37, 41, 42, 43, 46, 52, 54, 58
7.028 ± 4	$2^+; 0$	$\tau_m = 5.4 \pm 0.5$ fsec	γ	6, 7, 17, 18, 19, 24, 30, 31, 37, 39, 40, 41, 42, 43, 45, 46, 52, 54, 57, 58
7.966 ± 1	$2^{(-)}; 0$	$\Gamma < 0.37$	γ, p	6, 7, 17, 18, 19, 24, 30, 31, 40, 42, 46, 52, 54, 58

Table 14.7: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
8.061 \pm 2	$1^-; 1$	30 ± 1	γ, p	17, 24, 25, 30, 31, 38, 39, 42, 52, 54
8.489 \pm 3	$(4^-); 0$	≤ 0.2	γ, p	6, 7, 17, 18, 19, 24, 31, 54, 58
8.617 \pm 3	$0^+; 1$	7 ± 1	γ, p	17, 24, 25, 31, 52, 54, 58
8.80 \pm 50	$0^-; 1$	≈ 500	γ, p	24, 25, 31, 33, 54
8.907 \pm 3	$3^-; 1$	16 ± 2	γ, p	24, 25, 30, 31, 33, 40, 52, 54
8.963 \pm 3	$5^+; 0$	< 1	γ, p	17, 18, 19, 20, 24, 31, 52
8.979 \pm 4	$2^+; (0)$	9 ± 2	γ, p	6, 7, 24, 25, 30, 31, 52
9.129 \pm 2	$2^-; 0$	< 1	γ, p	6, 7, 24, 31
9.172 \pm 1.5	$2^+; 1$	0.074 ± 0.008	γ, p	17, 19, 24, 31, 38, 39, 40, 42, 52, 54, 57
9.388 \pm 4	$2^-; 0$	13 ± 3	p	6, 7, 17, 18, 19, 25, 52, 54, 58
9.508 \pm 3	$2^-; 1$	41 ± 2	γ, p	24, 25, 52, 54, 58
9.702 \pm 4	1^+	15 ± 3	p	6, 7, 17, 18, 19, 25, 52, 54, 58
10.063 \pm 15	$T = 0$			17
10.10 \pm 15	$(1, 2)^+$	5	γ, p	6, 7, 17, 18, 19, 24, 25, 52, 58
10.228 \pm 10	$1^{(-)}; 0$	80 ± 15	γ, p	24, 25, 52, 58
10.434 \pm 8	$2^+; 1$	33 ± 3	γ, p	6, 7, 17, 24, 25, 37, 38, 40, 52, 54, 58
10.56 \pm 10	1^-	140	p, d	12, 17, 25
10.809 \pm 12	$4^+; 0$			17, 18, 19, 20, 52, 58
11.04 \pm 32	$1^+; 0$	95	γ, n, p, d	12, 17, 18, 19, 24, 25, 26, 40, 52, 58
11.051 \pm 14		< 30	γ, p	6, 7, 24, 52
11.246 \pm 12	$3^-; 1$	11	n, p, d	12, 17, 19, 25, 26, 42, 43, 45, 52, 58
11.30	$2^-; 0$	180	γ, n, p, d	12, 17, 19, 24, 25, 26, 52, 58
11.374 \pm 12	$1^+; 0$	32	n, p, d	12, 13, 17, 19, 25, 26, 52, 58

Table 14.7: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
11.516 \pm 13	3^+	5	p, d	12, 13, 17, 19, 52, 58
11.66 \pm 40	$(1^+, 2^-)$	≈ 95	n, p, d	12, 17, 52
11.75	1^+	100	γ, n, p, d	12, 17, 24, 25, 26, 52
11.81	(2^+)	≈ 100	n, p	25, 26, 52
11.95 \pm 30	2^+			17, 19, 52
12.23	3^-		n, p, d	12, 26, 52
12.29 \pm 15			p, α	3, 52
12.414 \pm 6	4^-	37 ± 3	n, p, d, α	3, 4, 12, 13, 17, 19
12.47 \pm 10		≈ 20	p, α	3, 24, 37
12.504 \pm 7		31 ± 5	n, p, d, α	3, 12, 17, 24, 37, 52
12.608 \pm 6	3^+	47 ± 3	p, d, α	3, 12, 13, 17, 19, 52
12.689 \pm 4	3^-	17 ± 3	n, p, d, α	2, 3, 4, 12, 13, 17, 19, 26
12.793 \pm 6	4^+	18 ± 3	n, p, d, α	2, 3, 4, 12, 13, 17, 18, 20, 52
12.825 \pm 7	4^-	7 ± 2	n, p, d, α	3, 4, 12, 13, 52
12.853 \pm 15		71 ± 7	n, p, d	12, 17, 26
12.942 \pm 7	4	24 ± 4	p, d, α	3, 4, 12, 13, 17, 18
13.03		190	γ, p	24
13.05 \pm 20	$T = 0$	< 70		18
13.164 \pm 8	$(0, 1)^-$	16 ± 6	n, p, d, α	2, 3, 4, 5, 12, 13, 17, 18, 52
13.23 \pm 10		97 ± 7	n, p, α	2, 3, 26, 52
13.3		≈ 1000	γ, p	24
13.66 \pm 10	$(2, 3)^+$	≈ 110	p, d, α	3, 5, 12, 13
13.71 \pm 10		≈ 100	n, p	2, 3
13.75 \pm 10	$1^+; 1$	210 ± 30	n, p, d	3, 12, 13, 26, 37, 52, 54
14.04			p, d	12, 13
14.17		290	n, p, d, α	3, 12, 13, 26
14.40		≈ 140	p, d, α	3, 12, 13
14.70			d	13, 18
14.84		140	n, p, d, α	2, 3, 12, 13, 15, 17, 26
15.0		≈ 1000	γ, p	24

Table 14.7: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
15.10	6^-	40	n, p, d, α	2, 3, 13, 17, 18, 20, 26
15.26			p, d	12, 13
15.47		≈ 70	n, p, d, α	2, 3, 12, 15, 18
15.7		320	n, p, d, α	12, 13, 17, 26
16.3		140	n, p, d, α	15, 18, 26
17.2 ± 200		≈ 400	γ, n, p, d, α	13, 15, 17, 18, 24, 26
17.7				18
≈ 18	3^-	≈ 2000	d, α	15
18.1		550	n, p	24, 26
19.6	(5^+)		p, d	12, 13, 24
20.5	(5^+)	≈ 1000	γ, n, p, d	12, 13, 24
21.15 ± 150	(4^-)	≈ 1400	d, α	13, 15
≈ 22.5		≈ 3000	γ, n, p	24, 38
22.70 ± 100		≈ 900	d, α	15
23.0		≈ 450	γ, p	24
23.7		≈ 450	γ, p	24
≈ 23.9		≈ 1000	$n, {}^3\text{He}, \alpha$	8
≈ 24		several MeV	γ	39
≈ 30			γ	39

^a See also Tables 14.9 and 14.13.

Table 14.8: Resonances in $^{10}\text{B} + \alpha$


E_α (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particle ^a (x)	Γ_x (keV)	θ_x^2	$^{14}\text{N}^*$ (MeV)	J^π ^b	Refs.
0.95		p ₀			12.29		1953MA42
1.13 ± 5	30 ± 5	p ₀ , p ₁ , p ₂ , p ₃ , d			12.42	4 ⁻	{ 1953MA42, 1953SH64, 1954ST20, 1969GA01
1.20 ± 5	≈ 20	p ₀ , (p ₂), p ₃			12.47		1953MA42, 1969GA01
1.23 ± 5	35 ± 5	p ₀ , p ₃			12.49		1953SH64, 1969GA01
1.40 ± 5	46 ± 4	p ₁ , p ₂ , (p ₃)			12.61	3 ⁺	1953SH64, 1969GA01
1.508 ± 4	17 ± 3	α	1.7	6.0	12.690	3 ⁻	1953SH64, 1953TA06, 1954ST20, 1957BR18, 1969GA01
		p ₀	0.62	0.012			
		p ₁	0.17	0.29			
		p ₂	0.70	0.31			
		p ₃	5.6	0.47			
		d	0.93	0.26			
		n	4.3	0.19			
1.64 ± 5	16 ± 3	α	1.0	8.2	12.79	4 ⁺	1953SH64, 1953TA06, 1954ST20, 1969GA01
		p ₀	0.18	0.012			
		p ₁	0.085	2.7			
		p ₂	0.44	3.0			
		p ₃	9.6	4.9			
		d	2.0	3.9			
		n	0.59	0.16			
1.68 ± 5	5 ± 2	p ₁ , p ₂ , p ₃ , d			12.81	4 ⁻	1953SH64, 1953TA06, 1954ST20, 1969GA01
1.83 ± 5	22 ± 4	p ₀ , p ₁ , p ₂ , p ₃ , d			12.92	4 ⁺	1953SH64, 1953TA06, 1954ST20, 1969GA01
2.16 ± 5	14 ± 4	p ₀ , p ₁ , p ₂ , p ₃ , d, α_1 , n			13.16		1953TA06, 1955SH46, 1969GA01
2.26 ± 5	97 ± 7	p ₀ , p ₁ , p ₂ , p ₃ , n			13.23		1953TA06, 1955SH46, 1969GA01
2.86 ± 5	≈ 110	p ₁ , p ₂ , α_1			13.66		1969GA01
2.94 ± 5	≈ 100	p ₀ , p ₁ , p ₃ , n, α_1			13.71		1953TA06, 1956BO61, 1959GI47, 1969GA01
2.99 ± 5		p ₂			13.75		1969GA01
3.6	290	p			14.2		1956BO61, 1959GI47

Table 14.8: Resonances in $^{10}\text{B} + \alpha$ (continued)


E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particle ^a (x)	Γ_x (keV)	θ_x^2	$^{14}\text{N}^*$ (MeV)	J^π ^b	Refs.
3.94	≈ 140	p			14.43		1956BO61
4.53	140	n, p			14.85		1956BO61, 1959GI47
4.85	≈ 35	n, p			15.08		1956BO61, 1959GI47
5.36	≈ 70	n, p			15.44		1956BO61

^a p_0, p_1, p_2, p_3 correspond to the ground state and the 3.1, 3.7, 3.9 MeV states of ^{13}C , and the corresponding γ -rays.


^b From angular distributions and (p- γ) correlations ([1953SH64](#), [1954ST20](#)).



Resonances are reported at $E_{\alpha} = 1.507 \pm 0.004$ MeV ([1961BA22](#)) and at 1.64, 2.16, 2.26, 2.95, 4.53, 4.85 and 5.36 MeV: see Table 14.8 ([1953SH64](#), [1955SH46](#), [1956BO61](#), [1959GI47](#)). See also ([1969ED1C](#)).



Observed resonances in the yield of 3.09, 3.68, 3.85 and 0.17 [3.85 → 3.68] MeV γ -rays and of various proton groups are displayed in Table 14.8 ([1953SH64](#), [1954ST20](#), [1969GA01](#)). Excitation functions have also been measured for $E_{\alpha} = 4.5$ to 10 MeV ([1969ED1C](#)), 12.4 to 16 MeV ([1967IV1B](#); p_0) and 9.5 to 26 MeV ([1966SP08](#)). See also ([1959AJ76](#)).



Observed resonances below $E_{\alpha} = 2$ MeV are exhibited in Table 14.8 ([1953SH64](#)). Excitation curves have also been determined for $E_{\alpha} = 3.2$ to 3.8 MeV ([1960ON01](#)), $E_{\alpha} = 12$ to 24 MeV (d_0) and 18 to 25 MeV (d_1) ([1967AL16](#)). Clear resonance structure is not observed at these higher energies. See also ([ED69D](#)).



The yield of 0.72 MeV γ -rays has been measured for $E_{\alpha} = 2.1$ to 3.5 MeV: observed resonances are shown in Table 14.8 ([1969GA01](#)).



At $E(^6\text{Li}) = 5$ MeV, deuteron groups are observed to the ground state of ^{14}N and to excited states at 3.95, 4.91, 5.10, 5.69, 5.83, 6.23, 6.44, 7.03, 7.97, 8.47, 9.00, 9.13, 9.41, 9.71, 10.09, 10.43 and 11.06 MeV.

The $T = 1$ state at 10.43 MeV is populated weakly (([1966MC05](#)) and private communication). See also ([1963MO1B](#), [1969CO1D](#)). Branching ratios for the γ -decay of ^{14}N states have been measured: see Table 14.9 ([1966CA07](#)). See also reaction 1 ([1969TH01](#)). See also ([1964CA18](#), [1965CA05](#), [1967CA1D](#)), ([1966BR1G](#)) and ([1965RO1M](#), [1966RO1E](#), [1968TA1N](#)).

Table 14.9: Radiative decays in ^{14}N

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
2.31	0 ⁺ ; 1	0	1 ⁺ ; 0	100	$(8.1 \pm 1.4) \times 10^{-3}$ $(\text{M1}) (5.8 \pm 1.2) \times 10^{-4}$ $(\text{E2}) (4.81 \pm 0.33) \times 10^{-3}$	1966CA07 , 1968RO1C et al. 1966CA07 1965RI02 1967OL02 , 1968RO1C 1956GO42 , 1957BR18 1966GO15 1969YO1B 1956LE28
				6 ± 3		
				3.8 ± 0.5		
				3.7 ± 0.3		
				3.7 ± 0.6		
				3.6 ± 0.6		
				4.9 ± 0.6		
				5.3 ± 1		
				3.9 ± 0.2	0.140 ± 0.013	mean 1966CA07 1967OL02 , 1968RO1C 1969YO1B 1966GO15
				94 ± 3		
4.91	(0, 1) ⁻ ; 0	0	1 ⁺ ; 0	96.3 ± 0.3		
				95.1 ± 0.6		
				96.4 ± 0.6		
				96.1 ± 0.3		
				100		
				0.4 ± 0.7		
				1.3 ± 1.0		
				≤ 0.5		
				≤ 1		
				70 ± 5	$(4.0 \pm 0.5) \times 10^{-5}$	1966CA07 1964WA09 1965WA06 1969YO1B 1966GO15 , 1968AL12
5.11	2 ⁻ ; 0	0	1 ⁺ ; 0	75 ± 3		
				71 ± 5		
				81.2 ± 1.0		
				79 ± 4		
				79.9 ± 1.0		
				30 ± 4		

Table 14.9: Radiative decays in ^{14}N (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
5.69	1 ⁻ ; 0	3.95	1 ⁺ ; 0	25 ± 3		1964WA09
				18.0 ± 1.0		1969YO1B
				29 ± 5		1965WA06
				21 ± 4		1966GO15
				19.7 ± 1.2 (0.7 ± 0.4)		mean
		2.31	0 ⁺ ; 1	0.8 ± 0.2		1966GO15
				37 ± 2		1969YO1B
				40 ± 4		1963AL21
				40 ± 3		1966CA07
				38.4 ± 1.7		1964WA09
5.83	3 ⁻ ; 0	3.95	1 ⁺ ; 0	36 ± 4		1969YO1B
				38.1 ± 1.1		mean
				60 ± 5		1966CA07
				60 ± 3		1964WA09
				64 ± 4		1966GO15
		4.91	(0, 1) ⁻ ; 0	61.6 ± 1.7		1969YO1B
				63 ± 2		1963AL21
				61.9 ± 1.1		mean
				< 2		1966GO15
				< 1		1969YO1B
5.83	3 ⁻ ; 0	0	1 ⁺ ; 0	< 2		1966GO15
				< 1		1969YO1B
				25 ± 4		1966CA07
		0	(0, 1) ⁻ ; 0	25 ± 5		1965WA06
				19.4 ± 1.5		1969YO1B
				29 ± 4		1966GO15

Table 14.9: Radiative decays in ^{14}N (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
6.20	1 ⁺ ; 0	0	1 ⁺ ; 0	21.3 ± 1.3 < 3 < 1 75 ± 5 75 ± 5 80.6 ± 1.5 71 ± 4 78.8 ± 1.3 25 ± 4 24 ± 3 24.1 ± 1.1 21 ± 3 23.8 ± 1.0 72 ± 5 76 ± 3 74.6 ± 1.2 79 ± 3 75.2 ± 1.0 < 1 1.3 ± 0.2 73 ± 5 65 ± 3	$(\text{M2}) = (3.4 \pm 1.4) \text{ } \mu\text{eV}$ $(\text{E3}) = (5 \pm 2) \text{ } \mu\text{eV}$ 2.7×10^{-5}	GA69J
						mean 1966GO15 1969YO1B 1966GO15, 1969YO1B 1966GO15, 1969YO1B 1966CA07 1965WA06 1969YO1B 1966GO15, 1968AL12
						mean 1966CA07 1964WA09 1969YO1B 1966GO15
						mean 1966CA07 1964WA09 1969YO1B 1966GO15
						mean 1966GO15, 1969YO1B 1969YO1B 1966GO15, 1969YO1B 1959RO54, 1966CA07 1964WA09
						1966GO15 1969YO1B 1966GO15, 1969YO1B 1964WA09
						1966GO15, 1969YO1B 1969YO1B 1966GO15, 1969YO1B 1964WA09
						1966GO15, 1969YO1B 1969YO1B 1966GO15, 1969YO1B 1964WA09
						1966GO15, 1969YO1B 1969YO1B 1966GO15, 1969YO1B 1964WA09
						1966GO15, 1969YO1B 1969YO1B 1966GO15, 1969YO1B 1964WA09
6.44	3 ⁺ ; 0					

Table 14.9: Radiative decays in ^{14}N (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
7.03	2 ⁺ ; 0	0	1 ⁺ ; 0	75.3 ± 1.6		1969YO1B 1966GO15
				74 ± 4		mean 1966CA07 1969YO1B 1964WA09 1966GO15
				73.1 ± 1.5		mean 1966CA07 1964WA09 1969YO1B 1966GO15
				20 ± 4		1966CA07 1964WA09 1969YO1B 1966GO15
				18.2 ± 1.1		1966CA07 1964WA09 1969YO1B 1966GO15
		5.11	2 ⁻ ; 0	21 ± 2		1966CA07 1964WA09 1969YO1B 1966GO15
				19 ± 4		1966CA07 1964WA09 1969YO1B 1966GO15
				18.9 ± 0.9		1966CA07 1964WA09 1969YO1B 1966GO15
				7 ± 3		1966CA07 1964WA09 1969YO1B 1966GO15
				14 ± 3		1966CA07 1964WA09 1969YO1B 1966GO15
2.31	0	5.83	3 ⁻ ; 0	6.5 ± 0.6		1966CA07 1964WA09 1969YO1B 1966GO15
				7 ± 2		1966CA07 1964WA09 1969YO1B 1966GO15
				6.8 ± 0.6		1966CA07 1964WA09 1969YO1B 1966GO15
				< 3		1966CA07 1964WA09 1969YO1B 1966GO15
				< 2		1966CA07 1964WA09 1969YO1B 1966GO15
		3.95	1 ⁺ ; 0	< 1		1966CA07 1964WA09 1969YO1B 1966GO15
				100		1960RO13, 1966CA07, 1966GO15, 1969YO1B
				91 ± 4		1965WA06
				98.6 ± 0.3		1967OL02, 1968RO1C ^c
				(M1)(9.1 ± 1.3) × 10 ⁻²		1965WA06
3.95	1 ⁺ ; 0	2.31	0 ⁺ ; 1	(E2)(3.3 ± 0.9) × 10 ⁻²		1966GO15
				< 5		1969YO1B
				< 3		1967OL02, 1968RO1C
				< 1		1967BL22
				0.5 ± 0.1		1965WA06
		3.95	1 ⁺ ; 0	2 ± 1		1967OL02, 1968RO1C
				9 ± 5		1965WA06
				0.9 ± 0.25		1967OL02, 1968RO1C
				< (11 ± 3) × 10 ⁻⁴		

Table 14.9: Radiative decays in ^{14}N (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
7.97	2 ⁻ ; 0	0	1 ⁺ ; 0	< 1		1969YO1B
				< 2		1966GO15
		3.95	1 ⁺ ; 0	≤ 0.4		1967OL02
				< 1		1969YO1B
				< 4		1966GO15
	8.06	55 ± 3			$\omega\Gamma_p\Gamma_\gamma/\Gamma = 0.012$	1960HE14
		45 ± 3			= 0.010	1960HE14
		other states		≤ 3		1960HE14
			1 ⁺ ; 0	82	$\Gamma_\gamma = 10.5 \pm 6$	1956GR17, 1958GR97
			0 ⁺ ; 1	≈ 2		1956LE28, 1957BR25, 1957WI27
			1 ⁺ ; 0	11		1960WA12
8.49	1 ⁻ ; 1	3.95	1 ⁺ ; 0	5		1956LE28, 1957WI27, 1959WA04
		5.69	1 ⁻ ; 0		≈ 0.7	1956DE19, 1966CA07, 1967GA12
	8.62	5.11	2 ⁻ ; 0	100	$(5.6 \pm 2.0) \times 10^{-3}$	1959WA16
		0	1 ⁺ ; 0	23		1959WA16
		3.95	1 ⁺ ; 0	24		1959WA16
		5.69	1 ⁻ ; 0	13		1956LE28, 1957WI27, 1959WA16
		6.20	1 ⁺ ; 0	40		1957WI27
	8.80	0	1 ⁺ ; 0		43 ± 9	1960WA12
		3.95	1 ⁺ ; 0		0.9 ± 0.3	1960WA12
8.91	3 ⁻ ; 1	0	1 ⁺ ; 0		$(6.6 \pm 2.2) \times 10^{-3}$	1968CL05
		5.11	2 ⁻ ; 0	5.5 ± 2.5	$\omega\Gamma_\gamma = 0.040$	1959WA04
		5.83	3 ⁻ ; 0	90 ± 3		1959WA04
		6.44	3 ⁺ ; 0	3 ± 1		1959WA04
		7.03	2 ⁺ ; 0	1.4 ± 0.8		1959WA04
		0	1 ⁺ ; 0	< 1	$\left\{ \begin{array}{l} \Gamma_p/\Gamma_\gamma = 4.1 \pm 0.5 \\ \Gamma_\gamma = (1.36 \pm 0.21) \times 10^{-3} \end{array} \right.$	1967BL22, 1967GA12
		6.44	3 ⁺ ; 0	100		1965DE19, 1966CA07, 1967GA12
9.13	2 ⁻ ; 0	0	1 ⁺ ; 0	> 80	$\omega\Gamma_\gamma \approx 0.03$	1965DE19

Table 14.9: Radiative decays in ^{14}N (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV) ^a	Refs.
9.17	2 ⁺ ; 1	0	1 ⁺ ; 0	79 ± 4	7.7 ± 0.9	1959HA11, 1960RO13, 1963PR03, 1967GA12, 1968CL05
		2.31	0 ⁺ ; 1	1.1 ± 0.4	0.11 ± 0.4	1963PR03, 1968CL05
		5.11	2 ⁻ ; 0	< 1		1960RO13
		5.69	1 ⁻ ; 0	< 6	0.3 ± 0.25	1960RO13, 1968CL05
		5.83	3 ⁻ ; 0	3 ± 2	0.33	1960RO13
		6.44	3 ⁺ ; 0	8 ± 2	0.78 ± 0.35	1960RO13, 1968CL05
		7.03	2 ⁺ ; 0	6.3 ± 0.5	0.85 ± 0.15	1963PR03
				3 ± 1	0.3 ± 0.15	1960RO13, 1968CL05
		3.95	1 ⁺ ; 0	3.5 ± 0.5	0.34 ± 0.05	1963PR03
				6 ± 1	0.37	1959WA04
9.51	2 ⁻ ; 1	5.11	2 ⁻ ; 0	78 ± 3	4.8	1959WA04
		5.83	3 ⁻ ; 0	16 ± 2	1.0	1959WA04
		2.31	0 ⁺ ; 1		4 ± 1.3	1963RO17
10.23	1 ⁽⁻⁾ ; 0	3.95	1 ⁺ ; 0	82 ± 6	12.1 ± 1.5	1964RO03, 1968CL05
				2.31		1964RO03
		5.11	0 ⁺ ; 1	< 1		1964RO03
		5.69	1 ⁻ ; 0	< 2		1964RO03
		5.83	2 ⁻ ; 0	2 ± 1	0.3 ± 0.2	1964RO03, 1968CL05
		6.44	1 ⁻ ; 0	< 3		1964RO03
		6.44	3 ⁻ ; 0	< 1		1964RO03
		7.03	3 ⁺ ; 0	8 ± 1	1.2 ± 0.4	1964RO03, 1968CL05
		2.31	2 ⁺ ; 0	6 ± 1	0.88 ± 0.31	1964RO03, 1968CL05

^a See also Table 14.13.^b See also (1969HA49).



Triton groups corresponding to a number of ^{14}N states have been observed at $E(^7\text{Li}) = 5$ MeV: see (1966MC05) and reaction 6. The angular distribution of the ground state tritons has been measured at $E(^7\text{Li}) = 4.5$ MeV by (1963MO1B). See also (1965CA05, 1967CA1D) and (1966RO1E).

8. (a) $^{11}\text{B}(^3\text{He}, \gamma)^{14}\text{N}$	$Q_{\text{m}} = 20.735$	$E_{\text{b}} = 20.735$
(b) $^{11}\text{B}(^3\text{He}, \text{n})^{13}\text{N}$	$Q_{\text{m}} = 10.182$	
(c) $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$	$Q_{\text{m}} = 13.185$	
(d) $^{11}\text{B}(^3\text{He}, \text{d})^{12}\text{C}$	$Q_{\text{m}} = 10.463$	
(e) $^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$	$Q_{\text{m}} = -1.999$	
(f) $^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}$		
(g) $^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B}$	$Q_{\text{m}} = 9.122$	
(h) $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$	$Q_{\text{m}} = 4.566$	

The capture γ -rays (reaction (a)) have been studied for $E(^3\text{He}) = 1$ to 3 MeV. The differential cross section (at 90°) for the γ_0 transition increases rapidly with energy from $\approx 0.002 \mu\text{b}/\text{sr}$ at 1 MeV to $\approx 0.5 \mu\text{b}/\text{sr}$ at 3.0 MeV. At the latter energy $I_\gamma/I_{\gamma_0} \approx 0.22 \pm 0.08$. There is some evidence also for transitions to higher excited states of ^{14}N (1965PU1B).

The excitation function at 0° in the range $E(^3\text{He}) = 1.5$ to 5.6 MeV for neutrons corresponding to $^{13}\text{N}(0)$ (reaction (b)) shows a broad peak at $E(^3\text{He}) = 4.15$ MeV which may indicate the existence of a ^{14}N state at $E_x \approx 24.0$ MeV, $\Gamma \approx 1$ MeV (1966DI04). The excitation function for reaction (b) has also been measured for $E(^3\text{He}) = 6$ to 18 MeV (1967HA20). See also (1964DI1C, 1965BR42).

Yield curves for protons (reaction (c)) have been measured for $E(^3\text{He}) = 3.0$ to 5.5 MeV ($p_0, p_1, p_1 + p_2 + p_3$): they are rather featureless (1959HO01). This is also true for the ground state deuterons of reaction (d) in the same energy interval (1959HO01). Yield curves for reaction (e) have been measured for $E(^3\text{He}) = 6$ to 18 MeV (1967HA20) and 10 to 30 MeV (1965BR42). See also ^{13}C , and ^{11}C and ^{12}C in (1968AJ02). For reaction (f), see (1968PA1Y).

The excitation functions of α -particle groups ($\alpha_0, \alpha_1, \alpha_2, \alpha_3$) (reaction (g)) have been measured for $E(^3\text{He}) = 2.2$ to 5.5 MeV. No significant resonance behavior is seen except for the α_2 group which, in the 15° excitation function, exhibits a resonance at $E(^3\text{He}) = 4$ MeV, $\Gamma \approx 1$ MeV (1965FO06). See also ^{10}B in (1966LA04). The excitation function for the reaction $^{11}\text{B}(^3\text{He}, ^6\text{Li}_{\text{g.s.}})^8\text{Be}_{\text{g.s.}}$ has been measured for $E(^3\text{He}) = 1.4$ to 5.8 MeV: no pronounced compound nucleus effects are observed (1967YO02, 1967YO1C).



Table 14.10: Resonances in $^{12}\text{C} + \text{d}$

E_{d} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$^{14}\text{N}^*$ ^a (MeV)	$J^\pi; T$	Refs. ^b
0.335		p	10.559		1956KO26, 1956VA17
0.92	95	n, p ₀ , p ₁	11.06	1 ⁺ ; 0	1957SA01, 1960KA06, 1969BO32
1.13		p ₀ , p ₁	11.24	; 1	1957SA01, 1963JE02, 1969BO32
1.19	190	n, p ₀ , p ₁ , d	11.29	2 ⁻ ; 0	1957SA01, 1960KA06, 1963JE02, 1966LA1Q, 1969BO32
1.23		p ₀	11.33	(3 ⁺)	1969BO32
1.30	30	n, p ₀ , p ₁ , d	11.38	1 ⁺ ; 0	1957SA01, 1960KA06, 1963JE02, 1965FI05, 1966LA1Q, 1969BO32, 1969JA06
1.39		p ₀	11.46	(2 ⁻)	1969BO32
1.446	5	p ₀ , p ₁ , d	11.51	3 ⁺	1957SA01, 1960KA06, 1963JE02, 1965FI05, 1966LA1Q, 1969BO32
1.55		p ₀	11.60	(2 ⁻)	1969BO32
1.62		n, p	11.65	(2 ⁻)	1969BO32, 1969JA06
1.68		p ₀ , p ₁	11.71	(1 ⁺ , 2 ⁻)	1963JE02, 1965FI05, 1966KL05, 1968BE2A
1.78	85	n, p ₀ , p ₁	11.80	1 ⁺	1960KA06, 1963JE02, 1965FI05, 1969BO32, 1969JA06
1.95		p ₀	11.94	(1 ⁻)	1969BO32
2.30		n, p ₁ , p ₂ , p ₃	12.24		1965FI05, 1969JA06
2.498 \pm 6	40 \pm 3	n, p ₀ , p ₁ , p ₂ , p ₃ , d	12.411	4 ⁻	1956MC88, 1958MC63, 1960CH12, 1963JE02, 1965FI05, 1969JA06
2.622 \pm 8	19 \pm 9	n, p ₁ , p ₂ , p ₃	12.517		1956MC88, 1958MC63, 1960CH12, 1965FI05, 1969JA06
2.726 \pm 10	47 \pm 3	(n), p ₁ , p ₂ , p ₃ , d	12.606	3 ⁺	1956MC88, 1958MC63, 1960CH12, 1963JE02, 1965FI05, 1969JA06
2.817 \pm 7	27 \pm 6	n, p ₁ , p ₂ , p ₃ , d	12.684		1956MC88, 1960CH12, 1963JE02, 1965FI05, 1966GU04, 1969JA06
2.946 \pm 6	22 \pm 6	p ₂ , p ₃ , d	12.794		1956MC88, 1960CH12, 1963JE02
2.982 \pm 6	11 \pm 3	n, p ₃ , d	12.825		1956MC88, 1960CH12, 1969JA06
3.015 \pm 15	71 \pm 7	n, p ₀	12.853		1966GU04, 1966SA05
3.119 \pm 6	29 \pm 9	p ₁ , p ₂ , p ₃ , d	12.942	(3 ⁻ , 4 ⁻)	1956MC88, 1960CH12, 1963OH02, 1965FI05
3.39 \pm 12	47 \pm 15	n, p ₂ , p ₃ , d	13.17	(0 ⁻ , 1 ⁻)	1955MA76, 1956MC88, 1963JE02, 1963OH02, 1966SA05,

Table 14.10: Resonances in $^{12}\text{C} + \text{d}$ (continued)

E_{d} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$^{14}\text{N}^*$ ^a (MeV)	$J^\pi; T$	Refs. ^b
3.97 \pm 30	< 200	p ₀ , p ₂ , p ₃ , (d)	13.67	(2 ⁺ , 3 ⁺)	1969JA06
		n, (p), d	13.71	(1 ⁺)	1963JE02, 1965FI05, 1966BR2B, 1967FU03
					1963OH02, 1966BR2B, 1966HO11, 1966SA05, 1967FU03,
					1969JA06
		p ₀ , p ₁ , p ₂ , p ₃ , d	14.04		1963JE02, 1966BR2B
		n, p ₂ , d	14.17		1966BR2B, 1966HO11
		p ₀ , p ₂ , d	14.38		1963OH02, 1966BR2B
		d	14.70		1966BR2B
		p ₀ , p ₁ , p ₂ , p ₃ , d, α	14.84		1956BO08, 1963OH02, 1966BR2B, 1966PA1J, 1969CO02
		d	15.11		1956BO08, 1963OH02, 1966BR2B
		p ₁ , p ₃ , d	15.26		1966BR2B
		p ₁ , p ₂ , α	15.47		1966BR2B, 1966PA1J, 1969CO02
		p ₀ , p ₃ , d, α	15.7		1966BR2B
		α	16.4		1966PA1J, 1969CO02
		p ₀ , p ₂ , d, α	17.2		1963OH02, 1966PA1J, 1969CO02
		p ₀ , d	19.6	(5 ⁺)	1963OH02, 1968KL06
		n, p ₀ , d	20.4	(5 ⁺)	1965BA06, 1968KL06
12.70 \pm 150	\approx 1400	d, α_2	21.15	(4 ⁻)	1968JA09, 1968KL06
14.52 \pm 100	\approx 900	α_2	22.70		1968JA09

^a (1963JE02) report 39 excited states of ^{14}N with $11.2 < E_x < 14.2$ MeV.

^b See also (1959AJ76) and (1959EL44, 1961CI08, 1961GR06, 1961JA08, 1961ST10, 1966KA05, 1966KL05, 1966LA1P, 1966LA1Q, 1968BE2A, 1969PA1K).

Table 14.11: Recent $^{12}\text{C} + \text{d}$ yield curves ^a

E_{d} (MeV)	Yield of	Refs.
0.7 – 1.4	n_0	(1961JA08)
1.2 – 4.5	σ_t	(1969JA06)
1.4 – 3.1	n_0	(1959EL44)
2.7 – 3.2	n_0	(1966GU04)
2.8 – 4.2	n_0	(1966SA05)
3.8 – 4.2	n_1	(1966SA05)
3.8 – 4.2	n_0	(1967FU03)
3.8 – 5.0	n_0, n_1	(1966HO11)
9.2 – 13.8	σ_t	(1965BA06)
0.7 – 2.0	p_0, p_1	(1960KA06)
0.8 – 1.1	p_0	(1961GR06)
0.8 – 1.2	p_0	(1962WE12)
0.9 – 1.8	p_0	(1966KL05)
0.9 – 2.0	p_0	(1969BO32)
0.9 – 5.0	p_0, p_1, p_2, p_3	(1963FR1D, 1963JE02)
1 – 2	p_0, p_1	(1966LA1P)
1.1 – 1.2	p_0	(1961ST10)
1.2 – 4.3	p_1, p_2, p_3	(1960GO19, 1965FI05)
1.2 – 4.4	p	(1965BA1W)
1.5 – 4.0	p_0, p_1, p_2, p_3	(1967LE1K)
1.7 – 3.1	p_3	(1960CH12)
2 – 3.3	p_1, p_2, p_3	(1966KA05)
2.6 – 4.0	p_0, p_1, p_2, p_3	(1966GE03)
2.7 – 3.2	p_0	(1966GU04)
4 – 7	p_0, p_1, p_2, p_3	(1966BR2B)
5 – 10	p_0, p_1, p_2, p_3, p_4	(1969CO02)
8.5 – 10	p_5, p_6, p_7	(1969CO02)
5.5 – 12	p_0, p_1	(1963EV04)
9 – 14	p_0	(1968KL06)

Table 14.11: Recent $^{12}\text{C} + \text{d}$ yield curves ^a (continued)

E_{d} (MeV) (MeV)	Yield of	Refs.
9.9	σ_t	(1962WI15, 1963WI1D)
0.9 – 2.0	d_0	(1969BO32)
0.9 – 5	d_0	(1963FR1D, 1963JE02)
1 – 2	d_0	(1966LA1P)
1.5 – 2.0	d_0	(1960KA06)
2.0 – 3.3	d_0	(1966KA05)
2 – 6	d_0	(1967PL1B)
2.6 – 4.0	d_0	(1966GE03)
3 – 12	d_0	(1963OH02)
4 – 7	d_0	(1966BR2B)
4.5 – 5.7	d_0	(1968CO04)
5 – 10	d_0	(1969CO02)
8.5 – 10	d_1	(1969CO02)
9 – 14	d_0	(1968KL06)
26.5	σ_t	(1965MA57)
5 – 9	α	(1966PA1J)
5 – 10	α_0, α_1	(1969CO02)
7 – 10	α_3	(1969CO02)
9 – 10	α_4	(1969CO02)
9 – 12.5	$\alpha_0, \alpha_1, \alpha_2, \alpha_3$	(1966ME09)
9.2 – 13.8	$\alpha_0, \alpha_1, \alpha_3, \alpha_4$	(1965BA06, 1966BA32)
11 – 14	α_2	(1969SM03)
13 – 21	α_2	(1968JA09)
15 – 20	$\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4$	(1963YA1B)
21 – 29	$\alpha_0, \alpha_1, \alpha_3, \alpha_4, \alpha_5$	(1969YA1C)

^a See also (1959AJ76).


Angular distributions have been measured for $E_\alpha = 2.4$ to 3.7 MeV (1966MA04; n_0), 3.4 to 3.7 MeV (1966MA04; n_1), 2.1 to 5.4 MeV (1962CA05; n_0) and 13.5 and 13.9 MeV (1962KJ03; n_0, n_2). See also (1959HE1B) and (1959AJ76).



At $E(^6\text{Li}) = 4.5$ MeV, the angular distributions of the triton groups to $^{14}\text{N}^*(0, 2.31, 3.95)$ have been measured ([1963MO1B](#)). See also ([1965CA05](#)).



The capture cross section is reported to be $\approx 0.6 \pm 0.2 \mu\text{b}$ at $E_{\text{d}} = 1$ MeV ([1961KN03](#)): at $E_{\text{d}} = 1.5$ MeV it is $< 1 \mu\text{b}$ ([1955AL16](#)).




Resonances in the yields of neutrons and protons are displayed in Table 14.10. Recent measurements of the yields of neutrons and protons are listed in Table 14.11. See also ([1965WI11](#), [1968CO04](#), [1969RO1R](#)) and ([1959AJ76](#)).

Angular distributions of neutrons and protons have been reported at many of the energies listed in Table 14.11. Except at very low energies, direct interaction is the predominant mechanism although resonances are observed for the first few MeV (see Table 14.10) and fluctuations persist to 12 MeV ([1963EV04](#)). See also ([1955WI43](#)), ([1959AJ76](#)), ^{13}C and ^{13}N . At $E_{\text{d}} = 2.50$ MeV ($^{14}\text{N}^* = 12.41$) the $(\text{d}, \text{p}_3\gamma)$ angular correlation is undistorted. This is an example of a resonant direct reaction due to a single-particle state ([1966KA05](#)).

Polarization measurements are summarized in Table 14.12. See also ([1968BA2R](#), [1968BA47](#); theor.) (reaction (a)), and ([1959AL08](#), [1962GR10](#), [1963BE1M](#), [1966SK1A](#), [1967GR1L](#); exp.), ([1967CI1A](#), [1967KH1A](#), [1968BA2R](#), [1968BA47](#), [1969PE1N](#), [1969PE1L](#); theor.) (reaction (b)). See also the reviews by ([1961GO1K](#), [1963HA1G](#), [1966DA1B](#), [1966MI1E](#)).

For reaction (a) see also ([1959BR75](#), [1966WY01](#), [1967SC43](#), [1967WY02](#); exp.) and ([1961MA1E](#), [1964CA1F](#), [1967HO1K](#), [1968NO1C](#); theor.). For reaction (b) see also ([1963GE03](#), [1967AU05](#), [1967MO1P](#), [1967TI1A](#), [1968NO1C](#), [1969PE09](#)). For a discussion of sequential decay, see ([1963PI04](#)) and ^{13}C and ^{13}N .



Reported resonances are displayed in Table 14.10. Recent measurements of yields of scattered deuterons are listed in Table 14.11. See also ([1961CI08](#), [1962GR10](#), [1963GE03](#), [1965BA1W](#), [1967AU05](#), [1968BA2P](#), [1968GO1N](#)), and ([1965SA1H](#), [1966BA60](#), [1967HO1K](#), [1969IW1D](#)). See also ^{12}C in ([1968AJ02](#)) and ([1959AJ76](#)).

Polarization measurements are summarized in Table 14.12. See also ([1963ZA1B](#), [1965CA05](#), [1967RU1A](#); theor.).

Table 14.12: $^{12}\text{C} + \text{d}$ polarization studies

E_{d} (MeV)	Groups	Refs.
1.5 – 1.8	n_0	(1968KA1K)
1.7 – 2.8	n_0	(1966ME16)
2.8	n_0	(1965GA1G)
2.8, 3.0	n_0	(1963HA1G)
2.8 – 4.0	n_0	(1966SA05)
3.8 – 4.2	n_0	(1968DO09)
4 – 7.5	n	(1965KE10)
4.2 – 5.0	n_0	(1966MO14)
5.2 – 6.2	n_0, n_1	(1968DO09)
6 – 6.5	n_0, n_1	(1963BA38, 1963BA66, 1966BA33)
11.8	n_0	(1961LE1E)
12.9	n_3	(1959BU1E, 1960BU15)
51.5	n_0	(1969BR1E)
0.8 – 1.1	p_0	(1961GR06)
0.9, 1.0	p_0	(1968GL1B)
1.1	p_0	(1959CI38, 1959JU1A, 1964AS03)
2.4	p_0	(1963MI1G)
2.8, 3.0, 3.2	p_1	(1966HE1G)
2.9, 3.1, 3.3	p_0	(1969SO07)
4.0	p_0	(1966ST1N)
5.5 – 9.5	p_0	(1963EV04)
7 – 11	p_1	(1963EV04)
7, 10	p_0, p_1	(1968YU01)
8.9	p_0	(1959HI1E)
10	p_0	(1959AL09, 1962AL03)
10.8	p_0, p_1	(1961JO14)
11.9	p_0, p_1	(1968BA19)
13.6	p_0	(1966JI1A)
13.8	p_0	(1966MA25)
15	p_1	(1962IS04, 1964RE04)
21	p_0	(1963BO1J)

Table 14.12: $^{12}\text{C} + \text{d}$ polarization studies (continued)

E_{d} (MeV)	Groups	Refs.
2.8 – 4.4	d_0	(1969PA1K)
3.5 – 7.1	d_0	(1968CO10)
6	d_0	(1960AL09)
6 – 10.5	d_0	(1967CL07)
6.5	d_0	(1963NE1H)
12	d_0	(1968BA17)
13.6	d_0	(1968ZA02)
13.8	d_0	(1966MA2H)
22	d_0	(1965AR1F, 1965GR1Q)
51	d_0	(1968SE1E)

$$14. \ ^{12}\text{C}(\text{d}, \text{t})^{11}\text{C} \quad Q_{\text{m}} = -12.462 \quad E_{\text{b}} = 10.272$$

The cross section rises from ≈ 0.1 mb at $E_{\text{d}} = 16$ MeV to ≈ 10 mb at 20 MeV (1955WI43). See also ^{11}C in (1968AJ02).

$$15. \ ^{12}\text{C}(\text{d}, \alpha)^{10}\text{B} \quad Q_{\text{m}} = -1.341 \quad E_{\text{b}} = 10.272$$

Excitation functions have been measured for $E_{\text{d}} = 5$ to 29 MeV: see Table 14.11.


Resonant structure has been reported by (1966PA1J) and (1968JA09): see Table 14.10. Angular distributions have been measured at many of the above energies and analyzed by DWBA: see (1968KL06, 1969YA1C) and ^{10}B in (1966LA04). See also (1963PE07, 1967HO1K).

The yield of the α_2 group (to $^{10}\text{B}^*(1.74)$ [$J^\pi = 0^+$; $T = 1$]) is typically $< 1\%$ of the intensity of the groups to the neighboring $T = 0$ states in the range $E_{\text{d}} = 9$ to 12.5 MeV. This is partly due to isospin conservation and partly to the J^π selection rule involved in this transition. When the latter effect is calculated and the corresponding factor removed, the intensities of the $T = 1$ α_2 -groups range from $\approx 10\%$ of the intensities of the $T = 0$ α -groups at $E_{\text{d}} = 9$ MeV to $\approx 1-2\%$ at $E_{\text{d}} = 11$ MeV. Above $E_{\text{d}} = 11.5$ MeV, the yield of the α_2 group increases slightly indicating perhaps a direct-interaction mechanism involving isospin mixing at the surface of the nucleus (1966ME09). [See, however, below]. The mixing might also occur through Coulomb excitation during the d-capture or the α -emission (1966ME09). Some fluctuations in the cross sections, with widths of a few hundred keV, are observed at forward and backward angles. Ericson fluctuations

may be involved ([1966ME09](#)). The data of ([1966ME09](#)) have been interpreted as indicating an intermediate structure resonance corresponding to $E_x \approx 18$ MeV, $\Gamma \approx 2$ MeV [$J^\pi = 3^-$], whose doorway state is a single-particle cluster resonance in either, or both, the entrance or exit channels ([1968NO1C](#)). However, ([1969SM03](#)) find that the broad structure reported by ([1966ME09](#)) at $E_d \approx 13$ MeV can be resolved into at least three separate peaks, in contradiction to the predictions of ([1968NO1C](#)). Both the angular distributions and the excitation functions can be interpreted in terms of a few resonant states of ^{14}N [$l = 4$ and 5 account for most of the cross section at $E_d \approx 12.5$ MeV], without introducing any large direct reaction amplitudes ([1969SM03](#)). See also ([1969NO1C](#)).



Angular distributions of the n_0 , n_1 and n_2 groups have been measured at $E_t = 1.7$ MeV ([1966MA2G](#)). At $E(^{12}\text{C}) = 12$ to 20 MeV, the lifetimes of $^{14}\text{N}^*(5.11, 5.83)$ have been determined using recoil distance method: $\tau_m(5.11) = 12.4 \pm 1.4$ psec, $\tau_m(5.83) = 18 \pm 2$ psec: see Tables [14.9](#) and [14.13](#). The $5.11 \rightarrow 0$ transition is enhanced by 2.2 ± 0.7 W.u. The allowed M2 $5.11 \rightarrow 2.31$ transition has $|M|^2 = 0.83 \pm 0.14$ W.u. while the isospin forbidden part of the $5.11 \rightarrow 0$ transition has $|M|^2 = (3.3 \pm 1.3) \times 10^{-3}$ W.u. ([1968AL12](#)). See also ([1966AL11](#)).



Many proton groups have been observed: see Table [14.14](#). At $E(^3\text{He}) = 20$ MeV angular distributions of the protons corresponding to states with $0 < E_x < 12.6$ MeV have been measured by ([1968MA29](#)) and analyzed using the distorted-wave calculations of ([1965GL09](#)) and spin-independent interaction potential. L -values have been assigned and are displayed in Table [14.14](#). It is pointed out that in this reaction unnatural parity states of $T = 1$ are not allowed: the proton groups corresponding to the 0^- 8.80-MeV state and the 2^- 9.51-MeV state, both of which are $T = 1$, are not observed ([1967MA1G](#), [1968MA29](#)). Angular distributions have also been obtained at many other energies: see ([1965GR1R](#): 1 – 1.8 MeV; p_0, p_1, p_2), ([1963LU01](#), [1963LU1F](#), [1964LU1B](#): 2.3 – 3 MeV; p_2), ([1964KU05](#): 2.5 – 4.9 MeV; $p_0 \rightarrow p_9$), ([1969HA49](#): 3.0 – 9.2 MeV; p_0, p_1 : 3.0 – 11 MeV; $p_2 \rightarrow p_4$: 5.1 – 11 MeV; $p_5 \rightarrow p_9$), ([1967CL1C](#): 3.5 MeV; $p_0 \rightarrow p_8$), ([1966BL01](#): 5.1 MeV; $p_2 \rightarrow p_6$), ([1968LA19](#): 5.3 – 5.5 MeV; p_1, p_2, p_3, p_5), ([1965FU16](#): 6.6 – 10.7 MeV), ([1960PR12](#): 13.9 MeV; p_0, p_1, p_2), ([1969HO23](#): 15 MeV: $0 < E_x < 12.8$ MeV), ([HO66](#),

Table 14.13: Lifetimes of some ^{14}N states ^a

E_x (MeV)	τ_m (psec)	Reaction	Refs.
2.31	0.083 ± 0.019	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1966LI07)
	0.083 ± 0.03	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1965LO07)
	0.097 ± 0.03	$^{14}\text{N}(\gamma, \gamma)$	(1964BO22)
	0.077 ± 0.018	$^{14}\text{N}(\gamma, \gamma)$	(1961SW01)
	0.033 ± 0.003	$^{14}\text{O}(\beta^+)$	(1966SI05)
3.95	0.083 ± 0.011		mean ^b
	≤ 0.087	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1966LI07)
	0.0045 ± 0.0004	$^{14}\text{N}(\text{e}, \text{e})$	(1964BI09, 1967OL02)
	≤ 0.025	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1965LO07)
	≤ 0.02	$^{14}\text{N}(\text{n}, \text{n})$	(1969NY1A)
4.91	< 0.050	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1967LI04)
	< 0.050	$^{13}\text{C}(\text{d}, \text{n})$	(1963AL21)
	< 0.06	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1970GA09)
5.11	12.4 ± 1.4	$^{12}\text{C}(\text{t}, \text{n})$	(1968AL12)
	4.5 < τ < 20	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1967LI04)
	> 0.6	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1965LO07)
	> 0.3	$^{13}\text{C}(\text{p}, \gamma)$	(1959WA04)
	≤ 0.036	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1967LI04)
5.69	≤ 0.05	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1970GA09)
	18 ± 2	$^{12}\text{C}(\text{t}, \text{n})$	(1968AL12)
	> 4.5	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1967LI04)
6.20	> 23	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1964BE12)
	< 0.04	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1970GA09)
	0.63 ± 0.08	reactions 1, 6, 19	(1969TH01)
6.44 ^c	0.59 ± 0.12		(1964BE12)
	0.65 ± 0.05		(1969GA16)
	0.63 ± 0.04		(1970GA09)
	0.63 ± 0.03		mean
7.03 ^c	5.4 ± 0.5 fsec	$^{14}\text{N}(\gamma, \gamma)$	(1966SW01)
	< 0.05 psec	$^{12}\text{C}({}^3\text{He}, \text{p})$	(1970GA09)

^a See also Table 14.9.

^b Not included the value of (1966SI05).

^c See also (1969NY1A).

[1967FO1E](#): 15 MeV), ([1968MA46](#): 25.3 MeV: $0 < E_x < 9.5$ MeV) and ([1959AJ76](#)) for a listing of the earlier work. See also ^{15}O . The parity of $^{14}\text{N}^*(6.44)$ is even from angular distribution measurements of protons and γ -rays at the $E(^3\text{He}) = 2.99$ MeV resonance ([1964KU05](#)).

The γ decay of many states has been studied: Table 14.9 displays observed branching ratios and radiative widths ([1964WA09](#), [1965NE06](#), [1965RI02](#), [1965WA06](#), [1966GO15](#), [1967CH19](#), [1967GA12](#), [1967OL02](#), [GA69J](#), [1969HA49](#)). A very accurate value of the excitation energy of $^{14}\text{N}^*(5.11)$ is derived from E_γ : $E_x = 5.10587 \pm 0.00018$ MeV ([1967CH19](#)). p- γ angular correlations have been measured at many energies. The results demand $J = 0$ or 1 for $^{14}\text{N}^*(4.91)$ and $J = 2$ for $^{14}\text{N}^*(5.11)$. The ground state transition for the latter contains E1, M2 and E3 components ([1959WA04](#), [1965BL04](#), [1965NE06](#), [1966GO15](#)), $J = 2$ for $^{14}\text{N}^*(7.03)$ and the angular correlation of the 2.51 MeV γ -ray ($8.96 \rightarrow 6.44$) indicates $J \leq 5$ for $^{14}\text{N}^*(8.96)$ ([1967BL22](#)). $J^\pi = 5^+$ ([1967GA12](#)). See also ([1963LU01](#), [1963LU1F](#), [1964LU1B](#)) and Table 14.9. Polarization measurements lead to $J^\pi = 0^-$, 1^- for $^{14}\text{N}^*(4.91)$ and odd parity for $^{14}\text{N}^*(5.11, 5.83)$ ([1963BE33](#), [1968BL09](#)). The parity of $^{14}\text{N}^*(6.21)$ is even ([1964WA09](#)). An analysis of elastic ^3He , the angular distribution of the protons to $^{14}\text{N}^*(6.44)$ and of the subsequent ground state γ -rays at $E(^3\text{He}) = 2.99$ MeV ($^{15}\text{O}^*(14.46)$) leads to even parity for $^{14}\text{N}^*(6.44)$ ([1964KU05](#)). The angular correlation of internal pairs is consistent with E2 radiation ([1964WA09](#)) for the $6.44 \rightarrow 0$ transition.

Measurements of mean lifetimes are displayed in Table 14.13 ([1964BE12](#), [1965LO07](#), [1966LI07](#), [1967LI04](#), [1969GA16](#), [1970GA09](#)). See also ([1963BE33](#)).

See also ([1959AL96](#), [1959FA1A](#), [1959HI69](#), [1960HA31](#), [1961CE02](#), [1967BE2G](#), [1967FO1E](#), [1967OG1A](#), [1969BA1Z](#), [1969GO11](#), [1969HA2D](#)) and ([1959EL43](#), [1960EL1C](#), [1960NE1A](#), [1962EL1C](#), [1967HA1T](#), [1969BO1G](#), [1969LI1D](#); theor.).



Angular distributions of deuterons corresponding to various states of ^{14}N (see Table 14.15) have been measured at $E_\alpha = 42$ to 53 MeV ([1959BO40](#), [1960HA32](#), [1962CE01](#), [1962HA40](#), [1965PE03](#), [1967ZA01](#)). See ([1962CE01](#), [1963GL1C](#), [1965GL09](#)) for discussions of the analysis. The known $T = 1$ states are not excited in this reaction: see Table 14.15 and ([1960HA32](#), [1965PE03](#), [1967ZA01](#), [1968NO1C](#)). [It should be noted that in addition to isospin conservation, angular momentum and parity considerations would also inhibit the excitation of the $J^\pi = 0^+$; $T = 1$ 2.31-MeV state; see, e.g., ([1960HA32](#)).] No evidence is seen at $E_\alpha = 42$ MeV for deuterons in the $J^\pi = 0^+$; $T = 1$ singlet state leading to the excitation of the 2.31-MeV state: $\bar{d}_1/d_0 = 5 \times 10^{-3}$ ([1967CR1G](#), [1969BR1N](#)). The deuteron spectrum is dominated by very strong groups corresponding to the $(d_{5/2})^2$, $J^\pi = 5^+$, state at 8.96 MeV, and to a state at 15.1 MeV ([1962HA40](#), [1966RI04](#)).

Comparison of the angular distribution of ground state deuterons ($E_\alpha = 41.7$ MeV) with that of the ground state alphas from the $^{14}\text{N}(d, \alpha)^{12}\text{C}$ reaction ($E_d = 20$ MeV) leads to an upper limit of 3% for the time reversal non-conserving fraction of the Hamiltonian ([1959BO40](#), [1959HE1C](#)).

See also ([1964NA1E](#), [1965PE17](#), [1966BR1G](#), [1967OG1A](#)) and ([1961EL1A](#), [1968ZE1B](#); theor.).

Table 14.14: Excited states of ^{14}N from $^{12}\text{C}({}^3\text{He}, \text{p})^{14}\text{N}$

E_x (MeV \pm keV)					
(1960HI07)	(1968GR1G)	(1959YO25)	(1961ER01) ^a	(1969HO23)	L ^b
0	0	0		0	2
2.314 ± 10		2.313 ± 17		2.319 ± 15	0
3.946 ± 10	3.948 ± 5	3.946 ± 17		3.952 ± 15	0
4.909 ± 10	4.913 ± 5	4.888 ± 18		4.927 ± 15	1
5.102 ± 10	5.107 ± 5	5.078 ± 18		5.117 ± 15	1
5.686 ± 10	5.693 ± 5		5.692 ± 8	5.713 ± 15	1
5.830 ± 10	5.834 ± 5	5.812 ± 18	5.835 ± 8	5.885 ± 15	3
	6.202 ± 5 ^f	6.21 ± 20 ^b	6.204 ± 8	6.224 ± 15	0
	6.450 ± 5	6.46 ± 18 ^b	6.441 ± 8	6.468 ± 15	2
E_x (MeV \pm keV)		E_x (MeV \pm keV)			
(1968MA29)	(1969HO23)	L	(1968MA29)	(1969HO23)	
7.01 ± 42	7.036 ± 15	2	11.27 ± 50	11.249 ± 15	
7.95 ± 26	7.974 ± 15	3	11.39 ± 40	11.357 ± 15	
8.05 ± 35	8.072 ± 15	1	11.51 ± 30	11.517 ± 15	
8.47 ± 30 ^c	8.493 ± 15	3	11.66 ± 40		
8.61 ± 34	8.625 ± 15	0	11.79 ± 110		
	8.912 ± 15		11.95 ± 30		
8.96 ± 19	8.97 ± 15	3, 4, 2		12.29 ± 15	
9.15 ± 18	9.126 ± 15		12.40 ± 30	12.425 ± 15	
	9.176 ± 15		12.50 ± 20	12.506 ± 15	
9.39 ± 26 ^d	9.389 ± 15		12.63 ± 25	12.608 ± 15	
9.70 ± 22	9.703 ± 15			12.69 ± 15	
10.08 ± 18	10.063 ± 15		12.74 ± 30	12.80 ± 15	
	10.101 ± 15		12.90 ± 25		
10.43 ± 20	10.441 ± 15		13.15 ± 40		
10.56 ± 28			14.91 ± 60		
10.81 ± 23 ^e	10.812 ± 15		15.8 ± 200		
11.06 ± 50	11.053 ± 15		17.4 ± 200		

^a Based on Q_m .

^b From (1968MA29).

^c 4^- ; 0 (1968MA29).

^d 2^- ; 0 (1968MA29).

^e 4^+ ; 0 (1968MA29).

^f Private communication.



At $E(^6\text{Li}) = 20$ MeV, α -groups corresponding to most of the $T = 0$ states with $E_x < 12.7$ MeV are reported. See Table 14.15. The spectrum is dominated by the α -group corresponding to the 5^+ state at 9.0 MeV (1968ME10). See also (1962HO06, 1967DZ01). The α_1 group to the $J^\pi = 0^+$; $T = 1$ state at 2.31 MeV has been observed. Its intensity (< 3% of α_0) decreases sharply from $E(^6\text{Li}) = 4$ to 5.5 MeV, while the intensities of the α_0 and α_2 groups (to $T = 0$ states) increase rapidly. It is suggested that broad levels at the higher excitation energies correspond to such short lifetimes that the Coulomb forces do not have enough time to change the relative amounts of various isospin components in the compound nucleus wave function (1965CA06). See also (1962HO06, 1967DZ01).

Angular distributions of α -particles have been reported for $E(^6\text{Li}) = 2.0$ and 2.4 MeV (1966BE07; α_0, α_2), 3.0 MeV (1963BA08; α_0), 3.2 to 4.0 MeV (1962HO06; $\alpha_0, \alpha_1, \alpha_2$), 4.0 to 5.5 MeV (1965CA06; α_1), 4.5 to 5.5 MeV (1966HE05; α_0, α_2) and 20 MeV (1968ME10: see Table 14.15).

Doppler-shift attenuation measurements have led to the determination of τ_m for the 6.44 MeV state: (0.63 ± 0.08) psec (1969TH01): see Table 14.13 and reaction 1. See also Table 14.9.

See also (1960SH01, 1960SH05, 1961MA1K, 1961MA33, 1962LI08, 1963OL1B, 1964CA18, 1967CH34, 1967OG1A, 1968DA20) and (1962IN1C, 1963TA1B, 1966RO1E, 1966RO1F, 1968HO1J, 1968RO1D; theor.).



At $E(^{11}\text{B}) = 115$ MeV, the ^9Be spectrum is dominated by groups corresponding to the ^{14}N 5^+ state at $E_x = 8.96$ MeV, the 4^+ state at $E_x = 12.79$ MeV and to the 6^- state at $E_x = 15.10$ MeV, the levels also strongly populated in the $^{12}\text{C}(\alpha, d)^{14}\text{N}$ reaction (see reaction 18) (1966PO1E, 1967PO1E) and private communication). Groups to $^{14}\text{N}^*(0, 3.95, 5.83, 6.44, 10.81)$ are also reported (J.E. Poth, private communication). See also (1967VO1A). The angular distributions of the two strong groups are smoothly varying, exponential functions of angle, and have been fitted, in terms of a surface-diffuseness parameter, using a diffraction model (1965SA07). See also (1965DA1E, 1965GR1F) and (1964FL1D, 1964NA1E, 1967OG1A, 1968RO1D, 1969RO1G, 1969BR1D). For reaction (a), see (1969RO1G).



See (1962CH01).

Table 14.15: Energy levels of ^{14}N from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$

E_x ^a (MeV)	σ ^a (mb)	$\sigma/\sigma_{\text{g.s.}}$ ^b	Angular distribution ^c
0	5.8	1.00	d, α
2.31 ^d	d		α
3.94	4.8	0.31	d, α
4.91	4	0.16	α
5.10	5	1.32	d, α
5.69	4	0.11	α
5.83	6	0.97	d, α
6.21	7.0	0.24	α
6.44	13.0	1.15	d
7.03	4.0	0.18	d, α
7.97	5.0	0.16	d, α
8.47	8.3	0.53	d, α
9.00 ± 0.05 ^e	28.0	3.67	d, α
9.17	medium	(0.30)	
9.41	8.1	0.30	d, α
9.71	6.9	0.18	α
10.09	13.8	0.5	α
10.76 ± 0.02 ^f	medium	0.62	
11.06	weak	very weak ^h	
11.23	medium		
11.29	medium		
11.39	medium	very weak, broad	
11.51	weak		
11.97	weak		
12.21	weak	weak	
12.41	medium		
12.61	strong		
12.69	strong	strong, broad ($E_x = 12.76$)	
13.05 ± 0.02 ^g		very weak	
14.7 ^h		weak, broad	

Table 14.15: Energy levels of ^{14}N from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$
(continued)

E_x ^a (MeV)	σ ^a (mb)	$\sigma/\sigma_{\text{g.s.}}$ ^b	Angular distribution ^c
15.1 ± 0.1 ^e		strong, sharp	
15.5		weak	
16.0		weak	
16.3		medium	
17.1		medium	
17.7		medium	

^a From $^{12}\text{C}(^6\text{Li}, \alpha)$. $E(^6\text{Li}) = 20$ MeV ([1968ME10](#)).

^b Cross section (relative to ground state) integrated for $0^\circ - 90^\circ$ (c.m.), from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$, $E_\alpha = 42$ MeV ([1967ZA01](#)). The $T = 1$ states at $E_x = 2.31, 8.06, 8.63, 9.51, 10.41$ MeV were not observed: $\sigma/\sigma_{\text{g.s.}} < 0.0027, 0.027, 0.018, 0.30, 0.03$, respectively. For absolute cross sections at $E_\alpha = 53$ MeV, see ([1965PE03](#)).

^c d and α refer to angular distributions of the corresponding group from the $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and the $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$ reactions: see text for E_α , $E(^6\text{Li})$ and references.

^d See text.

^e ([1966RI04](#)).

^f $E_x = 10.85 \pm 0.02$ MeV ([1967ZA01](#)).

^g From $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ ([1967ZA01](#)): $\Gamma < 70$ keV.

^h Comments here and below are from ([1965PE03](#)).

22. $^{12}\text{C}(^{14}\text{N}, ^{12}\text{C})^{14}\text{N}$

At $E(^{14}\text{N}) = 28$ MeV, excitation of $^{14}\text{N}^*(0, 3.95, 4.91, 5.11)$ and some higher states is reported. The total cross section for excitation of the $T = 1$ state at $E_x = 2.31$ MeV is $< 60 \mu\text{b}$ ([1961HA04](#)). See also ([1968HU1H](#), [1969BR1D](#), [1969HE06](#)).

23. $^{12}\text{C}(^{18}\text{F}, ^{16}\text{O})^{14}\text{N}$ $Q_m = 2.745$

See ([1968RO1D](#)).

24. $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$ $Q_m = 7.550$

Table 14.16: Levels of ^{14}N from $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$ and $^{13}\text{C}(\text{p}, \text{p})^{13}\text{C}$ ^a

E_{p} (MeV \pm keV)	Γ_{lab} (keV)	l_{p}	$\omega\Gamma_{\gamma}$ (eV)	$J^{\pi}; T$	$^{14}\text{N}^*$	Refs.
0.4485 \pm 0.5	< 0.4	2	0.022	2^-	7.966	1960HE14
0.551 \pm 1	32.5 ± 1	0	9.2	$1^-; 1$	8.061	A, 1960HE14 , 1961KA04
1.012 \pm 2	≤ 0.21	4	$\cong 0.01$	$(4^-); 0$	8.489	1965DE19
1.150 \pm 2	7.5 ± 1	1	1.3	$0^+; 1$	8.617	A, 1961KA04 , 1966LA03
1.34 \pm 50	≈ 500	0	12.8	$0^-; 1$	8.80	A, 1961KA04 , 1966LA03
1.462 \pm 3	17 ± 2	2	0.72	$3^-; 1$	8.907	A, 1959WA04 , 1961KA04 , 1966LA03 , 1969WA1H
1.523 \pm 2	< 1		$\cong 0.003$	$5^+; 0$	8.963	1965DE19
1.540 \pm 3	9 ± 2	1, (3)	0.13	2^+	8.979	A, 1961KA04 , 1966LA03
1.701 \pm 1	< 1	2	$\cong 0.03$	$2^-; 0$	9.129	1965DE19
1.7476 \pm 0.9 ^b	0.075 ± 0.050		14.8	$2^+; 1$	9.172	B, 1959BO14 , 1959RO54 , 1959WA04 , 1960RO13 , 1960RO23 , 1963PR03
1.980 \pm 3	14 ± 3	2		$3^-, 2^-$	9.388	C, 1966LA03
2.110 \pm 3	44 ± 2	2	6.2	$2^-; 1$	9.508	D, 1966LA03
2.319 \pm 4	16 ± 3	1		1^+	9.702	C, 1966LA03
2.743 ^c	5.5	1	^d	$1^+, (2^+)$	10.096	1960RO23 , 1961KA04
2.885 \pm 10 ^c	85 ± 15	0, 2		$1(^-); 0$	10.228	C, 1961KA04 , 1963RO17
3.105 \pm 7 ^c	36 ± 3	1	17	$2^+; 1$	10.432	1957WI30 , 1960RO23 , 1961KA04 , 1964RO03
(3.12 \pm 30)	80 ± 10				(10.45)	C
3.20 ^c	150	0, 2		1^-	10.52	1961KA04
3.77	< 33		^f		11.05	1967RI1D
3.79	100			1^+	11.07	1960BA35 , 1961KA04 , 1967RI1D
3.98 ^c	12	2		3^-	11.24	1961KA04

Table 14.16: Levels of ^{14}N from $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$ and $^{13}\text{C}(\text{p}, \text{p})^{13}\text{C}$ ^a (continued)

E_{p} (MeV \pm keV)	Γ_{lab} (keV)	l_{p}	$\omega\Gamma_{\gamma}$ (eV)	$J^{\pi}; T$	$^{14}\text{N}^*$	Refs.
4.04 ^c	190	2	f	2^-	11.30	1960BA35, 1961KA04, 1967RI1D
4.14 ^c	30	1		1^+	11.39	1960BA35, 1961KA04
4.52	130		f	1^+	11.75	1960BA35, 1961BA09, 1967RI1D
4.60	≈ 100				11.82	1961BA09
5.31	55		f		12.48	1967RI1D
5.90	200		f		13.03	1967RI1D
6.20	≈ 1100		f		13.30	1967RI1D
8.0	e		f		15.0	1968BL1E, 1968RI1R
10.4	e		f,g		17.2	1968BL1E, 1968RI1R
12.6	e		f		19.2	1968BL1E, 1968RI1R
14.0	e		f		20.5	1968BL1E, 1968RI1R
16.1	≈ 500		f		22.5	1968BL1E, 1968RI1R
16.6	≈ 500		f		23.0	1968BL1E, 1968RI1R
17.4	≈ 500		g		23.7	1968BL1E, 1968RI1R

A: (1952SE01, 1953WO41, 1954MI05).

B: (1952SE01, 1953WO41, 1956MA87).

C: (1957ZI09, 1958ZI17).

D: (1952SE01, 1953WO41, 1957ZI09, 1958ZI17, 1959WA04).

^a See also Table 14.9.^b See also (1963BO07, 1966MA60).^c Reduced width for proton emission is of the order of 1% of the Wigner limit (1961KA04).^d $(2J + 1) \Gamma_{\gamma} = 0.5 \pm 0.2 \text{ eV}$, $\Gamma = 12 \pm 3 \text{ keV}$ (1960RO23).^e $\Gamma \approx 0.8$ to 1.5 MeV for these resonances (1968RI1R).^f Resonance in γ_0 .^g Resonance in γ_1 .

Observed resonances are displayed in Table 14.16. The decay schemes of various levels of ^{14}N , as derived from the γ -spectra in this and other reactions are exhibited in Table 14.9 ([1953WO41](#), [1959WA04](#), [1960HE14](#), [1960RO13](#), [1960RO23](#), [1963PR03](#), [1963RO17](#), [1964RO03](#), [1965DE19](#), [1967RI1D](#), [1968BL1E](#), [1968RI1R](#)).

The low-energy capture cross-section yields an extrapolated σ -factor at $E_p = 25$ keV (cm), $S_0 = 6.0 \pm 0.8$ keV · b ([1960HE14](#)). See also ([1963BR14](#), [1964FO1A](#), [1966BA2P](#), [1967FO1B](#)). The capture cross section rises from $(7.7 \pm 1.8) \times 10^{-10}$ b at $E_p = 100$ keV to $(9.8 \pm 1.2) \times 10^{-9}$ b at $E_p = 140$ keV ([1961HE02](#)).

Direct radiative capture of protons into the lowest three odd-parity states $^{14}\text{N}^*(4.91, 5.11, 5.69, 5.83)$ has been studied by ([1964TR04](#)). See also ([1963RO17](#)).

The angular distribution of the γ -rays at the $E_p = 0.45$ MeV resonance ($^{14}\text{N}^* = 7.97$ MeV) is consistent with $J^\pi = 2^-$ ([1960HE14](#)). The width of the $E_p = 0.55$ MeV resonance and the isotropy of the γ -rays ([1949DE1A](#), [1953WO41](#)) indicate s-wave formation of $^{14}\text{N}^*(8.06)$: $J^\pi = 1^-$ from $^{13}\text{C}(p, p)^{13}\text{C}$. The decay properties of this state (see Table 14.9) show that the ground state transition is an uninhibited E1 transition, and thus that $^{14}\text{N}^*(8.06)$ has $T = 1$ ([1953CL39](#)) but with a strong $T = 0$ admixture [as shown by a 2% $(8.06 \rightarrow 2.31)$ transition] ([1956LE28](#), [1957BR25](#), [1957WI27](#)). The strong transition $8.06 \rightarrow 5.69$ admits either E1, $\Delta T = 1$ or M1, $\Delta T = 1$. Since the transition $5.69 \rightarrow 2.31$ is observed, $^{14}\text{N}^*(5.69)$ cannot have $J^\pi = 0^+$, and 2^+ is excluded by the strength of the $8.62 \rightarrow 5.69$ transition. It appears then that $^{14}\text{N}^*(5.69)$ has $J = 1$: see ([1956LE28](#), [1957WI27](#), [1959WA04](#)).

Anomalies in the capture cross section (“midget resonances”) are observed at $E_p = 1.01, 1.52$ and 1.70 MeV: see Table 14.16. The predominant γ -decay at the first resonance, $^{14}\text{N}^*(8.49)$, is to the 5.10 MeV state. The angular distribution of the γ -rays and $\omega\Gamma_\gamma$ are consistent with $J^\pi = 4^-$; $T = 0$ for $^{14}\text{N}^*(8.49)$. The γ -decay of the second state, $^{14}\text{N}^*(8.96)$, is predominantly to the 6.44 MeV state: it’s $J^\pi = 5^+$. The third state decays primarily to the ground state: the angular distribution of the γ -rays is consistent with $J^\pi = 2^-$; $T = 0$ for $^{14}\text{N}^*(9.13)$ ([1965DE19](#)).

The narrow $E_p = 1.16$ MeV resonance, $^{14}\text{N}^*(8.62)$ [$J = 0^+$ from $^{13}\text{C}(p, p)^{13}\text{C}$] shows strong transitions to $^{14}\text{N}^*(0, 3.95, 5.69)$: hence $T = 1$ ([1959WA16](#)). The strong transition $8.62 \rightarrow 3.95$ requires dipole radiation and hence $J = 1$ for the latter ([1959WA04](#)) while the strength of the transition $8.62 \rightarrow 6.21$ and the angular correlation in the cascade $8.62 \rightarrow 6.21 \rightarrow \text{g.s.}$ is consistent with $J^\pi = 1^+$; $T = 0$ for $^{14}\text{N}^*(6.21)$: see ([1956GO1L](#), [1956GO39](#), [1957GA1B](#), [1957GO30](#), [1957WI27](#), [1959GA05](#)).

The $E_p = 1.25$ MeV resonance, $^{14}\text{N}^*(8.80)$ [$J^\pi = 0^-$ from $^{13}\text{C}(p, p)^{13}\text{C}$] has a large γ -width consistent with E1 radiation and $T = 1$ ([1953WI1A](#)). At $E_p = 1.47$ MeV, the plane polarization of 0.73 MeV γ -rays (from $5.83 \rightarrow 5.11$) suggests odd parity for both these states ([1962RO21](#)).

Angular correlation and angular distributions of γ -rays at the $E_p = 1.75$ MeV resonance, $^{14}\text{N}^*(9.17)$, indicates $J = 2^+$ for that state, $J = 3$ for $^{14}\text{N}^*(6.44)$ and $J = 2$ for $^{14}\text{N}^*(7.03)$ ([1959RO54](#), [1959WA04](#), [1960RO13](#), [1960RO23](#), [1963PR03](#)). The angular distribution of the 2.73 MeV γ -rays ($9.17 \rightarrow 6.44$) suggests odd parity for $^{14}\text{N}^*(6.44)$ ([1961SE03](#), [1963PR03](#)).

Marked variations are observed in the $(\cos \theta)$ term at $E_p = 2.75$ and 2.90 MeV [$^{14}\text{N}^*(10.09)$ and (10.23)] ([1960RO23](#)). The angular distribution of the γ -rays from the $10.23 \rightarrow 2.31$ transition ($E_p = 2.88$ MeV resonance) is consistent with $J^\pi = 1^+$, assuming a single state at 10.23

MeV: $M^2(M1)$ leads to a $T = 0$ assignment ([1963RO17](#)). At the $E_p = 3.11$ MeV resonance [$^{14}\text{N}^*(10.43)$], the angular distribution of ground state γ -rays is consistent with $J = 2$ ([1959WA16](#), [1960RO23](#)): the similar decay characteristics of $^{14}\text{N}^*(10.43)$ and of the $J^\pi = 2^+$; $T = 1$ state at $E_x = 9.17$ MeV suggest that the 10.43 MeV state is in fact also a $J^\pi = 2^+$; $T = 1$ level ([1964RO03](#)).

The yield of γ_0 for $E_p > 3.5$ MeV shows a steady rise marked at first by pronounced resonances and then by broad structures most of which are related to levels seen in the inverse reaction ($^{14}\text{N}(\gamma, p)^{13}\text{C}$). The γ_1 yield (to $^{14}\text{N}^*(2.31)$) is on the average only about a third the γ_0 yield with roughly constant cross section for $E_p = 7$ to 18 MeV. Only weak structure is observed ([1967RI1D](#), [1968BL1E](#), [1968RI1R](#)): see Table [14.16](#).

See also ([1959WA02](#), [1962WA1C](#)), ([1965FA1E](#), [1965MA1H](#); theor.) and ([1959AJ76](#)).

25. (a) $^{13}\text{C}(p, p)^{13}\text{C}$	$E_b = 7.550$
(b) $^{13}\text{C}(p, p')^{13}\text{C}^*$	

The elastic scattering has been studied for $E_p = 0.14$ to 0.75 MeV ([1960HE14](#)), 0.45 to 1.60 MeV ([1954MI05](#)), 1.0 to 2.6 MeV ([1966LA03](#)), 1.5 to 3.4 MeV ([1957ZI09](#), [1958ZI17](#)), 2.6 to 5.0 MeV ([1961KA04](#)), 5 to 8.1 MeV ([1965BA1W](#)), and 7 to 11 MeV ([1966SH1H](#)): parameters of resonances observed in this reaction and in the $^{13}\text{C}(p, \gamma)^{14}\text{N}$ reaction are displayed in Table [14.16](#). Angular distributions of elastically scattered protons have been measured at many of these energies.

The yield of γ -rays in reaction (b) has been measured for $E_p = 3.6$ to 5.0 MeV: the 3.1 MeV γ -yield shows broad resonances at $E_p = 3.80, 4.1, 4.14$ and 4.60 MeV while the 3.7 MeV γ -yield shows one resonance at $E_p = 4.52$ MeV ($^{14}\text{N}^* = 11.07, 11.30, 11.39, 11.82$ and 11.75 MeV, respectively) ([1960BA35](#), [1961BA09](#)): see Table [14.16](#). The angular distribution of inelastic protons to $^{13}\text{C}^*(3.68)$ at the $E_p = 4.52$ MeV resonance leads to the assignment $J^\pi = 1^+$ for $^{14}\text{N}^*(11.75)$ ([1961BA09](#)). The excitation functions for proton groups to $^{13}\text{C}^*(3.09, 3.68, 3.85)$ have been measured for $E_p = 5$ to 11 MeV ([1965BA1W](#), [1966SH1H](#)). See also ([1962BE04](#)).

Polarization measurements for elastically scattered protons are reported at $E_d = 7$ MeV by ([1969GU02](#)) and at $E_p = 14.5$ MeV by ([1962RO20](#), [1965RO22](#), [1966RO1B](#), [1966RO1R](#)). At $E_p = 32.9$ MeV, the polarization and asymmetry in the elastic scattering have been compared. They are equal to within 1%, and there is therefore no evidence for violation of time reversal invariance for strong interactions, at least for that part of the force which flips the spin of the proton ([1968GR1K](#), [1969MA2D](#)).


See also ([1962WA1C](#), [1963GE03](#), [1964FO02](#), [1967AR1D](#)), ([1967HO1L](#), [1969WA11](#)), ([1964PF1A](#), [1966AM1B](#); theor.) and ([1959AJ76](#)).

26. $^{13}\text{C}(p, n)^{13}\text{N}$	$Q_m = -3.003$	$E_b = 7.550$
--	----------------	---------------


The yield of neutrons has been measured from threshold to $E_p = 13.7$ MeV: see (1959AJ76) and (1959BR06, 1960BA35, 1961DA09, 1961WO03, 1966DI03). Observed resonances are displayed in Table 14.17. Angular distributions have been measured at many energies: over much of the energy range (below $E_p = 10$ MeV) there is pronounced backward peaking with a secondary maximum near 50° and no forward peak (1961DA09). The polarization of neutrons corresponding to $^{13}\text{N}(0)$ has been studied for $E_p = 6.9$ to 12.3 MeV (1964WA1G, 1965WA02). See also (1962BE04, 1964CA1F, 1965VA23, 1969BA1N) and ^{13}N .



The yield of ground state deuterons has been determined for $E_p = 5$ to 11 MeV (1965BA1W, 1966SH1H). See also (1962BE04). A polarization measurement at $E_p = 7$ MeV is reported by (1969GU02). See also (1968TA1V) and ^{12}C in (1968AJ02).



See (1968TA1V).



Alpha-particle yields have been measured for $E_p = 7$ to 12 MeV (1963PE07, 1964FO02, 1966SH1H).



Observed neutron groups are exhibited in Table 14.18. Recent angular distribution measurements are reported by (1969CH04: 0.5 to 0.8 MeV; n_3), (1961JA09: 1.2 MeV; $n_0 \rightarrow n_6$), (1961ZD01, 1964WI03: 1.3 to 2.5 MeV; n_0, n_1, n_2), (1963BE05: 3.9 MeV; $n_0, n_1, n_2, n_3 + n_4$), (1966FU10, 1966SI02, 1967FU04: 5.5 and 6 MeV; many groups), (1968CO24: 7 to 12 MeV; n_0, n_1, n_2), and (1969VE1D: 11.8 MeV; n_0, n_1, n_2). See also (1960VA11, 1962SH19, 1963DE19). Comparison of relative spectroscopic factors here and in the $^{13}\text{C}(^3\text{He}, \text{d})^{14}\text{N}$ reaction [see Table 14.19] shows smaller values for the $T = 1$ state [$^{14}\text{N}^*(2.31)$] in this reaction than in the ($^3\text{He}, \text{d}$) reaction (1966SI02, 1967FU04). Simple DWBA calculations would suggest that the factors would be the same in both proton pickup reactions. However, the dependence of the cross section magnitude on the T of the final state [the $t \cdot T$ term] appears to be energy dependent: see Table 14.19 (1968CO24). See also (1967LE1F).

Table 14.17: Resonances in $^{13}\text{C}(\text{p}, \text{n})^{13}\text{N}$
^a(1961DA09)

E_{p} (MeV)	Γ (keV)	$^{14}\text{N}^*$ (MeV)
3.76 ± 0.05	100	11.04
3.98 ± 0.02	30	11.24
4.05		11.31
4.15 ± 0.02	40	11.40
4.5 ± 0.1	100	11.7
4.7 ± 0.1	150	11.9
5.03 ^b		12.22
(5.44 ± 0.03)	(60)	(12.60)
5.53 ± 0.03	50	12.68
5.72 ± 0.03	60	12.86
6.20 ± 0.04	70	13.30
6.67 ± 0.13	250	13.74
7.0 ± 0.1	150	14.1
7.3		14.3
7.85 ± 0.08	150	14.84
7.93 ± 0.03	50	14.91
8.03 ± 0.03	50	15.00
8.7 ± 0.2	350	15.6
9.3 ± 0.1	150	16.2
10.2 ± 0.2	400	17.0
11.4 ± 0.3	600	18.1

^a See also (1960BA35) and Table 14.9 in (1959AJ76).

^b (1959GI47).

Table 14.18: ^{14}N levels from $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$ and $^{13}\text{C}(^3\text{He}, \text{d})^{14}\text{N}$

E_x (MeV)	$J^\pi; T$	l_p ^a	l ^c
0	$1^+; 0$	1	1
2.31	$0^+; 1$	1	1
3.95	$1^+; 0$	1	1
4.91	$(0^-); 0$	0	0
5.11	$2^-; 0$	2	2
5.69	$1^-; 0$	0	0
5.83	$3^-; 0$	2	2
6.21	$1^+; 0$	unspec.	1
6.44	$3^+; 0$	iso.	1
7.03	$2^+; 0$	1	1
7.97	$2^-; 0$	b	d
8.06	$1^-; 1$	0	
8.91	$3^-; 1$	(1, 2)	d
8.98	$2^+; (0)$	(1, 2)	

^a $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$: $E_{\text{d}} = 5.5$ and 6 MeV ([1966FU10](#)).

See also Table 14.10 in ([1959AJ76](#)) and ([1961JA09](#), [1963BE05](#), [1967FU04](#), [1968CO24](#)).

^b Group was masked by contaminant.

^c $^{13}\text{C}(^3\text{He}, \text{d})^{14}\text{N}$: $E(^3\text{He}) = 15$ MeV ([1966HO15](#)).

^d See ([1969HO23](#)).

In the range $E_{\text{d}} = 0.4$ to 4.2 MeV, a single strong neutron threshold occurs at $E_{\text{d}} = 0.422 \pm 0.005$ MeV ($^{14}\text{N}^* = 5.691 \pm 0.006$) ([1955MA76](#)). See also ([1965MA1K](#)).

Observed γ -rays attributed to transitions in ^{14}N are shown in Table 14.20 ([1952TH24](#), [1955BE62](#), [1955MA36](#), [1958RA13](#), [1966AL10](#)). The decay of $^{14}\text{N}^*(5.69)$ is via $^{14}\text{N}^*(2.31)$ [$63 \pm 2\%$] and directly to $^{14}\text{N}(0)$ [$37 \pm 2\%$]. τ_m [$^{14}\text{N}^*(4.91)$] < 0.5 psec ([1963AL21](#)) [see also Tables 14.9 and 14.13]. The angular correlation of internal pairs conclusively establish the parities of $^{14}\text{N}^*(4.91, 5.10, 5.69)$ as odd ([1964WA05](#)).

See also ([1959CH28](#), [1959GO78](#), [1966EV1B](#)), ([1959NE1B](#), [1960AB02](#), [1964BA1G](#), [1966BA2R](#), [1966ST1L](#), [1967BA2J](#); theor.) and ([1959AJ76](#)). See also ^{15}N .

$$\begin{aligned}
Q_0 &= 2.070 \pm 0.015 \text{ (1962SH21)}; \\
Q_0 &= 2.048 \pm 0.014 \text{ (1963SP1B)}; \\
Q_0 &= 2.050 \pm 0.015; \\
Q_1 &= -0.265 \pm 0.015 [E_x = 2.315 \pm 0.022] \text{ (1959YO25).}
\end{aligned}$$

Table 14.19: Spectroscopic factors for ^{14}N states ^a

E_x (MeV)	$J^\pi; T$	$^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$				$^{13}\text{C}({}^3\text{He}, \text{d})^{14}\text{N}$			Theory ^f
		E_{d} (MeV)	3.9 ^b	5.5 ^c	6.0 ^d	7.0 to 12.0 ^a	$E({}^3\text{He})$ (MeV)		
		13 ^c	15 ^e	17 ^c					
0	$1^+; 0$	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1
2.31	$0^+; 1$	0.97	1.1	1.15	0.920 ± 0.052	1.93	1.53	1.6	1.24
3.95	$1^+; 0$	0.72	0.8	0.57	0.551 ± 0.032	0.52	0.39	0.5	0.29

^a Normalized to 1.0 for the ground state (1968CO24).

^b (1963BE05).

^c (1966SI02).

^d (1966FU10).

^e (1966HO15).

^f (1967CO32).

Angular distributions have been obtained at $E({}^3\text{He}) = 13$ and 17 MeV (1966SI02: $^{14}\text{N}^*(0, 2.31, 3.95)$), 15 MeV (1966HO15, 1969HO23: $^{14}\text{N}^*(0 \rightarrow 9.17)$), 17.8 MeV (1966EC1B: $^{14}\text{N}^*(0, 2.31, 3.95)$) and 22 MeV (1969MA1R: $^{14}\text{N}^*(0, 2.31)$). Relative spectroscopic factors for the first three states have been compared with those obtained in the $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$ reaction (1966HO15, 1966SI02): see reaction 30 and Table 14.19. See also (1967LE1F). Spectroscopic factors for the higher states have also been obtained by (1966HO15), from a DWBA analysis. l -values for the observed groups are in agreement with those obtained by (1966FU10) in the (d, n) reaction: see Table 14.18. See also (1965HE01, 1967BA1D).



Angular distributions have been obtained at $E_\alpha = 46$ MeV of the tritons corresponding to the first seven states of ^{14}N : the data are consistent with the assumption of a direct interaction in which a proton is transferred from the incident α -particle (1969FO1C).

Table 14.20: Gamma rays from $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$

(1952TH24) ^a E_γ ^e (keV)	(1955BE62) ^b E_γ ^f (keV)	(1955MA36) ^c E_γ ^e (keV)	(1958RA13) ^d E_γ ^f (keV)	Assignment $^{14}\text{N}^*$
725 \pm 4		729 \pm 3	728.3 \pm 1.0 ^g	5.83 \rightarrow 5.11
1638 \pm 8				3.95 \rightarrow 2.31
2310 \pm 12				2.31
3381 \pm 13	3410 \pm 40			5.69 \rightarrow 2.31
	3920 \pm 70 ^g	3910 \pm 50		3.95
	4940 \pm 40	4930 \pm 40	4897 \pm 25	4.91
5052 \pm 25	5100 \pm 50	5130 \pm 30	5102 \pm 25	5.11
5690 \pm 50	5720 \pm 40	5730 \pm 30	5669 \pm 25	5.69
			5833 \pm 30 ^e	5.83
	6490 \pm 60	6450 \pm 50	6419 \pm 30	6.44
	7050 \pm 40		7012 \pm 25	7.03

^a $E_{\text{d}} = 1.2, 1.6 \text{ MeV}$.

^b $E_{\text{d}} = 2, 4 \text{ MeV}$.

^c $E_{\text{d}} = 1.4 \text{ MeV}$.

^d $E_{\text{d}} = 4.5 \text{ MeV}$.

^e Not corrected for Doppler shift.

^f Includes $\approx 0.5\%$ Doppler correction.

^g (1966AL10).

$$33. \ ^{13}\text{C}(^{11}\text{B}, ^{10}\text{Be})^{14}\text{N} \quad Q_{\text{m}} = -3.678$$

At $E(^{11}\text{B}) = 113.5 \text{ MeV}$, ^{10}Be groups are observed corresponding to $^{14}\text{N}^*(5.69+5.83)$, $(8.80+8.91)$ and possibly to $^{14}\text{N}^*(12.2)$. The two lower transitions correspond to the addition of $s_{1/2}$ and $d_{5/2}$ protons to the ^{13}C target core. States which were not excited would have required the excitation of the target core in addition to the transfer of a nucleon (1967PO13). See also (1969BR1D).

$$34. \ ^{13}\text{C}(^{14}\text{N}, ^{13}\text{C})^{14}\text{N}$$

See (1968HU1H).

$$35. \ ^{14}\text{C}(\beta^-)^{14}\text{N} \quad Q_{\text{m}} = 0.156$$

See ^{14}C .



Neutron thresholds have been observed at $E_p = 671.5 \pm 0.5$ and 3149.6 ± 1.1 keV ([1956SA06](#)) and at $E_p = 4910 \pm 8$ keV ([1960BA34](#)) corresponding to the ground state of ^{14}N and to excited states at 2.3119 ± 0.0012 and 3.952 ± 0.008 MeV. Angular distributions of the neutrons corresponding to $^{14}\text{N}^*(0, 2.31, 3.95)$ have been obtained for $E_p = 6$ to 14 MeV. At the highest energies these have been analyzed by using finite range DWBA, taking into account various isospin and spin factors. The ground state transition is not inhibited, whereas the ^{14}C β -decay is. This requires spin-flip mechanisms such as a tensor interaction or particle exchange ([1967WO05](#)). See also ([1969MA1G](#)). Angular distributions are also reported at $E_p = 20$ and 30 MeV ([1969SA1M](#); n_0, n_1, n_2). See also ^{15}N .



At $E(^3\text{He}) = 44.8$ MeV, triton groups are observed corresponding to all the known levels of ^{14}N with $E_x < 7.1$ MeV. Triton groups were also seen to unresolved states with $E_x = 8.0 - 9.5$ MeV, to $^{14}\text{N}^*(10.43)$ and to excited states with $E_x = 12.49 \pm 0.04, 12.83 \pm 0.05$ and 13.70 ± 0.04 MeV. Angular distributions were obtained for nine of the triton groups and analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture. Dominant $L = 0$ transitions are found to $^{14}\text{N}^*(2.31, 3.95, 13.7)$, $L = 1$ to $^{14}\text{N}^*(5.11)$, $L = 2$ to $^{14}\text{N}^*(0, 7.03, 10.43)$ and $L = 3$ to $^{14}\text{N}^*(5.83)$ ([1967BA13](#), [1968BA1E](#), [1969BA06](#)). See also ([1969MA1G](#)) and reaction 45.

38. (a) $^{14}\text{N}(\gamma, \text{n})^{13}\text{N}$	$Q_m = -10.553$
(b) $^{14}\text{N}(\gamma, \text{p})^{13}\text{C}$	$Q_m = -7.550$
(c) $^{14}\text{N}(\gamma, \text{d})^{12}\text{C}$	$Q_m = -10.272$
(d) $^{14}\text{N}(\gamma, \text{pn})^{12}\text{C}$	$Q_m = -12.497$
(e) $^{14}\text{N}(\gamma, \alpha)^{10}\text{B}$	$Q_m = -11.613$
(f) $^{14}\text{N}(\gamma, \text{n}\alpha)^9\text{B}$	$Q_m = -20.051$
(g) $^{14}\text{N}(\gamma, \text{p}\alpha)^9\text{Be}$	$Q_m = -18.201$

The total absorption over the range $E_\gamma = 9$ to 31 MeV is dominated by a single peak at 22.5 MeV [estimated $\sigma \approx 29$ mb, $\Gamma \approx 2 - 3$ MeV] and appreciable strength extending beyond 30 MeV. The cross section cannot be accounted solely by the (γ, n) and (γ, p_0) processes: particle unstable excited states of ^{13}C , ^{13}N are believed involved ([1969BE92](#)). Over the interval $E_\gamma = 20.0$ to 20.5

MeV, the average cross section is 10.5 ± 4 mb ([1959CA1C](#), [1960CA09](#)). See also ([1968CO13](#)). The cross section for neutron production, reactions (a) and (d), exhibits maximum at $E_\gamma = 22.5$ MeV, $\Gamma = 3.2$ MeV, $\sigma = 15.3$ mb ([1954FE16](#)) [$\Gamma = 3.8$ MeV, $\sigma = 14.5$ mb ([1960FA06](#))]. The cross section for reaction (a) shows a maxima at $E_\gamma = 11.7, 13.2, 15.2, 19.5$ and 22.8 MeV ([1960KI02](#)). [The peak cross sections for the two highest energy maxima are ≈ 1.8 and 2.8 mb, respectively; the widths are $\approx 2 - 3$ MeV, estimated from the published curve.] Breaks in the (γ , n) activation curve are reported at $E_\gamma = 11.07$ MeV ([1960GE06](#)) and $11.49, 11.61, 12.39, 12.92, 13.28, 13.87, 14.62, 15.25, 16.35, 18.05$ and 19.10 MeV ([1959MU08](#), [1960MU02](#): ± 50 keV). See also ([1959FU1A](#), [1960BA15](#), [1960KO05](#), [1960SA09](#), [1962GO1E](#), [1962KO23](#), [1963CO1D](#), [1963FU06](#), [1970LO1A](#)).

Studies of reaction (b) indicate the involvement of $^{14}\text{N}^*(8.06, 9.17, 10.43)$ ([1956WR22](#), [1960WA17](#)) as well as of ^{14}N states with $E_x = 11.8, 13.0$ and 15.2 MeV ([1960WA17](#)). See also ([1959FU1A](#), [1960BA15](#), [1960KO05](#), [1962GO1E](#), [1962KO23](#), [1963FI1B](#), [1964ED01](#), [1964KO1D](#)). For reaction (c), see ([1964ED01](#)). For reaction (d) see also ([1959RE1A](#), [1960BA15](#), [1962GO1E](#), [1962KO23](#), [1963KO1B](#), [1964KO1D](#)).

For reactions (e), (f) and (g) see ([1956LI05](#), [1958MA1A](#), [1960BA15](#), [1960KO05](#), [1962GO1E](#), [1962MO16](#), [1964ED01](#), [1964TO1B](#)). See also ([1959AJ76](#)).

39. $^{14}\text{N}(\gamma, \gamma)^{14}\text{N}$

Resonant absorption measurements give $\Gamma = 72 \pm 10$ eV ([1965LU05](#)), 77 ± 12 eV ([1959HA11](#)) for $^{14}\text{N}^*(9.17)$: $\omega\Gamma_\gamma = 14.5 \pm 2$ eV consistent with dipole radiation ([1959HA11](#)). ([1966SW01](#)) finds $E_x = 7.029 \pm 0.006$ MeV for $^{14}\text{N}^*(7.03)$. The angular distribution of the scattered radiation is consistent with $J = 2$. Assuming that $^{14}\text{N}^*(7.03)$ decays entirely to the ground state [actually 98%: see Table 14.13], $\tau_m = 5.4 \pm 0.5$ fsec ([1966SW01](#)). The mean lifetime for $^{14}\text{N}^*(2.31)$ is 77 ± 18 fsec ([1961SW01](#)), 97 ± 30 fsec ([1964BO22](#)). For $^{14}\text{N}^*(8.06)$, ([1956GR17](#), [1958GR97](#)) finds $\Gamma_\gamma = 10.5 \pm 6$ eV. See Tables 14.9, 14.13 and 14.21.

An elastic scattering study shows a broad (several MeV wide) maximum centered around $E_\gamma = 24$ MeV and indicates a secondary maximum around 30 MeV ([1967LO1B](#)).

40. $^{14}\text{N}(e, e)^{14}\text{N}$

The r.m.s. radius of ^{14}N at $E_e = 400$ MeV is 2.58 ± 0.05 fm ([1968DA1Q](#)). See also ([1964BI04](#)). See also ([1959ME24](#)). Measurements of magnetic form factors at $E_e = 100$ to 180 MeV favor jj -coupling for $^{14}\text{N}(0)$ ([1966RA29](#)).

Inelastic scattering ($\theta = 180^\circ$) gives evidence for the excitation of $^{14}\text{N}^*(8.91, 9.17, 10.43)$: the ground state Γ_γ are given in Table 14.21 ([1962ED02](#), [1963BA19](#), [1966KO08](#), [1968CL05](#)). In addition ([1968CL05](#)) report the excitation of a state with $E_x = 11.01 \pm 0.07$ MeV and ([1966KO08](#)) report structure at $E_x = 11.7$ and 12.7 MeV. Partial Γ_γ [for cascade transitions of $^{14}\text{N}^*(9.17,$

Table 14.21: Gamma widths ^a of unbound levels from $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$, $^{14}\text{N}(\gamma, \gamma)^{14}\text{N}$ and $^{14}\text{N}(\text{e}, \text{e})^{14}\text{N}$

E_x (MeV)	$J^\pi; T$	Γ_γ to g.s. (eV)	Multipol.	Reaction	Refs.
8.06	$1^-; 1$	10.5 ± 6	E1	(γ, γ)	(1956GR17, 1958GR97)
8.91	$3^-; 1$	$(6.6 \pm 2.2) \times 10^{-3}$	M2	(e, e)	(1968CL05)
		$(16 \pm 4) \times 10^{-3}$		(e, e)	(1966KO08)
9.17	$2^+; 1$	7.7 ± 0.9	M1	(e, e)	(1968CL05)
		8.7 ± 1.5		(γ, γ)	(1959HA11)
		9 ± 3		(e, e)	(1962ED02)
		13 ± 3		(e, e)	(1966KO08)
		4.6 ± 1.4		(e, e)	(1963BA19)
10.43	$2^+; 1$	12.1 ± 1.5	M1	(e, e)	(1968CL05)
		16.4 ± 3.0		(e, e)	(1966KO08)
		17 ± 5		(p, γ)	(1957WI30)
		20 ± 6		(e, e)	(1962ED02)
		4.9 ± 1.5		(e, e)	(1963BA19)

^a See also Table 14.9.

10.43)] have been obtained by (1968CL05) and are shown in Table 14.9. See also (1963GO04). Inelastic scattering is also reported to $^{14}\text{N}^*(2.3, 3.95, 5.1, 5.85, 7.05, 8.0)$ (1964BI09). See also (1964BI04). See (1962BA1D) for a general discussion. See also (1958CA1B, 1960PA08, 1963GU1A, 1967KA1A, 1969VI02).

41. $^{14}\text{N}(\text{n}, \text{n}')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered neutrons are displayed in Table 14.23. See also (1959AJ76) and ^{15}N .

Observed gamma rays are displayed in Table 14.23 (1969DI1B, 1969NY1A). See also (1959HA13, 1968CO1W). Lifetime measurements are reported in Table 14.22 (1969NY1A).

See also ^{15}N , (1961AS1B, 1963MO04, 1963OP1A, 1964EN1B, 1964MO1D, 1964PE20, 1965VA1K, 1966MO1C) and (1965JO07, 1965TA07, 1968CA1A, 1969WA11).

42. (a) $^{14}\text{N}(\text{p}, \text{p}')^{14}\text{N}^*$

$$\begin{array}{ll} (\text{b}) \quad ^{14}\text{N}(\text{p}, 2\text{p})^{13}\text{C} & Q_m = -7.550 \\ (\text{c}) \quad ^{14}\text{N}(\text{p}, \text{pd})^{12}\text{C} & Q_m = -10.272 \end{array}$$

Table 14.22: Gamma rays from $^{14}\text{N}(\text{n}, \text{n}')^{14}\text{N}^*$

E_γ^{a} (MeV \pm keV)	E_γ^{d} (MeV)	Transition
$0.727 \pm 3^{\text{b}}$	0.728	$5.83 \rightarrow 5.11$
$1.634 \pm 5^{\text{b}}$	1.622	$3.95 \rightarrow 2.31$
$2.312 \pm 3^{\text{b}}$	2.313	$2.31 \rightarrow \text{g.s.}$
	2.49	$6.44 \rightarrow 3.95$
2.792 ± 3	2.792	$5.11 \rightarrow 2.31$
	3.372	$5.69 \rightarrow 2.31$
	3.895	$6.20 \rightarrow 2.31$
$4.919 \pm 10^{\text{c}}$	4.91	$4.91 \rightarrow \text{g.s.}$
5.106 ± 3	5.105	$5.11 \rightarrow \text{g.s.}$
	5.685	$5.69 \rightarrow \text{g.s.}$
$5.835 \pm 5^{\text{b}}$	5.833	$5.83 \rightarrow \text{g.s.}$
$6.426 \pm 10^{\text{c}}$	6.44	$6.44 \rightarrow \text{g.s.}$
$7.021 \pm 10^{\text{c}}$	7.03	$7.03 \rightarrow \text{g.s.}$

^a (1969NY1A): $E_n = 15$ MeV.

^b $E_x(3.95)$ is then 3.947 ± 0.005 MeV and $E_x(5.83)$ is 5.833 ± 0.003 MeV.

^c These γ -ray energies have been corrected for Doppler shift.

^d (1969DI1B): $E_n = 8.6$ MeV: gamma-ray peaks located in terms of the 2.313 and 5.105 MeV γ -rays. No attempt was made to obtain accurate γ -ray energies.

Angular distributions of elastically and inelastically scattered protons have been measured and analyzed at a number of energies: see Table 14.23. See also ^{15}O and (1968OD1B). Observed inelastic proton groups are shown in Table 14.24. The proton groups corresponding to $^{14}\text{N}^*(7.40, 7.60)$ reported by (1956BU16) are spurious: see (1964DO03, 1964EA04). See also (1963BR14, 1966ME1L). (1965DE21) reports the excitation of ^{14}N states with $E_x = 11.2 \pm 0.2$ and 12.8 ± 0.4 MeV, and the measurement of the angular distribution of the protons from the decay of $^{14}\text{N}^*(11.2)$. See also (1957HO34, 1961CL09, 1969CU1D) and (1962KA1E, 1962WA1D, 1965TA07, 1969MA1G, 1969WA11). See also (1959AJ76) and (1960WA12).

Reaction (b) at $E_p = 19$ MeV proceeds at least in part through an intermediate state in ^{14}N at $E_x \approx 11.2$ MeV (1965DE21). See also (1961CL09, 1965RI1A, 1966TY01) and (1965BE1E, 1965DE1P, 1967JA1E, 1967KO1P, 1969KO1J; theor.) and ^{13}C . For reaction (c) see (1961CL09) and (1964BA1P, 1966JA1A; theor.).

43. $^{14}\text{N}(\text{d}, \text{d}')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered deuterons have been obtained at

Table 14.23: $^{14}\text{N}(\text{n}, \text{n})$, (p, p) , (d, d) , $(^3\text{He}, ^3\text{He})$ and (α, α) angular distribution studies

(a) *Angular distribution of neutrons*

E_{n} (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.
1.6 – 3.2	g.s.	(1966FO1D, 1966FO1E)
5.0 – 6.5	g.s.	(1960SM02)
6.8 – 14.0	g.s.	(1967BA03)
14	g.s.	(1961ST22, 1963BA46)
14	2.31	(1961BO28)
14.1	g.s.	(1966BE1P)
14.8	g.s.	(1967LU1B)

(b) *Angular distribution of protons*

E_{p} (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.
6 – 9	2.31	(1968SH11)
6 – 11.4	g.s., 3.95	(1968BO36, 1968SH11)
7.7, 8.5, 10.5, 14.1	g.s., 2.31	(1960OD01)
9.5	g.s., 3.95	(1954FR38, 1957GI14)
9.8	g.s.	(1957HI56)
9.9 – 11.4	5.11, 5.83	(1968BO36)
10.2	3.95 → 7.03	(1964DO03)
10.5, 14.1	3.95	(1960OD01)
12.2	g.s.	(1968RI1Q)
19.4	g.s.	(1956VA1B, 1957VA1B)
20	g.s.	(1955CH1A)
21	g.s.	(1969BA23)
24.8	2.31	(1967AU1E)
31	g.s.	(1964KI06)
142	g.s.	(1961TA06)
155	g.s., 2.31, 3.95, 4.91 + 5.11, 5.69 + 5.83	(1968GE04)

(c) *Angular distribution of deuterons*

E_{d} (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.

Table 14.23: $^{14}\text{N}(\text{n}, \text{n})$, (p, p) , (d, d) , $(^3\text{He}, ^3\text{He})$ and (α, α) angular distribution studies (continued)

0.7 – 2.1	g.s.	(1963SE05)
1.2 – 3.1	g.s.	(1969GO14)
1.3 – 3.2	g.s.	(1969BE08)
5.5 – 10.2	2.31, 3.95	(1968DU1E)
8	g.s.	(1952GI01)
11	g.s.	(1960TA08)
11.8	g.s.	(1967FI07)
13.6	g.s.	(1963NE1C, 1964NE1D)
14.2	g.s., 3.95, 4.91, 5.11	(1966NG01)
21	g.s.	(1959FI30)
27	g.s.	(1962ER03)
28	g.s.	(1968GA13)
28.5	g.s.	(1966VI1A)
52	g.s., 3.95, 4.91 + 5.11, 5.69 + 5.83, 7.03	(1968HI09, 1968HI14)

(d) Angular distribution of ^3He particles

$E(^3\text{He})$ (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.
17.4, 25.4, 36.6	g.s.	(1965AR1E)
25.7	g.s.	(1964SE05)
29	g.s., 3.95, 4.91 + 5.11	(1962SE13)
44.6	g.s., 2.31, 3.95, 4.91, 5.11, 5.69, 5.83, 7.03, 11.24, 12.77	(1968BA1E, 1969BA06)

(e) Angular distribution of α particles

E_α (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.
11.4 – 12.7	2.31	(1966CH1E)
19.2	g.s., 3.95, 4.91 + 5.11	(1961PL01)
20.1 – 23.3	g.s.	(1967BO1M)
25.7 – 32.9	g.s., 3.95	(1963NO1B)
37.9	g.s.	(1960AG01)
40	g.s.	(1959YA01)
40.5	g.s., 3.95, 4.91, 5.11, 5.69, 5.83, 6.21, 6.44, 7.03	(1966HA19)
56	g.s.	(1969GA11)
65	g.s.	(1964HA16)

Table 14.23: $^{14}\text{N}(\text{n}, \text{n})$, (p, p) , (d, d) , $(^3\text{He}, ^3\text{He})$ and (α, α) angular distribution studies (continued)

104	g.s.	(1968HA1D, 1969HA14)
-----	------	-----------------------------

a number of energies: see Table 14.23. See also ^{16}O in (1971AJ02). Inelastic deuteron groups are displayed in Table 14.24. The deuteron group to the 0^+ ; $T = 1$ state at $E_x = 2.31$ MeV is isospin forbidden: its intensity is 1 – 3 % of the deuteron group to $^{14}\text{N}^*(3.95)$ for $E_d = 7.7$ to 10.2 MeV (1968DU1E). See also (1953BO70). The deuteron group to the $T = 1$ state $^{14}\text{N}^*(8.06)$ is also not seen: see Table 14.24. See also general discussions in (1960WA12, 1966BR1G, 1968NO1C). See also (1968ME1E, 1969CU08).

44. $^{14}\text{N}(\text{t}, \text{t})^{14}\text{N}$

See (1964SC09).

45. $^{14}\text{N}(^3\text{He}, ^3\text{He}')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered ^3He particles have been determined by (1962SE13, 1964SE05, 1965AR1E, 1969BA06): see Table 14.23. See also (1967KN1B).

At $E(^3\text{He}) = 44.6$ MeV, twelve ^3He groups are reported corresponding to states in ^{14}N : see Table 14.24 (1969BA06). The angular distributions were analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture. A comparison of the cross sections of the reactions $^{14}\text{N}(^3\text{He}, \text{t})^{14}\text{O}_{\text{g.s.}}$, $^{14}\text{N}(^3\text{He}, ^3\text{He}')^{14}\text{N}(2.31)$ and $^{14}\text{C}(^3\text{He}, \text{t})^{14}\text{N}(0)$ [which all correspond to transitions between identical initial and final states] shows that they are roughly equal, as would be expected from charge independence, once detailed-balance, isospin coupling and phase-space corrections have been applied (1969BA06). See also (1968HO1C, 1968LE1G, 1969RA1B).

46. (a) $^{14}\text{N}(\alpha, \alpha')^{14}\text{N}^*$

$$(b) \quad ^{14}\text{N}(\alpha, \alpha\text{p})^{13}\text{C} \qquad Q_m = -7.550$$

$$(c) \quad ^{14}\text{N}(\alpha, \alpha\text{d})^{12}\text{C} \qquad Q_m = -10.272$$

Angular distributions of elastically and inelastically scattered α -particles have been measured for $E_\alpha = 11$ to 104 MeV: see Table 14.23. The intensity of the isospin-forbidden α_1 group to $^{14}\text{N}^*(2.31)$ is low. The highest intensities reported are 0.18 of the α_0 group and 0.4 of the α_2 group for $E_\alpha = 11.4$ to 12.7 MeV (1966CH1E). Generally the intensity of the α_1 group is much smaller than that, typically a few percent of the α_0 or α_2 group: see (1959AJ76, 1966HA19): see also Table 14.23.

Table 14.24: ^{14}N levels from $^{14}\text{N}(\text{p}, \text{p}')$, (d, d') , $(^3\text{He}, ^3\text{He}')$ and (α, α')

$^{14}\text{N}^*(\text{MeV} \pm \text{keV})$				L^k
(p, p')	(d, d')	$(^3\text{He}, ^3\text{He}')$	(α, α')	
$2.313 \pm 5^{\text{a}}$	see text	h	see text	
$3.945 \pm 5^{\text{a}}$	f	h	j	2
$4.910 \pm 10^{\text{a}}$	f	h	j	1, 3
$5.104 \pm 10^{\text{a}}$	f	h	j	1, 3
$5.69 \pm 30^{\text{b}}$	f	h	j	1, 3
$5.83 \pm 30^{\text{b}}$	f	h	j	1, 3
$6.23 \pm 20^{\text{b}}$	f	h	j,l	
$6.46 \pm 20^{\text{b}}$	f		j,l	
$(6.60 \pm 40)^{\text{b}}$				
$7.03 \pm 20^{\text{b}}$	f	h	j	2
7.97 ^c			m	
8.06 ^c		$8.0 \rightarrow 11.0^{\text{i}}$		
9.2 ^d				
$11.2 \pm 200^{\text{e}}$	f,g	$11.22 \pm 50^{\text{h}}$		
$12.8 \pm 400^{\text{e}}$	f,g	$12.77 \pm 50^{\text{h}}$		
17 ^d				
21.5 ^d				

^a (1953BO70). See also (1952AR29, 1964EA04).

^b (1956BU16). See also text, and (1964EA04).

^c (1964EA04).

^d (1958TY46).

^e (1965DE21).

^f Observed: see (1953BO70, 1966NG01, 1968DU1E, 1968HI09, 1968HI14).

^g $E_x = 11.3$ and 13 MeV (1968HI14).

^h (1969BA06).

ⁱ Unresolved structure (1969BA06).

^j (1966HA19).

^k From (α, α') : (1966HA19).

^l Relatively low cross section due to two-nucleon transition (1966HA19).


^m See also (1956MI17).

Reduced transition probabilities are reported by (1966HA19): $B(E2)\downarrow/e^2 = 6.5$ and 3.3 fm^4 , respectively for $^{14}\text{N}^*(3.95, 7.03)$; $B(E3)\downarrow/e^2 = 40$ and 60 fm^6 , for $^{14}\text{N}^*(5.11, 5.83)$. See also (1962HA40, 1962JO14, 1963MI1C, 1969BA06), (1968FA1A), (1960WA12), (1968NO1C, 1968RA1C; theor.) and ^{18}F in (1972AJ02).

At $E_\alpha = 22.9 \text{ MeV}$, reaction (b) to $^{13}\text{C}(0)$ appears to involve twelve states of ^{14}N with $E_x = 8.4$ to 13.2 MeV , while reaction (c) proceeds via five states with $E_x = 11.5$ to 12.9 MeV (1969BA17). See also (1967BE30, 1968KU1C).



See (1964KU1D, 1969BR1D).



See (1961NE04, 1962SM02, 1964KU1D).



See (1962SM02, 1968JA1N) and (1967BR1M, 1969VO1E).



See (1961NE04, 1968JA1P).



The decay proceeds almost entirely to the $J^\pi = 0^+$; $T = 1$ state of ^{14}N at 2.31 MeV : see ^{14}O . Measurement of the γ -ray energy from the decay of $^{14}\text{N}^*(2.31)$ leads to $E_x = 2.31287 \pm 0.00010 \text{ MeV}$ for this state (1968FR08), $2.31289 \pm 0.00010 \text{ MeV}$ (1967CH19). The spectrum shape for the transition to $^{14}\text{N}(0)$ differs markedly from the statistical shape. τ_m for $^{14}\text{N}^*(2.31)$ extracted from these data is $33 \pm 3 \text{ fsec}$ (1966SI05): see, however, Table 14.13.



Angular distributions have been obtained for the deuterons corresponding to $^{14}\text{N}^*(0, 2.31, 3.95)$ (**1961BE12**: $E_p = 18.6$ MeV) and to $^{14}\text{N}^*(0 - 8.06, 8.62, 8.91, 8.96 + 8.98, 9.17 - 10.43, 10.81, 11.04, 11.24 + 11.30, 11.39 - 11.66, 11.75 + 11.81, 11.95, 12.23 + 12.29, 12.50, 12.61, 12.79 + 12.83, 13.16 + 13.23, 13.72)$ (**1969SN04**: $E_p = 39.8$ MeV). Spectroscopic factors were extracted by DWBA analysis of the $l_n = 1$ pickup angular distributions. $\Gamma = 210 \pm 30$ keV for $^{14}\text{N}^*(13.75)$. Weak deuteron groups to ^{14}N states at $E_x = 6.70, 7.40$ and 7.60 MeV are reported [see, however, reaction 54] (**1969SN04**).



Not reported.



At $E(^3\text{He}) = 2.8$ MeV, α -particle groups are observed to ^{14}N states at $E_x = 3.95, 4.91, 5.113 \pm 0.008, 5.691 \pm 0.008, 5.832 \pm 0.008, 6.048 \pm 0.012$ [see, however, (**1969HO23**)], $6.224 \pm 0.012, 6.436 \pm 0.012, 7.032 \pm 0.010, 7.97$ and 8.06 MeV (**1962CL12**, **1962CL1D**). The previously reported states at $E_x = 6.70, 7.40$ and 7.60 MeV are unobserved (**1962CL12**, **1969HO23**). Angular distributions have been obtained at $E(^3\text{He}) = 2.8$ MeV (**1962CL12**: to $^{14}\text{N}^*(3.95, 5.69, 5.83)$), 15 MeV (**1969HO23**: to $^{14}\text{N}^*(0 - 8.91, 9.17 - 9.70)$ and 39.8 MeV (**1966BA13**: to $^{14}\text{N}^*(0, 2.31, 3.95, 7.03, 9.17, 10.43)$ and to a state at $E_x = 13.72 \pm 0.04$ MeV). See also (**1965SE01**).



Not reported.



See (**1964BA1C**).



At $E_p = 40$ MeV, angular distributions have been measured for the ^3He particles corresponding to $^{14}\text{N}^*(0, 2.31, 3.95)$ (**1966BR1X**, **1966HO1F**). The excitation of $^{14}\text{N}^*(7.03, 9.17)$ is also reported (**1965PE17**). At $E_p = 43.7$ MeV, a comparison has been made between the angular distributions

of the ${}^3\text{He}$ particles to ${}^{14}\text{N}^*(2.31)$ and the tritons (from the ${}^{16}\text{O}(\text{p}, \text{t}){}^{14}\text{O}$ reaction) to ${}^{14}\text{O}_{\text{g.s.}}$. As would be expected from charge independence, the shape of the distributions and the cross sections are approximately the same to the two analog states: $\sigma(\text{p}, \text{t})/\sigma(\text{p}, {}^3\text{He}) = 1.12/1$ ([1964CE02](#)).



$$Q_m = 3.111$$


$$Q_0 = 3.110 \pm 0.006 \quad (\textcolor{red}{1964MA57}).$$

Angular distribution of α -particles have been measured at many energies: see Table [14.25](#). See also ([1959FI30](#)).

Alpha particle groups have been seen corresponding to most known states of ${}^{14}\text{N}$ with $E_x \leq 11.51$ MeV (in some cases, the identification of the groups is inconclusive), with the exception of previously reported states at $E_x = 6.05, 6.70, 7.40$ and 7.60 MeV ([1968JO07](#): $E_d = 5$ to 9 MeV). See also ([1965IS04](#)). The yield of the isospin-forbidden α_1 group [to ${}^{14}\text{N}^*(2.31)$] has been studied for $E_d = 3$ to 15 MeV. The intensity of the α_1 group, relative to the α_0 and α_2 groups [to the $T = 0$ states of ${}^{14}\text{N}^*(0, 3.95)$] depends on the deuteron energy and on the angle of observation [the isospin impurity in the compound nucleus is a function of the excitation energy in ${}^{18}\text{F}$ and of J] ([1969JO09](#)). Studies of the α_1 yield have also been conducted for $E_d = 5.5$ to 7.5 MeV ([1956BR36](#)) and 6.8 to 8.9 MeV ([1958DA16](#)). For further discussions, see ([1963CE02](#), [1966BR1G](#), [1968NO1C](#), [1969NO1B](#), [1969NO1C](#)). See also ([1960HU10](#), [1961PE09](#), [1961YA08](#), [1963JA03](#), [1963YA1B](#)) and ${}^{18}\text{F}$ in ([1972AJ02](#)).

Preliminary results on the absolute cross sections of this reaction and its inverse [${}^{14}\text{N}(\alpha, \text{d}){}^{16}\text{O}$] are in agreement, to $\pm 0.56\%$, with the principle of detailed balance ([1967TH1E](#), [1968TH1J](#), [1969PL1C](#)).

See also ([1961TE02](#), [1963DO1B](#), [1963VA1E](#), [1964RI1C](#), [1966BE1E](#), [1966JA05](#), [1966KL1E](#), [1968KO24](#)), ([1960EL1E](#), [1961BA1J](#), [1961JA1H](#), [1964HO1E](#), [1966ME1E](#), [1967JO1H](#); theor.) and ([1959AJ76](#)).



$$Q_m = -19.264$$

See ([1965PE17](#)).



$$Q_m = -2.822$$

See ([1965SH10](#), [1965SH14](#)).



$$Q_m = 1.193$$

Table 14.25: $^{16}\text{O}(\text{d}, \alpha)^{14}\text{N}$ angular distribution studies

E_{d} (MeV)	Angular distribution to $^{14}\text{N}^*$	Refs.
1.3 – 1.9	g.s.	(1964AM1A)
1.8 – 2.4	g.s.	(1965MA59)
1.9 – 3.6	g.s.	(1968DI06)
3.3 – 11.0	2.31	(1969JO09)
3.8, 3.9	g.s.	(1963JA03)
3.9	2.31	(1963JA03)
4.4, 4.6, 4.7	g.s.	(1967TH1E)
5.5 – 6.7	g.s.	(1965SA18)
5.6 – 6.9	g.s., 3.95	(1968JO07)
5.8 – 6.6	3.95	(1965SA18)
6.3	g.s.	(1969CO12)
6.6	3.95	(1964GR19)
6.8, 6.9	g.s., 3.95	(1958DA16)
7	g.s., 2.31, 3.95	(1956BR36)
9	g.s., 3.95	(1956GR37)
10.5 – 11.4	g.s.	(1960HU10)
11.4, 12.4	g.s., 3.95	(1965DO08)
14.5	g.s., 3.95	(1961YA07)
14.7	g.s.	(1961YA08)
14.9 – 19.6	g.s., 3.95	(1963YA1B)
15	g.s., 3.95	(1961PE09, 1965PE09)
15	g.s. → 7.97, 8.49, 8.91, 9.17 → 9.51, 10.09	(1969HO23)
19	g.s.	(1953FR23)
24	g.s., 3.95, 4.91 + 5.11, 5.69 + 5.83, 6.21 + 6.44, 7.03	(1965PE17)

See ^{18}F in (1972AJ02).



See (1967DE03).



See (1966KU1D).

¹⁴O
(Figs. 7 and 8)

GENERAL:

See (1959OT1A, 1960PI1C, 1960TA1C, 1961FR1C, 1963BL1B, 1964LO1B, 1965KO1D, 1966BO1R, 1966KE16, 1966MI1G, 1967AU1B, 1968EI1C, 1968FA1B, 1968NE1C, 1969GA1G, 1969SO08).



The decay proceeds primarily to the $J^\pi = 0^+; T = 1$ first excited state of ^{14}N : see Table 14.27. Weak branches are also observed to the ground state of ^{14}N and to the 3.95 MeV state.

The ground-state decay is considerably faster than the mirror transition $^{14}\text{C} \rightarrow ^{14}\text{N}$ supporting the hypothesis that both decays are inhibited by accidental cancellation of matrix elements: see ^{14}C and (1968RO1C).

The decay of $^{14}\text{N}^*(2.31)$ is an example of pure Fermi $0 \rightarrow 0$ transition and provides a precise determination of the fundamental coupling constant G_V (see surveys by (1964WU01, 1966FR15, 1968MO1F)). $G_V = 1.4025 \times 10^{-49} \text{ erg} \cdot \text{cm}^3$ (1962BA26), $1.409 \times 10^{-49} \text{ erg} \cdot \text{cm}^3$ (1968BR1L: see (1969DA06)), $1.407 \times 10^{-49} \text{ erg} \cdot \text{cm}^3$ (1969DA06). See also (1959KI1C, 1960BA36, 1967BL24, 1968BE03). According to the hypothesis of a universal weak interaction, with conserved vector current, $G_V = G_\mu$, the coupling constant for muon decay. The mean for seven cases cited by (1966FR15) is $(G_\mu - G_V)/G_\mu = (2.2 \pm 0.1 \pm 0.5)\%$, where the second error reflects the uncertainty of radiative corrections. See also (1962BA26, 1968BR1L, 1969BL1E, 1969DA06).

The transition to $^{14}\text{N}^*(3.95)$ appears to be super allowed. See also (1962BR16, 1964GA1B, 1966FR15, 1966CH1J, 1966MAZY, 1969LO1F).

The polarization of the positrons has been studied by (1959GE36, 1961HO04). See also (1959AJ76) and (1958AL1E, 1959JA1B, 1960DU1B, 1961BL1A, 1962AL1F, 1962WE09, 1962WE1H, 1962BL1D, 1963LO04, 1964BA1T, 1965BL1E, 1964NA1C, 1965MI1B, 1966BA1A, 1966MA57, 1967CH24, 1967SA1H, 1968BH1C, 1968DA1J, 1968HA38, 1968LA1J, 1968YA1F).



See (1957NO17).



Table 14.26: Energy levels of ^{14}O

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$0^+; 1$	$\tau_{1/2} = 70.98 \pm 0.06$ sec	β^+	1, 2, 3, 4, 5, 6, 7
5.17 ± 40	(1^-)	$\Gamma \leq 150$ keV		3, 5
5.905 ± 12	(0^+)	≤ 60		3, 5
6.29 ± 25	(3^-)	≈ 120		3, 4, 5
6.586 ± 12	2^+	≤ 60		3, 5
6.79 ± 30	$\pi = (-)$			5
7.78 ± 30	2^+			3, 4, 5
8.74 ± 60				5
$(9.74 \pm 30)^{\text{a}}$				3, 4, 5
10.89 ± 50				5
11.24 ± 50				5
$(11.97)^{\text{a}}$				5
12.84 ± 50				5
13.01 ± 50				3, 5
14.15 ± 40				5
14.64 ± 60				5
17.40 ± 60				5

^a Possibly more than one level.

Reported values of the threshold energy are displayed in Table 14.28. Neutron groups have been observed corresponding to ^{14}O states at 5.17 ([1967AD1D](#), [1968TO09](#)), 5.905 ± 0.012 , 6.30 ± 0.03 and 6.586 ± 0.012 MeV, with widths ≤ 60 , 120 and ≤ 60 keV, respectively ([1961TO03](#)). Higher excited states at 7.78 ([1967AD1D](#)), 8.0 ± 0.4 , 10.1 ± 0.4 and 13.3 ± 0.4 MeV are reported by ([1964BR13](#)). Angular distributions are reported at many energies from 1.9 to 25 MeV: see ([1957BR18](#), [1960AJ03](#), [1960GA13](#), [1963MA22](#), [1964DE1C](#), [1964DI02](#), [1964KU05](#), [1967AD1D](#), [1967MC03](#), [1967SC27](#), [1968TO09](#)).

The angular distributions for n_0 and for p_1 , to $^{14}\text{N}^*(2.3)$, show strong forward peaking in the range $E(^3\text{He}) = 6.5$ to 11 MeV. The distributions are similar in detail, and well fitted by plane wave direct interaction theory with $L = 0$. Excitation functions at $\theta = 10^\circ$ show marked energy dependence, but $(d\sigma_n/d\sigma_p)$ is nearly constant at the expected value of 2.0 ([1965FU16](#)). Angular distributions (at $E(^3\text{He}) = 9.2$ and 10.2 MeV) of the ^3He groups to $^{14}\text{O}^*(0, 5.17, 5.91, 6.29)$,

Table 14.27: Positron decay of ^{14}O

	(1954GE38, 1955SH84)	(1961HE03)	(1961BU04)	(1962BA26)	(1963FR10, 1965FR09)	(1965KAZX, 1966SI05)
E_0^{a} (keV)	4145 ± 20			4124 ± 2	4085 ± 30	
E_1^{a} (keV)	1835 ± 8	$[1810.6 \pm 1.5]^{\text{b}}$	1809.7 ± 1.5	$1812.6 \pm 1.4^{\text{e}}$	1821 ± 7	
$\tau_{1/2}$ (sec)	72.1 ± 0.4	70.91 ± 0.04	$[71.4 \pm 0.2]$	71.00 ± 0.13	71.3 ± 0.1	
branch ₀ (%) ^a	0.60 ± 0.10				0.65 ± 0.05	0.61 ± 0.01
branch ₁ (%) ^a	99.4 ± 0.1	$[99.4 \pm 0.1]$	$[99.4 \pm 0.1]$	$[99.4 \pm 0.1]$	99.35 ± 0.05	
branch ₂ (%) ^a						$0.062 \pm 0.007^{\text{d}}$
ft_0 (sec)	$(2.0 \pm 0.3) \times 10^7$				$(1.7 \pm 0.2) \times 10^7$	$(2.14 \pm 0.03) \times 10^7^{\text{f}}$
ft_1 (sec)	3275 ± 75	3061 ± 10	3057 ± 20	$3074 \pm 10^{\text{c}}$	3137 ± 70	$3076 \pm 7^{\text{g}}$
ft_2 (sec)						$1200 \pm 150^{\text{d}}$

^a End-point energies and branches, to $^{14}\text{N}(\text{g.s.: } 1^+; 2.31: 0^+; 3.95: 1^+)$, respectively.^b Square brackets indicate values used for ft -calculations but not determined in present experiment.^c Using $\tau(\text{partial}) = [71.36 \pm 0.009]$ sec; includes form-factor and screening corrections. Radiative corrections of (1959KI1C) increase ft to 3126 sec: see also (1966BA1A, 1966FR15, 1967SU1E).^d And private communication: 0.025 β^+ (1969KA1B).^e 1809.1 ± 1.5 keV is obtained from measurements of the threshold energy of the $^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$ reaction and the energy of $^{14}\text{N}^*(2.31)$. This value leads to a 0.8% decrease in the ft value of (1962BA26, 1968FR08).^f $(1.9 \pm 0.2) \times 10^7$ (1969KA1B).^g (1969KA1B).

Table 14.28: Threshold energy of the $^{12}\text{C}(^{3}\text{He}, \text{n})^{14}\text{O}$ reaction ^a

$E_{\text{thresh.}}$ (keV)	Refs.
1449.6 ± 2.8	(1957BR18)
1436.2 ± 0.9	(1961BU04)
1434.8 ± 1.6	(1961TO06)
1437.5 ± 0.7	(1962BA26)
1436.7 ± 0.5	recommended value: (1966MA60)

^a See also (1965RY01).

analyzed by DWBA double stripping theory, give $L = 0$; $L = 1$ or (0); $L = 0$ or (1); $L = 3$ or 4, respectively. These fits, taken together with the relatively sharp widths of the three excited states (see Table 14.25), lead to probable J^π for $^{14}\text{O}^*(5.17, 5.91, 6.29)$ of $1^-, 0^+, 3^-$, respectively (1968TO09). See also (1960NE1A, 1961GI1B, 1963GL1C, 1964HE06, 1965BR1H, 1966HA1Q, 1966SH1F, 1967CH19) and ^{15}O .



$$E_{\text{thresh.}} = 6.3544 \pm 0.0016 \text{ (1968FR08);}$$

$$E_{\text{thresh.}} = 6.3588 \pm 0.0030 \text{ (1956KU1A; see also (1964KU06)).}$$

The angular distribution of ground state neutrons has been determined at $E_p = 14.8$ and 31.5 MeV (1960DA06, 1961AD02). Neutron groups are reported corresponding to broad or unresolved states at $E_x \approx 6.2, 7.5$ and 9.3 MeV (1954AJ11). See also (1966SI05, 1968RI1Q, 1969MA1G, 1969PA1J, 1969VE02).



Triton groups observed at $E(^3\text{He}) = 44.6$ MeV are displayed in Table 14.29. Comparisons of angular distributions of the tritons to the first eight states of ^{14}O with those of triton groups in other reactions involving known l -transfers lead to the J^π assignments quoted in the table (1967BA13, 1969BA06). See also (1968BA1E) and reaction 45 in ^{14}N .



Table 14.29: ^{14}O levels from $^{14}\text{N}(^3\text{He}, \text{t})^{14}\text{O}$
 (1967BA13)

E_x (MeV \pm keV)	J^π ^a
0	0^+
5.17 ± 40	(1^-)
5.91 ± 40	(0^+)
6.28 ± 30	(3^-)
6.60 ± 30	2^+
6.79 ± 30	(2^-)
7.78 ± 30	2^+
8.74 ± 60	
9.74 ± 30	
10.89 ± 50	
11.24 ± 50	
11.97 ^b	
12.84 ± 50	
13.01 ± 50	
14.15 ± 40	
14.64 ± 60	
17.40 ± 60	

^a See text.

^b Unresolved states.

See (1959OC07, 1962BR16, 1964MO26).

$$7. \ ^{16}\text{O}(\text{p}, \text{t})^{14}\text{O} \quad Q_m = -20.406$$

The angular distribution of ground state protons has been determined at $E_p = 43.7$ (1964CE02).
 See also (1966CE05, 1969SO08).

References

(Closed 31 December 1969)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1947HU03 D.J. Hughes, C. Eggler and C.M. Huddleston, Phys. Rev. 71 (1947) 269
1949DE1A Devons and Hine, Proc. Roy. Soc. A199 (1949) 56, 73
1952AR29 J.C. Arthur, A.J. Allen, R.S. Bender, H.J. Hausman and C.J. McDole, Phys. Rev. 88 (1952) 1291
1952GI01 W.M. Gibson and E.E. Thomas, Proc. Roy. Soc. A210 (1952) 543
1952SE01 J.D. Seagrave, Phys. Rev. 85 (1952) 197
1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
1953BO70 C.K. Bockelman, C.P. Browne, W.W. Buechner and A. Sperduto, Phys. Rev. 92 (1953) 665
1953CL39 A.B. Clegg and D.H. Wilkinson, Phil. Mag. 44 (1953) 1269
1953FR23 R.G. Freemantle, W.M. Gibson, D.J. Prowse and J. Rotblat, Phys. Rev. 92 (1953) 1268
1953MA42 G. Manning and B. Singh, Proc. Phys. Soc. (London) A66 (1953) 842
1953SH64 E.S. Shire, J.R. Wormald, G. Lindsay-Jones, A. Lundan and A.G. Stanley, Phil. Mag. 44 (1953) 1197
1953TA06 F.L. Talbott and N.P. Heydenburg, Phys. Rev. 90 (1953) 186
1953WI1A Wilkinson, Phil. Mag. 44 (1953) 450
1953WO41 H.H. Woodbury, R.B. Day and A.V. Tollestrup, Phys. Rev. 92 (1953) 1199
1954AJ11 F. Ajzenberg and W. Franzen, Phys. Rev. 94 (1954) 409
1954FE16 G.A. Ferguson, J. Halpern, R. Nathans and P.F. Yergin, Phys. Rev. 95 (1954) 776
1954FR38 R.G. Freemantle, D.J. Prowse and J. Rotblat, Phys. Rev. 96 (1954) 1268
1954GE38 J.B. Gerhart, Phys. Rev. 95 (1954) 288
1954JA1A B. Jancovici and I. Talmi, Phys. Rev. 95 (1954) 289
1954MI05 E.A. Milne, Phys. Rev. 93 (1954) 762
1954MO84 A. Moljk and S.C. Curran, Phys. Rev. 96 (1954) 395
1954SA68 P. Savel and M.E. Nahmias, Compt. Rend. 238 (1954) 2155

- 1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, Phys. Rev. 96 (1954) 1316
- 1954ST20 A.G. Stanley, Phil. Mag. 45 (1954) 430
- 1955AL16 H.R. Allan and N. Sarma, Proc. Phys. Soc. (London) A68 (1955) 535
- 1955BE62 R.D. Bent, T.W. Bonner and R.F. Sippel, Phys. Rev. 98 (1955) 1237
- 1955CH1A Chow and Wright, Phys. Rev. 99 (1955) 640
- 1955MA36 R.J. Mackin, Jr., W.B. Mims and W.R. Mills, Jr., Phys. Rev. 98 (1955) 43
- 1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847
- 1955MC75 J.N. McGruer, E.K. Warburton and R.S. Bender, Phys. Rev. 100 (1955) 235
- 1955SH46 E.S. Shire and R.D. Edge, Phil. Mag. 46 (1955) 640
- 1955SH84 R. Sherr, J.B. Gerhart, H. Horie and W.F. Hornyak, Phys. Rev. 100 (1955) 945
- 1955WI43 D.H. Wilkinson, Phys. Rev. 100 (1955) 32
- 1956BO08 T.W. Bonner, J.T. Eisinger, A.A. Kraus, Jr. and J.B. Marion, Phys. Rev. 101 (1956) 209
- 1956BO61 T.W. Bonner, A.A. Kraus, Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956BR36 C.P. Browne, Phys. Rev. 104 (1956) 1598
- 1956BU16 E.J. Burge and D.J. Prowse, Phil. Mag. 1 (1956) 912
- 1956EL1B Elliott, Phil. Mag. 1 (1956) 503
- 1956FR1A J.B. French, Phys. Rev. 103 (1956) 1391
- 1956GO1L S. Gorodetzky, A. Gallman and M. Croissiaux, Physica 22 (1956) 1160A
- 1956GO39 S. Gorodetzky, A. Gallman and M. Croissiaux, J. Phys. Rad. 17 (1956) 550
- 1956GO42 H.E. Gove, A.E. Litherland, E. Almqvist and D.A. Bromley, Phys. Rev. 103 (1956) 835
- 1956GR17 G.M. Griffiths, Can. J. Phys. 34 (1956) 339
- 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28
- 1956KO26 B. Koudijs, Ph.D. Thesis, Univ. of Utrecht (1956)
- 1956KU1A Kurath, Phys. Rev. 101 (1956) 216
- 1956LE28 P. Lehmann, A. Leveque and R. Pick, Compt. Rend. 243 (1956) 743
- 1956LI05 D.L. Livesey, Can. J. Phys. 34 (1956) 216
- 1956MA87 J.B. Marion and F.B. Hagedorn, Phys. Rev. 104 (1956) 1028
- 1956MC88 M.T. McEllistrem, K.W. Jones, R. Chiba, R.A. Douglas, D.F. Herring and E.A. Silverstein, Phys. Rev. 104 (1956) 1008

- 1956MI17 D.W. Miller, B.M. Carmichael, U.C. Gupta, V.K. Rasmussen and M.B. Sampson, Phys. Rev. 101 (1956) 740
- 1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434
- 1956VA17 F.P.G. Valckx, Ph.D. Thesis, Univ. of Utrecht (1956)
- 1956VA1B Vanetsian and Fedchenko, Physica 22 (1966) A1124
- 1956WR22 I.F. Wright, D.R.O. Morrison, J.M. Reid and J.R. Atkinson, Proc. Phys. Soc. (London) A69 (1956) 77
- 1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson, Phys. Rev. 105 (1957) 957
- 1957BR25 C. Broude, L.L. Green, J.J. Singh and J.C. Willmott, Phil. Mag. 2 (1957) 499
- 1957GA1B Gallmann, Ph.D. Thesis, Univ. of Strasbourg (1957)
- 1957GI14 W.M. Gibson, D.J. Prowse and J. Rotblat, Proc. Roy. Soc. A243 (1957) 237
- 1957GO30 S. Gorodetzky, A. Gallmann, M. Croissiaux and R. Armbruster, Compt. Rend. 244 (1957) 62
- 1957HI56 N.M. Hintz, Phys. Rev. 106 (1957) 1201
- 1957HO34 A. Hossain and A.N. Kamal, Ind. J. Phys. 31 (1957) 553
- 1957HU1C Huper, Z. Naturforsch. A12 (1957) 295
- 1957NO14 E. Norbeck, Jr., Phys. Rev. 105 (1957) 204
- 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1957SA01 N. Sarma, M. Govindjee and H.R. Allan, Proc. Phys. Soc. (London) A70 (1957) 68
- 1957VA11 D.M. Van Patter and W. Whaling, Rev. Mod. Phys. 29 (1957) 757
- 1957VA1B Vanetsian and Fedchenko, Sov. J. At. Energy 2 (1957) 141
- 1957VI1A W.M. Visscher and R.A. Ferrell, Phys. Rev. 107 (1957) 781
- 1957WI27 D.H. Wilkinson and S.D. Bloom, Phil. Mag. 2 (1957) 63
- 1957WI30 H.B. Willard, J.K. Bair, H.O. Cohn and J.D. Kington, Phys. Rev. 105 (1957) 202
- 1957ZI09 D. Zipoy, G. Freier and K. Famularo, Phys. Rev. 106 (1957) 93
- 1958AL1E Altman and MacDonald, Phys. Rev. Lett. 1 (1958) 456
- 1958BA1A E. Baranger and S. Meshkov, Phys. Rev. Lett. 1 (1958) 30
- 1958CA1B Cazzola and Foglia, Nuovo Cim. 10 (1958) 913
- 1958CH1A Chevallier, Thesis, Univ. of Strasbourg (1958)
- 1958DA16 A.W. Dalton, S. Hinds and G. Parry, Proc. Phys. Soc. (London) A71 (1958) 252
- 1958GO81 S. Gorodetzky, P. Chevallier, R. Armbruster, G. Sutter and A. Gallman, Nucl. Phys. 8 (1958) 412

- 1958GR97 G.M. Griffiths, Proc. Phys. Soc. (London) A72 (1958) 337
- 1958MA1A Maikov, Zh. Eksp. Teor. Fiz. 34 (1958) 1406; Sov. Phys. JETP 7 (1958) 973
- 1958MC63 M.T. McEllistrem, Phys. Rev. 111 (1958) 596
- 1958RA13 W.A. Ranken, T.W. Bonner, J.M. McCrary and T.A. Rabson, Phys. Rev. 109 (1958) 917
- 1958TY46 H. Tyren and T.A.J. Maris, Nucl. Phys. 6 (1958) 82
- 1958WA02 E.K. Warburton and H.J. Rose, Phys. Rev. 109 (1958) 1199
- 1958ZI17 D.M. Zipoy, Phys. Rev. 110 (1958) 995
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AL08 M.A. Al-Jeboori, M.S. Bokhari, B. Hird and A. Strzalkowski, Proc. Phys. Soc. (London) A74 (1959) 705
- 1959AL09 R.G. Allas and F.B. Shull, Phys. Rev. 116 (1959) 996
- 1959AL96 E. Almqvist, D. A. Bromley, A. J. Ferguson, H. E. Gove, and A. E. Litherland, Phys. Rev. 114 (1959) 1040
- 1959AR1A Armstrong, Moore and Blair, Bull. Amer. Phys. Soc. 4 (1959) 17
- 1959BA1F Balashov and Tulinov, Zh. Eksp. Teor. Fiz. 36 (1959) 615; Sov. Phys. JETP 9 (1959) 426
- 1959BO14 R.O. Bondelid and C.A. Kennedy, Phys. Rev. 115 (1959) 1601
- 1959BO40 D. Bodansky, S.F. Eccles, G.W. Farwell, M.E. Rickey and P.C. Robinson, Phys. Rev. Lett. 2 (1959) 101
- 1959BR06 D.A. Bromley, A.J. Ferguson, H.E. Gove, J.A. Kuehner, A.E. Litherland, E. Almqvist and R. Batchelor, Can. J. Phys. 37 (1959) 1514
- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959BR75 O.D. Brill and L.V. Sumin, Atomnaya Energ. 7 (1959) 377
- 1959BU1E Budzanowski, Bull. Acad. Pol. Sci. 7 (1959) 961
- 1959CA1C Carroll, Thesis, Univ. of Pennsylvania (1959)
- 1959CH28 P. Chevallier, Ann. Phys. 4 (1959) 1389
- 1959CI38 S.D. Cirilov and M.K. Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 9 (1959) 39
- 1959EL43 M. El Nadi and M. El Khishin, Proc. Phys. Soc. (London) A73 (1959) 705
- 1959EL44 A. Elwyn, J.V. Kane, S. Ofer and D.H. Wilkinson, Phys. Rev. 116 (1959) 1490
- 1959FA1A Fagg and Hanna, Rev. Mod. Phys. 31 (1959) 711
- 1959FA1C Fallieros and Ferrell, Phys. Rev. 116 (1959) 660
- 1959FI30 G.E. Fischer and V.K. Fischer, Phys. Rev. 114 (1959) 533

- 1959FU1A Fujii, Prog. Theor. Phys. 21 (1959) 511
- 1959GA05 A. Gallmann, Ann. Phys. 4 (1959) 185
- 1959GA14 F. Gabbard, H. Bischel and T.W. Bonner, Nucl. Phys. 14 (1959) 277
- 1959GE36 J.B. Gerhart, F.H. Schmidt, H. Bichsel and J.C. Hopkins, Phys. Rev. 114 (1959) 1095
- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959GO78 S. Gorodetzky, P. Chevallier, R. Armbruster and G. Sutter, Nucl. Phys. 12 (1959) 349
- 1959HA11 S.S. Hanna and L. Meyer-Schutzmeister, Phys. Rev. 115 (1959) 986
- 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
- 1959HE1B Hess, Ann. Phys. 6 (1959) 115
- 1959HE1C Henley and Jacobsohn, Phys. Rev. 113 (1959) 225
- 1959HI1E Hird, Cookson and Bokhari, Congres Int. Phys. Nucl., Dunod, Paris, 1958 (1959) 470
- 1959HI69 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 196
- 1959HO01 H.D. Holmgren, E.A. Wolicki and R.L. Johnston, Phys. Rev. 114 (1959) 1281
- 1959JA1B Jancovici, Ann. Phys. 4 (1959) 689
- 1959JU1A Juric and Cirilov, Congres Int. Phys. Nucl., Dunod, Paris, 1958 (1959) 473
- 1959KA07 Y. Kato and O. Nakahara, J. Phys. Soc. Jpn. 14 (1959) 690
- 1959KI1C Kinoshita and Sirlin, Phys. Rev. 113 (1959) 1652
- 1959KU1C Kurepin and Neudachin, Zh. Eksp. Teor. Fiz. 36 (1959) 1725; Sov. Phys. JETP 9 (1959) 1229
- 1959ME24 U. Meyer-Berkhout, K.W. Ford and A.E.S. Green, Ann. Phys. (N.Y.) 8 (1959) 119
- 1959MU08 N. Mutsuro, Y. Ohnuki, K. Sato, K. Kageyama and M. Kimura, J. Phys. Soc. Jpn. 14 (1959) 1457
- 1959NE1B Neudachin, Teplov and Shevchenko, Zh. Eksp. Teor. Fiz. 36 (1959) 850; Sov. Phys. JETP 9 (1959) 5999
- 1959OC07 J. O'Connell, P. Dyal and J. Goldemberg, Phys. Rev. 116 (1959) 173
- 1959OT1A Ott, Z. Naturforsch. A14 (1959) 769
- 1959RE1A Reid et al., Congres Int. Phys. Nucl., Dunod, Paris, 1958 (1959) 700
- 1959RO54 H.J. Rose, W. Trost and F. Riess, Nucl. Phys. 12 (1959) 510
- 1959SK1A Skyrme, Nucl. Phys. 9 (1959) 641
- 1959WA02 E.K. Warburton, W.T. Pinkston, H.J. Rose and E.N. Hatch, Bull. Amer. Phys. Soc. 4 (1959) 219, AB9
- 1959WA04 E.K. Warburton, H.J. Rose and E.N. Hatch, Phys. Rev. 114 (1959) 214
- 1959WA16 E.K. Warburton, Phys. Rev. 113 (1959) 595

- 1959YA01 A.I. Yavin and G.W. Farwell, Nucl. Phys. 12 (1959) 1
- 1959YO25 T.E. Young, G.C. Phillips, R.R. Spencer and D.A.A.S.N. Rao, Phys. Rev. 116 (1959) 962
- 1960AB02 T.L. Abelishvili, T.G. Gachechiladze and O.M. Mdivani, Zh. Eksp. Teor. Fiz. 38 (1960) 631; Sov. Phys. JETP 11 (1960) 453
- 1960AG01 J. Aguilar, W.E. Burcham, J. Catala, J.B.A. England, J.S.C. McKee and J. Rotblat, Proc. Roy. Soc. A254 (1960) 395
- 1960AJ03 F. Ajzenberg-Selove and K.L. Dunning, Phys. Rev. 119 (1960) 1681
- 1960AL09 M.A. Al Jeboori, M.S. Bokhari, A. Strzalkowski and B. Hird, Proc. Phys. Soc. (London) 75 (1960) 875
- 1960BA15 D. Balfour and D.C. Manizies, Proc. Phys. Soc. (London) A75 (1960) 543
- 1960BA34 J.K. Bair, R.D. Edge and H.B. Willard, Phys. Rev. 119 (1960) 1948
- 1960BA35 J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 119 (1960) 2026
- 1960BA36 R.K. Bardin, C.A. Barnes, W.A. Fowler and P.A. Seeger, Phys. Rev. Lett. 5 (1960) 323
- 1960BU15 A. Budzanowski, K. Grotowski, H. Niewodniczanski and J. Nurzynski, J. Phys. Rad. 21 (1960) 366
- 1960CA09 E.E. Carroll, Jr. and W.E. Stephens, Phys. Rev. 118 (1960) 1256
- 1960CH12 L.F. Chase, Jr., R.G. Johnson and E.K. Warburton, Phys. Rev. 120 (1960) 2103
- 1960DA06 W.W. Daehnick, R. Sherr and M.K. Banerjee, Bull. Amer. Phys. Soc. 5 (1960) 246, HA6
- 1960DU1B Durand, Landovitz and Marr, Phys. Rev. Lett. 4 (1960) 620
- 1960EL1C El-Nadi, Phys. Rev. 120 (1960) 1360
- 1960EL1E El-Nadi, Phys. Rev. 119 (1960) 242
- 1960FA06 R.W. Fast, P.A. Flournoy, R.S. Tickle and W.D. Whitehead, Phys. Rev. 118 (1960) 535
- 1960GA13 N.H. Gale, J.B. Garg, J.M. Calvert and K. Ramavataram, Nucl. Phys. 20 (1960) 313
- 1960GE06 K.N. Geller, J. Halpern and E.G. Muirhead, Phys. Rev. 119 (1960) 716
- 1960GO19 S. Gorodetzky, A. Gallmann, P. Fintz and J. Samuel, J. Phys. Rad. 21 (1960) 358
- 1960HA31 B.G. Harvey, J. Cerny and R.H. Pehl, Bull. Amer. Phys. Soc. 5 (1960) 493, C4
- 1960HA32 B.G. Harvey and J. Cerny, Phys. Rev. 120 (1960) 2162
- 1960HE14 D.F. Hebbard and J.L. Vogl, Nucl. Phys. 21 (1960) 652
- 1960HI07 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 745
- 1960HU10 C. Hu, J. Phys. Soc. Jpn. 15 (1960) 1741

- 1960JA17 A.A. Jaffe, F. de S. Barros, P.D. Forsyth, J. Muto, I.J. Taylor and S. Ramavataram, Prog. Phys. Soc. 76 (1960) 914
- 1960KA06 E. Kashy, R.R. Perry and J.R. Risser, Phys. Rev. 117 (1960) 1289
- 1960KI02 J.D. King, R.N.H. Haslam and R.W. Parsons, Can. J. Phys. 38 (1960) 231
- 1960KO05 A.P. Komar, J. Krzhemenek and I.P. Yavor, Dokl. Akad. Nauk SSSR 131 (1960) 283; Sov. Phys. Dokl. 5 (1960) 295
- 1960LI10 C.C. Lin, Phys. Rev. 119 (1960) 1027
- 1960MA32 R.D. Macfarlane and J.B. French, Rev. Mod. Phys. 32 (1960) 567
- 1960MU02 N. Mutsuro, K. Sato and M. Mishina, J. Phys. Soc. Jpn. 15 (1960) 358
- 1960MU07 J. Muto, F. de S. Barros and A.A. Jaffe, Proc. Phys. Soc. (London) 75 (1960) 929
- 1960NE1A Newns, Proc. Phys. Soc. (London) 76 (1960) 489
- 1960OD01 Y. Oda, M. Takeda, N. Takano, T. Yamazaki, C. Hu, K. Kikuchi, S. Kobayashi, K. Matsuda and Y. Nagahara, J. Phys. Soc. Jpn. 15 (1960) 760
- 1960ON01 K. Ono and K. Kuroda, Phys. Rev. 117 (1960) 214
- 1960PA08 M.L. Pal, Phys. Rev. 117 (1960) 566
- 1960PI1C Pinajiom, Nucl. Phys. 17 (1960) 44
- 1960PR12 J.R. Priest, D.J. Tendam and E. Bleuler, Phys. Rev. 119 (1960) 1295
- 1960RO13 H.J. Rose, Nucl. Phys. 19 (1960) 113
- 1960RO23 H.J. Rose, F. Riess and W. Trost, Nucl. Phys. 21 (1960) 367
- 1960SA09 D. Sadeh, Compt. Rend. 250 (1960) 1632
- 1960SH01 S.M. Shafroth, Bull. Amer. Phys. Soc. 5 (1960) 55, RA1
- 1960SH05 S.M. Shafroth, J. Phys. Rad. 21 (1960) 353
- 1960SM02 R.V. Smith, L.F. Chase, Jr., R.H. Abramson, J.B. Reagan and M. Walt, Bull. Amer. Phys. Soc. 5 (1960) 19, E8
- 1960TA08 M. Takeda, J. Phys. Soc. Jpn. 15 (1960) 557
- 1960TA1C Talmi and Unna, Ann. Rev. Nucl. Sci. 10 (1960) 353
- 1960VA11 F.J. Vaughn, L.F. Chase, Jr. and R.G. Johnson, Bull. Amer. Phys. Soc. 5 (1960) 404, B2
- 1960WA12 E.K. Warburton and W.T. Pinkston, Phys. Rev. 118 (1960) 733
- 1960WA17 I. Wahlstrom and B. Forkman, Ark. Fys. 18 (1960) 83
- 1961AD02 H.E. Adelson and C.N. Waddell, Bull. Amer. Phys. Soc. 6 (1961) 375, X2
- 1961AS1B Ashe, McGrary, Morgan and Prud'Homme, Bull. Amer. Phys. Soc. 6 (1961) 61
- 1961BA09 J.K. Bair, Phys. Rev. 122 (1961) 897

- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. (London) 77 (1961) 853
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165
- 1961BA1F Barker, Phys. Rev. 122 (1961) 572
- 1961BA1J Bashkin, SUI-61-6 (1961)
- 1961BA22 J.K. Bair and H.B. Willard, Bull. Amer. Phys. Soc. 6 (1961) 440, P3
- 1961BE12 E.F. Bennett, Phys. Rev. 122 (1961) 595
- 1961BL1A Blin-Stoyle and Le Tourneau, Phys. Rev. 123 (1961) 627
- 1961BO28 V.V. Bobyr, L.Ya. Grona and V.I. Strizhak, Zh. Eksp. Teor. Fiz. 41 (1961) 24; JETP (Sov. Phys.) 14 (1962) 18
- 1961BR13 H.R. Brooker, P.J. Haigh and T.A. Scott, Phys. Rev. 123 (1961) 2143
- 1961BU04 J.W. Butler and R.O. Bondelid, Phys. Rev. 121 (1961) 1770
- 1961CE02 J. Cerny, N.K. Glendenning, B.G. Harvey, R.H. Pehl and E. Rivet, Bull. Amer. Phys. Soc. 6 (1961) 507, E11
- 1961CI08 S.D. Cirilov, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 781
- 1961CL09 A.B. Clegg, K.J. Foley, G.L. Salmon and R.E. Segel, Proc. Phys. Soc. (London) 78 (1961) 681
- 1961CO05 H.O. Cohn, J.K. Bair and H.B. Willard, Phys. Rev. 122 (1961) 534
- 1961DA09 P. Dagley, W. Haeberli and J.X. Saladin, Nucl. Phys. 24 (1961) 353
- 1961EL1A El Nadi and Sherif, Nucl. Phys. 28 (1961) 331
- 1961ER01 J.R. Erskine and C.P. Browne, Phys. Rev. 123 (1961) 958
- 1961FR1C J.B. French, Nucl. Phys. 26 (1961) 161
- 1961GI1B Gibbs and Tobocman, Bull. Amer. Phys. Soc. 6 (1961) 236
- 1961GO1K Goldfarb, Proc. Rutherford Jub. Int. Conf. (1961) 479
- 1961GR06 A.G. Gregory and P.B. Treacy, Proc. Phys. Soc. (London) 77 (1961) 499
- 1961HA04 M.L. Halbert and A. Zucker, Phys. Rev. 121 (1961) 236
- 1961HE02 R.E. Hester and W.A.S. Lamb, Phys. Rev. 121 (1961) 584
- 1961HE03 D.L. Hendrie and J.B. Gerhart, Phys. Rev. 121 (1961) 846
- 1961HO04 J.C. Hopkins, J.B. Gerhart, F.H. Schmidt and J.E. Stroth, Phys. Rev. 121 (1961) 1185
- 1961JA08 A.N. James, Nucl. Phys. 23 (1961) 648
- 1961JA09 A.N. James, Nucl. Phys. 24 (1961) 132

- 1961JA1H Jacobson and Ryndin, Nucl. Phys. 24 (1961) 505
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, Rev. Mex. Fisica 10 (1961) 247
- 1961JO14 W.P. Johnson and D.W. Miller, Phys. Rev. 124 (1961) 1190
- 1961KA04 E. Kashy, R.R. Perry, R.L. Steele and J.R. Risser, Phys. Rev. 122 (1961) 884
- 1961KN03 H.H. Knitter and H. Waffler, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 823
- 1961LE1E Levintov and Trostin, Zh. Eksp. Teor. Fiz. 40 (1961) 1570; Sov. Phys. JETP 13 (1961) 1102
- 1961MA1E Maksimov, Sov. J. At. Energ. 10 (1962) 250; Atomn. Energ. (USSR) 10 (1961) 260
- 1961MA1K L. Marquez and P.D. Lien, Centro Brasiliero De Pesquisas Fisicas 8, No. 11 (1961)
- 1961MA1L W.B. Mann, W.F. Marlow and E.E. Hughes, National Bureau of Standards (U.S.) Tech. News Bull. 45 (1961) 21
- 1961MA32 W.B. Mann, W.F. Marlow and E.E. Hughes, Int. J. Appl. Rad. Isotopes 11 (1961) 57
- 1961MA33 L. Marquez and P.D. Lien, J. Phys. Rad. 22 (1961) 589
- 1961NE04 E. Newman, P.G. Roll and F.E. Steigert, Phys. Rev. 122 (1961) 1842
- 1961PE09 F. Pellegrini, Nucl. Phys. 24 (1961) 372
- 1961PL01 W.D. Plough, Phys. Rev. 122 (1961) 1232
- 1961SE03 R.E. Segel, J.W. Daughtry and J.W. Olness, Phys. Rev. 123 (1961) 194
- 1961ST10 D.M. Stanojevic, M.K. Jric and B.S. Marsicanin, Bull. Inst. Nucl. Sci., Horis Kidrich, 11 March 1961 (1961) 11
- 1961ST22 V.I. Strizhak, V.V. Bobyr and L.Y. Grona, Zh. Eksp. Teor. Fiz. 41 (1961) 313; Sov. Phys. JETP 14 (1962) 225
- 1961SW01 C.P. Swann, V.K. Rasmussen and F.R. Metzger, Phys. Rev. 121 (1961) 242
- 1961TA06 A.E. Taylor and E. Wood, Nucl. Phys. 25 (1961) 642
- 1961TA10 Tamers and Delibrias, Compt. Rend. 253 (1961) 1202
- 1961TE02 A. Tejera, M. Mazari, A. Jaidar and G. Lopez, Rev. Mex. Fis. 10 (1961) 229
- 1961TO03 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. (London) 77 (1961) 399
- 1961TO06 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. (London) 77 (1961) 1217
- 1961TR1B True and Warburton, Nucl. Phys. 22 (1961) 426
- 1961WA16 D.E. Watt, D. Ramsden and H.W. Wilson, Int. J. Appl. Rad. Isotopes 11 (1961) 68; Nucl. Sci. Abs. 15, 4221, Abs. 32740 (1961)
- 1961WI1A Wilkinson, Wollan and Koehler, Ann. Rev. Nucl. Sci. 11 (1961) 303
- 1961WO03 C. Wong, J.D. Anderson, S.D. Bloom, J.W. McClure and B.D. Walker, Phys. Rev. 123 (1961) 598

- 1961YA07 T. Yanabu, S. Yamashita, T. Nakamura, K. Takamatsu, A. Masaike, S. Kakigi, D.C. Nguyen and K. Takimoto, J. Phys. Soc. Jpn. 16 (1961) 2594
- 1961YA08 T. Yanabu, J. Phy. Soc. Jpn. 16 (1961) 2118
- 1961ZD01 R. Zdanis, G.E. Owen and L. Madansky, Phys. Rev. 121 (1961) 854
- 1962AL03 R.G. Allas and F.B. Shull, Phys. Rev. 125 (1962) 941
- 1962AL1F Altman and MacDonald, Nucl. Phys. 35 (1962) 593
- 1962BA1D Barber, Ann. Rev. Nucl. Sci. 12 (1962) 1
- 1962BA1K Baird, J. Chem. Phys. 37 (1962) 1879
- 1962BA26 R.K. Bardin, C.A. Barnes, W.A. Fowler and P.A. Seeger, Phys. Rev. 127 (1962) 583
- 1962BE04 T.A. Belote, B. Mainsbridge and J.R. Risser, Bull. Amer. Phys. Soc. 7 (1962) 112, A7
- 1962BE24 E. Berkowitz, S. Bashkin, R.R. Carlson, S.A. Coon and E. Norbeck, Phys. Rev. 128 (1962) 247
- 1962BL1D Blin-Stoyle and Le Tourneux, Ann. Phys. 18 (1962) 12
- 1962BR16 H. Breuer and W. Pohlit, Nucl. Phys. 30 (1962) 417
- 1962CA05 J.H. Calvert, N.H. Gale, J.B. Garg and K. Ramavataram, Nucl. Phys. 31 (1962) 471
- 1962CE01 J. Cerny, B.G. Harvey and R.H. Pehl, Nucl. Phys. 29 (1962) 120
- 1962CH01 C. Chasman and D.A. Bromley, Bull. Amer. Phys. Soc. 7 (1962) 36, JA4
- 1962CL12 D.D. Clayton, Phys. Rev. 128 (1962) 2254
- 1962CL1D Clayton, Thesis, CalTech, Unpublished (1962)
- 1962ED02 R.D. Edge and G.A. Peterson, Phys. Rev. 128 (1962) 2750
- 1962EL1C El Nadi and Sherif, Proc. Phys. Soc. (London) 80 (1962) 1041
- 1962ER03 H.J. Erramuspe and R.J. Slobodrian, Nucl. Phys. 34 (1962) 532
- 1962GO1E Gorbunov et al., Zh. Eksp. Teor. Fiz. 42 (1962) 747; Sov. Phys. JETP 15 (1962) 520
- 1962GO33 H. Godwin, Nature 195 (1962) 984
- 1962GR10 A.G. Gregory, G.D. Symons and P.B. Treacy, Proc. Phys. Soc. (London) 80 (1962) 315
- 1962GU01 G.D. Gutsche, H.D. Holmgren, L.M. Cameron and R.L. Johnston, Phys. Rev. 125 (1962) 648
- 1962HA40 B.G. Harvey, J. Cerny, R.H. Pehl and E. Rivet, Nucl. Phys. 39 (1962) 160
- 1962HE1A Henning and Waard, Phys. Lett. 3 (1962) 139
- 1962HO06 R.K. Hobbie and F.F. Forbes, Phys. Rev. 126 (1962) 2137
- 1962IN1C Inglis, Phys. Rev. 126 (1962) 1789
- 1962IS04 A. Isoya, S. Micheletti and L. Reber, Phys. Rev. 128 (1962) 806

- 1962JO14 J.C. Jodogne, P.C. Macq and J. Steyaert, Phys. Lett. 2 (1962) 325
- 1962KA1E Kawai, Terasewa and Izumo, Prog. Theor. Phys. 27 (1962) 404
- 1962KJ03 J. Kjellman, Ark. Fys. 21 (1962) 543
- 1962KO23 A.P. Komar, Ya. Krzhemenek and I.P. Yavor, Nucl. Phys. 34 (1962) 551
- 1962KU09 B. Kuhn, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teopr. Fiz. 43 (1962) 1660; Sov. Phys. JETP 16 (1963) 1171
- 1962LA15 N.O. Lassen, Nucl. Phys. 38 (1962) 442
- 1962LI08 Pham-dinh-Lien and L. Marquez, Nucl. Phys. 33 (1962) 202
- 1962MA1P Majling, Zh. Eksp. Teor. Fiz. 42 (1962) 831; Sov. Phys. JETP 15 (1962) 579
- 1962MO16 W.T. Morton and T.G. Walker, Phil. Mag. 7 (1962) 741
- 1962MO1A Morpurgo, Nucl. Spectroscopy, Ed., Racah (1962)
- 1962OL04 I.U. Olsson, I. Karlen, A.H. Turnbull and N.J.D. Prosser, Ark. Fys. 22 (1962) 237
- 1962RO20 L. Rosen and W.T. Leland, Phys. Rev. Lett. 8 (1962) 379
- 1962RO21 H.J. Rose, F. Uihlein, F. Riess and W. Trost, Nucl. Phys. 36 (1962) 583
- 1962SE13 H.M. Sen Gupta, J. Rotblat, P.E. Hodgson and J.B.A. England, Nucl. Phys. 38 (1962) 361
- 1962SH19 E.F. Shrader and R.L. Zimmerman, Bull. Amer. Phys. Soc. 7 (1962) 550, T5
- 1962SH21 R.D. Sharp, Bull. Amer. Phys. Soc. 7 (1962) 622, W1
- 1962SM02 A.M. Smith and F.E. Steigert, Phys. Rev. 125 (1962) 988
- 1962TA1E Talmi, Nucl. Spectroscopy; Ed., Racah (1962)
- 1962UN1A Unna, Nucl. Spectrosc. (1962) 254
- 1962VA1F Vakselj, Nucl. Phys. 31 (1962) 525
- 1962WA1C Wakburton, Electromag. Lifetime and Properties of Nucl. States; N.A.S.-N.R.C. Pub. 974 (1962) 180
- 1962WA1D Wang Tieh-Yu et al., Acta Phys. Sin. (China) 18 (1962) 227
- 1962WE09 H.A. Weidenmuller, Phys. Rev. 128 (1962) 841
- 1962WE12 C. Weddigen, Z. Phys. 170 (1962) 436
- 1962WE1B Weidenmuller, Nucl. Phys. 36 (1962) 151
- 1962WE1C Wegner, Rev. Sci. Instrum. 33 (1962) 271
- 1962WE1H H.A. Weidenmuller, Phys. Rev. 127 (1962) 537; Erratum Phys. Rev. 127 (1962) 2287
- 1962WI15 B. Wilkins and G. Igo, Phys. Lett. 3 (1962) 48
- 1963AL21 D.E. Alburger and E.K. Warburton, Phys. Rev. 132 (1963) 790
- 1963BA08 S. Bashkin, V.P. Hart and W.A. Seale, Phys. Rev. 129 (1963) 1750

- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963BA38 NN.P. Babenko, B.A. Bibichev, I.O. Konstantinov and Y.A. Nemilov, Zh. Eksp. Teor. Fiz. 44 (1963) 135; Sov. Phys. JETP 17 (1963) 92
- 1963BA46 R.W. Bauer, J.D. Anderson and L.J. Christensen, Nucl. Phys. 47 (1963) 241
- 1963BA66 N.P. Babenko, I.O. Konstantinov and Y.A. Nemilov, Zh. Eksp. Teor. Fiz. 45 (1963) 1389; Sov. Phys. JETP 18 (1964) 959
- 1963BE05 R.E. Benenson and B. Yaramis, Phys. Rev. 129 (1963) 720
- 1963BE1M Beurtey et al., J. Phys. 24 (1963) 1038
- 1963BE33 J.A. Becker, Phys. Rev. 131 (1963) 322
- 1963BL1B Blin-Stoyle and Nair, Phys. Lett. 7 (1963) 161
- 1963BO07 R.O. Bondelid and J.W. Butler, Phys. Rev. 130 (1963) 1078
- 1963BO1J Boschitz, in Padua (1963) 640
- 1963BR14 R.E. Brown, Astrophys. J. 137 (1963) 338
- 1963CA09 R.R. Carlson and E. Norbeck, Phys. Rev. 131 (1963) 1204
- 1963CE02 J. Cerny, R.H. Pehl, E. Rivet and B.G. Harvey, Phys. Lett. 7 (1963) 67
- 1963CO1D Costa et al., Phys. Lett. 6 (1963) 226
- 1963DE19 V.K. Deshpande, Nucl. Phys. 47 (1963) 257
- 1963DO1B Douglas, Sala, Gomes and Polga, in Padua (1963) 558A
- 1963DR1B Drisko, Satchler and Dassel, 3rd Conf. on Reactions between Complex Nuclei (1963) 85
- 1963EV04 J.E. Evans, J.A. Kuehner and E. Almqvist, Phys. Rev. 131 (1963) 1632
- 1963FI1B Finck et al., Z. Phys. 174 (1963) 337
- 1963FR10 G. Frick, A. Gallmann, D.E. Alburger, D.H. Wilkinson and J.P. Coffin, Phys. Rev. 132 (1963) 2169
- 1963FR1D Freeman, Jeronymo, Mani and Sadeghi, J. Phys. 24 (1963) 868
- 1963FU06 H. Fuchs, Z. Phys. 171 (1963) 416
- 1963GE03 D.G. Gerke, D.R. Tilley, N.R. Fletcher and R.M. Williamson, Bull. Amer. Phys. Soc. 8 (1963) 302, D5
- 1963GL1C Glendenning, Ann. Rev. Nucl. Sci. 13 (1963) 191
- 1963GO04 J. Goldemberg and Y. Torizuka, Phys. Rev. 129 (1963) 312
- 1963GU1A Gupta and Waghmare, Nucl. Phys. 48 (1963) 321
- 1963HA1G Haeberli, Fast Neutron Phys., Eds., Marion and Fowler (1963) 1379

- 1963HO1E Hortig, Werner and Gentner, 3rd Conf. on Reactions between Complex Nuclei (1963) 178
- 1963HU02 M.N. Huberman, M. Kamegai and G.C. Morrison, Phys. Rev. 129 (1963) 791
- 1963JA03 J. Jastrzebski, F. Picard, J.P. Schapira and J.L. Picou, Nucl. Phys. 40 (1963) 400
- 1963JE02 J.M.F. Jeronymo, G.S. Mani, F. Picard and A. Sadeghi, Nucl. Phys. 43 (1963) 417
- 1963KO1B Kopaleishvili and Jibuti, Nucl. Phys. 44 (1963) 34
- 1963KU03 D. Kurath, Phys. Rev. 130 (1963) 1525
- 1963KU1B Kunz, Can. J. Phys. 41 (1963) 2187
- 1963LI09 J.C. Lisle, J.O. Newton, W.R. Phillips and F.H. Read, Nucl. Phys. 47 (1963) 56
- 1963LO04 L. Lovitch, Nucl. Phys. 46 (1963) 353
- 1963LU01 C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, Bull. Amer. Phys. Soc. 8 (1963) 12, BA10
- 1963LU1F C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, in Padua (1963) 850A
- 1963MA22 J.H. Manley, Phys. Rev. 130 (1963) 1475
- 1963MA28 G.S. Mani, P.D. Forsyth and R.R. Perry, Nucl. Phys. 44 (1963) 625
- 1963MI02 P.D. Miller, G.C. Morrison, N. Gale and G. Dearnaley, Bull. Amer. Phys. Soc. 8 (1963) 292, A16
- 1963MI1C Mikumo, Nonaka, Yamaguchi and Maki, in Padua (1963) 1088A
- 1963MI1G Mingay and Sellschop, in Padua (1963) 664
- 1963MO04 I.L. Morgan, D.O. Nellis, R. Benjamin and J.B. Ashe, Bull. Amer. Phys. Soc. 8 (1963) 120, M6
- 1963MO1B Morrison, Gale, Hussain and Murray, 3rd Conf. on Reactions between Complex Nuclei (1963) 168
- 1963MO1C Motz and Jurney, Wash-1044 (1963)
- 1963NA04 M.A. Nagarajan, Nucl. Phys. 42 (1963) 454
- 1963NE1C Nemets, Picard, Slyusarenko and Tokarevskii, Zh. Eksp. Teor. Fiz. 45 (1963) 850; Sov. Phys. JETP 18 (1964) 583
- 1963NE1H Nemilov and Pobedonotsev, Zh. Eksp. Teor. Fiz. 45 (1963) 103; Sov. Phys. JETP 18 (1964) 76
- 1963NO1B Nonaka et al., INSJ-57 (1963)
- 1963OH02 G.G. Ohlsen and R.E. Shamu, Nucl. Phys. 45 (1963) 523
- 1963OL1B Ollerhead, Chasman and Bromley, in Padua (1963) 984
- 1963OP1A Oparin, Saukov and Shuvalov, Atomn. Energ. (USSR) 15 (1963) 411; J. Nucl. Energ. 18 (1964) 596

- 1963PE07 R.R. Perry, I.J. Taylor and P.D. Forsyth, Bull. Amer. Phys. Soc. 8 (1963) 302, D4
- 1963PI04 A.E. Pitts, J.D. Bronson, T.A. Belote and G.C. Phillips, Nucl. Phys. 48 (1963) 75
- 1963PR03 F.W. Prosser, Jr., R.W. Krone and J.J. Singh, Phys. Rev. 129 (1963) 1716
- 1963RO17 H.J. Rose, W. Trost and F. Riess, Nucl. Phys. 44 (1963) 287
- 1963SE05 R.F. Seiler, W.J. Ansick, D.F. Herring and K.W. Jones, Bull. Amer. Phys. Soc. 8 (1963) 304, D14
- 1963SE19 T. Sebe, Prog. Theor. Phys. 30 (1963) 290
- 1963SO04 P.C. Sood and Y.R. Waghmare, Nucl. Phys. 46 (1963) 181
- 1963SP1B Sperduto and Bufchner, 2nd Int. Conf. on Nucl. Masses, Vienna, July 1963 (1963)
- 1963TA1B Takeda and Nakasima, 3rd Conf. on Reactions between Complex Nuclei (1963) 159
- 1963TR02 W.W. True, Phys. Rev. 130 (1963) 1530
- 1963VA1E Van Der Zwan, Porterfield and Ritter, Nucl. Instrum. Meth. 24 (1963) 329
- 1963VL1A Vlasov, Zh. Eksp. Teor. Fiz. 45 (1963) 160; Sov. Phys. JETP 18 (1964) 160
- 1963WA15 H.G. Wahsweiler, Z. Phys. 175 (1963) 370
- 1963WI1D Wilkins, UCR-10783 (1963)
- 1963YA1B Yanabu et al., J. Phys. Soc. Jpn. 18 (1963) 747
- 1963ZA1B Zamick, Ann. Phys. 21 (1963) 550
- 1964AB1B Abul-Magd and El-Nadi, Phys. Lett. 13 (1964) 328
- 1964AM1A Amsel, Ann. Phys. 9 (1964) 297
- 1964AM1D Amit and Latz, Nucl. Phys. 58 (1964) 297
- 1964AS03 F. Asfour, I. Bondouk, V.J. Gontchar, V.A. Lutsik, F. Machali and I.I. Zaloubovsky, Nuovo Cim. 32 (1964) 1107
- 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
- 1964BA1G Barz, Ann. Phys. 13 (1964) 164
- 1964BA1P Balashov and Boyarkina, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1964) 359
- 1964BA1T Barker, 5/C32, Paris (1964)
- 1964BE12 J.A. Becker and E.K. Warburton, Phys. Rev. 134 (1964) B349
- 1964BI04 G.R. Bishop and M. Bernheim, Phys. Lett. 8 (1964) 128
- 1964BI09 G.R. Bishop, M. Bernheim and P. Kossanyi-Demay, Nucl. Phys. 54 (1964) 353
- 1964BL1C Bliden, Phys. Lett. 9 (1964) 176
- 1964BO22 E.C. Booth, B. Chasan and K.A. Wright, Nucl. Phys. 57 (1964) 403
- 1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97

- 1964BR1L Bromley, Symp. on Nucl. Spectrosc. with Direct Reactions, ANL 6878 (1964)
- 1964CA05 R.R. Carlson, E. Norbeck and V. Hart, Bull. Amer. Phys. Soc. 9 (1964) 419, DA9
- 1964CA18 R.R. Carlson and M. Throop, Phys. Rev. 136 (1964) B630
- 1964CA1F Carter, Bull. Amer. Phys. Soc. 9 (1964) B630
- 1964CE02 J. Cerny and R.H. Pehl, Phys. Rev. Lett. 12 (1964) 619
- 1964DE1C Deshpande, Fulbright and Verba, Nucl. Phys. 52 (1964) 457
- 1964DI02 G.U. Din, H.M. Kuan and T.W. Bonner, Nucl. Phys. 50 (1964) 267
- 1964DI1C Din, Unpublished Thesis, Rice Univ. (1964)
- 1964DO03 P.F. Donovan, J.F. Mollenauer and E.K. Warburton, Phys. Rev. 133 (1964) B113
- 1964DZ1A Dzubay and Blair, Phys. Rev. 134 (1964) B586
- 1964EA04 L.G. Earwaker and D.F. Hebbard, Nucl. Phys. 53 (1964) 252
- 1964ED01 R.D. Edge, J.R. Stewart and R.C. Morrison, Bull. Amer. Phys. Soc. 9 (1964) 429, EA2
- 1964EN1B Engesser, Thompson and Ferguson, USNRDL TR-791 (1964)
- 1964FE02 J.M. Ferguson, Nucl. Phys. 59 (1964) 97
- 1964FL1D Flerov and Karnaukhov, Congres Int. Phys. Nucl., Paris, 1964 (1964) 373
- 1964FO02 P.D. Forsyth and I.J. Taylor, Bull. Amer. Phys. Soc. 9 (1964) 57, FA15
- 1964FO1A Fowler and Vogl, Lectures in Theor. Phys., Vol. 6 (1964) 379
- 1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49
- 1964GA1B Gallman, Nucl. Instrum. Meth. 28 (1964) 33
- 1964GR19 K.A. Gridnev, A.E. Denisov, Y.A. Nemilov, V.S. Sadkovskii and E.D. Teterin, Zh. Eksp. Teor. Fiz. 46 (1964) 1473; Sov. Phys. JETP 19 (1964) 994
- 1964HA16 B.G. Harvey, E.J-M. Rivet, A. Springer, J.R. Meriwether, W.B. Jones, J.H. Elliott and P. Darriulat, Nucl. Phys. 52 (1964) 465
- 1964HE06 E.M. Henley and D.U.L. Yu, Phys. Rev. 133 (1964) B1445
- 1964HE1F Herczeg, 5/C267, Paris (1964)
- 1964HO1E Honda, Horie, Kudo and Ui, Prog. Theor. Phys. 31 (1964) 424
- 1964HU09 E.E. Hughes and W.B. Mann, Int. J. App. Rad. Isotopes 15 (1964) 97; Nucl. Sci. Abs. 18, 2271, Abs.16961 (1964)
- 1964KI06 C.C. Kim, S.M. Bunch, D.W. Devins and H.H. Forster, Nucl. Phys. 58 (1964) 32
- 1964KO10 R. Kosiek, Z. Phys. 179 (1964) 544
- 1964KO1D Kosiek, Maier and Schlupmann, Phys. Lett. 9 (1964) 260
- 1964KU05 H.-M. Kuan, T.W. Bonner and J.R. Risser, Nucl. Phys. 51 (1964) 481

- 1964KU06 H.-M. Kuan and J.R. Risser, Nucl. Phys. 51 (1964) 518
- 1964KU1D Kuehner and Almqvist, Phys. Rev. 134 (1964) B1229
- 1964KU1F Kurath, Bull. Amer. Phys. Soc. 9 (1964) 628
- 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 379
- 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 411
- 1964LO1B Lovitch, Nucl. Phys. 53 (1964) 477
- 1964LU1B Ludemann, Holmgren and Hornyak, Bull. Amer. Phys. Soc. 9 (1964) 171
- 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
- 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed., W.H. Johnson Jr. (1964) 305
- 1964MC1C McIntosh, Park and Rawitscher, Phys. Rev. 134 (1964) B1016
- 1964MI05 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 63
- 1964MO1D Morgan, Bull. Amer. Phys. Soc. 9 (1964) 653
- 1964MO26 Y. Moriceau and A. Bertin, J. Phys. (Paris) Suppl. 25 (1964) 129A
- 1964NA1C Nataf, Congres Int. Phys. Nucl., Paris, 1964, Vol. 1 (1964) 421
- 1964NA1E Nagatani and Bromley, Bull. Amer. Phys. Soc. 9 (1964) 679
- 1964NE1D Nemets, Pikar, Slyusarenko and Tokarevskyi, Ukr. Fiz. Zh. 9 (1964) 599
- 1964NE1E Neudachin, Shevchenko and Yudin, Phys. Lett. 10 (1964) 180
- 1964PE20 J.L. Perkin, Nucl. Phys. 60 (1964) 561
- 1964PF1A Pfitzner and Riedel, Nucl. Phys. 60 (1964) 672
- 1964RE04 L.H. Reber and J.X. Saladin, Phys. Rev. 133 (1964) B1155
- 1964RI1C Ritter and Porterfield, Bull. Amer. Phys. Soc. 9 (1964) 67
- 1964RO03 H.J. Rose, F. Riess and W. Trost, Nucl. Phys. 52 (1964) 481
- 1964SC09 R.B. Schwartz, H.D. Holmgren, L.M. Cameron and A.R. Knudson, Phys. Rev. 134 (1964) B577
- 1964SE05 H.M. Sen Gupta, J. Rotblat, E.A. King and J.B.A. England, Nucl. Phys. 50 (1964) 549
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TH03 M.J. Throop and R.R. Carlson, Bull. Amer. Phys. Soc. 9 (1964) 406, CA3

- 1964TO1B Toms, Nucl. Phys. 54 (1964) 625
- 1964TR04 W. Trost, H.J. Rose and F. Riess, Phys. Lett. 10 (1964) 83
- 1964UL1A Ullah and Nesbet, Phys. Rev. 134 (1964) B308; Erratum Phys. Rev. 139 (1965) AB2
- 1964WA05 E.K. Warburton, D.E. Alburger, A. Gallmann, P. Wagner and L.F. Chase, Jr., Phys. Rev. 133 (1964) B42
- 1964WA09 E.K. Warburton, J.W. Olness, D.E. Alburger, D.J. Bredin and L.F. Chase, Jr., Phys. Rev. 134 (1964) B338
- 1964WA1G Walker, UCRL 7676 (1964)
- 1964WI03 E.H. Willen, P.W. Keaton and G.E. Owen, Phys. Rev. 133 (1964) B930
- 1964WU01 C.S. Wu, Rev. Mod. Phys. 36 (1964) 618
- 1964YO1B Young, Nucl. Phys. 55 (1964) 84
- 1965AR1E Artemov et al., Yad. Fiz. 1 (1965) 629; Sov. J. Nucl. Phys. 1 (1965) 450
- 1965AR1F Arieux et al., Phys. Lett. 16 (1965) 149
- 1965BA06 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klabes, B. Kuhn and G. Stiller, Nucl. Phys. 64 (1965) 55
- 1965BA1W Barnes, Kuan, Carter and Risser, Bull. Amer. Phys. Soc. 10 (1965) 440
- 1965BA1X Balashov et al., Izv. Akad. Nauk SSSR. Ser. Fiz. 29 (1965) 1177
- 1965BE1B Becker and McIntyre, Phys. Rev. 138 (1965) B339
- 1965BE1E Berggren, Ark. Fys. 30 (1965) 508
- 1965BE1N Benisz, Jasielska and Panek, Acta Phys. Pol. 28 (1965) 763
- 1965BI10 G.R. Bishop, Phys. Lett. 17 (1965) 311
- 1965BL04 R.S. Blake, D.J. Jacobs, J.O. Newton and J.P. Schapira, Phys. Lett. 14 (1965) 219
- 1965BL1E Blin-Stoyle, Nair and Papageorgiou, Proc. Phys. Soc. (London) 85 (1965) 477
- 1965BR1H Bryant and Flynn, Bull. Amer. Phys. Soc. 10 (1965) 515
- 1965BR42 O.D. Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965CA05 R.R. Carlson and R.L. McGrath, Phys. Rev. Lett. 15 (1965) 173
- 1965CA06 R.R. Carlson and D.W. Heikkinen, Phys. Lett. 17 (1965) 305
- 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707
- 1965DA1E Dar, Phys. Rev. 139 (1965) B1193
- 1965DE19 R.W. Detenbeck, J.C. Armstrong, A.S. Figuera and J.B. Marion, Nucl. Phys. 72 (1965) 552
- 1965DE1P Detenbeck, Nucl. Phys. 74 (1965) 199
- 1965DE21 R.W. Detenbeck, Nucl. Phys. 74 (1965) 184

- 1965DO08 V.K. Dolinov, Y.V. Melikov and A.F. Tulinov, *Pisma Zh. Eksp. Teor. Fiz.* 2 (1965) 120; *JETP Lett. (USSR)* 2 (1965) 74
- 1965FA1E Faessler, *Nucl. Phys.* 65 (1965) 329
- 1965FI05 P. Fintz, *Ann. Phys. (Paris)* 10 (1965) 435
- 1965FO06 P.D. Forsyth, I.J. Taylor and R.R. Perry, *Nucl. Phys.* 66 (1965) 376
- 1965FR09 G. Frick, *Ann. Phys. (Paris)* 10 (1965) 155
- 1965FU16 H.W. Fulbright, W.P. Alford, O.M. Bilaniuk, V.K. Deshpande and J.W. Verba, *Nucl. Phys.* 70 (1965) 553
- 1965GA1G Garber and Shrader, *Bull. Amer. Phys. Soc.* 10 (1965) 510
- 1965GI1B Giraud, *Nucl. Phys.* 71 (1965) 373
- 1965GL07 R.N. Glover and A.D.W. Jones, *Phys. Lett.* 18 (1965) 165
- 1965GL09 N.K. Glendenning, *Phys. Rev.* 137 (1965) B102
- 1965GR1F Greider, *Ann. Rev. Nucl. Sci.* 15 (1965) 291
- 1965GR1Q Griffith and Roman, *Phys. Lett.* 14 (1965) 42
- 1965GR1R Gromov et al., *Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P-2184 (1965)*
- 1965HE01 H.J. Hennecke and H.A. Enge, *Bull. Amer. Phys. Soc.* 10 (1965) 9, AC7
- 1965IS04 T. Ishimatsu, S. Morita, T. Tohei, N. Kawai, N. Takano, N. Kato and Y. Yamanouchi, *J. Phys. Soc. Jpn.* 20 (1965) 1112
- 1965IS1A Ishiware, *J. Phys. Soc. Jpn.* 20 (1965) 658
- 1965JO07 B. Johansson, *Nucl. Phys.* 67 (1965) 289
- 1965KAZX R.W. Kavanagh and A. Knipper, *Bull. Amer. Phys. Soc.* 10 (1965) 715, DG11
- 1965KE10 C.A. Kelsey and A.S. Mahajan, *Nucl. Phys.* 71 (1965) 157
- 1965KO1D Koo, *Rev. Mex. Fis.* 14 (1965) 182A
- 1965LA09 J.M. Lacambra, D.R. Tilley, N.R. Roberson and R.M. Williamson, *Nucl. Phys.* 68 (1965) 273
- 1965LO07 J.A. Lonergan and D.J. Donahue, *Nucl. Phys.* 74 (1965) 318
- 1965LU05 A. Luukko, *Soc. Sci. Fennica, Comment. Phys. Math. (Finland)* 31 No. 6 (1965)
- 1965MA1H Mahaux, *Nucl. Phys.* 71 (1965) 241
- 1965MA1K Marion, *Nucl. Phys.* 68 (1965) 463
- 1965MA1N Malvano and Ricco, *Nuovo Cim.* 35 (1965) 484
- 1965MA57 S. Mayo, W. Schimmerling, M.J. Sametband and R.M. Eisberg, *Nucl. Phys.* 62 (1965) 393

- 1965MA59 N.A. Mansour, H.R. Saad, Z.A. Saleh, E.M. Sayed, I.I. Zaloubovsky and V.I. Gontchar, Nucl. Phys. 65 (1965) 433
- 1965MI1B Migdal and Khodel, Yad. Fiz. 8 (1965) 28
- 1965NE06 J.O. Newton, R.S. Blake, D.J. Jacobs and J.P. Schapira, Nucl. Phys. 71 (1965) 113
- 1965PE03 R.H. Pehl, E. Rivet, J. Cerny and B.G. Harvey, Phys. Rev. 137 (1965) B114
- 1965PE09 F. Pellegrini, Nuovo Cim. 38 (1965) 655
- 1965PE17 R.H. Pehl, J. Cerny, E. Rivet and B.G. Harvey, Phys. Rev. 140 (1965) B605
- 1965PU1B Puttaswamy and Kohler, Bull. Amer. Phys. Soc. 10 (1965) 1194
- 1965RI02 F. Riess, W. Trost, H.J. Rose and E.K. Warburton, Phys. Rev. 137 (1965) B507
- 1965RI1A Riou, Rev. Mod. Phys. 37 (1965) 375
- 1965RO1M Rotter, Joint Inst. Nucl. Res. Lab. Theor. Phys., USSR, Rept. No. E-2244 (1965)
- 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96
- 1965RY01 A. Rytz, Nucl. Phys. 70 (1965) 369
- 1965SA07 M.W. Sachs, C. Chasman and D.A. Bromley, Phys. Rev. 139 (1965) B92
- 1965SA18 V.S. Sadkovskii, E.D. Teterin, K.A. Gridnev, A.E. Denisov, R.P. Kolalis and Y.A. Nemilov, Yad. Fiz. 2 (1965) 843; Sov. J. Nucl. Phys. 2 (1966) 601
- 1965SA1H Sawada, Nucl. Phys. 74 (1965) 289
- 1965SE01 K.K. Seth, G. Walter, P.D. Miller and J.A. Biggerstaff, Bull. Amer. Phys. Soc. 10 (1965) 10, AC8
- 1965SH10 S.N. Shumilov, A.P. Klyucharev and N.Y. Rutkevich, Pisma Zh. Eksp. Teor. Fiz. 2 (1965) 213; JETP Lett. (USSR) 2 (1965) 135
- 1965SH14 S.N. Shumilov, A.P. Klyucharev and N.Y. Rutkevich, Zh. Eksp. Teor. Fiz. 49 (1965) 1754; Sov. Phys. JETP 22 (1966) 1198
- 1965SH1E Shapiro and Timashev, Yad. Fiz. 2 (1965) 459
- 1965TA07 T. Takemiya, Prog. Theor. Phys. 34 (1965) 433
- 1965VA13 P.I. Vatset, L.Y. Kolesnikov and S.G. Tonapetyan, Yad. Fiz. 1 (1965) 809; Sov. J. Nucl. Phys. 1 (1965) 579
- 1965VA1K Vaucher et al., Helv. Phys. Acta 38 (1965) 371
- 1965VA23 L. Valentin, Nucl. Phys. 62 (1965) 81
- 1965WA02 B.D. Walker, C. Wong, J.D. Anderson, J.W. McClure and R.W. Bauer, Phys. Rev. 137 (1964) B347
- 1965WA06 E.K. Warburton, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and M.F. Thomas, Phys. Rev. 138 (1965) B104
- 1965WA1J Waghmare and Majumdar, Phys. Lett. 14 (1965) 144

- 1965WI11 J.H. Williamson, Nucl. Phys. 69 (1965) 481
- 1965ZA1B Zamick, Phys. Lett. 19 (1965) 580
- 1965ZH1A Zhdanov, Kuzmin and Yakovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 239
- 1966AL10 D.E. Alburger, A. Gallmann, J.B. Nelson, J.T. Sample and E.K. Warburton, Phys. Rev. 148 (1966) 1050
- 1966AL11 K.W. Allen, T.K. Alexander and D.C. Healey, Phys. Lett. 22 (1966) 193
- 1966AM1B Amos, Nucl. Phys. 77 (1966) 225
- 1966BA13 G.C. Ball and J. Cerny, Phys. Lett. 21 (1966) 551
- 1966BA1A Bahcall, Nucl. Phys. 75 (1966) 10
- 1966BA2P Barnes, 2nd Symp. on the Struct. of Low-Medium Mass Nucl., April 1966 (1966) 242
- 1966BA2R Barz, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintmeister, ZFK 122 (1966) 207
- 1966BA32 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klubes, B. Kuhn and G. Stiller, Nucl. Phys. 85 (1966) 171
- 1966BA33 N.P. Babenko, B.A. Bibichev and Y.A. Nemilov, Yad. Fiz. 3 (1966) 663; Sov. J. Nucl. Phys. 3 (1966) 486
- 1966BA42 F.J. Bartis, Nuovo Cim. B45 (1966) 113
- 1966BA60 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klubes, B. Kuhn and G. Stiller, Nucl. Phys. 84 (1966) 305
- 1966BE07 C. Bergman and R.K. Hobbie, Phys. Rev. 142 (1966) 575
- 1966BE1E H. Beaumeville et al., J. Phys. 27 (1966) C1-150
- 1966BE1P Beach, Finlay, Koshel and Cassola, Bull. Amer. Phys. Soc. 11 (1966) 471
- 1966BL01 R.S. Blake, D.J. Jacobs, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 254
- 1966BO1R O. Bohigas, J. Phys. 27 (1966) C1-39
- 1966BR1G Browne, F. S. U. Isobaric Spin Conf. (1966) 136
- 1966BR1X Brown et al., Bull. Amer. Phys. Soc. 11 (1966) 316
- 1966BR2B G. Bruno, J. Decharge, A. Perrin and G. Surget, J. Phys. 27 (1966) C1-151
- 1966CA07 R.R. Carlson, Phys. Rev. 148 (1966) 991
- 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, Phys. Rev. 152 (1966) 950
- 1966CH1E Chesterfield and Spicer, Bull. Amer. Phys. Soc. 11 (1966) 628; F.S.U. Isobaric Spin Conf. (1966) 433
- 1966CH1J Chen, Durand and McGee, Phys. Rev. 146 (1966) 638
- 1966DA1B Darden, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (1966) 433

- 1966DA1C Daehnick and Denes, Bull. Amer. Phys. Soc. 11 (1966) 30
- 1966DA1E Dalidchik and Sayasov, Yad. Fiz. 3 (1966) 820
- 1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097
- 1966DI03 O. Dietzsch, Nucl. Phys. 85 (1966) 689
- 1966DI04 G.U. Din and J.L. Weil, Nucl. Phys. 86 (1966) 509
- 1966EC1B Eccles, Lutz and Rohn, Bull. Amer. Phys. Soc. 11 (1966) 735
- 1966EV1B Evans, Brown and Marion, Rev. Sci. Instrum. 37 (1966) 991
- 1966FO1D Fowler and Johnson, Bull. Amer. Phys. Soc. 11 (1966) 510; ORNL P-2026 (1966)
- 1966FO1E Fowler, Johnson and Kernal, Bull. Amer. Phys. Soc. 11 (1966) 653
- 1966FR15 J.M. Freeman, J.G. Jenkin, G. Murray and W.E. Burcham, Phys. Rev. Lett. 16 (1966) 959
- 1966FU10 H. Fuchs, K. Grabisch, P. Kraaz and G. Roschert, Phys. Lett. 23 (1966) 363
- 1966GA25 G.T. Garvey and I. Kelson, Phy. Rev. Lett. 16 (1966) 197
- 1966GE03 D.G. Gerke, D.R. Tilley, N.R. Fletcher and R.M. Williamson, Nucl. Phys. 75 (1966) 609
- 1966GL01 R.N. Glover and A.D.W. Jones, Nucl. Phys. 84 (1966) 673
- 1966GL1C R.N. Glover and A.D.W. Jones, Nucl. Phys. 81 (1966) 277
- 1966GO15 S. Gorodetzky, R.M. Freeman, A. Gallman and F. Hass, Phys. Rev. 149 (1966) 801
- 1966GU04 T. Gudehus, M. Cosack, R. Felst and H. Wahl, Nucl. Phys. 80 (1966) 577
- 1966GU08 R.K. Gupta and P.C. Sood, Phys. Rev. 152 (1966) 917
- 1966HA18 E.C. Halbert, Y.E. Kim and T.T.S. Kuo, Phys. Lett. 20 (1966) 657
- 1966HA19 B.G. Harvey, J.R. Meriwether, J. Mahoney, A. Bussiere de Nercy and D.J. Horen, Phys. Rev. 146 (1966) 712
- 1966HA1Q Hahn and Ricci, Bull. Amer. Phys. Soc. 11 (1966) 724
- 1966HA31 O. Hausser, H.J. Rose, J.S. Lopes and R.D. Gill, Phys. Lett. 22 (1966) 604
- 1966HE05 D.W. Heikkinen, Phys. Rev. 141 (1966) 1007; Erratum Phys. Rev. 149 (1966) 990
- 1966HE1E Hehl and Reidel, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 322
- 1966HE1G Herbert, Bull. Amer. Phys. Soc. 11 (1966) 903
- 1966HO11 C.E. Hollandsworth, F.O. Purser, Jr., J.R. Sawers, Jr. and R.L. Walter, Phys. Rev. 150 (1966) 825
- 1966HO15 C.H. Holbrow, R. Middleton and B. Rosner, Phys. Rev. 152 (1966) 970
- 1966HO1F Hoot, Bull. Amer. Phys. Soc. 11 (1966) 316

- 1966JA05 R. Jahr, K. Kayser, A. Kostka and J.P. Wurm, Nucl. Phys. 76 (1966) 79
- 1966JA1A Jacob and Maris, Rev. Mod. Phys. 38 (1966) 121
- 1966JI1A Jiang et al., Chin. J. Phys. 22 (1966) 439
- 1966KA05 T.S. Katman, N.R. Fletcher, D.R. Tilley and R.M. Williamson, Nucl. Phys. 80 (1966) 449
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KL05 A.P. Klyucharev and Y.I. Titov, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 224; Bull. Acad. Sci. USSR Phys. Ser. 30 (1966) 229
- 1966KL1E Klyucharev and Titov, Bull. Acad. Sci. USSR 30 (1966) 234
- 1966KO08 P. Kossanyi-Demay and G.J. Vanpraet, Nucl. Phys. 81 (1966) 529
- 1966KU1D Kuehner, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 146
- 1966LA03 V.A. Latorre and J.C. Armstrong, Phys. Rev. 144 (1966) 891
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LA1P Lambert, Dumazet, Benendetti and Gresillon, Compt. Rend. B262 (1966) 1459
- 1966LA1Q Lambert and Foucou, Compt. Rend. B262 (1966) 1547
- 1966LI07 K.P. Lieb, Nucl. Phys. 85 (1966) 461
- 1966MA04 G.S. Mani and G.C. Dutt, Nucl. Phys. 78 (1966) 613; Erratum Nucl. Phys. A119 (1968) 691
- 1966MA1P MacFarlane, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 411
- 1966MA25 A.I. Malko, N.N. Pucherov and L.S. Saltykov, Yad. Fiz. 3 (1966) 307; Sov. J. Nucl. Phys. 3 (1966) 221
- 1966MA2G Malushinska, Przytula and Sizov, Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR, Rept. No. P3-3079 (1966)
- 1966MA2H Malko and Soroka, Ukr. Fiz. Zh. 11 (1966) 904
- 1966MA57 J.J. Matese and W.R. Johnson, Phys. Rev. 150 (1966) 846
- 1966MA60 J.B. Marion, Rev. Mod. Phys. 38 (1966) 660
- 1966MAZY W.M. MacDonald, Univ. of Maryland, Dept. Phys. Astron., Tech. Rept. 555 (1966)
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966ME09 L. Meyer-Schutzmeister, D. Von Ehrenstein and R.G. Allas, Phys. Rev. 147 (1966) 743
- 1966ME16 M.M. Meier, L.A. Schaller and R.L. Walter, Phys. Rev. 150 (1966) 821
- 1966ME1E Melikov, Vestnik Mosk. Univ. Fiz. Astron. Nov-Dec., No. 6 (1966) 102
- 1966ME1L Messelt, Bull. Amer. Phys. Soc. 11 (1966) 317

- 1966MI1E Miller, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (1966) 410
- 1966MI1G Mikulinsky, Yad. Fiz. 3 (1966) 245
- 1966MO14 G.L. Morgan, R.L. Walter, C.S. Soltesz and T.R. Donoghue, Phys. Rev. 150 (1966) 830
- 1966MO1C Morgan et al., Antwerp 1965 Neutron Conf. (1966) 537
- 1966NG01 D.-C. Nguyen, J. Phys. Soc. Jpn. 21 (1966) 2462; Erratum J. Phys. Soc. Jpn. 22 (1967) 684
- 1966PA1J Parish, Rawlins and Shin, Bull. Amer. Phys. Soc. 11 (1966) 27
- 1966PI02 L.L. Pinsonneault and J.M. Blair, Phys. Rev. 141 (1966) 961
- 1966PO09 A.M. Poskanzer, S.W. Cosper, E.K. Hyde and J. Cerny, Phys. Rev. Lett. 17 (1966) 1271
- 1966PO1E Poth and Bromley, Bull. Amer. Phys. Soc. 11 (1966) 317
- 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246
- 1966RI02 F. Riess and W. Trost, Nucl. Phys. 78 (1966) 385
- 1966RI04 E. Rivet, R.H. Pehl, J. Cerny and B.G. Harvey, Phys. Rev. 141 (1966) 1021
- 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (1966) 253
- 1966RO1E Rotter and Zhusupov, Ann. Phys. 17 (1966) 57
- 1966RO1F Rotter, Ann. Phys. 17 (1966) 247
- 1966RO1R Rosen, Antwerp 1965 Neutron Conf. (1966) 379
- 1966SA05 J.R. Sawers, Jr., F.O. Purser, Jr. and R.L. Walter, Phys. Rev. 141 (1966) 825
- 1966SH1F I.S. Shapiro and S.F. Timashev, Nucl. Phys. 79 (1966) 46
- 1966SH1H Shin, Bull. Amer. Phys. Soc. 11 (1966) 27
- 1966SI02 R.H. Siemssen, G.C. Morrison, B. Zeidman and H. Fuchs, Phys. Rev. Lett. 16 (1966) 1050
- 1966SI05 G.S. Sidhu and J.B. Gerhart, Phys. Rev. 148 (1966) 1024
- 1966SK1A Skakun and Strashinsky, Ukr. Fiz. Zh. 11 (1966) 1265
- 1966SP08 A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 3 (1966) 652; Sov. J. Nucl. Phys. 3 (1966) 477
- 1966ST1L G.L. Strobel, Nucl. Phys. 86 (1966) 535
- 1966ST1N Stout, Blue and Marr, Bull. Amer. Phys. Soc. 11 (1966) 316
- 1966SW01 C.P. Swann, Phys. Rev. 148 (1966) 1119

- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacsson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692
- 1966VI1A J.L. Vidal et al., J. Phys. 27 (1966) C1-128
- 1966WA1C Warburton and Alburger, Nucl. Spin-Parity Assign., Ed., Gove (1966) 114
- 1966WA1E Warburton, F. S. U. Isobaric Spin Conf. (1966) 90
- 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97
- 1966WY01 W.R. Wylie, J.L. Binney, R.M. Bahnsen and H.W. Lefevre, Bull. Amer. Phys. Soc. 11 (1966) 903, P5
- 1966ZA03 L. Zamick, Phys. Lett. 21 (1966) 194
- 1967AD1D Adelberger and McDonald, Bull. Amer. Phys. Soc. 12 (1967) 1143
- 1967AL16 A.U. Aldzhauakhiri, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 6 (1967) 248; Sov. J. Nucl. Phys. 6 (1968) 180
- 1967AN08 B. Antolkovic, G. Paic, D. Rendic and P. Tomas, Izv. Akad. Nauk SSSR Ser. Fiz. 31 (1967) 110; Bull. Acad. Sci. USSR Phys. Ser. 31 (1968) 107
- 1967AR1D Armstrong, Nucl. Research with Low Energy Accelerators, Eds., Marion and Van Patter (1967) 247
- 1967AU05 E.G. Auld, D.G. Crabb, J.G. McEwen, L. Bird, C. Whitehead and E. Wood, Nucl. Phys. A101 (1967) 65
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967AU1E Austin, Beneson and Crawley, Bull. Amer. Phys. Soc. 12 (1967) 1199
- 1967BA03 P.W. Bauer, J.D. Anderson, H.F. Lutz, C. Wong, J.W. McClure and B.A. Pohl, Nucl. Phys. A93 (1967) 673
- 1967BA13 G.C. Ball and J. Cerny, Phys. Rev. 155 (1967) 1170
- 1967BA1D Bastawros, Dissertation Abs. B28 (1967)
- 1967BA2H Backenstoss et al., Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 484
- 1967BA2J H.W. Barz, Nucl. Phys. A91 (1967) 262
- 1967BA78 G. Backenstoss, S. Charalambus, H. Daniel, H. Koch, G. Poelz, H. Schmitt and L. Tauscher, Phys. Lett. B25 (1967) 547
- 1967BE2G Beyea, Nessin, Patton and Lidofsky, Bull. Amer. Phys. Soc. 12 (1967) 502
- 1967BE30 T. Becker, K. Bahr, R. Jahr and W. Kuhlmann, Phys. Lett. B24 (1967) 458
- 1967BI06 J. Birnbaum, J.C. Overley and D.A. Bromley, Phys. Rev. 157 (1967) 787
- 1967BL22 R.S. Blake, E.B. Paul, C.H. Sinex and S.T. Emerson, Nucl. Phys. A102 (1967) 305
- 1967BL24 R.J. Blin-Stoyle and S.C.K. Nair, Nucl. Phys. A105 (1967) 640
- 1967BO1M Bolta, Garcia and Senent, An. Real. Soc. Espan. Fis. y Quim. 63 (1967) 35

- 1967BR1M Breit, Polak and Torchia, Phys. Rev. 161 (1967) 993
- 1967BU1D Bukhvostov and Popov, Yad. Fiz. 6 (1967) 1241
- 1967CA1D Carlson, Nucl. Research with Low Energy Accelerators, Eds., Marion and van Patter (1967) 475
- 1967CH19 C. Chasman, K.W. Jones, R.A. Ristinen and D.E. Alburger, Phys. Rev. 159 (1967) 830
- 1967CH24 B. Chern, T.A. Halpern and L. Logue, Phys. Rev. 161 (1967) 1116
- 1967CH34 V.I. Chuev, V.V. Davidov, A.A. Ogleblin and S.B. Sakuta, Ark. Fys. 36 (1967) 263
- 1967CI1A Cirelli, Marini and Gulmanelli, Nuovo Cim. B47 (1967) 39
- 1967CL07 G. Clausnitzer, R. Fleischmann and H. Wilsch, Phys. Lett. B25 (1967) 466
- 1967CL1C Cloud, Leonard, Gibson and Wells, Bull. Amer. Phys. Soc. 12 (1967) 894
- 1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1
- 1967CR1G Cramer, Braitwaite, Eidson and Slee, Bull. Amer. Phys. Soc. 12 (1967) 1198
- 1967DE03 L.J. Denes and W.W. Daehnick, Phys. Rev. 154 (1967) 928
- 1967DZ01 T.G. Dzubay, Phys. Rev. 158 (1967) 977
- 1967EV1C J.A. Evans and R.P.J. Perazzo, Nucl. Phys. A103 (1967) 225
- 1967FE06 P. Fessenden and D.R. Maxson, Phys. Rev. 158 (1967) 948
- 1967FI07 W. Fitz, R. Jahr and R. Santo, Nucl. Phys. A101 (1967) 449
- 1967FO1A Fowler and Mayes, Proc. Phys. Soc. (London) 92 (1967) 377
- 1967FO1B Fowler, Caughlan and Zimmerman, Ann. Rev. Astron. Astrophys. 5 (1967) 525
- 1967FO1E Focht, Zurmuhle and Fou, Bull. Amer. Phys. Soc. 12 (1967) 35
- 1967FU03 H.W. Fulbright, J.A. Robbins, R. West, D.P. Saylor and J.W. Verba, Nucl. Phys. A94 (1967) 214
- 1967FU04 H. Fuchs and R. Santo, Phys. Lett. B24 (1967) 234
- 1967GA12 A. Gallmann, F. Haas and B. Heusch, Phys. Rev. 164 (1967) 1257
- 1967GR1D L. Grunbaum and N.K. Ganguly, Nucl. Phys. A100 (1967) 645
- 1967GR1L Gray, Fortune, Trost and Fletcher, Bull. Amer. Phys. Soc. 12 (1967) 34
- 1967HA10 K.B. Haque and J.G. Valatin, Nucl. Phys. A95 (1967) 97
- 1967HA1T J.C. Hardy and I.S. Towner, Phys. Lett. B25 (1967) 98
- 1967HA20 R.L. Hahn and E. Ricci, Nucl. Phys. A101 (1967) 353
- 1967HO1K Hodgson and Wilmore, Proc. Phys. Soc. (London) 90 (1967) 361
- 1967HO1L Hodgson, Ann. Rev. Nucl. Sci. 17 (1967) 1
- 1967IN1B Ing and Gupta, Bull. Amer. Phys. Soc. 12 (1967) 666

- 1967IV1B Ivascu, Dumitrescu and Semenescu, Rev. Roum. Phys. 12 (1967) 279
- 1967JA1E B.K. Jain and D.F. Jackson, Nucl. Phys. A99 (1967) 113
- 1967JO1H Jolivette, Bull. Amer. Phys. Soc. 12 (1967) 1172
- 1967KA1A Kabachnik and Grishanova, Sov. J. Nucl. Phys. 4 (1967) 583
- 1967KE1J Keaton et al., Bull. Amer. Phys. Soc. 12 (1967) 1198
- 1967KH1A Khomyakov, Skakun and Strashinsky, Ukr. Fiz. Zh. (USSR) 12 (1967) 375
- 1967KN1B Knudson and Young, Bull. Amer. Phys. Soc. 12 (1967) 502
- 1967KO1D D.S. Koltun, Phys. Rev. 162 (1967) 963
- 1967KO1F Kohman and Bender, High Energy Nucl. Reactions in Astrophys., Ed., B.S.P. Chen (1967) 169
- 1967KO1N Kolata and Galonsky, Bull. Amer. Phys. Soc. 12 (1967) 540
- 1967KO1P Kolibasov and Smorodinskaya, Proc. Prob. Symp. on Nucl. Phys., Tbilisi, April 1967 (1967) 374
- 1967KU1E D. Kurath and R.D. Lawson, Phys. Rev. 161 (1967) 915
- 1967LE1F Lee, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 31
- 1967LE1K Leonard, Cloud, Gibson and Wells, Bull. Amer. Phys. Soc. 12 (1967) 1143
- 1967LI04 K.P. Lieb and R. Hartmann, Z. Phys. 200 (1967) 432
- 1967LI06 R.H. Lindsay and J.J. Veit, Phys. Rev. 157 (1967) 933
- 1967LO1B Loiseaux, Maison and Langevin, J. Phys. 28 (1967) 11
- 1967LU1B Lundberg, Schwarz and Zetterstrom, Ark. Fys. 34 (1967) 247
- 1967MA1G Magelson, Thesis, Univ. of California, Berkeley (1967)
- 1967MC03 W.R. McMurray, P. Van Der Merwe and I.J. Van Heerden, Nucl. Phys. A92 (1967) 401
- 1967MI1B Miller, Thesis, Princeton Univ. (1967)
- 1967MO1P Moore et al., Bull. Amer. Phys. Soc. 12 (1967) 33
- 1967NE06 J.B. Nelson and W.R. Smith, Nucl. Phys. A96 (1967) 671
- 1967OD01 F.H. O'Donnell and C.P. Browne, Phys. Rev. 158 (1967) 957
- 1967OG1A Ogloblin, Proc. Problem Symp. on Nucl. Phys., Tbilisi, April 1967 (1967) 169
- 1967OL02 J.W. Olness, A.R. Poletti and E.K. Warburton, Phys. Rev. 154 (1967) 971
- 1967PA05 J.C. Parikh and N. Ullah, Nucl. Phys. A99 (1967) 529
- 1967PL1B Plough, Springer and Kness, Bull. Amer. Phys. Soc. 12 (1967) 214
- 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, Phys. Rev. 164 (1967) 1295
- 1967PO1E Poth and Bromley, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 94

- 1967PO1J A.R. Poletti, E.K. Warburton and D. Kurath, Phys. Rev. 155 (1967) 1096
- 1967RI1D Riess, O'Connell and Paul, Bull. Amer. Phys. Soc. 12 (1967) 51
- 1967RU1A M.L. Rustgi, Phys. Lett. B24 (1967) 229
- 1967SA1H Sapershtein and Khodel, Proc. Prob. Symp. on Nucl. Phys., Tbilisi, April 1967 (1967) 720
- 1967SC27 L.A. Schaller, R.S. Thomason, N.R. Roberson, R.L. Walter and R.M. Drisko, Phys. Rev. 163 (1967) 1034
- 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
- 1967SC43 G.W. Schweimer, Nucl. Phys. A100 (1967) 537
- 1967SE08 W.A. Seale, Phys. Rev. 160 (1967) 809
- 1967SH05 Y.Y. Sharon, Nucl. Phys. A99 (1967) 321
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SI1F Silverstein and Herling, Bull. Amer. Phys. Soc. 12 (1967) 33
- 1967SO1A Soga, Bull. Amer. Phys. Soc. 12 (1967) 501
- 1967SU1E Suslov, Sov. J. Nucl. Phys. 4 (1967) 854
- 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, Nucl. Instrum. Meth. 56 (1967) 325
- 1967TH1E Thornton, ORNL TM 1917 (1967)
- 1967TI1A Tilley, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 389
- 1967TI1B Timashev, Sov. J. Nucl. Phys. 4 (1967) 192
- 1967VO1A Volkov, Proc. Problem Symp. on Nucl. Phys., Tbilisi, April 1967 (1967) 226
- 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators, Eds., Marion and van Patter (1967) 43
- 1967WO05 C. Wong, J.D. Anderson, J. McClure, B. Pohl, V.A. Madsen and F. Schmittroth, Phys. Rev. 160 (1967) 769
- 1967WY02 W.R. Wylie, R.M. Bahnsen and H.W. Lefevre, Bull. Amer. Phys. Soc. 12 (1967) 894, H10
- 1967YO02 F.C. Young, P.D. Forsyth and J.B. Marion, Nucl. Phys. A91 (1967) 209
- 1967YO1C Young, Nucl. Research with Low Energy Accelerators, Eds., Marion and van Patter (1967) 109
- 1967ZA01 C.D. Zafiratos, J.S. Lilley and F.W. Slee, Phys. Rev. 154 (1967) 887
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968AL12 K.W. Allen, T.K. Alexander and D.C. Healey, Can. J. Phys. 46 (1968) 1575

- 1968BA17 A.M. Baxter, J.A.R. Griffith, S.W. Oh and S. Roman, Nucl. Phys. A112 (1968) 209
- 1968BA19 A.M. Baxter, J.A.R. Griffith and S. Roman, Phys. Rev. Lett. 20 (1968) 1114
- 1968BA1E Ball, UCRL-18263 (1968)
- 1968BA1M Batusov, Bunyatov, Sidorov and Yarba, Sov. J. Nucl. Phys. 6 (1968) 836
- 1968BA2G Backenstoss et al., Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968BA2P Baldweg, Zentralinstitut Kerf. Ross. Bei Dresden, Rept. No. ZFK 140 (1968)
- 1968BA2R Babenko, Yad. Fiz. 7 (1968) 1037
- 1968BA47 N.P. Babenko, K.A. Gridnev and Y.A. Nemilov, Yad. Fiz. 7 (1968) 22; Sov. J. Nucl. Phys. 7 (1968) 15
- 1968BA48 Y.A. Batusov, S.A. Bunyatov, V.M. Sidorov and V.A. Yarba, Yad. Fiz. 7 (1968) 28; Sov. J. Nucl. Phys. 7 (1968) 20
- 1968BE03 H. Behrens and W. Buhring, Nucl. Phys. A106 (1968) 433
- 1968BE2A Beaumevielle, Yaker and Lambert, in Tokyo (1968) 268
- 1968BE30 R.A.I. Bell, R.D. Gill, B.C. Robertson, J.S. Lopes and H.J. Rose, Nucl. Phys. A118 (1968) 481
- 1968BE47 F. Bella, M. Alessio and P. Fratelli, Nuovo Cim. B58 (1968) 232
- 1968BH1C Bhalla, in Tokyo (1968) 402
- 1968BL09 R.S. Blake, G. Johnson, H. Laurent, J.P. Schapira and F. Picard, Nucl. Phys. A117 (1968) 561
- 1968BL1E Black et al., in Tokyo (1968) 375
- 1968BO36 F. Boreli, P.N. Shrivastava, B.B. Kinsey and V.C. Mistry, Phys. Rev. 174 (1968) 1221
- 1968BR1L Brene, Roos and Sirlin, Nucl. Phys. B6 (1968) 255
- 1968CA1A Cassola and Koshel, Nuovo Cim. B55 (1968) 83
- 1968CE1C Cerny and Ball, Bull. Amer. Phys. Soc. 13 (1968) 632
- 1968CH1F Cheon, Proc. Phys. Soc. (London) A1 (1968) 350
- 1968CL05 H.-G. Clerc and E. Kuphal, Z. Phys. 211 (1968) 452
- 1968CO04 R.L.A. Cottrell, J.C. Lisle and J.O. Newton, Nucl. Phys. A109 (1968) 288
- 1968CO10 H. Cords, G.U. Din, M. Ivanovich and B.A. Robson, Nucl. Phys. A113 (1968) 608
- 1968CO13 B.S. Cooper and J.M. Eisenberg, Nucl. Phys. A114 (1968) 184
- 1968CO1W Conde, Bergqvist and Nystrom, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 763
- 1968CO24 J.A. Cookson, Phys. Lett. B27 (1968) 619

- 1968DA1J Daniel, Rev. Mod. Phys. 40 (1968) 659
- 1968DA1Q Dally, Croissiaux and Schweitz, Bull. Amer. Phys. Soc. 13 (1968) 607
- 1968DA20 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Yad. Fiz. 7 (1968) 758; Sov. J. Nucl. Phys. 7 (1968) 463
- 1968DE13 N. de Takacsy, Can. J. Phys. 46 (1968) 2091
- 1968DI06 O. Dietzsch, R.A. Douglas, E.F. Pessoa, V.G. Porto, E.W. Hamburger, T. Polga, O. Sala, S.M. Perez and P.E. Hodgson, Nucl. Phys. A114 (1968) 330
- 1968DO09 T.R. Donoghue, W.L. Baker, P.L. Beach, D.C. DeMartini and C.R. Soltesz, Phys. Rev. 173 (1968) 952
- 1968DU1E Duray and Browne, Bull. Amer. Phys. Soc. 13 (1968) 84
- 1968EI1C Eisenberg and Cooper, Bull. Amer. Phys. Soc. 13 (1968) 18
- 1968FA1A J.C. Faivre, H. Krivine and A.M. Papiau, Nucl. Phys. A108 (1968) 508
- 1968FA1B Faessler, Sauer and Stingl, Z. Phys. 212 (1968) 1
- 1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1
- 1968FR03 N. Freed and P. Ostrander, Nucl. Phys. A111 (1968) 63
- 1968FR08 J.M. Freeman, J.G. Jenkin, D.C. Robinson, G. Murray and W.E. Burcham, Phys. Lett. B27 (1968) 156
- 1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957
- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161
- 1968GE04 B. Geoffrion, N. Marty, M. Morlet, B. Tatischeff and A. Willis, Nucl. Phys. A116 (1968) 209
- 1968GL1B Gleyvod et al., Helv. Phys. Acta 41 (1968) 442
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GO1N Gofman et al., Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 690
- 1968GO1Q Goldring, Loebenstein, Plessner and Sachs, in Tokyo (1968) 206
- 1968GO1T Golovanova and Zelenskaya, Yad. Fiz. 8 (1968) 274
- 1968GR1G Greene, Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968GR1K E.E. Gross, J.J. Malanify, A. van der Woude and A. Zucker, Phys. Rev. Lett. 21 (1968) 1476
- 1968HA1D G. Hauser, R. Lohken, H. Rebel, G. Schatz, W. Schweimer and J. Specht, Phys. Lett. B27 (1968) 220
- 1968HA38 T.A. Halpern and B. Chern, Phys. Rev. 175 (1968) 1314

- 1968HI09 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, G.J. Wagner and P. Turek, Nucl. Phys. A115 (1968) 570
- 1968HI14 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, G.J. Wagner and P. Turek, Nucl. Phys. A111 (1968) 265
- 1968HI1H Hinterberger et al., in Tokyo (1968) 274
- 1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, Sept. 1967 (1968) 41
- 1968HO1H Horie and Hsieh, in Tokyo (1968) 51
- 1968HO1J Honda, Horie and Yoshida, in Tokyo (1968) 210
- 1968HU1H Huber et al., Helv. Phys. Acta 41 (1968) 438
- 1968JA09 J. Janecke, T.F. Yang, R.M. Polichar and W.S. Gray, Phys. Rev. 175 (1968) 1301
- 1968JA1N Jacobson, Bull. Amer. Phys. Soc. 13 (1968) 650
- 1968JA1P Jacobson, Bull. Amer. Phys. Soc. 13 (1968) 1466
- 1968JO07 J.E. Jobst, Phys. Rev. 168 (1968) 1156
- 1968KA1K Katori, Uchida, Imaizumi and Kobayashi, in Tokyo (1968) 269
- 1968KL06 R. Klubes, F. Baldeweg and G. Stiller, Nucl. Phys. A121 (1968) 113; Erratum Nucl. Phys. A127 (1969) 693
- 1968KO1C Kopaleishvili, Machabeli, Gogsadze and Krupennikova, Yad. Fiz. 7 (1968) 292
- 1968KO24 R.P. Kolalis, Y.A. Nemilov, V.S. Sadkovskii, E.D. Teterin and A.E. Denisov, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1739; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1603
- 1968KU1C Kuhlmann, BMWF FBK 68 06 (1968)
- 1968KU1E Kuriyama, Nagata and Bando, in Tokyo (1968) 58
- 1968LA19 C.M. Lamba, N. Sarma and N.S. Thampi, Nucl. Phys. A122 (1968) 390
- 1968LA1J Laird and Huffaker, Bull. Amer. Phys. Soc. 13 (1968) 18
- 1968LE1G Leonhardt, Univ. Lyon. Rept. No. Lycen 6811 (1968)
- 1968MA29 N.F. Mangelson, B.G. Harvey and N.K. Glendenning, Nucl. Phys. A117 (1968) 161
- 1968MA46 K. Matsuda, N. Nakanishi, S. Takeda and T. Wada, J. Phys. Soc. Jpn. 25 (1968) 1207
- 1968ME10 K. Meier-Ewert, K. Bethge and K.-O. Pfeiffer, Nucl. Phys. A110 (1968) 142
- 1968ME1E Meyer, Audebert, Elbaz and Lafoucriere, Compt. Rend. B266 (1968) 969
- 1968MO1F Morita, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 419
- 1968NE1C Nemirovskii, Sov. J. Nucl. Phys. 6 (1968) 29
- 1968NO1A M.E. Nordberg, K.F. Kinsey and R.L. Burman, Phys. Rev. 165 (1968) 1096

- 1968NO1C J.V. Noble, Phys. Rev. 173 (1968) 1034
- 1968OD1B Oda et al., in Tokyo (1968) 278
- 1968PA1Y Park et al., Bull. Amer. Phys. Soc. 13 (1968) 1447
- 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
- 1968RA10 S. Radhakant and N. Ullah, Nucl. Phys. A116 (1968) 43
- 1968RA1C Rahman, Khan and Sen Gupta, Nuovo Cim. B54 (1968) 260
- 1968RI1H A. Richter and R.A. Chatwin, Phys. Lett. B27 (1968) 181
- 1968RI1Q A. Richter and L.J. Parish, Phys. Rev. Lett. 21 (1968) 1824
- 1968RI1R Riess, O'Connell and Paul, Bull. Amer. Phys. Soc. 13 (1968) 885
- 1968RO1C Rose, Hausser and Warburton, Rev. Mod. Phys. 40 (1968) 591
- 1968RO1D I. Rotter, Nucl. Phys. A122 (1968) 567
- 1968SC1H G. Schiffner, Nucl. Phys. A113 (1968) 367
- 1968SE1E E. Seibt, Ch. Weddigen and K. Weigle, Phys. Lett. B27 (1968) 567
- 1968SH11 P.N. Shrivastava, F. Boreli and B.B. Kinsey, Phys. Rev. 169 (1968) 842
- 1968SO1B Soga, in Tokyo (1968) 50
- 1968ST1U Stokes and Young, Bull. Amer. Phys. Soc. 13 (1968) 652
- 1968TA1C Tanner, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 91
- 1968TA1N Takeuchi and Sanada, TUENS-4 (1968)
- 1968TA1V Taketani, Muto, Yamaguchi and Kokame, in Tokyo (1968) 277
- 1968TH04 T.D. Thomas, G.M. Raisbeck, P. Boerstling, G.T. Garvey and R.P. Lynch, Phys. Lett. B27 (1968) 504
- 1968TH1J S.T. Thornton, C.M. Jones, J.K. Bair, M.D. Mancusi and H.B. Willard, Phys. Rev. Lett. 21 (1968) 447
- 1968TO09 J.H. Towle and G.J. Wall, Nucl. Phys. A118 (1968) 500
- 1968WI1B Wilkinson, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 469
- 1968YA1F Yamada and Kodama, in Tokyo (1968) 400
- 1968YU01 T.J. Yule and W. Haeberli, Nucl. Phys. A117 (1968) 1
- 1968ZA02 N.I. Zaika, Y.V. Kibkalo, A.V. Mokhnach, O.F. Nemets, P.L. Shmarin and A.M. Yasnogorodskii, Yad. Fiz. 7 (1968) 754; Sov. J. Nucl. Phys. 7 (1968) 460
- 1968ZE1B Zelenskaya et al., Sov. J. Nucl. Phys. 6 (1968) 47

- 1968ZH05 M.A. Zhusupov, O. Lkhagva and I. Rotter, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1714; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1579
- 1968ZU1A Zupancic, Proc. Symp. on Use of Nimrod, 1968; RHEL/R166 (1968) 67
- 1969AL04 W.P. Alford and K.H. Purser, Nucl. Phys. A132 (1969) 86
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969AT1A Atkinson and Bloom, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 799
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969BA17 K. Bahr, T. Becker, R. Jahr and W.R. Kuhlmann, Nucl. Phys. A129 (1969) 388
- 1969BA1N Bahcall and Fowler, Astrophys. J. 157 (1969) 659
- 1969BA1P Bacon et al., Acta Cryst. A25 (1969) 391
- 1969BA1Z Barnes, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 179
- 1969BA23 N. Baron, R.F. Leonard and D.A. Lind, Phys. Rev. 180 (1969) 978
- 1969BE08 H. Beaumevieille, M. Lambert, M. Yaker, A. Amokrane and Nguyen van Sen, Nucl. Phys. A125 (1969) 568
- 1969BE92 N. Bezic, D. Brajnik, D. Jamnik and G. Kernel, Nucl. Phys. A128 (1969) 426
- 1969BL1E R.J. Blin-Stoyle, Phys. Lett. B29 (1969) 12
- 1969BO1G S. Boffi and F.D. Pacati, Nucl. Phys. A129 (1969) 673
- 1969BO32 I. Borbely, T. Dolinszky, J. Ero and G. Hrehuss, Acta Phys. Acad. Sci. Hung. 26 (1969) 269
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como 1967 (1969) 242
- 1969BR1E Bruckmann, Kluge and Schanzler, Z. Phys. 221 (1969) 379
- 1969BR1N Braithwaite, Cameron, Cramer and Rudy, Bull. Amer. Phys. Soc. 14 (1969) 1220
- 1969CA1A R.R. Carlson and H.W. Wyborny, Phys. Rev. 178 (1969) 1529
- 1969CH04 P.W. Chudleigh, C.K. Gowers and E.G. Muirhead, Nucl. Phys. A123 (1969) 114
- 1969CH1C D.T. Chivers, E.M. Rimmer, B.W. Allardyce, R.C. Witcomb, J.J. Domingo and N.W. Tanner, Nucl. Phys. A126 (1969) 129
- 1969CO02 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A127 (1969) 95
- 1969CO12 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A134 (1969) 561
- 1969CO1D Comfort, Fortune, Morrison and Zeidman, Bull. Amer. Phys. Soc. 14 (1969) 507
- 1969CU08 J.R. Curry, W.R. Coker and P.J. Riley, Phys. Rev. 185 (1969) 1416
- 1969CU1D Curtis, Lutz, Bartolini and Heikkinen, Bull. Amer. Phys. Soc. 14 (1969) 508

- 1969DA06 J. Damgaard, Nucl. Phys. A130 (1969) 233
- 1969DA1P Danielopoulos et al., Bull. Amer. Phys. Soc. 14 (1969) 506
- 1969DE16 J. Dey, J.P. Elliott, A.D. Jackson, H.A. Mavromatis, E.A. Sanderson and B. Singh, Nucl. Phys. A134 (1969) 385
- 1969DI1B Dickens and Perey, Nucl. Sci. Eng. 36 (1969) 280
- 1969ED1C Edge, Private Communication (1969)
- 1969EL1B Elliott, Proc. Int. Conf., Montreal (1969) 277
- 1969ET01 K. Etoh, T. Murata, N. Kawai, R. Chiba and S. Takayanagi, J. Phys. Soc. Jpn. 26 (1969) 1335, and Private Communication (1969)
- 1969FO1C Fortune, Dehnhard, Siemssen and Zeidman, Bull. Amer. Phys. Soc. 14 (1969) 487
- 1969FR1E Freed and Ostrander, Contrib., Montreal (1969) 178
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA01 A. Gallmann, F. Hibou and P. Fintz, Nucl. Phys. A123 (1969) 27
- 1969GA11 P. Gaillard, R. Bouche, L. Feuvrais, M. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
- 1969GA16 A. Gallmann, F. Haas, B. Heusch and M. Toulemonde, Rev. Phys. Appl. 4 (1969) 216
- 1969GA1G Garvey, Ann. Rev. Nucl. Sci. 19 (1969) 433
- 1969GO11 S. Gorodetzky, F.A. Beck, P. Engelstein, T. Byrski and A. Knipper, Nucl. Instrum. Meth. 69 (1969) 163
- 1969GO14 V. Gomes Porto, N. Ueta, R.A. Douglas, O. Sala, D. Wilmore, B.A. Robson and P.E. Hodgson, Nucl. Phys. A136 (1969) 385
- 1969GU02 H. Guratzsch, G. Hofmann, H. Muller and G. Stiller, Nucl. Phys. A129 (1969) 405
- 1969HA14 G. Hauser, R. Lohken, H. Rebel, G. Schatz, G.W. Schweimer and J. Specht, Nucl. Phys. A128 (1969) 81
- 1969HA2D Haas, Thesis, Univ. of Strasbourg (1969)
- 1969HA49 F. Haas, B. Heusch, A. Gallmann and D.A. Bromley, Phys. Rev. 188 (1969) 1625
- 1969HE06 H.-D. Helb, H. Voit, G. Ischenko and W. Reichardt, Phys. Rev. Lett. 23 (1969) 176
- 1969HO1Y Houk, Bull. Amer. Phys. Soc. 14 (1969) 494
- 1969HO23 C.H. Holbrow, R. Middleton and W. Focht, Phys. Rev. 183 (1969) 880
- 1969IW1D Iwao, DPKU-028-68 (1969)
- 1969JA06 R.J. Jaszczak, R.L. Macklin and J.H. Gibbons, Phys. Rev. 181 (1969) 1428
- 1969JO09 J. Jobst, S. Messelt and H.T. Richards, Phys. Rev. 178 (1969) 1663
- 1969KA1B R.W. Kavanagh, Nucl. Phys. A129 (1969) 172

- 1969KO1J V.M. Kolybasov and N.Ya. Smorodinskaya, Nucl. Phys. A136 (1969) 165
- 1969LI1D T.K. Lim, Nucl. Phys. A129 (1969) 259
- 1969LO1F L.J. Logue, Phys. Lett. B29 (1969) 642
- 1969MA1G Madsen, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 149
- 1969MA1R Mairle et al., Nucl. Isospin. Proc. 1969 Asilomar Conf. (1969) 291
- 1969MA2D Malanify, Gross, Van Der Woude and Zucker, Bull. Amer. Phys. Soc. 14 (1969) 506
- 1969MO1E L. Moyer and D.S. Koltun, Phys. Rev. 182 (1969) 999
- 1969NO1B J.V. Noble, Phys. Rev. Lett. 22 (1969) 473, Erratum Phys. Rev. Lett. 22 (1969) 1028
- 1969NO1C Noble, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 233
- 1969NY1A Nyberg, Jonsson and Bergqvist, Research Inst. Nat. Defence, Stockholm, NP 6902 (1969)
- 1969PA1J Parish, Richter and Love, Bull. Amer. Phys. Soc. 14 (1969) 1207
- 1969PA1K Paetz Gen. Schieck et al., Bull. Amer. Phys. Soc. 14 (1969) 507
- 1969PE09 C.A. Pearson and J.C. Wilcott, Phys. Rev. 181 (1969) 1477
- 1969PE1L Pearson, Bang and Pocs, Ann. Phys. 52 (1969) 83
- 1969PE1N C.A. Pearson, J.C. Wilcott and L.C. McIntyre, Nucl. Phys. A125 (1969) 111
- 1969PL1C Pledger, Thesis, Univ. of Wisconsin (1969)
- 1969RA1B M. Rahman, N.C. Dutta Banik and H.M. Sen Gupta, Nucl. Phys. A125 (1969) 449
- 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
- 1969RO1R Roughton and Taylor, Bull. Amer. Phys. Soc. 14 (1969) 1242
- 1969SA1L Y. Sakamoto, Phys. Lett. B29 (1969) 88
- 1969SA1M Saltmarsh, Goodman and Rapaport, Bull. Amer. Phys. Soc. 14 (1969) 1207
- 1969SH1A Sharon, Bull. Amer. Phys. Soc. 14 (1969) 35
- 1969SM03 H.V. Smith, Jr. and H.T. Richards, Phys. Rev. Lett. 23 (1969) 1409
- 1969SN02 F.D. Snyder and M.A. Waggoner, Phys. Rev. 186 (1969) 999
- 1969SN04 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1259
- 1969SO07 M. Sosnowski, Nucl. Phys. A133 (1969) 266
- 1969SO08 B. Sorensen, Nucl. Phys. A134 (1969) 1
- 1969SO1E Sorensen, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 239
- 1969SU1E Suzuki, Prog. Theor. Phys. 41 (1969) 695
- 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
- 1969UL03 N. Ullah and S.S.M. Wong, Phys. Rev. 188 (1969) 1645

- 1969VA1C S. Varma and P. Goldhammer, Nucl. Phys. A125 (1969) 193
- 1969VE02 V.V. Verbinski and W.R. Burrus, Phys. Rev. 177 (1969) 1671
- 1969VE1D Velkley et al., Bull. Amer. Phys. Soc. 14 (1969) 1200
- 1969VI02 D. Vinciguerra and T. Stovall, Nucl. Phys. A132 (1969) 410
- 1969VO1E Vogt, Proc. Int. Conf., Montreal (1969) 5
- 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
- 1969WA1F Waghmare, Joshi and Mehta, Contrib., Montreal (1969) 186
- 1969WA1H Warburton and Weneser, Private Communication (1969)
- 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
- 1969YA1C Yang, Janecke, Gray and Polichar, Bull. Amer. Phys. Soc. 14 (1969) 1228
- 1969YO1B Young, Phillips and Marion, Private Communication (1969)
- 1970GA09 A. Gallman, F. Haas, N. Balaux, B. Heusch and M. Toulemonde, Can. J. Phys. 48 (1970) 1595
- 1970LO1A Lokan et al., Bull. Amer. Phys. Soc. 15 (1970) 85
- 1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1
- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- DE65C Unknown Source
- ED69D Unknown Source
- GA69J Unknown Source
- HO66 Unknown Source

