

Energy Levels of Light Nuclei

$A = 14$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 13\text{--}15$ was published in *Nuclear Physics A268* (1976), p. 1. This version of $A = 14$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed January 31, 1976)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 14$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [¹⁴Li](#), [¹⁴Be](#), [¹⁴B](#), [¹⁴C](#), [¹⁴N](#), [¹⁴O](#), [¹⁴F](#)

B. Tables of Recommended Level Energies:

[Table 14.1](#): Energy levels of ¹⁴B

[Table 14.3](#): Energy levels of ¹⁴C

[Table 14.11](#): Energy levels of ¹⁴N

[Table 14.29](#): Energy levels of ¹⁴O

C. References

D. Figures: [¹⁴B](#), [¹⁴C](#), [¹⁴N](#), [¹⁴O](#), Isobar diagram

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{14}Li

(Not illustrated)

^{14}Li has not been observed: it is calculated to be particle unstable with a binding energy of -2.66 MeV for decay into $^{13}\text{Li} + \text{n}$ and of -3.23 MeV for decay into $^{12}\text{Li} + 2\text{n}$. The calculated mass excess is 72.29 MeV ([1974TH01](#)).

^{14}Be

(Figs. 5 and 9)

^{14}Be has been observed with a production cross section of $\approx 10 \mu\text{b}$ in the 4.8 GeV proton bombardment of uranium ([1973BO30](#), [1974BO05](#)). It is particle stable: its atomic mass excess is calculated to be 40.69 MeV. ^{14}Be is then bound by 2.73 and 0.41 MeV, respectively, with respect to decay into $^{13}\text{Be} + \text{n}$ and $^{12}\text{Be} + 2\text{n}$ ([1974TH01](#)). See, however, ([1975JE02](#)). ^{14}Be had not been observed in the bombardment of ^{232}Th with 145 MeV ^{15}N ions ([1970AR27](#)). See also ([1971AR1P](#), [1973VO1D](#), [1973KO1D](#), [1973TA30](#)) and ([1971ST40](#), [1972TH13](#), [1975BE31](#); theor.).

^{14}B

(Figs. 5 and 9)

GENERAL: (See also ([1970AJ04](#))).

See ([1970AR1D](#), [1970AR27](#), [1971AR02](#)), ([1973KO1D](#), [1973TO16](#)) and ([1972TH13](#), [1973WI15](#), [1974TH01](#), [1975JE02](#), [1975BE31](#), [1975MI12](#); theor.).

Mass of ^{14}B : From the Q -value of the $^{14}\text{C}(^{7}\text{Li}, ^{7}\text{Be})^{14}\text{B}$ reaction the atomic mass excess of ^{14}B is 23.657 ± 0.030 MeV. ^{14}B is then stable with respect to breakup into $^{13}\text{B} + \text{n}$ and $^{12}\text{B} + 2\text{n}$ by 0.98 and 5.86 MeV, respectively ([1973BA34](#)). The lowest $T = 2$ states in ^{14}C , ^{14}N and ^{14}O are estimated to lie at $E_x = 22.5$, 24.8 and 22.5 MeV. These states would then be stable against allowed neutron decay by ≈ 0.7 MeV and unstable to proton decay by < 3.0 MeV ([1973BA34](#)).



^{14}B has a half-life of 16.1 ± 1.2 msec. The β -decay is primarily to $^{14}\text{C}^*(6.09)$ [$J^\pi = 1^-$] and (6.73) [$J^\pi = 3^-$]: see Table [14.2](#). The measured γ -ray intensity ratios are $I_{6.09}/I_{6.73}/I_{7.34} = 100/(10.0 \pm 2.0)/< 2.2$. The nature of the decay fixes the J^π of ^{14}B to be 2^- ([1974AL11](#)). See also ([1975MI12](#); theor.).



Table 14.1: Energy levels of ^{14}B

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ (msec)	Decay	Reactions
g.s.	$2^-; 2$	16.1 ± 1.2 msec	β^-	1, 2, 3, 4
0.74 ± 40	$(1^-); 2$			4
1.38 ± 30	$(3^-); 2$			4
1.82 ± 60	$(2^-); 2$			4
2.08 ± 50	$(4^-); 2$			4
(2.32 ± 40)				4
2.97 ± 40	$T = 2$			4

 Table 14.2: Beta decay of ^{14}B ^a

Decay to $^{14}\text{C}^*$ (MeV)	J^π	Branch (%)	$\log ft$
0	0^+	(5 ± 3) ^d	
6.09 ^b	1^-	81 ± 9	4.22 ± 0.05
6.73	3^-	$8.6^{+1.7}_{-4.0}$	$5.10^{+0.30}_{-0.08}$
7.34	2^-	< 11 ^e	> 4.9
c			

^a (1974AL11).

^b E_{β^-} -(max) = 14.0 ± 0.7 MeV to this state.

^c A search for possible delayed neutrons following the population of higher states has not yet been carried out.

^d This branch has not been observed. It is assumed to be $5 \pm 3\%$ in the calculation of the branching ratios to $^{14}\text{C}^*(6.09, 6.73)$. However, (1975MI12; theor.) suggest that the branch may be as small as $\approx 0.3\%$. The errors shown for the branching ratios reflect this uncertainty (1974AL11).

^e This branch has not been observed: the upper limit is shown. In the calculations of the branching ratios to $^{14}\text{C}^*(6.09, 6.73)$ a value $5 \pm 5\%$ was used.

^{14}B has been populated in this reaction at $E(^6\text{Li}) = 31$ MeV ([1974AL11](#)).



See ([1974AL11](#)).



^{14}B states with $0 < E_x < 3$ MeV have been populated in this reaction at $E(^7\text{Li}) = 52$ MeV: see Table [14.1](#). Similarities in the relative intensities of $^{14}\text{B}^*(0, 0.74, 1.38, 1.82, 2.08)$ and of $^{12}\text{B}^*(1.67, 2.62, 3.39, 4.37, 4.52)$ [populated in $^{12}\text{C}(^7\text{Li}, ^7\text{Be})^{12}\text{B}$], and the similarity in the ΔE_x of these ^{12}B states with the E_x of the ^{14}B states suggest that they have the same J^π . The differential cross section for producing $^{14}\text{B}_{\text{g.s.}}$ at $\theta = 15^\circ$ is $\approx 4 \mu\text{b}/\text{sr}$ ([1973BA34](#)).

¹⁴C
(Figs. 6 and 9)

GENERAL: (See also (1970AJ04).)

Nuclear models: (1970FR13, 1970FR1C, 1970KH01, 1971BA2Y, 1972LI06, 1973KU03, 1973SA30, 1975MI12, 1975VE01).

Special reactions involving ¹⁴C: (1969GA18, 1969KR21, 1971AR02, 1972VO06, 1973KO1D, 1973KU03, 1973VO1G, 1973WI15, 1974BE16, 1974KO25, 1975HU14, 1975KU01, 1975UD01).

Reactions involving muons: (1970DM01, 1970VA24, 1973BE16, 1973KI12, 1973MU05, 1973MU11, 1973MU1B, 1974FU1B).

Reactions involving pions (See also reactions 17, 33, 41.): (1970DO04, 1971KO23, 1973DA37, 1973KA1D, 1974LI15, 1975EI1B, 1975HU1D, 1975RE01).

Other topics: (1970FO1B, 1970GU13, 1970KH01, 1970SU1B, 1971BA2Y, 1971JA14, 1971ST40, 1972AN05, 1972CA37, 1973AL1B, 1973GO1H, 1973HO32, 1973KI05, 1973KU03, 1973MA48, 1973PA1F, 1974CA1H, 1974DZ03, 1974HA1C, 1975KU01, 1975MO1M, 1975VE01, 1976MA04).

Ground state: (1974DE1E, 1975BE31).

Mass of ¹⁴C: 14.003241983 (12) amu (1975SM02).



Measured $\tau_{1/2}$ are 5745 ± 50 y (1961MA1L, 1961MA32, 1964HU09), 5780 ± 65 y (1961WA16), 5680 ± 40 y (1962OL04), 5660 ± 30 y (1968BE47), 5736 ± 56 y (1972EM01): the recommended value of $\tau_{1/2} = 5730 \pm 40$ y [†]. Using Q_m , $\log ft = 9.02$ (1969KA1B). The spectrum does not deviate from the allowed shape down to 3 keV (1954MO84). The large $\log ft$ difference between this and the mirror transition in $^{14}\text{O}(\beta^+)$ cannot be accounted for by the differences in the binding energies of ^{14}C and ^{14}O (1972WI05). For other discussions of the long lifetime of ^{14}C see (1959AJ76, 1970AJ04) and (1976GOZX). See also (1970SO08) and (1969EL1A, 1970FR13, 1971HU10, 1973HA49, 1973KU1J, 1974ZA1G; theor.).

2. (a) $^7\text{Li}(^7\text{Li}, 2n)^{12}\text{C}$	$Q_m = 13.674$	$E_b = 26.797$
(b) $^7\text{Li}(^7\text{Li}, n)^{13}\text{C}$	$Q_m = 18.620$	
(c) $^7\text{Li}(^7\text{Li}, p)^{13}\text{B}$	$Q_m = 5.966$	
(d) $^7\text{Li}(^7\text{Li}, d)^{12}\text{B}$	$Q_m = 3.311$	

[†] W. Mann, private communication.

- (e) ${}^7\text{Li}({}^7\text{Li}, \text{t}){}^{11}\text{B}$ $Q_m = 6.199$
 (f) ${}^7\text{Li}({}^7\text{Li}, \alpha){}^{10}\text{Be}$ $Q_m = 14.784$
 (g) ${}^7\text{Li}({}^7\text{Li}, {}^8\text{Be}){}^6\text{He}$ $Q_m = 7.278$
 (h) ${}^7\text{Li}({}^7\text{Li}, {}^7\text{Li}){}^7\text{Li}$

Table 14.3: Energy levels of ${}^{14}\text{C}$ ^a

E_x in ${}^{14}\text{C}$ (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
g.s.	$0^+; 1$	$\tau_{1/2} = 5730 \pm 40$ y	β^-	1, 2, 3, 4, 8, 9, 10, 12, 13, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 42, 44, 45, 46, 47
6.0942 ± 1.6	1^-	$\tau_m < 10$ fsec	γ	2, 3, 4, 5, 9, 10, 13, 17, 18, 20, 24, 27, 34, 39
6.5898 ± 1.6	0^+	> 0.6 psec	γ	2, 3, 4, 9, 10, 17, 18, 27
6.7282 ± 1.3	3^-	97 ± 15 psec $ g = 0.272 \pm 0.007$	γ	2, 3, 4, 5, 9, 10, 11, 18, 20, 21, 24, 27, 34, 39, 41
6.9023 ± 1.8	0^-	36 ± 4 fsec	γ	2, 3, 4, 5, 10, 18, 41
7.0120 ± 4.2	2^+	13 ± 2 fsec	γ	3, 4, 9, 10, 26, 27, 33, 39
7.3414 ± 3.1	2^-		γ	3, 4, 9, 10, 18, 21, 27, 39
8.3183 ± 0.9	2^+	$\Gamma_{\text{c.m.}} = 3.4 \pm 0.6$	γ, n	3, 4, 9, 10, 13, 14, 17, 18, 26, 27, 33, 39
9.799 ± 7	$(3, 1)$	45 ± 12	n	3, 9, 14, 18, 26, 27, 39
10.437 ± 9	$(2^+, 3)$	16	n	3, 9, 14, 18, 39
10.509 ± 13	(4^+)	26 ± 8	n	3, 9, 12, 14, 18, 27
10.743 ± 5		20 ± 7		3, 9, 10
11.306 ± 15	$1^{(-)}$	46 ± 12		3, 26, 33, 39
11.397 ± 15	$(2^+, 3)$	22 ± 7		3
11.667 ± 15	(5^-)	20 ± 7		3, 9, 39
11.740 ± 20				3
11.9 ± 300		950 ± 300	n	14, 18
12.589 ± 16		105 ± 15		3, 18, 39
12.860 ± 14	$(4, 5)$	30 ± 10		3, 18

Table 14.3: Energy levels of ^{14}C ^a (continued)

E_x in ^{14}C (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
12.964 \pm 14	(3, 4)	30 \pm 10		3, 18
14.667 \pm 20	(2 ⁺ , 3, 4, 5, 6 ⁺)	57 \pm 15		3, 9
14.867 \pm 25				3, 39
15.19 \pm 30				3, 9
(15.37 \pm 30)				3
15.44 \pm 40				3
(16.02 \pm 50)				3
16.411 \pm 20				3
(16.57 \pm 40)				3
16.72 \pm 30				3
(17.28 \pm 40)				3
17.95 \pm 40				3
18.10 \pm 40				3
b				
23.624 \pm 36	($T = 2$)	120 $^{+120}_{-60}$		6

^a See also Tables 14.6 and 14.10.

^b See also reactions 7 and 33.

These reactions have been studied with $E(^7\text{Li})$ to 6.5 MeV: see (1970AJ04) for the earlier references. For $E(^7\text{Li})$ = 2.3 to 5.8 MeV, the cross section for emission of α_0 , α_1 and α_{2+3+4} is found to increase monotonically with energy (1964DZ1A, 1969CA1A). However, (1971WY01) reports several broad structures in the 0° yield of α_0 and α_1 for $E(^7\text{Li})$ = 2 to 20 MeV: it is suggested that these are neither compound nucleus resonances nor Ericson-type fluctuations but that they are due to a forward-direction cluster transfer process (1971WY01).



Observed proton groups are displayed in Table 14.4 (1973AJ01). Angular distributions have been measured at $E(^6\text{Li})$ = 1.9 MeV (1969KO1A: $p_0, p_1, p_{2+3+4+5}, p_6, p_7, p_8$) and at 20 MeV ((1973AJ01) and unpublished results: to most states shown in Table 14.4). See also ^{15}N .

Table 14.4: Levels of ^{14}C from $^9\text{Be}(^6\text{Li}, \text{p})^{14}\text{C}$ ^a

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	σ ^b (μb)	$2J_f + 1$ ^c	J^π ^d	θ_n^2 ^e
6.089 \pm 10		71	2.5 [2]		
6.588 \pm 10		30	1.0 [1]		
6.726 \pm 10		220	7.6 [7]		
6.899 \pm 10		31	1.1 [1]		
7.016 \pm 10		130	4.5 [5]		
7.341 \pm 10		151	5.2 [5]		
8.318 \pm 10	22 \pm 6	146	5.1	2 ⁺	0.08 \pm 0.02
9.796 \pm 10	45 \pm 12	223	7.7	3	0.4 \pm 0.1 if $\pi = +$ 0.04 \pm 0.01 if $\pi = -$
10.441 \pm 15		312	10.8	2 ^{+, 3}	
10.512 \pm 15	26 \pm 8	262	9.1	4	0.08 \pm 0.02 if $\pi = +$ 1.3 \pm 0.4 if $\pi = -$
10.743 \pm 15	20 \pm 7	444	15.4		
11.306 \pm 15	46 \pm 12	70	2.4	1 ⁻	0.012 \pm 0.003
11.397 \pm 15	22 \pm 7	179	6.2	2 ^{+, 3}	
11.667 \pm 15	20 \pm 7	358	12.4	5 ⁻	0.20 \pm 0.07
11.74 \pm 20					
12.57 \pm 25	80 \pm 20	435	15.1		
12.867 \pm 20	30 \pm 10	300	10.4	4, 5	
12.970 \pm 20	30 \pm 10	225	7.8	3, 4	
14.667 \pm 20	57 \pm 15			2 ^{+, 3, 4, 5, 6⁺}	
14.867 \pm 25					
15.19 \pm 30					
(15.37 \pm 30)					
15.44 \pm 40					
(16.02 \pm 50)					
16.411 \pm 20					
(16.57 \pm 40)					
16.715 \pm 30					
(17.28 \pm 40)					

Table 14.4: Levels of ^{14}C from ${}^9\text{Be}({}^6\text{Li}, \text{p}){}^{14}\text{C}$ ^a (continued)

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	σ ^b (μb)	$2J_f + 1$ ^c	J^π ^d	θ_n^2 ^e
17.95 ± 40					
18.10 ± 40					

^a (1973AJ01): $E({}^6\text{Li}) = 20$ MeV.

^b Total cross section for formation of this state; absolute value: $\pm 20\%$ except for the last six values, $\pm 30\%$.

^c The first number gives $2J_f + 1$, based on a best fit to the experimentally determined values for the cross section of the states with known spins. These $2J_f + 1$ values are determined to $\pm 10\%$, except for the last six values which are determined to $\pm 20\%$. The second number, in brackets, gives $2J_f + 1$ derived from the J_f assignments shown in Table 14.3.

^d Suggested from the $2J_f + 1$ rule and comparison of predicted neutron width with observed $\Gamma_{\text{c.m.}}$ assuming $0.01 < \theta_n^2 < 1.0$.

^e Calculated from neutron penetration of the centrifugal barrier, assuming that the total width is the neutron decay width Γ_n .

$$4. {}^9\text{Be}({}^7\text{Li}, \text{d}){}^{14}\text{C} \quad Q_m = 10.101$$

Angular distributions of the deuterons to ${}^{14}\text{C}^*(0, 6.09, 6.59 + 6.73, 6.90 + 7.01, 7.34, 8.32)$ have been measured at $E({}^7\text{Li}) = 5.6, 5.8, 6.0$ and 6.2 MeV (1969SN02). Angular distributions for the d_0 group have also been measured at $E({}^7\text{Li}) = 1.7$ to 1.9 MeV (1969KA1C). The γ -decay of six bound excited states has been studied by (1966CA07). (1969TH01) report $E_\gamma = 6094.5 \pm 3.2$, 6728.1 ± 1.4 and 7011.7 ± 5.2 keV for the ground state transitions of these states and $\tau_m < 120$ fsec for ${}^{14}\text{C}^*(7.01)$.

$$5. {}^9\text{Be}({}^{13}\text{C}, {}^8\text{Be}){}^{14}\text{C} \quad Q_m = 6.512$$

At $E({}^{13}\text{C}) = 9$ MeV ${}^{14}\text{C}^*(6.09, 6.73, 6.90)$ have been populated: $\tau_m < 10, > 5 \times 10^4$ and 36 ± 4 fsec, respectively (1975SE04). See also (1975SE03).

$$6. {}^{10}\text{Be}(\alpha, \text{p}){}^{13}\text{B} \quad Q_m = -8.818 \quad E_b = 12.013$$

Table 14.5: ^{14}C levels from $^{12}\text{C}(\text{t}, \text{p})^{14}\text{C}$

E_x^a (MeV \pm keV)	(1964MI05) ^b L	(1960JA17) ^c L
0	0	0
6.090 ± 10	(2)	1
6.582 ± 10	1	1
6.725 ± 10	(2)	3
6.893 ± 10	e	e
7.009 ± 10	(2)	0
7.335 ± 10	e	e
8.32 ^a	2	
10.74 ± 20^d		

^a The excitation energies of the first six excited states have been determined by (1960JA17); the seventh comes from the $^{13}\text{C}(\text{n}, \text{n})^{13}\text{C}$ work of (1961CO05); the eighth has been determined by (1964MI05).

^b $E_t = 11$ MeV; except ground state $E_t = 8$ to 13 MeV.

^c $E_t = 5.5$ MeV.

^d $\Gamma < 15$ keV. No states are observed between this level and the broad state at 11.9 MeV (1964MI05).

e Weak group.

The yield of 3.7 MeV γ -rays coincident with $E_\beta > 4$ MeV (from the ^{13}B decay) has been measured for $E_\alpha = 13.6$ to 17 MeV: a strong interference anomaly is observed which appears to correspond to $E_x = 23.624 \pm 0.036$ MeV, $\Gamma_{\text{c.m.}} = 120^{+120}_{-60}$ keV. It is suggested that it is a $T = 2$ state whose analog is either $^{14}\text{B}^*(0.74)$ or (1.38) (1974GO1T).

$$7. \ ^{11}\text{B}(\text{t}, \text{n})^{13}\text{C} \quad Q_m = 12.4214 \quad E_b = 20.5984$$

Resonant structure has been reported in the yield of neutrons with $E_t = 1.20$ to 2.32 MeV: these are thought to correspond to ten excited states in ^{14}C in the range $21.54 < E_x < 22.42$ MeV (1965VA13).

$$8. \ ^{11}\text{B}(\alpha, \text{p})^{14}\text{C} \quad Q_m = 0.7837$$

Angular distributions of p_0 have been measured at $E_\alpha = 1.43$ to 2.94 MeV ([1976DA1U](#)), 2.5 to 5.0 MeV ([1963MA28](#)) and 10.8 to 25.1 MeV ([1969SP02](#)). See also ([1967SP09](#)), ([1971BU1K](#); theor.) and ^{15}N .



At $E(^7\text{Li}) = 5$ MeV, α -particle groups are observed to the known states of ^{14}C with $E_x < 10$ MeV except $^{14}\text{C}^*(6.90)$, and to the unresolved $10.4 - 10.5$ MeV states. There is some indication also of ^{14}C states at $(10.71), (11.35), (11.66), (14.15), (14.73)$ and (15.07) MeV (\pm approx. 50 keV). Angular distributions have been obtained for the α -particles to $^{14}\text{C}^*(0, 6.09, 8.32)$ ([1966MC05](#)).



Observed proton groups are displayed in Table [14.5](#). Angular distributions have been measured at $E_t = 5.5$ MeV ([1960JA17](#)), 8 to 13 MeV ([1964MI05](#)) [see Table [14.5](#)] and at 16 and 20 MeV ([1972KE02](#): p_0). Particle-gamma correlations have been studied by ([1968BE30](#)): see Table [14.6](#). The lifetime of $^{14}\text{C}^*(6.73)$ is 97 ± 15 psec; the $(6.73 \rightarrow 0)$ E3 transition is enhanced by 3.3 ± 0.8 W.u. ([1968AL12](#)). The nuclear gyromagnetic ratio for $^{14}\text{C}^*(6.73)$ is $|g| = 0.272 \pm 0.007$ showing that the state is not of a pure $(\text{p}_{1/2}^{-3}\text{d}_{5/2})$ configuration ([1974AL07](#)).

See also ([1969AR1B](#)), ([1974BA65](#); theor.) and ^{15}N .



At $E(^{12}\text{C}) = 114$ MeV $^{14}\text{C}^*(6.73, 10.25)$ [$J^\pi = 3^-$ and 4^+ , respectively] are populated ([1972SC21](#), [1974AN36](#)). See also ([1971SC1F](#), [1972MO1E](#), [1973SC1J](#)).



For reaction (b) see ([1970BA1J](#), [1974CH1Q](#)). See also ([1970AJ04](#)), ([1974GO1L](#)) and ([1970AN1D](#), [1971AL1D](#), [1971RO1G](#), [1972RO35](#), [1975TA02](#); theor.). For reaction (a) see ([1975NA11](#)).




Table 14.6: Branching ratios of γ -rays in ^{14}C

E_i (MeV)	J_i^π	E_f (MeV)	Branch (%)	
			A	B
6.09	1^-	0		
6.59	0^+	0	1.0 ± 0.4 ^c	
		6.09	99.0 ± 0.4 ^d	
6.73	3^-	0	97.3 ± 1	93 ± 1
		6.09	7 ± 2	2.7 ± 1
		6.09	100 ^{a,e}	
7.01	2^+	0		98.6 ± 0.7
		6.09		1.4 ± 0.7
		6.09		
7.34	2^-	0	18 ± 4	14 ± 4
		6.09	47 ± 4 ^f	52 ± 5 ^b
		6.73	35 ± 7	34 ± 4 ^b

A: (1966AL10): $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$.

B: (1968BE30): $^{12}\text{C}(\text{t}, \text{p})^{14}\text{C}$.

^a (1958WA02).

^b $\delta(M2/E1) = -0.04 \pm 0.09$ and $+0.07 \pm 0.30$, respectively (1968BE30).


^c Internal pairs.

^d $E_\gamma = 495.8 \pm 0.4$ keV (1966AL10).

^e $E_\gamma = 808.7 \pm 1.0$ keV (1966AL10).

^f $E_\gamma = 1248 \pm 3$ keV (1966AL10).

The thermal capture cross section is 0.9 ± 0.2 mb (1973MU14). The neutron capture yield has been measured for $E_n = 95$ to 235 keV. The parameters for $^{14}\text{C}^*(8.32)$ are $E_n = 152 \pm 1$ keV, $\Gamma_n = 5 \pm 1$ keV, $\Gamma_\gamma = 4.0 \pm 1.6$ eV [see Table 14.7]. The calculated cross section for $kT = 30$ keV is 0.12 ± 0.04 mb suggesting that ^{13}C is not significantly involved as a neutron poison in s-process nucleosynthesis (1971AL09). See also (1970CL1C, 1973CL1E; astrophys. questions). Two γ -rays are observed with $E_\gamma = 8.174 \pm 0.002$ and 6.093 ± 0.002 MeV [$E_x = 6.094 \pm 0.002$ MeV], with intensities of 87 ± 5 and $13 \pm 1\%$. Intensities of transitions via other ^{14}C states are $< 2\%$ (1967TH05).



$$E_b = 8.170$$

Table 14.7: Resonances in $^{13}\text{C}(\text{n}, \gamma)$ and (n, n)

E_{res} (keV)	E_{x} (MeV ± keV)	Γ_{lab} (keV)	Γ_{γ} (eV)	σ^{a} (bn)	l_{n}	J^{π}
$152 \pm 1^{\text{b}}$	8.3183 ± 0.9	5 ± 1	$4.0 \pm 1.6^{\text{f}}$			
$152.9 \pm 1.4^{\text{c}}$		$3.7 \pm 0.7^{\text{f}}$		21 ± 2	1	2^{+}
$1751 \pm 8^{\text{d}}$	9.802 ± 8	20		$[\approx 1.3]$	> 0	1
$2432 \pm 10^{\text{d}}$	10.434 ± 10	17		$[\approx 1.3]$		2
$2454 \pm 10^{\text{d}}$	10.454 ± 10	10		$[\approx 0.7]$		≥ 1
3800 ^e	11.7					

^a Corrected peak cross section above background.

^b (1971AL09): (n, γ) .

^c (1975HE02).

^d (1961CO05).

^e Broad resonance structure.

^f Recommended by (1973MU14). I am indebted to S.F. Mughabghab for a very useful discussion.

The coherent scattering length (thermal, bound) is 6.0 ± 0.2 fm (1973MU14). The total cross section has been measured for $E_{\text{n}} = 0.10$ to 9 MeV and 16 to 23 MeV: observed resonances are displayed in Table 14.7 (1961CO05, 1975HE02). See also (1973MU14).

$$15. \ ^{13}\text{C}(\text{n}, \text{t})^{11}\text{B} \quad Q_{\text{m}} = -12.4214 \quad E_{\text{b}} = 8.1770$$

See (1973BI1B).

$$16. \ ^{13}\text{C}(\text{n}, \alpha)^{10}\text{Be} \quad Q_{\text{m}} = -3.836 \quad E_{\text{b}} = 8.1770$$

See (1970AJ04).

$$17. \ ^{13}\text{C}(\text{p}, \pi^{+})^{14}\text{C} \quad Q_{\text{m}} = -132.174$$

At $E_{\text{p}} = 185$ MeV, an angular distribution has been measured for the transition to $^{14}\text{C}_{\text{g.s.}}$: it is forward peaked and goes through a minimum at 90° [while the π^- to $^{14}\text{O}_{\text{g.s.}}$ show a flat distribution: see reaction 6 in ^{14}O]. There is evidence also for the population of $^{14}\text{C}^*(6.09, 8.32, 14.9 \pm 0.2)$ as well as some other structure (1973DA37). See also (1970DO04).

Table 14.8: Proton groups from $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

E_x (MeV \pm keV)	(1954SP01, 1955MC75) l_n ^d	J^π	(1958WA02, 1959WA04) J^π
0	1	$0^+, 1^+, 2^+$	0^+
6.091 ± 10	0	$0^-, 1^-$	1^-
6.589 ± 20	$1, 2, 3$ ^e	$(1^-, 2, 3^-)$	
6.723 ± 10 ^a	2	$1^-, 2^-, 3^-$	$3^-(2^-)$
6.894 ± 10 ^a	$0, 1$ ^e	$0, 1, 2^+$	0^-
7.346 ± 20	2	$1^-, 2^-, 3^-$	$2^-, 3^-$
8.321 ± 20			
9.800 ± 20			
10.433 ± 20			
10.505 ± 20			
11.9 ± 300 ^b			
12.601 ± 20 ^c			
12.854 ± 20			
12.958 ± 20			

^a The spacing of these two levels is 171 ± 3 keV (1954SP01).

^b $\Gamma_{\text{lab}} = 1.10 \pm 0.30$ MeV.

^c $\Gamma_{\text{lab}} = 0.130 \pm 0.020$ MeV.

^d See also (1959AJ76).

^e See footnotes 18 and 31 in (1958WA02).

$$18. \ ^{13}\text{C}(\text{d}, \text{p})^{14}\text{C} \quad Q_m = 5.9524$$

Observed proton groups are displayed in Table 14.8. Angular distributions have been measured at a number of energies up to $E_d = 14.8$ MeV: recent measurements have been carried out for the p₀ group at $E_p = 0.41$ to 0.81 MeV (1971PU01) and 12.1 to 14.0 MeV (1969GU1C; prelim. results). See also (1970LI1E, 1971LI1K, 1975ZA06). For listings of θ_n^2 see Table 14.5 in (1970AJ04).

Gamma rays are exhibited in Table 14.6: studies of these, of the angular distributions analyzed by DWBA, and of p γ correlations lead to the following J^π assignments [see reaction 14 in (1970AJ04) for a full discussion of the evidence and of the relevant references]. $^{14}\text{C}^*(6.09)$ is 1^- (decay is E1); $^{14}\text{C}^*(6.59)$ is 0^+ (internal pairs only); $^{14}\text{C}^*(6.73)$ is 3^- (γ_0 is E3; $l_n = 2$); $^{14}\text{C}^*(6.90)$ is 0^- (no γ_0 ; 0.81 MeV cascade via 6.09 is predominantly dipole; $\gamma_{0.8} + \gamma_{6.1}$ correlation is consistent with $J = 0$, and plane polarization leads to negative parity); $^{14}\text{C}^*(7.34)$ is 2^- (strength of cascade

decay and angular correlation results): see, in particular, ([1958WA02](#), [1965LA09](#), [1966AL10](#), [1966RI02](#), [1966WA1C](#)). $\tau_m < 15$ fsec ([1974RU1B](#)), < 40 fsec ([1973RE1E](#), [1973RE1F](#); prelim. results) for $^{14}\text{C}^*(6.09)$. See also ([1967SP09](#)) and ^{15}N .



See ([1961BA10](#)).




At $E(^7\text{Li}) = 34$ MeV the angular distribution has been measured for the ground state transition. $^{14}\text{C}^*(6.09, 6.73)$ and possibly other states are also populated ([1973SC26](#)).



At $E(^{11}\text{B}) = 114$ MeV $^{14}\text{C}_{\text{g.s.}}$ is weakly excited but $^{14}\text{C}^*(6.73, 7.34)$ [$J^\pi = 3^-$ and 2^- , respectively] are strongly populated ([1974AN36](#)). See also ([1972MO1E](#), [1973SC1J](#)).



See ([1971GA05](#)) and ([1973TO05](#); theor.) for reaction (a). For reaction (b) see ^{14}N , reaction 60 ([1975GA17](#)).



See ([1974CH1Q](#)).




^{14}B decays primarily to $^{14}\text{C}^*(6.09, 6.73)$: see Table 14.2. $\tau_{1/2} = 16.1 \pm 1.2$ msec ([1974AL11](#)).



See ([1973KI05](#); theor.).



The elastic scattering has been studied at $E_e = 374.6$ MeV: the charge radius $r_{\text{rms}} = 2.56 \pm 0.05$ fm ([1973KL12](#)). At $E_e = 101.2$ MeV, $^{14}\text{C}^*(7.01, 8.32)$ [both $J^\pi = 2^+$] are excited strongly: $\Gamma_\gamma = 50.8 \pm 7.1$ and 4.1 ± 1.7 meV ([1972CRZN](#); prelim. results). There is some evidence also for excitation of other states of ^{14}C including $^{14}\text{C}^*(9.8, 11.3)$ ([1973CR1B](#); prelim. results; see also ([1975FA1A](#))), however, there is no evidence of sharp structure in the region $E_x = 19$ to 24 MeV where the $T = 2$ analog state of $^{14}\text{B}_{\text{g.s.}}$ would be expected ([1974KL12](#): $E_e = 61.5$ and 80.7 MeV). See also ([1973FA1K](#), [1974DE1E](#)).




The elastic angular distribution has been measured by ([1971CU01](#)) at $E_p = 14.5$ MeV (reaction (a)) and by ([1967NE06](#)) at $E_d = 3.4$ MeV (reaction (b)). The population of $^{14}\text{C}^*(6.09, 6.58, 6.72, 7.01 \pm 0.02, 7.34, 8.32, 9.80, 10.5)$ has been reported by ([1959AR1A](#)) at $E_d = 14.9$ MeV. Angular distributions have also been studied at $E_d = 17$ MeV to $^{14}\text{C}^*(0, 6.73, 7.01)$ ([1975CE04](#)).



The elastic angular distribution has been measured at $E(^3\text{He}) = 10, 12, 15$ and 18 MeV ([1970DU07](#)). See also ([1971CO14](#)).



Elastic angular distributions have been studied at $E_\alpha = 22, 24$ and 28 MeV ([1972OE01](#), [1973OE01](#)).



The elastic scattering for reaction (a) has been studied at $E(^{12}\text{C}) = 12, 15, 18$ and 20 MeV and for reaction (b) at 15 MeV ([1972BO68](#)). See also ([1973FE03](#)), ([1974GO1L](#)) and ([1973MC1J](#), [1973VO04](#), [1975DE09](#), [1975VO1B](#); theor.).



See ([1971BO1U](#)) and ([1971VO1D](#), [1975DE09](#), [1975VO1B](#); theor.).



The elastic scattering angular distributions have been measured at $E(^{16}\text{O}) = 20, 25$ and 30 MeV ([1974KO1P](#)).



Branching ratios have been measured for the capture of stopped pions to $^{14}\text{C}^*(0, 7.0 \pm 0.1, 8.3, 11.3, 20.0 \pm 1.0)$ ([1975BA52](#)): the total radiative capture branching is $2.1 \pm 0.2\%$.



$^{14}\text{C}^*(6.09)$ decays with $E_\gamma = 6098 \pm 5 \text{ keV}$, $\tau_m < 0.1 \text{ psec}$ ([1972NY02](#)), and $E_\gamma = 6732 \pm 5 \text{ keV}$ from the decay of $^{14}\text{C}^*(6.73)$ ([1971NY03](#)). The p_0 angular distribution has been measured at $E_n = 14 \text{ MeV}$ ([1971CO32](#)). See also ([1970AJ04](#)), ([1970DI1A](#)), ([1973CL1E](#); astrophys. questions) and ^{15}N .



See ([1970AJ04](#)).



See ([1971BAWQ](#)).

Table 14.9: Levels of ^{14}C from the $^{15}\text{N}(\text{d}, ^3\text{He})$ reaction ^a

E_x (MeV)	l	J^π ^a	C^2S
g.s.	1		1.10
6.12 ± 0.04	b		d
6.73 ± 0.04	b		(< 0.11)
7.02 ± 0.03	1		1.13
7.34 ± 0.04	b		(< 0.04)
8.32	1	2^+	0.86
9.78 ± 0.04	b		
10.44 ± 0.03		$((0, 1, 2)^+$	0.39
11.29 ± 0.04	1	1^+	1.23
11.67 ± 0.05	b		
12.08 ± 0.10	b		
12.60 ± 0.08	b		
14.81 ± 0.08			
c			

^a (1971KA18).

^b Dominant non-stripping reaction mechanism.

^c Several other weak transitions occur up to $E_x \approx 18$ MeV.

^d C^2S for $^{14}\text{C}^*(6.59) < 0.01$.



For $E_{bs} = 19.6$ to 38.3 MeV ^{14}C is reported to be formed preferentially in the ground state and in excited states with $E_x = 7.0 \pm 0.2$ and 10.7 MeV (1971DE42). See also (1973DE30) and ^{15}N . For reaction (b) see (1975MU07).



The angular distributions of n_0 have been measured at $E_n = 14.1$ and 14.8 MeV (1967FE06, 1971MI12).




^{14}C states populated in this reaction at $E_{\text{d}} = 52$ MeV are displayed in Table 14.9. The table also shows the l -values obtained from the analysis of the angular distributions ([1971KA35](#)).



Angular distributions of the α_0 group have been measured at $E_{\text{t}} = 0.82$ to 1.62 MeV ([1973AB1E](#)). See also ^{18}O in ([1978AJ03](#)).



At $E_{\pi^-} = 230$ MeV, γ -rays from the population of $^{14}\text{C}^*(6.73, 6.90)$ are observed ([1974LI15](#)). See also ([1971KO23](#)).



See ([1973CE1B](#)).



See ([1972KU1H](#); theor.).



Angular distributions are reported at $E(^{18}\text{O}) = 28$ and 32 MeV ([1971BA68](#)).



See ^{18}O in ([1978AJ03](#)) and ([1973CL1E](#); astrophys. relevance).




Table 14.10: Lifetimes of bound excited states of ^{14}C

$^{14}\text{C}^*$	τ_m	Reaction	Refs.
6.09 ^a	< 15 fsec	$^{13}\text{C}(\text{d}, \text{p})$	(1974RU1B)
	< 10 fsec ^A	$^9\text{Be}(^{13}\text{C}, ^8\text{Be})$	(1975SE04)
6.59	> 0.6 psec	$^{13}\text{C}(\text{d}, \text{p})$	(1966AL10)
6.73	> 50 psec	$^9\text{Be}(^{13}\text{C}, ^8\text{Be})$	(1975SE04)
	97 \pm 15 psec ^A	$^{12}\text{C}(\text{t}, \text{p})$	(1968AL12)
6.90	36 \pm 4 fsec ^A	$^9\text{Be}(^{13}\text{C}, ^8\text{Be})$	(1975SE04)
7.01	< 120 fsec	$^9\text{Be}(^7\text{Li}, \text{d})$	(1969TH01)
	13 \pm 2 fsec ^A	$^{14}\text{C}(\text{e}, \text{e})$	(1972CRZN)

^A = adopted value.

^a See also (1972NY02).

See (1975BE01), (1970AJ04) and (1974DO03; theor.).



See (1967DE03).



Not observed: see (1962LA15).

¹⁴N
(Figs. 7 and 9)

GENERAL: (See also (1970AJ04).)

Shell model: (1970CO1H, 1970FR13, 1970HS02, 1970UL01, 1971NO02, 1972LE1L, 1972LI06, 1973IG02, 1973KU03, 1973SA30, 1974KI1B, 1975DI04, 1975MI12, 1975VE01).

Collective and deformed models: (1972LA12).

Cluster model: (1969BA1J, 1969KU1B, 1970KO26, 1971NO02, 1972LE1L, 1973KU03, 1975CH1V).

Special levels: (1969HA1F, 1971HS05, 1971JA13, 1971MC15, 1971NO02, 1972LA12, 1972LI06, 1973SA30, 1974BO22, 1974NI1A, 1974VA24, 1975CH1V, 1975DI04, 1975MI12).

Giant resonance: (1971JA14, 1973KI05, 1974FA1A, 1974HA1C, 1974MU13, 1974VE10, 1975VE01).

Electromagnetic transitions: (1969HA1F, 1969WA1C, 1970HS02, 1971JA13, 1972BE1E, 1972LI06, 1972LO1D, 1972TA21, 1973HA49, 1973SA30, 1974HA1C, 1974MU13, 1974PI1C, 1974VE10, 1975VE01).

Special reactions: (1969GA18, 1971AR02, 1971HE24, 1973KU03, 1973WI15, 1974KO25, 1975AR14, 1975HU14, 1975KA07, 1975KU01, 1975ME1F, 1975RE08, 1976YOZZ).

Astrophysical questions: (1972CL1A, 1973AR1E, 1973AU1B, 1973AU1D, 1973AU1C, 1973BO1R, 1973CA1B, 1973CO1B, 1973IB1B, 1973MI1D, 1973RA1D, 1973SA1J, 1973SC1T, 1973SM1A, 1973TA1E, 1973TA1D, 1973TR1B, 1974BE1R, 1974SN1B, 1974WI1F, 1975AU1D, 1975CO1Q, 1975DW1A, 1975EN1A, 1975GO1Y, 1975IB1A, 1975JA1F, 1975KA1L, 1975KE1A, 1975NO1D, 1975PE1E, 1975PR1B, 1975RA1M, 1975RY1A, 1975SC1H, 1975SH1N, 1975SN1A, 1975TR1A, 1975UL1A).

Muon and neutrino capture and reactions: (1970BU1D, 1970DM01, 1970VA24, 1971BA96, 1972BU29, 1973BA68, 1973KI12, 1973MU05, 1973MU11, 1973MU1B, 1974DU02, 1974EN10, 1974FU1B, 1975CA1H, 1975CH22).

Pion capture and reactions (See also reaction 60.): (1969BA1L, 1969GO1C, 1969MA1C, 1970BA44, 1970BA1E, 1970DO04, 1970KO26, 1970NE1F, 1971AM1A, 1971FA09, 1971KO23, 1971MA38, 1971MA1T, 1971MA2A, 1972AG01, 1972FA14, 1972HU1A, 1972KO31, 1972SW1A, 1972YO1C, 1973AG05, 1973AG06, 1973AR1B, 1973BA2G, 1973BU1B, 1973CH20, 1973EI01, 1973GR1F, 1973KA19, 1973KA1D, 1973NY04, 1973ST1K, 1974HU14, 1974KO07, 1974LE25, 1974LI15, 1974NE18, 1974ST1G, 1974TA18, 1974ZA05, 1975AR02, 1975BA52, 1975DE1D, 1975EI1B, 1975VE05, 1976JA04).

Kaon capture: (1969KU1B, 1971MA1T, 1972WA1F, 1974CA1H).

Applied work: (1975BE1U, 1975OL1A).

Table 14.11: Energy levels of ^{14}N ^a

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$1^+; 0$	stable		1, 6, 7, 9, 10, 18, 19, 20, 21, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 57, 59, 60, 61, 63, 64, 65, 66, 67, 68, 70, 71, 72
2.31287 ± 0.07	$0^+; 1$	$\tau_m = 87 \pm 6$ fsec	γ	1, 9, 10, 18, 19, 21, 26, 32, 33, 34, 35, 38, 39, 41, 42, 43, 44, 47, 56, 57, 59, 60, 61, 64, 65, 66
3.9478 ± 0.4	$1^+; 0$	4.5 ± 0.4 fsec	γ	1, 6, 7, 9, 10, 18, 19, 20, 21, 26, 32, 33, 34, 35, 38, 39, 42, 43, 44, 45, 47, 48, 57, 59, 61, 64, 65, 66
4.9150 ± 1.3	$0^-; 0$	7.6 ± 1.4 fsec	γ	1, 6, 7, 18, 19, 20, 21, 32, 33, 34, 39, 41, 42, 43, 44, 45, 47, 48, 57, 59, 65
5.10587 ± 0.18	$2^-; 0$	(12.4 ± 1.4) psec	γ	6, 7, 18, 19, 20, 21, 26, 32, 33, 34, 39, 42, 43, 44, 45, 47, 48, 57, 59, 61, 64, 65
5.6896 ± 1.1	$1^-; 0$	10.0 ± 2.0 fsec	γ	6, 7, 18, 19, 20, 21, 32, 33, 39, 42, 43, 44, 45, 47, 48, 57, 59, 65
5.8324 ± 1.4	$3^-; 0$	18 ± 2 psec	γ	6, 7, 11, 18, 19, 20, 21, 24, 32, 33, 39, 42, 43, 44, 45, 47, 48, 57, 59, 61, 65
6.2035 ± 0.6	$1^+; 0$	124 ± 15 fsec	γ	1, 6, 7, 18, 19, 20, 21, 26, 32, 33, 39, 44, 45, 47, 48, 57, 59, 65
6.4444 ± 1.1	$3^+; 0$	627 ± 33 fsec	γ	1, 6, 7, 18, 19, 20, 21, 32, 33, 39, 43, 44, 45, 48, 57, 59, 65
7.0279 ± 1.4	$2^+; 0$	5.4 ± 0.5 fsec	γ	6, 7, 18, 19, 20, 21, 32, 33, 39, 41, 43, 44, 45, 47, 48, 57, 59, 64, 65
7.9666 ± 0.6	$2^-; 0$	$\Gamma = (2.5 \pm 0.7) \times 10^{-3}$	γ, p	6, 7, 18, 19, 20, 21, 26, 32, 33, 44, 48, 57, 59, 65

Table 14.11: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
8.062 ± 1.0	$1^-; 1$	30 ± 1	γ, p	18, 19, 26, 27, 32, 41, 44, 57, 59
8.4877 ± 1.2	$4^-; 0$	27×10^{-6}	γ, p	6, 7, 18, 19, 20, 21, 26, 32, 33, 44, 65
8.618 ± 2	$0^+; 1$	7 ± 1	γ, p	18, 19, 26, 27, 32, 33, 44, 57, 59, 65
8.79 ± 50	$0^-; 1$	≈ 460	γ, p	26, 27
8.9091 ± 1.6	$3^-; 1$	19.7 ± 1.9	γ, p	19, 26, 27, 32, 33, 42, 44, 57, 59
8.9612 ± 1.3	$5^+; 0$	7×10^{-6}	γ, p	11, 19, 20, 21, 23, 24, 26, 52, 57
8.979 ± 3	$2^+; (0)$	8 ± 2	γ, p	6, 19, 26, 27, 32, 57
9.1241 ± 1.5	3^+			6, 19
9.129 ± 1.0	$2^-; 0$	< 1	γ, p	26, 32
9.1708 ± 1.6	$2^+; 1$	0.074 ± 0.008	γ, p	19, 21, 26, 32, 41, 42, 44, 57, 59, 64
9.3860 ± 1.4	$2^-, 3^-; 0$	15.6 ± 2.0	γ, p	6, 19, 20, 21, 26, 27, 32, 44, 57, 59, 65
9.509 ± 3	$2^-; 1$	41 ± 2	γ, p	19, 26, 27, 32, 44, 57, 59, 65
9.703 ± 4	$1^+; 0$	15 ± 3	p	6, 19, 20, 21, 27, 32, 44, 57, 59, 65
10.063 ± 15	$3^+, \geq 4$	< 10	p	19, 57
10.101 ± 15	$2^+, 1^+; 0$	5	γ, p	6, 19, 20, 21, 26, 27, 44, 57, 65
10.228 ± 10	$1^{(-)}; 0$	80 ± 15	γ, p	19, 26, 27, 57, 65
10.434 ± 6	$2^+; 1$	33 ± 3	γ, p	6, 19, 26, 27, 39, 42, 57, 59, 64, 65
(10.54)	(1)	(140)	p	19, 27
10.811 ± 7	$5^+; 0$	$(0.39 \pm 0.16) \times 10^{-3}$	γ, p	19, 20, 21, 26, 57, 65
11.00 ± 30		165 ± 30	γ, p	26, 42
11.050 ± 5	$(2, 3)^+$	≤ 2	γ, p	6, 19, 26, 42, 57
11.07	$1^+; 0$	100	n, p, d	6, 13, 20, 21, 26, 28, 64, 65
11.24 ± 20	; 1	220 ± 30	γ, p, d	13, 19, 24, 26, 44, 45, 47, 57, 65

Table 14.11: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
11.24	(3 ⁻)	≈ 20	n, p	27, 28, 57
11.29	2 ⁻ ; 0	180	n, p, d	13, 14, 21, 27, 28, 57
11.357 \pm 15	1 ⁺ ; 0	30	γ , n, p, d	13, 14, 19, 20, 21, 27, 28, 57, 65
11.5136 \pm 1.5	2 ⁺ , 3 ⁺	7.0 ± 0.5	p, d	13, 14, 19, 21, 57, 65
11.68 \pm 20	1 ⁻ , 2 ⁻	150 ± 20	n, p, d	13, 19, 57
11.741 \pm 6	1 ⁻ , 2 ⁻	40 ± 9	$(\gamma, n), p, d$	13, 19, 26, 27, 28, 57
11.761 \pm 6	3 ⁻ , 4 ⁻	78 ± 6	$(\gamma, n), p, d$	13, 19, 26, 27, 28, 57
11.807 \pm 7	(2 ⁻)	119 ± 9	n, p, d	13, 14, 19
11.874 \pm 6	(2 ⁻)	101 ± 9	p, d	13, 19, 28
12.20 \pm 20	1 ⁻ , 2 ⁻	300 ± 30	n, p, d	13, 21, 28, 57
(12.29 \pm 15)				19
12.408 \pm 3	(3 ⁺)	37 ± 4	n, p, d	13, 14, 19, 21
12.418 \pm 3	(4 ⁻)	41 ± 4	p, d	3, 4, 13, 19
(12.47 \pm 5)		≈ 20	p, α	3
12.497 \pm 4		35 ± 10	γ, n, p, d, α	3, 13, 19, 26, 39, 57, 64
12.594 \pm 3	3 ⁺	48 ± 2	n, p, d, α	3, 13, 14, 19, 21, 28, 57
12.688 \pm 4	3 ⁻	22 ± 4	n, p, d, α	2, 3, 4, 5, 13, 14, 19, 20, 21, 28
(12.708 \pm 9)		43 ± 15	p, d	13
12.792 \pm 4	4 ⁺	18 ± 3	p, d, α	3, 4, 5, 11, 13, 14, 19, 23, 44, 45, 47, 57
(12.813 \pm 5)		37 ± 6	p, d	13
12.819 \pm 4	4 ⁻	8 ± 3	n, p, d, α	3, 4, 13, 14, 57
12.857 \pm 6		78 ± 10	n, p, d	13, 28
12.923 \pm 5	(4 ⁺)	25 ± 3	p, d, α	3, 4, 13, 14, 19
13.03 \pm 15		≈ 100	γ, p	20, 26
13.166 \pm 5	(1 ⁺)	15 ± 5	n, p, d, α	2, 3, 4, 5, 13, 14, 19, 57
13.192 \pm 10	3 ⁺	65 ± 10	α	5, 9, 57
13.243 \pm 10	2 ⁻	92 ± 5	n, p, α	2, 3, 28, 57
13.30 \pm 40	(2 ⁻); 1	1000 ± 150	γ, p	26
13.656 \pm 5	(2 ⁺ , 3 ⁺)	≈ 90	n, p, d, α	2, 3, 5, 13
13.714 \pm 5	2, 3 ⁺	105 ± 25	n, p, d, α	2, 3, 4, 28, 57

Table 14.11: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
13.71 \pm 30	1 $^+$; 1	180 \pm 20	γ, n, p, α	2, 3, 5, 13, 14, 26, 28, 39, 57, 59
13.72 \pm 20		110	p, α	3
14.04 \pm 30		100	n, p, d, α	2, 3, 13, 14, 28
14.16 \pm 30		230	n, p, d, α	2, 3, 13, 14
14.25 \pm 50	3 $^+$	420 \pm 100	p, α	3, 5
14.30 \pm 20		150	p, d, α	3, 13, 14
14.56 \pm 20		100	n, p, α	2, 3
14.59 \pm 30		50	n, p, α	2, 3
14.66 \pm 10	2 $^-$	100 \pm 20	α	5
14.73 \pm 30		125	n, p, α	2, 3
14.86 \pm 30		140	n, p, d, α	2, 3, 13, 14, 16, 19, 28
14.92 \pm 25		43 \pm 8	n, p, α	2, 3, 19, 28
15.015 \pm 15		\approx 60	n, d, α	2, 14, 20, 28
15.24 \pm 20		100	p, d, α	3, 13, 14, 23
15.43 \pm 20		100	n, p, d, α	2, 3, 13, 16, 20
15.7 \pm 150		350	n, p, d, α	13, 14, 16, 19, 20, 28
16.21 \pm 20		125	n, p, α	2, 3, 20, 23, 28
16.40 \pm 20		150	p, α	3, 16, 20, 23
16.8	4 $^+$; 0 + 1	\approx 300	d, α	16
16.91 \pm 20	(5 $^-$); 0 + 1	\approx 100	p, d, α	3, 16
17.17 \pm 20	4 $^+$; 0 + 1	\approx 300	n, p, d, α	3, 13, 14, 16, 19, 20, 23, 28
17.7				20, 23
18.1	(1 $^-, 2^+$); 0 + 1	(\approx 300)	d, α	16
18.1	4 $^+$; 0 + 1	\approx 600	d, α	16, 28
18.2	3 $^-$; 0 + 1	(\approx 400)	d, α	16
18.4	3 $^-$; 0 + 1	(\approx 300)	d, α	16
18.50	5 $^-$; 0 + 1	\approx 60	d, α	16
18.8	4 $^+$; 0 + 1	(\approx 400)	d, α	16
20.1	1 $^-$; 0 + 1	(\approx 500)	d, α	16
20.6		500 \pm 100	γ, d	12
20.8	5 $^-$; 0 + 1	\approx 600	d, α	16

Table 14.11: Energy levels of ^{14}N ^a (continued)

E_x in ^{14}N (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
20.8	$(3^-, 4^+); 0 + 1$	(≈ 500)	d, α	16
21.3	$4^+; 0 + 1$	$(\approx 1 \text{ MeV})$	d, α	16
21.5	$3^-; 0 + 1$	(≈ 500)	d, α	16, 44
21.7	$5^-; 0 + 1$	≈ 200	d, α	16
21.8		650	$\gamma, {}^3\text{He}$	8
22.5 ± 100	$5^-; 0 + 1$	610 ± 100	d, α	16
22.5	$(2^-); 1$		γ, p	26
23.0	$(0, 1, 2)^-; 1$	$\approx 3 \text{ MeV}$	γ, n	40
23.0	$2^-; 1$		γ, p	26
23.3 ± 100	$5^-; 0 + 1$	500 ± 100	d, α	16
24.0		≈ 1000	$n, {}^3\text{He}, \alpha$	8

^a See also Table 14.7 in (1970AJ04) and Tables 14.12 and 14.13 here.

Other topics: (1968JO1C, 1970BO1M, 1970CO1H, 1970FO1B, 1970FR13, 1971BO1F, 1971HS05, 1971JA13, 1971MC15, 1972CA37, 1972LA12, 1972LE1L, 1972WA1F, 1973CL09, 1973CO16, 1973GO1H, 1973IG02, 1973KE1C, 1973KU03, 1973MA48, 1973PA1F, 1973PO1D, 1973RO1R, 1973RO1Q, 1973SH14, 1973SI21, 1974BO22, 1974CA1H, 1974DU11, 1974FA1A, 1974MU13, 1974SI04, 1974VA24, 1974ZU1A, 1975CA1N, 1975KU01, 1976MA04, 1976PA03).

Ground state: (1969LE1B, 1971SC29, 1971SH26, 1971TA1A, 1972LE1L, 1972VA36, 1973CH38, 1973HO32, 1973SU1B, 1974AD1C, 1974DE1E, 1974DU02, 1974EN10, 1974MU13, 1975AL19, 1975BE31, 1975CA1N, 1976PA03).

$$\mu = +0.403562 \pm 0.000010 \text{ nm} \quad (1964BA11); \text{ see also } (1969FU11);$$

$$= +0.40376077 \pm 0.00000006 \quad (1974SHYR);$$

$$Q = +10 \text{ mb} \quad (1969FU11); \text{ see also } (1968OK1C, 1969BO1D, 1974SHYR);$$

$$r_{\text{rms}} = 2.55 \pm 0.03 \text{ fm} \quad (1974DU02).$$

Mass of ^{14}N : 14.003 074 001 (11) amu (1975SM02).

See also (1971KE01, 1971SM01, 1972WA1G).

1. (a) ${}^9\text{Be}({}^6\text{Li}, n){}^{14}\text{N}$ $Q_m = 14.500$
- (b) ${}^9\text{Be}({}^7\text{Li}, 2n){}^{14}\text{N}$ $Q_m = 7.250$

Table 14.12: Radiative decays in ^{14}N ^a

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
2.31	$0^+; 1$	0	$1^+; 0$	100	$(7.6 \pm 0.5) \times 10^{-3}$	Table 14.13
	$1^+; 0$	0	$1^+; 0$	^A 3.9 ± 0.2 ^b	$(\text{M1})(5.8 \pm 1.2) \times 10^{-4}$	(1967OL02, 1968RO1C)
					$(\text{E2})(4.81 \pm 0.33) \times 10^{-3}$	(1972RE10)
		2.31	$0^+; 1$	^A 96.1 ± 0.3 ^b	0.140 ± 0.013	(1967OL02, 1968RO1C)
				4.2 ± 1.4		(1972RE10)
				95.8 ± 1.4		(1972RE10)
	$0^-; 0$	0	$1^+; 0$	97 ± 3	$(8.4 \pm 1.6) \times 10^{-2}$	(1972RE10), Table 14.13)
		2.31	$0^+; 1$	0.4 ± 0.7		(1965NE06; see also 1966GO15)
				≤ 2		(1972RE10)
		3.95	$1^+; 0$	1.3 ± 1.0		(1965NE06)
4.92				≤ 2		(1972RE10)
				≤ 0.5		(1966GO15)
	$2^-; 0$	0	$1^+; 0$	^A 79.9 ± 1.0 ^b	$(4.0 \pm 0.5) \times 10^{-5}$ ^d	(1966GO15, 1968AL12)
		2.31	$0^+; 1$	^A 19.7 ± 1.2 ^b		(1966GO15)
		3.95	$1^+; 0$	(0.7 ± 0.4)		(1963AL21)
	$1^-; 0$	0	$1^+; 0$	37 ± 2		(1964WA09)
				40 ± 3		(1966CA07)
				40 ± 4		(1966GO15)
				36 ± 4		(1972RE10)
				31.6 ± 1.9		mean
5.69		2.31	$0^+; 1$	35.6 ± 1.2	$(4.2 \pm 0.8) \times 10^{-2}$ ^d	(1963AL21); Table 14.13
				63 ± 2		

Table 14.12: Radiative decays in ^{14}N ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
5.83	3 ⁻ ; 0	0	1 ⁺ ; 0	60 ± 3		(1964WA09)
				60 ± 5		(1966CA07)
				64 ± 4		(1966GO15)
				64.7 ± 1.9		(1972RE10)
				63.1 ± 1.2		mean
				≤ 0.4 ^e		(1972RE10)
				≤ 0.3 ^e		(1972RE10)
				25 ± 5		(1965WA06)
				25 ± 4		(1966CA07)
				29 ± 4		(1966GO15)
6.20	1 ⁺ ; 0	0	1 ⁺ ; 0	26.5 ± 2.5		mean
				< 3		(1966GO15)
				< 1		(1966GO15)
				< 1		(1966GO15)
				75 ± 5	$(2.7 \pm 0.3) \times 10^{-5}$ ^d	(1965WA06, 1966CA07)
				71 ± 4		(1966GO15, 1968AL12)
				73 ± 3		mean
				24 ± 3		(1964WA09)
				25 ± 4		(1966CA07)
				21 ± 3		(1966GO15)
29		2.31	0 ⁺ ; 1	23.0 ± 1.9	$(4.1 \pm 0.5) \times 10^{-3}$ ^d	mean
				76 ± 3		(1964WA09); Table 14.13

Table 14.12: Radiative decays in ^{14}N ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
30	6.44	3 ⁺ ; 0	3.95	72 ± 5		(1966CA07)
				79 ± 3		(1966GO15)
				76.7 ± 2.0		mean (1966GO15)
			5.11	< 1		(1966GO15)
			0	2 ⁻ ; 0		(1966GO15)
	7.03	2 ⁺ ; 0	3.95	65 ± 3	$(7.3 \pm 0.4) \times 10^{-4}$ ^d	(1964WA09)
				73 ± 5		(1959RO54, 1966CA07)
				74 ± 4		(1966GO15)
				69.4 ± 2.2		mean (1964WA09)
				21 ± 2		(1966CA07)
30	7.03	2 ⁺ ; 0	5.11 ^g	20.5 ± 1.6		(1966GO15)
				7 ± 3		(1966CA07)
				^A 7 ± 2		(1966GO15)
			5.83	3 ⁻ ; 0		(1966GO15)
			0	1 ⁺ ; 0		(1965WA06)
	7.03	2 ⁺ ; 0	3.95	91 ± 4	$(\text{M1})(9.1 \pm 1.3) \times 10^{-2}$	(1967OL02, 1968RO1C)
				^A 98.6 ± 0.3	$(\text{E2})(3.3 \pm 0.9) \times 10^{-2}$	(1967OL02, 1968RO1C)
			2.31	0 ⁺ ; 1	$(\text{E2})(6.2 \pm 1.4) \times 10^{-4}$	(1967OL02, 1968RO1C)
			3.95	1 ⁺ ; 0	$< (11 \pm 3) \times 10^{-4}$	(1967OL02, 1968RO1C)
			other states	≤ 0.4		(1967OL02)

Table 14.12: Radiative decays in ^{14}N ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
31	7.97 ^h	2 ⁻ ; 0	0	1 ⁺ ; 0	55 ± 3	$\omega\Gamma_p\Gamma_\gamma/\Gamma = 0.012$
			3.95	1 ⁺ ; 0	45 ± 3	= 0.010
			other states	≤ 3		(1960HE14)
	8.06	1 ⁻ ; 1	0	1 ⁺ ; 0	80.3 ± 0.6	10.3
			2.31	0 ⁺ ; 1	1.40 ± 0.14	0.2
			3.95	1 ⁺ ; 0	12.7 ± 0.4	1.6
			4.92	0 ⁻ ; 0	1.86 ± 0.14	0.3
			5.11	2 ⁻ ; 0	0.25 ± 0.14	0.03
			5.69	1 ⁻ ; 0	3.5 ± 0.4	0.5
	8.49 ^j	4 ⁻ ; 0	5.11	2 ⁻ ; 0	100	$(5.61 \pm 2.0) \times 10^{-3}$
	8.62	0 ⁺ ; 1	0	1 ⁺ ; 0	23	1.20
			3.95	1 ⁺ ; 0	24	1.26
			5.69	1 ⁻ ; 0	13	0.69
			6.20	1 ⁺ ; 0	40	(1957WI27)
	8.79	0 ⁻ ; 1	0	1 ⁺ ; 0		43 ± 9
			3.95	1 ⁺ ; 0		0.9 ± 0.3
	8.91	3 ⁻ ; 1	0	1 ⁺ ; 0		$(6.6 \pm 2.2) \times 10^{-3}$
			5.11	2 ⁻ ; 0	5.5 ± 2.5	$\omega\Gamma_\gamma = 0.040$
			5.83	3 ⁻ ; 0	89 ± 3	= 0.65
			6.44	3 ⁺ ; 0	3 ± 1	= 0.022
			7.03	2 ⁺ ; 0	1.4 ± 0.8	= 0.010
	8.96 ^j	5 ⁺ ; 0	0	1 ⁺ ; 0	< 1	(1967BL22, 1967GA12)

Table 14.12: Radiative decays in ^{14}N ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
		6.44	$3^+; 0$	100	$\left\{ \begin{array}{l} \Gamma_p/\Gamma_\gamma = 4.1 \pm 0.5 \\ \Gamma_\gamma = (1.36 \pm 0.21) \times 10^{-3} \end{array} \right.$	(1965DE19, 1966CA07, 1967GA12)
9.13	$2^-; 0$	0	$1^+; 0$	> 80	$\omega\Gamma_\gamma \approx 0.03$	(1965DE19)
9.17	$2^+; 1$	0	$1^+; 0$	79 ± 4	7.7 ± 0.9	(1959HA11, 1960RO13, 1963PR03, 1967GA12, 1968CL05)
		2.31	$0^+; 1$	1.1 ± 0.4	0.11 ± 0.4	(1963PR03, 1968CL05)
		5.11	$2^-; 0$	< 1		(1960RO13)
		5.69	$1^-; 0$	< 6	0.3 ± 0.25	(1960RO13, 1968CL05)
		5.83	$3^-; 0$	3 ± 2	0.33	(1960RO13)
		6.44	$3^+; 0$	8 ± 2	0.78 ± 0.35	(1960RO13, 1968CL05)
				6.3 ± 0.5	0.85 ± 0.15	(1963PR03)
		7.03	$2^+; 0$	3 ± 1	0.3 ± 0.15	(1960RO13, 1968CL05)
				3.5 ± 0.5	0.34 ± 0.05	(1963PR03)
9.39	$2^-; 3^-; 0$	ⁱ				(1975NO1F)
9.51	$2^-; 1$	3.95	$1^+; 0$	6 ± 1	0.37	(1959WA04)
		5.11	$2^-; 0$	78 ± 3	4.8	(1959WA04)
		5.83	$3^-; 0$	16 ± 2	1.0	(1959WA04)
10.23	$1^{(-)}; 0$	2.31	$0^+; 1$		4 ± 1.3	(1963RO17)
10.43	$2^+; 1$	0	$1^+; 0$	82 ± 6	12.1 ± 1.5	(1960RO13, 1968CL05)
		2.31	$0^+; 1$	< 1		(1964RO03)
		3.95	$1^+; 0$	< 2		(1964RO03)
		5.11	$2^-; 0$	2 ± 1	0.3 ± 0.2	(1960RO13, 1968CL05)

Table 14.12: Radiative decays in ^{14}N ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	$J_f^\pi; T$	Branch (%)	Γ_γ (eV)	Refs.
10.81 ^k	5 ⁺ ; 0	5.69	1 ⁻ ; 0	< 3		(1964RO03)
		5.83	3 ⁻ ; 0	< 1		(1964RO03)
		6.44	3 ⁺ ; 0	8 ± 1	1.2 ± 0.4	(1960RO13, 1968CL05)
		7.03	2 ⁺ ; 0	6 ± 1	0.88 ± 0.31	(1960RO13, 1968CL05)
		6.44	3 ⁺ ; 0	100	$\Gamma_\gamma/\Gamma = 4.1 \pm 0.8\%$	(1972NO08)

A = Adopted.

^a See also Table 14.9 in (1970AJ04) and Tables 14.13 and 14.20 here.

^b Means of branching ratio values quoted in Table 14.9 (1970AJ04).

^c See also (1972NY02, 1974RAZD).

33

^d Corrected for branching: see Table 14.13.

^e See also (1966GO15).

^f 5.83 → 5.11: $E_\gamma = 728.3 \pm 1.0$ keV (1966AL10): the plane polarization of the γ -rays leads to odd parity for $^{14}\text{N}^*(5.83)$ (1962RO21).

^g See also (1964WA09).

^h $\Gamma_\gamma/\Gamma = 0.7 \pm 0.2\%$; $(2J+1)\Gamma_p = 12.6 \pm 3.6$ eV (1972BA56); $\Gamma = 2.5 \pm 0.7$ eV (J.W. Noe, private communication).

ⁱ Decays predominantly to $^{14}\text{N}^*(5.11, 5.83, 8.9)$: ≈ 5 meV each (1975NO1F).

^j $\Gamma = 27 \times 10^{-3}$ and 7×10^{-3} eV, respectively for $^{14}\text{N}^*(8.49, 8.96)$ (J.W. Noe, private communication).

^k $\Gamma = 0.39 \pm 0.16$ eV (J.W. Noe, private communication).

Table 14.13: Lifetimes of some ^{14}N states ^a

E_x (MeV)	τ_m (fsec)	Reaction	Refs.
2.31 ^b	83 ± 19	$^{12}\text{C}(^3\text{He}, \text{p})$	(1966LI07)
	83 ± 30	$^{12}\text{C}(^3\text{He}, \text{p})$	(1965LO07)
	114 ± 30	$^{12}\text{C}(^3\text{He}, \text{p})$	(1973HA40)
	70 ± 10	$^{13}\text{C}(\text{p}, \gamma)$	(1971BI03)
	75 ± 19	$^{13}\text{C}(\text{p}, \gamma)$	(1972RE10)
	106 ± 10	$^{14}\text{N}(\gamma, \gamma)$	(1975RA22)
3.95 ^c	87 ± 6		mean
	10.5 ± 3.5	$^{13}\text{C}(\text{p}, \gamma)$	(1971BI03)
	4.5 ± 0.4 ^A	$^{14}\text{N}(\text{e}, \text{e})$	(1964BI09, 1967OL02)
4.92 ^d	7.05 ± 1.5	$^{13}\text{C}(\text{d}, \text{n})$	(1974RU1B)
	10 ± 3	$^{14}\text{N}(\gamma, \gamma)$	(1974RAZD)
	7.6 ± 1.4		mean
5.11 ^e	(12.4 ± 1.4) psec ^A	$^{12}\text{C}(\text{t}, \text{n})$	(1968AL12)
	< 21 fsec	$^{12}\text{C}(^3\text{He}, \text{p})$	(1973HA40)
	< 15 fsec	$^{13}\text{C}(\text{p}, \gamma)$	(1971BI03, 1972RE10)
5.83	10.0 ± 2.0 ^A	$^{13}\text{C}(\text{d}, \text{n})$	(1974RU1B)
	18 ± 2 psec ^A	$^{12}\text{C}(\text{t}, \text{n})$	(1968AL12)
	200 ± 45	$^{12}\text{C}(^3\text{He}, \text{p})$	(1973HA40)
6.20 ^f	118 ± 13	$^{13}\text{C}(\text{p}, \gamma)$	(1971BI03)
	124 ± 15		mean
	630 ± 80	reactions 1, 6, 21	(1969TH01)
6.44	590 ± 120		(1964BE12)
	630 ± 100		(1973HA40)
	630 ± 40		(1970GA09)
7.03	627 ± 33		mean
	5.4 ± 0.5	$^{14}\text{N}(\gamma, \gamma)$	(1966SW01)

A = adopted.

^a See also Table 14.12, and also Table 14.13 in (1970AJ04).

^b See also (1974RU1B).

^c See also (1974RU1B, 1973HA40).

^d See also (1973HA40).

^e See also Table 14.13 in (1970AJ04) and (1974RU1B).

^f See also (1970GA09).

Table 14.14: Resonances in $^{10}\text{B} + \alpha$ ^a

E_α (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particle ^b (x)	Γ_x ^c (keV)	$^{14}\text{N}^*$ (MeV)	J^π	Refs.
0.95		p_0		12.29		(1953MA42)
1.13 ± 5	30 ± 5	$p_0 \rightarrow p_3, d$		12.42	4 ⁻	A
1.20 ± 5	≈ 20	$p_0, (p_2), p_3$		12.47		(1953MA42, 1969GA01)
1.23 ± 5	35 ± 5	p_0, p_3		12.49		(1953SH64, 1969GA01)
1.40 ± 5	46 ± 4	$p_1, p_2, (p_3)$		12.61	3 ⁺	(1953SH64, 1969GA01)
1.507 ± 5	18 ± 5	n_0	4.3	12.690	3 ⁻	A, (1973VA25)
		p_0	0.62			
		p_1	0.17			
		p_2	0.70			
		p_3	5.6			
		d	0.93			
		α	1.7			
		n_0	≤ 0.6	12.789	4 ⁺	A, (1973VA25)
		p_0	0.18			
		p_1	0.085			
1.645 ± 5	16 ± 3	p_2	0.44			
		p_3	9.6			
		d	2.0			
		α	1.0			
		p_1, p_2, p_3, d		12.814	4 ⁻	A
1.68 ± 5	5 ± 2	$p_0 \rightarrow p_3, d$		12.921	4 ⁺	A
1.83 ± 5	22 ± 4					
2.174 ± 5	15 ± 5	$n_0, p_0 \rightarrow p_3, d, \alpha_1$		13.166	1 ⁺	A, (1973VA25, 1975WI04)

Table 14.14: Resonances in $^{10}\text{B} + \alpha$ ^a (continued)

E_α (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particle ^b (x)	Γ_x ^c (keV)	$^{14}\text{N}^*$ (MeV)	J^π	Refs.
2.21 ± 10	65 ± 10	α_0		13.192	3 ⁺	(1973MO15)
2.281 ± 10	92 ± 5	$n_0, p_0 \rightarrow p_3$		13.243	2 ⁻	A, (1973VA25, 1975WI04)
2.86 ± 5	≈ 90	n_0, p_1, p_2, α_1		13.656		(1969GA01, 1973VA25)
2.94 ± 5	105 ± 25	$n_0, p_0 \rightarrow p_3, d$		13.714	2, 3 ⁺	A, (1973VA25, 1975WI04)
2.95 ± 50	180 ± 20	$n_0, p_0, (p_2), \alpha_0$		13.72	1 ⁽⁺⁾	(1973MO15, 1973VA25, 1975WI04) ^d
2.95 ± 20	110	p_1, p_3		13.72		(1969GA01, 1975WI04) ^d
3.40 ± 30	100	n_0, p_1		14.04		(1973VA25, 1975WI04)
3.56 ± 30	230	$n_0, (p_0), p_3$		14.16		(1973VA25, 1975WI04)
3.69 ± 50	420 ± 100	$p\gamma, \alpha_0$		14.25	3 ⁺	(1956BO61, 1959GI47, 1973MO15)
3.76 ± 20	150	p_1		14.30		(1975WI04)
3.98 ± 20	100	n_0, p_0, p_2		14.56		(1956BO61, 1973VA25, 1975WI04)
4.16 ± 30	50	n_0, p_0, p_3		14.59		(1973VA25, 1975WI04)
4.26 ± 10	100 ± 20	α_0		14.66	2 ⁻	(1973MO15)
4.36 ± 30	125	$n_0, p_0, p_1, (p_2)$		14.73		(1973VA25, 1975WI04)
4.54 ± 30	140	n_0, p_2, p_3		14.86		(1956BO61, 1959GI47, 1973VA25, 1975WI04)
4.633 ± 30	43 ± 8	n_0, n_{2+3}, p_0		14.923		(1973VA25, 1975WI04)
4.77 ± 20	≈ 60	n_0, n_1		15.02		(1973VA25)
5.08 ± 20	100	p_3		15.24		(1975WI04)
5.35 ± 20	100	n_1, p_2, p_3		15.43		(1956BO61, 1975WI04)
6.44 ± 20	125	n_0, p_0, p_2		16.21		(1975WI04)
6.70 ± 20	150	p_2		16.40		(1975WI04)
7.42 ± 20		p_0		16.91		(1975WI04)

Table 14.14: Resonances in $^{10}\text{B} + \alpha$ ^a (continued)

E_α (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particle ^b (x)	Γ_x ^c (keV)	$^{14}\text{N}^*$ (MeV)	J^π	Refs.
7.78 ± 20	50	p ₃		17.17		(1975WI04)

A: See references quoted for this state in (1970AJ04).


^a See Table 1 in (1975WI04) for a display of the resonance data obtained both in $^{12}\text{C} + \text{d}$, $^{13}\text{C} + \text{p}$ and $^{10}\text{B} + \alpha$.

^b n₀, n₁, n₂₊₃ correspond to the g.s. and $^{13}\text{N}^*(2.37, 3.51 + 3.55)$; p₀, p₁, p₂, p₃ correspond to the g.s. and $^{13}\text{C}^*(3.09, 3.68, 3.85)$ and the corresponding γ -rays; α_1 corresponds to the transition to $^{10}\text{B}^*(0.7)$.


^c For θ_x^2 see Table 14.8 in (1970AJ04).

^d See reference^f to Table 1 of (1975WI04).


In reaction (a) $E_\gamma = 1631.3 \pm 1.3$ [3.95 → 2.31], 4913.8 ± 3.0 [4.92 → g.s.], 3883.0 ± 1.9 [6.20 → 2.31] and 6443.7 ± 1.8 keV [6.44 → g.s.] have been observed [transitions shown in brackets]. The τ_m of $^{14}\text{N}^*(6.44)$ is 620 ± 80 fsec [see also Table 14.13] (1969TH01). For reaction (b) see (1957NO17).




Recent measurements have been carried out at $E_\alpha = 1.0$ to 5.0 MeV (1973VA25: n_0, n_1, n_{2+3}) and at 2.0 to 7.0 MeV (1975WI04: n_0). Observed resonances in these and in previous experiments are displayed in Table 14.14. See also reaction 3.



Excitation functions have been measured to $E_\alpha = 26$ MeV: see (1970AJ04) for a listing of the earlier references and (1975WI04: $E_\alpha = 2.0$ to 10.7 MeV (p_0), to 7.5 MeV (p_1) and to 8 MeV (p_2, p_3); see also reaction 2). Observed resonances are displayed in Table 14.14. (1975WI04) has expanded the angular distributions of the $p_0 \rightarrow p_3$ groups into Legendre polynomials and fitted the coefficients at the resonances corresponding to $^{14}\text{N}^*(13.16, 13.24, 13.67, 13.76)$ obtaining $J^\pi = 1^+, 2^-, 2$ or 3^+ , and 1 , respectively, for these states. (1975WI04) also finds that a surprising proportion of states have a higher cross section for neutron than for proton emission: the fluctuations of σ_n/σ_p at low E_α suggest sizable isospin impurities in the ^{14}N states (1975WI04).



Excitation curves have been measured for E_α up to 27 MeV: see (1970AJ04) for the earlier references and (1975SP04: $E_\alpha = 18$ to 27 MeV; yields of 12.7 and 15.1 MeV γ). The low-energy resonances ($E_d < 2.2$ MeV) are exhibited in Table 14.14. At the higher energies the yield curves are fairly smooth (see (1967AL16, 1975SP04)), although (1975SP04) report a sharp rise in the 15.1 MeV γ -yield ≈ 1 MeV above the $^{12}\text{C}^*(15.1) + p + n$ threshold, a reaction which is not isospin forbidden.



The yield of 0.72 MeV γ -rays has been measured for $E_\alpha = 2.1$ to 3.5 MeV by (1969GA01). Excitation functions for elastically scattered α -particles have been measured for $E_\alpha = 2.0$ to 4.3

MeV ([1973MO15](#)) and for 5 to 30 MeV ([1972DA04](#)). In addition to two strong resonances in the α_0 yields at $E_\alpha = 2.21$ and 4.26 MeV ($^{14}\text{N}^*(13.19, 14.66)$), two other states ($^{14}\text{N}^*(13.72, 14.25)$) are required to fit the data: an R -matrix calculation leads to $J^\pi = 3^+, 1^+$ [see, however, ([1975WI04](#))], 3^+ and 2^- , respectively for $^{14}\text{N}^*(13.19, 13.72, 14.25, 14.66)$ ([1973MO15](#)). For reaction (b) see ([1974JE1A](#)).



At $E(^6\text{Li}) = 5$ MeV $^{14}\text{N}^*(0, 3.95, 4.92, 5.11, 5.69, 5.83, 6.20, 6.44, 7.03, 7.97, 8.49, 8.98, 9.12, 9.39, 9.70, 10.10, 10.43 (T = 1, weakly populated), 11.06)$ are populated (([1966MC05](#)), and private communication). ([1975FO01](#)) have examined $^{14}\text{N}^*(9.13)$ in detail and conclude that it is a closely spaced doublet of which one member is populated in $^{13}\text{C}(\text{p}, \gamma)$ and has been assigned $J^\pi = 2^-$, and the other member, populated here, has $J^\pi = 3^+$. This assignment is based on a DWBA analysis of the angular distributions at $E(^6\text{Li}) = 16.5$ and 21.0 MeV which shows the contributions of several L values and, in particular, $L = 0$. See also reaction 19. Branching ratios for the γ -decay of ^{14}N states are displayed in Table 14.12 ([1966CA07](#)). See also reaction 1 ([1969TH01](#)).



Triton groups corresponding to a number of ^{14}N states have been observed at $E(^7\text{Li}) = 5$ MeV: see ([1966MC05](#)) and reaction 6. See also ([1974KO1G](#)).

8. (a) $^{11}\text{B}(^3\text{He}, \gamma)^{14}\text{N}$	$Q_m = 20.7359$	$E_b = 20.7359$
(b) $^{11}\text{B}(^3\text{He}, \text{n})^{13}\text{N}$	$Q_m = 10.182$	
(c) $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$	$Q_m = 13.1852$	
(d) $^{11}\text{B}(^3\text{He}, \text{d})^{12}\text{C}$	$Q_m = 10.4634$	
(e) $^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$	$Q_m = -2.001$	
(f) $^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}$		
(g) $^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B}$	$Q_m = 9.1225$	
(h) $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$	$Q_m = 4.570$	

The capture γ -rays (reaction (a)) have been studied at $E(^3\text{He}) = 0.9$ to 2.6 MeV ([1970BL10](#): $\gamma_0; \theta = 0^\circ$ and 90°) and 1.5 to 5.8 MeV (γ_0) and 2.0 to 5.8 MeV (γ_1) ([1973MA1W](#): 90°). When the barrier penetration factor has been removed the low-energy work shows a single resonance at $E(^3\text{He}) \approx 1.4$ MeV ($^{14}\text{N}^*(21.8)$), $\Gamma_{\text{c.m.}} = 0.65$ MeV ([1970BL10](#)). ([1973MA1W](#): prelim. results)

report two broad resonances at $E_x = 23.4$ and 25.0 MeV in the γ_0 yield and at 24.0 MeV in the γ_1 yield.

The excitation function at 0° in the range $E(^3\text{He}) = 1.5$ to 5.6 MeV for n_0 (reaction (b)) shows a broad peak at $E(^3\text{He}) = 4.15$ MeV which may indicate the existence of a ^{14}N state at $E_x \approx 24.0$ MeV, $\Gamma \approx 1$ MeV ([1966DI04](#)). The excitation function for reaction (b) has also been measured for $E(^3\text{He}) = 6$ to 18 MeV ([1967HA20](#)).

Yield curves for protons (reaction (c)) have been measured for $E(^3\text{He}) = 3.0$ to 5.5 MeV ($p_0, p_1, p_1 + p_2 + p_3$): they are rather featureless ([1959HO01](#)). This is also true for the ground state deuterons of reaction (d) in the same energy interval ([1959HO01](#)). Yield curves for reaction (e) have been measured for $E(^3\text{He}) = 6$ to 18 MeV ([1967HA20](#)) and 10 to 30 MeV ([1965BR42](#)). See also ^{13}C and ^{13}N . For reactions (d, e, f) see ^{12}C , ^{11}C and ^{11}B in ([1975AJ02](#)).

The excitation functions for α -particle groups (reaction (g)) have been measured for $E(^3\text{He}) = 0.9$ to 2.15 MeV ([1966LO15](#): $\alpha_0 \rightarrow \alpha_4$) and 2.2 to 5.5 MeV ([1965FO06](#): $\alpha_0 \rightarrow \alpha_3$). No significant resonance behavior is seen except for the α_2 group which, in the 15° excitation function, exhibits a resonance at $E(^3\text{He}) = 4$ MeV, $\Gamma \approx 1$ MeV ([1965FO06](#)). See also ^{10}B in ([1974AJ01](#)).

The excitation function for reaction (h) to $^6\text{Li}_{\text{g.s.}} + ^8\text{Be}_{\text{g.s.}}$ has been measured for $E(^3\text{He}) = 1.4$ to 5.8 MeV: no pronounced structure is observed ([1967YO02](#)). At $E(^3\text{He}) = 25.20$ to 26.25 MeV the excitation functions for the transitions to $^8\text{Be}^*(0, 16.63, 16.91, 17.64)$ are smooth, indicating a predominantly direct reaction mechanism ([1974DE25](#)).



Angular distributions have been measured for E_α to 13.9 MeV [see ([1970AJ04](#)) for the earlier references], $E_\alpha = 3.7$ to 7.9 MeV ([1975VA06](#): n_0, n_1, n_2) and at $E_\alpha = 12.6$ to 14.8 MeV ([1972CI01](#), [1973CI02](#): n_0).



See ([1970AJ04](#)).



At $E(^{11}\text{B}) = 114$ MeV the relatively strongly populated states are $^{14}\text{N}^*(5.83, 8.96, 12.8)$ [$J^\pi = 3^-, 5^+$ and 4^+ , respectively] ([1974AN36](#)).




Table 14.15: Resonances in $^{12}\text{C} + \text{d}$

E_{d} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$^{14}\text{N}^*$ ^a (MeV)	$J^\pi; T$	Refs. ^b
0.92	95	n, p ₀ , p ₁	11.06	1 ⁺ ; 0	A, (1972SE09, 1975SE07)
1.13		p ₀ , p ₁	11.24	; 1	A
1.19	190	n, p ₀ , p ₁ , d	11.29	2 ⁻ ; 0	A, (1972SE09, 1975SE07)
1.23		p ₀	11.33	(3 ⁺)	(1969BO32)
1.30	30	n, p ₀ , p ₁ , d	11.39	1 ⁺ ; 0	A
1.39		p ₀	11.46	(2 ⁻)	(1969BO32)
1.4495 \pm 1.5	7.0 \pm 0.5	p ₀ , p ₁ , d	11.5136	2 ⁺ , 3 ⁺	A, (1970AL26, 1973TR02, 1975TR07)
1.55		p ₀	11.60	(2 ⁻)	(1969BO32)
1.640 \pm 20	150 \pm 20	n, p ₁	11.68	1 ⁻ , 2 ⁻	(1969BO32, 1969JA06, 1975TR07)
1.715 \pm 6	40 \pm 9	p ₂	11.741	1 ⁻ , 2 ⁻	(1975TR07)
1.738 \pm 6	78 \pm 6	p ₁	11.761	3 ⁻ , 4 ⁻ , (2 ⁻)	(1975TR07)
1.792 \pm 7	119 \pm 9	n, p ₀ , p ₁ , p ₂ , d ₀	11.807	2 ⁻ , (1 ⁺)	A, (1970AL26, 1971ME18, 1975TR07)
1.870 \pm 6	101 \pm 9	p ₀ , p ₁ , p ₂	11.874	2 ⁻ , (1 ⁻)	(1969BO32, 1974GM01, 1975TR07)
		c			
2.250 \pm 19	300 \pm 30	n, p ₀ \rightarrow p ₃	12.20	1 ⁻ , 2 ⁻	A, (1974GM01, 1975TR07)
2.494 \pm 3 ^d	37 \pm 4	n, p ₀ \rightarrow p ₃ , d ₀	12.408	3 ⁺ , (3 ⁻ , 4 ⁻)	A, (1971ME18, 1974DA06, 1974ST05, 1975TR07)
2.506 \pm 3	41 \pm 4	p ₁	12.418	3 ⁻ , 4 ⁻ , (2 ⁺ , 3 ⁺)	(1975TR07)
2.610 \pm 20	30 \pm 20	n, p ₁ , p ₂ , p ₃	12.507		A, (1975TR07)
2.712 \pm 3	48 \pm 2	(n), p ₀ \rightarrow p ₃ , d	12.594	3 ⁺	A, (1975TR07)
(2.817 \pm 7)	27 \pm 6	n, p ₁ , p ₂ , p ₃ , d	(12.684)		A
2.844 \pm 9	43 \pm 15	p ₂ , p ₃	12.708		(1975TR07)
2.940 \pm 10	30 \pm 10	p ₂ , p ₃ , d	12.790		A, (1975TR07)

Table 14.15: Resonances in $^{12}\text{C} + \text{d}$ (continued)

E_{d} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$^{14}\text{N}^*$ ^a (MeV)	$J^\pi; T$	Refs. ^b
2.967 \pm 5	37 \pm 6	p ₁	12.813		(1975TR07)
2.982 \pm 6	11 \pm 3	n, p ₃ , d	12.826		A
3.018 \pm 6	78 \pm 10	n, p ₀ , p ₁	12.857		A, (1975TR07)
3.049 \pm 8	134 \pm 11	p ₁	12.883		(1975TR07)
3.100 \pm 10	20 \pm 14	p ₁ , p ₂ , p ₃ , d	12.927	(3 ⁻ , 4 ⁻)	A, (1975TR07)
3.39 \pm 12	47 \pm 15	n, p ₂ , p ₃ , d	13.17	(0 ⁻ , 1 ⁻)	A
3.97 \pm 30	< 200	p ₀ , p ₂ , p ₃ , (d)	13.67	(2 ⁺ , 3 ⁺)	A
4.02 ⁺²⁰ ₋₁₀	\approx 235	n, (p), d	13.71	(1 ⁺)	A
4.40		p ₀ \rightarrow p ₃ , d	14.04		A
4.55		n, p ₂ , d	14.17		A
4.80		p ₀ , p ₂ , d	14.38		A
5.17		d	14.70		A
5.34	\approx 100	p ₀ \rightarrow p ₃ , d, α	14.84		A
5.65		d	15.11		A
5.83		p ₁ , p ₃ , d	15.26		A
6.07		p ₁ , p ₂ , α	15.47		A
6.3		p ₀ , p ₃ , d, α	15.7		A
7.2		α	16.4		A
7.6 ^e	\approx 300	α_2	16.8	4 ⁺ ^f	(1972SM07) ^e
7.8	\approx 100	α_2	16.9	(5 ⁻)	(1972SM07)
8.1	\approx 300	p ₀ , p ₂ , d, α_2	17.2	4 ⁺	A, (1972SM07)
9.1	(\approx 300)	α_2	18.1	(1 ⁻ , 2 ⁺)	(1972SM07)

Table 14.15: Resonances in $^{12}\text{C} + \text{d}$ (continued)

E_{d} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$^{14}\text{N}^*$ ^a (MeV)	$J^\pi; T$	Refs. ^b
9.2	≈ 600	α_2	18.1	4^+	(1972SM07)
9.3	(≈ 400)	α_2	18.2	3^-	(1972SM07)
9.5	(≈ 300)	α_2	18.4	3^-	(1972SM07)
9.61	≈ 60	α_2	18.50	5^-	(1972SM07)
10.0 ^g	(≈ 400)	α_2	18.8	4^+	(1972SM07)
11.5	(≈ 500)	α_2, α_3	20.1	1^-	(1972SM07)
12.3	≈ 600	α_2	20.8	5^-	(1972SM07)
12.3	(≈ 500)	α_2	20.8	($3^-, 4^+$)	(1972SM07)
12.9	(≈ 1000)	α_2	21.3	4^+	A, (1972SM07)
13.1	(≈ 500)	α_2	21.5	3^-	(1972SM07)
13.4	≈ 200	α_2	21.7	5^-	(1972SM07)
14.29 ± 100	610 ± 100	α_2	22.5	5^- ^h	(1974JO01) ⁱ
15.28 ± 100	500 ± 100	α_2	23.3	5^- ^h	(1974JO01)

A: See references quoted for this state in (1970AJ04).

^a (1963JE02) report 39 excited states of ^{14}N with $11.2 < E_x < 14.2$ MeV.

^b See also (1959AJ76, 1970AJ04).

^c (1974GM01) also report a state at 12.05 MeV with $\Gamma = 190$ keV in the p_0 yield.

^d A study of this resonance shows that either f-shell components are present in the wave function or that the coupling is very strong or that both effects are present (1974DA06).

^e For all states reported by (1972SM07) see their discussion of the S -matrix analysis of their α_2 data: the resonances shown below correspond to one possible (albeit the most reasonable) solution. The α_2 channel is sensitive only to ^{14}N states that have natural parity and are isospin mixed.

^f These, and all the states shown below, are isospin mixed. (1972SM07) notes that the average spacing of these states, as observed via the α_2 channel, $D \approx 400$ keV, as is the average width of the states, so $\Gamma \approx D$.

^g See also Table 14.10 in (1970AJ04) for states reported at $^{14}\text{N}^*(17.2, 19.6, 20.4)$ by (1963OH02, 1965BA06, 1968KL06, 1969CO02).

^h These states appear to be an isospin mixed pair with $\langle H_c \rangle \geq 40$ keV (1974JO01).

ⁱ See also (1968JA09).

Table 14.16: Recent $^{12}\text{C} + \text{d}$ yield curves ^a

E_{d} (MeV)	Yield of	Refs.
3.6 – 12.0	n_0	(1972DA02)
12 – 17	n_0	(1970GA07)
0.4 – 1.35	p_0	(1972HU13)
0.80 – 2.20	$p_1\gamma$	(1973TR02)
1.4 – 2.3	p_0, p_1	(1970AL26)
1.4 – 3.2	$p_1\gamma, p_2\gamma, p_3\gamma$	(1975TR07)
1.82 – 2.50	p_0	(1974GM01)
2.61 – 2.82	$p_0 \rightarrow p_3$	(1974DA06)
3.3 – 3.7	p_0	(1970LE20)
12.3 – 14.7	p_0, p_1	(1968HO23)
1.5 – 2.0	d_0	(1970AL26)
2.61 – 2.82	d_0	(1974DA06)
7.19 – 13.99	$\alpha_0 - \alpha_3$	(1971RI15, 1972SM07)
13.8 – 16	α_2	(1974JO01)
16.0 – 17.0	α_2	(1971VO04)
26.2 – 29.5	α_2	(1971JA04)

^a See Table 14.11 in (1970AJ04) for a listing of the earlier measurements to $E_{\text{d}} = 13.8$ MeV for neutrons, to 14 MeV for protons, to 26.5 MeV for deuterons and to 29 MeV for α -particles. See also (1959AJ76).

At $E_{\text{d}} = 1.5$ MeV the capture cross section is $< 1 \mu\text{b}$ (1955AL16). The yield of γ_0 is nearly flat from $E_{\text{d}} = 9$ to 11.7 MeV and then exhibits broad resonance structures at $E_{\text{d}} = 12.10$ and ≈ 12.8 MeV, with $\Gamma = 0.5 \pm 0.1$ and $1.5 - 2$ MeV, respectively [$^{14}\text{N}^*(20.6, 21.2)$]. The γ_1 and γ_2 yields have also been measured [1972DI1B: prelim. results].

$$\begin{array}{lll}
 13. \text{ (a)} & ^{12}\text{C}(\text{d}, \text{n})^{13}\text{N} & Q_{\text{m}} = -0.281 \\
 & (\text{b}) & E_{\text{b}} = 10.2725 \\
 & ^{12}\text{C}(\text{d}, \text{p})^{13}\text{C} & Q_{\text{m}} = 2.7218
 \end{array}$$

Resonances in the yields of neutrons and protons are displayed in Table 14.15. Recent measurements of the yields of these particles are listed in Table 14.16. For angular distributions of neutrons see ^{13}N . See also (1971MU18, 1973BI1G), (1971DU1B; theor.) and (1972LU1B, 1973WE19, 1975LE1K; applied work). For angular distributions of protons see ^{13}C . See also (1971PU01,

Table 14.17: $^{12}\text{C} + \text{d}$ polarization studies ^a

E_{d} (MeV)	Groups	Refs.
2.71, 2.96	n_0	(1971JA17)
5.7 – 9.8	n_0, n_1	(1975TE1A)
6 – 14	n_0, n_1	(1974LI1K)
6.4	n_0, n_1	(1970BA63)
8.5	n_0	(1971HI09)
1.13, 1.17	p_0	(1973LE26)
1.40 – 2.40	p_0	(1970BO20)
1.4 – 3.0	p_0, p_1	(1971BO44, 1972BO56, 1972MA77)
3.90 – 5.80	p_0	(1972BL04)
5 – 12	p_0	(1973ME22)
7.7 – 10	p_0, p_1	(1969CU10)
12.1, 12.3	p_0	(1971BR44, 1971GR20, 1973JO10)
12.4	p_0, p_1	(1971BU03)
13.6	p_0	(1967GO27)
15	$p_0 \rightarrow p_4, p_8 \rightarrow p_{11}$	(1973DA17)
51	p_0	(1971FE1D)
1.6 – 2.7	d_0	(1971ME18)
1.6 – 3.0	d_0	(1971BO39)
9, 10, 11, 12	d_0	(1971WI02)
12.1	d_0	(1971GR20)
15.0	d_0	(1974BU06)
19.8 – 29.6	d_0	(1972PE09, 1972PE15, 1974AR16)
28.0	d_0, d_1	(1970BU08)
41, 46, 51	d_0	(1971SE06)


^a See Table 14.12 in (1970AJ04) for a listing of previous studies with E_{d} to 51.5 MeV for neutron groups, 21 MeV for proton groups and 51 MeV for the d_0 group.

[1973BE25](#)) and ([1971DU1B](#), [1974ST05](#), [1975GR12](#), [1975MO1L](#); theor.). For spallation measurements see ([1971BA70](#), [1971JA02](#), [1975FU01](#)).


Polarization measurements are summarized in Table [14.17](#) [and in Table 14.12 in ([1970AJ04](#))]. The polarization transfer coefficient has been measured for $E_d = 6$ to 14 MeV for \bar{n}_0 and \bar{n}_1 . The \bar{n}_0 dependence shows some influence from the compound nucleus ([1974LI1K](#)). See also ([1975TE1A](#)). For neutron polarization studies see also ([1971TH1E](#), [1975KA26](#)), ([1971WA1D](#)) and ([1974BO51](#), [1974BO53](#), [1975BO1W](#); theor.). For proton polarization studies see also ([1971GO1E](#), [1971HA1R](#)) and ([HA70P](#), [1970PE1B](#), [1972GO27](#), [1972KO30](#), [1972SE09](#), [1974BO52](#), [1974SA11](#), [1974SA20](#), [1975SE07](#); theor.).



Reported resonances are displayed in Table [14.15](#). Recent measurements of yields of d_0 are listed in Table [14.16](#). For a discussion of the relative yields of $^{12}\text{C}^*(12.7, 15.1)$ see ^{12}C in ([1975AJ02](#)). See also ([1970VE06](#), [1971PU01](#), [1971ZA04](#), [1975JA1A](#)) and ([1969IW1A](#), [1969KO1B](#), [1970SO11](#), [1974IN07](#), [1974ST05](#), [1975GU10](#); theor.). The (d, np) processes are discussed in reaction 38 of ^{13}C and in reaction 21 of ^{13}N .



The cross section rises from ≈ 0.1 mb at $E_d = 16$ MeV to ≈ 10 mb at 20 MeV ([1955WI43](#)).



Reported resonances are displayed in Table [14.15](#). Recent measurements of the yields of α -groups are listed in Table [14.16](#) [see also Table 14.11 in ([1970AJ04](#))]. For angular distributions see ^{10}B in ([1974AJ01](#)).

The major interest in this reaction has been the study of the yield of the α_2 group to the $J^\pi = 0^+$, isospin “forbidden” $T = 1$ state. In particular, the work of ([1971RI15](#), [1972SM07](#)) has shown that while the α_0 , α_1 and α_3 yields show only weak fluctuations, the α_2 “forbidden” yield shows narrow resonances which implies that the source of the isospin mixing (at least in the region which they, and the subsequent work of ([1974JO01](#)) studied: $E_d = 7.2$ to 16 MeV) is due to states in the ^{14}N compound nucleus. The ratio of the σ_t for the α_2 group compared to the σ_t for the “allowed” groups is $\approx 1\%$, an order of magnitude greater than predicted by direct or multistep processes ([1972SM07](#)). Partial wave analyses lead to the resonance parameters shown in Table [14.15](#) ([1972SM07](#), [1974JO01](#)). For an earlier discussion of these problems see ([1970AJ04](#)). See also ([1970WE03](#)).

Table 14.18: Excited states of ^{14}N from $^{12}\text{C}(^3\text{He}, \text{p})^{14}\text{N}$ ^a

E_x (MeV \pm keV)			L ^d	$J^\pi; T$
(1971DU03)	(1969HO23)	(1968MA29)		
	0		2	
	2.319 \pm 15		0	
3.9502 \pm 1.5	3.952 \pm 15		0	
4.9153 \pm 1.4	4.927 \pm 15		1	
$\equiv 5.10587 \pm 0.18$ ^b	5.117 \pm 15		1	
5.6888 \pm 1.4	5.713 \pm 15		1	
5.8324 \pm 1.4	5.885 \pm 15		3	
6.2025 \pm 1.4	6.224 \pm 15	6.21 \pm 20	0	
6.4449 \pm 1.4	6.468 \pm 15	6.46 \pm 18	2	
7.0279 \pm 1.4	7.036 \pm 15	7.01 \pm 42	2	
7.9649 \pm 1.4	7.974 \pm 15	7.95 \pm 26	3	^l
	8.072 \pm 15	8.05 \pm 35	1	
8.4864 \pm 1.5 ^g	8.493 \pm 15	8.47 \pm 30	3	$4^-; 0$ ^{d,f}
8.6174 \pm 4	8.625 \pm 15	8.61 \pm 34	0	$(0^+; 1)$ ^h
8.9099 \pm 1.9 ^c	8.912 \pm 15			$(3^-; 1)$ ^h
8.9598 \pm 1.4				
	8.97 \pm 15	8.96 \pm 19		
8.9773 \pm 4				$(2^+; 0)$ ^h
9.1241 \pm 1.5	9.126 \pm 15			ⁱ
		9.15 \pm 18		
9.1674 \pm 1.4	9.176 \pm 15		j	$(2^+; 1)$ ^h
9.3854 \pm 1.6 ^c	9.389 \pm 15	9.39 \pm 26		$2^-; 0$ ^{d,k}
		9.51 ^f		$(2^-; 1)$ ^h
	9.703 \pm 15	9.70 \pm 22		$(1^+; 0)$ ^h
	10.063 \pm 15 ^m	10.08 \pm 18		$3^+, \geq 4$ ^f
	10.101 \pm 15			$1^+, 2^+$ ^f
		10.23 ^f		1 ^f
	10.441 \pm 15	10.43 \pm 20	j	$(2^+, 1)$ ^h
		10.56 ^f		1, 2 ^f
	10.812 \pm 15	10.81 \pm 23		$5^+, 0$ ^{e,f}

Table 14.18: Excited states of ^{14}N from $^{12}\text{C}(^3\text{He}, \text{p})^{14}\text{N}$ ^a (continued)

E_x (MeV \pm keV)			L ^d	$J^\pi; T$
(1971DU03)	(1969HO23)	(1968MA29)		
	11.053 \pm 15	11.06 \pm 50		
	11.249 \pm 15	11.27 \pm 50		
	11.357 \pm 15	11.39 \pm 40		
	11.517 \pm 15	11.51 \pm 30		
		11.66 \pm 40		
		11.79 \pm 110		
		11.95 \pm 30		
	12.29 \pm 15			
	12.425 \pm 15	12.40 \pm 30		
	12.506 \pm 15	12.50 \pm 20		
	12.608 \pm 15	12.63 \pm 25		
	12.69 \pm 15	12.74 \pm 30		
	12.80 \pm 15			
		12.90 \pm 25		
		13.15 \pm 40		
		14.91 \pm 60		
		15.8 \pm 200		
		17.4 \pm 200		

^a See also Table 14.14 in (1970AJ04).

^b All E_x shown by (1971DU03) are measured relative to this energy obtained by (1967CH19) from E_γ .

^c The widths of $^{14}\text{N}^*(8.91, 9.39)$ are, respectively, 19.7 ± 1.9 and 15.6 ± 2.0 keV.

^d (1968MA29).

^e (1972NO08): from study of angular correlations. See also (1968MA29).

^f (1974NO01): from a study of decay proton correlation ($^{12}\text{C}(^3\text{He}, \text{p}')^{14}\text{N}^*(\text{p})^{13}\text{C}_{\text{g.s.}}$) with the relevant p' group.

^g $\Gamma_{\text{p}}/\Gamma = 0.73 \pm 0.10$ (1974NO01).

^h Known from other data: consistent with results of (1974NO01).

ⁱ Unresolved doublet: see reactions 21 and 26.

^j $\theta_{\text{p}}^2(l=3) = (2.3 \pm 1.1) \times 10^{-3}$ and $< 1.6 \times 10^{-3}$ for $^{14}\text{N}^*(9.17, 10.43)$ (1974NO01).

^k The results of (1974NO01) are consistent with either $J^\pi = 2^-$ or 3^- for this state.

^l $\Gamma_\gamma/\Gamma = 0.7 \pm 0.2\%$; $(2J+1)\Gamma_{\text{p}} = 12.6 \pm 3.6$ eV: see Table 14.12 (1972BA56).

^m $\Gamma < 10$ keV (J.W. Noe, private communication).

17. (a) $^{12}\text{C}(\text{d}, ^6\text{Li})^8\text{Be}$	$Q_m = -5.893$	$E_b = 10.2725$
(b) $^{12}\text{C}(\text{d}, ^7\text{Li})^7\text{Be}$	$Q_m = -17.543$	

For reaction (a) see (1970AN1E) and ^8Be in (1974AJ01). For reaction (b) see (1975FU01) and ^7Be in (1974AJ01).



Angular distributions have been measured at $E_t = 1.12$ to 1.68 MeV (1971MA46: n_0, n_1, n_2) and at 8 MeV (1972CO01: to ^{14}N states with $E_x < 8.7$ MeV). In the latter experiment L -assignments are suggested (1972CO01). For τ_m measurements see Table 14.13 (1968AL12).



Many proton groups have been observed: see Table 14.18. Angular distributions have been measured at many energies up to $E(^3\text{He}) = 25.3$ MeV: see (1970AJ04) for a listing of the earlier work and (1969SC1G: $2.22 - 3.60$ MeV; p_0, p_1, p_2), (1970CL1D: 3.0 to 4.05 MeV; p_0, p_1, p_2), (1971DU03: 11.95 MeV; to states shown in Table 14.18) and (1972BR60: 13.3 MeV; to states with $E_x < 8.7$ MeV). The reaction mechanism involves the compound nucleus, at least for $E(^3\text{He}) < 12$ MeV: see (1973HE09, 1973SO04) and ^{15}O .

Extensive studies of $\text{p}'\gamma$ and $\text{p}'\text{p}$ correlations (the latter from $^{12}\text{C}(^3\text{He}, \text{p}')^{14}\text{N}*(\text{p})^{13}\text{C}_{\text{g.s.}}$) have led to the confirmation and determination of J^π of many of the unbound states [see Table 14.18 and (1972NO08, 1974NO01, 1974NO1G)] and of Γ_γ/Γ [see Table 14.12 and (1972BA56, 1972NO08)]. Table 14.12 also displays branching ratios and radiative widths studied earlier. These studies led to $J^\pi = 0^-$ or 1^- for $^{14}\text{N}^*(4.92)$, $J^\pi = 2^-$ for $^{14}\text{N}^*(5.11)$, odd parity for $^{14}\text{N}^*(5.83)$, even parity for $^{14}\text{N}^*(6.20, 6.44)$, $J = 2$ for $^{14}\text{N}^*(7.03)$: see (1970AJ04) for a fuller discussion and a listing of the relevant references. For τ_m measurements see Table 14.13 (1970AJ04, 1973HA40). (1973BEVW) find $0.5 \leq |g| \leq 0.85$ for the 3^- state $^{14}\text{N}^*(5.83)$. See also (1970CA28), (1970AN1D) and (1970LK1A, 1973LI15, 1974GR39, 1974NE18; theor.).



Angular distributions of deuterons corresponding to $T = 0$ states in ^{14}N have been measured at $E_\alpha = 42$ to 55 MeV: see Table 14.19 and (1970AJ04) for a listing of the references. The deuteron spectrum is dominated by very strong groups corresponding to the $(\text{d}_{5/2})^2, J^\pi = 5^+$, state at 8.96 MeV, and to a state at 15.1 MeV (1962HA40, 1966RI04). See also (1974VAZK, 1975VA1H), (1970AN1D) and (1971BU1K; theor.).

Table 14.19: Energy levels of ^{14}N from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$

E_x ^a (MeV)	σ ^a (mb)	$\sigma/\sigma_{\text{g.s.}}$ ^b	Angular distribution ^c
0	5.8	1.00	d, α ⁱ
2.31 ^d	d		α
3.95	4.8	0.31	d, α ⁱ
4.92	4	0.16	α
5.11	5	1.32	d, α
5.69	4	0.11	α
5.83	6	0.97	d, α
6.20	7.0	0.24	α
6.44	13.0	1.15	d, α
7.03	4.0	0.18	d, α
7.97	5.0	0.16	d, α
8.49	8.3	0.53	d, α
9.00 ± 0.05 ^e	28.0	3.67	d, α
$9.12 + 9.13 + 9.17$	medium	(0.30)	
9.39	8.1	0.30	d, α
9.70	6.9	0.18	α
10.10	13.8	0.5	α
10.76 ± 0.02 ^f	medium	0.62	
11.06	weak	very weak ^h	
11.24	medium		
11.29	medium		
11.39	medium	very weak, broad	
11.51	weak		
11.95	weak		
12.24	weak		
		weak	
12.41	medium		
12.61	strong		
12.69	strong	strong, broad ($E_x = 12.76$)	
13.05 ± 0.02 ^g		very weak	

Table 14.19: Energy levels of ^{14}N from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$
(continued)

E_x^{a} (MeV)	σ^{a} (mb)	$\sigma/\sigma_{\text{g.s.}}^{\text{b}}$	Angular distribution ^c
14.7 ^h		weak, broad	
$15.1 \pm 0.1^{\text{e}}$		strong, sharp	
15.5		weak	
16.0		weak	
16.3		medium	
17.1		medium	
17.7		medium	

^a From $^{12}\text{C}(^6\text{Li}, \alpha)$. $E(^6\text{Li}) = 20$ MeV ([1968ME10](#)).

^b Cross section (relative to ground state) integrated for $0^\circ - 90^\circ$ (c.m.), from $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$, $E_\alpha = 42$ MeV ([1967ZA01](#)). The $T = 1$ states at $E_x = 2.31, 8.06, 9.51, 10.43$ MeV were not observed: $\sigma/\sigma_{\text{g.s.}} < 0.0027, 0.027, 0.018, 0.30, 0.03$, respectively. For absolute cross sections at $E_\alpha = 53$ MeV; see ([1965PE03](#)).

^c d and α refer to angular distributions of the corresponding group from the $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ and the $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$ reactions: see text for E_α , $E(^6\text{Li})$ and references.

^d The excitation of $^{14}\text{N}^*(2.3)$ is inhibited by angular momentum and parity consideration as well as by isospin in the (α, d) reaction.

^e ([1966RI04](#)).

^f $E_x = 10.85 \pm 0.02$ MeV ([1967ZA01](#)).

^g From $^{12}\text{C}(\alpha, \text{d})^{14}\text{N}$ ([1967ZA01](#)): $\Gamma < 70$ keV.

^h Comments here and below are from ([1965PE03](#)).

ⁱ Angular distributions reported at $E_\alpha = 55$ MeV ([1974VAZO](#)).

$$21. \ ^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N} \quad Q_m = 8.7987$$

At $E(^6\text{Li}) = 20$ MeV, α -groups corresponding to most of the $T = 0$ states with $E_x < 12.7$ MeV are reported: see Table 14.19. The spectrum is dominated by the α -group corresponding to the 5^+ state at 9.0 MeV ([1968ME10](#)). The α_1 group to the 0^+ ; $T = 1$ state at 2.31 MeV state has an intensity (< 3% of α_0) which decreases sharply from $E(^6\text{Li}) = 4$ to 5.5 MeV, while the intensities of the α_0 and α_2 groups increase rapidly ([1965CA06](#)). Angular distributions of α -particles have been reported for $E(^6\text{Li}) = 2.0$ to 20 MeV: see ([1970AJ04](#)) for the earlier references, ([1970JO09](#), [1970JO1D](#): 5.6 to 14.0 MeV; $\alpha_0, \alpha_2, \alpha_3 + \alpha_4$) and ([1973WH03](#), [1975WH03](#): 33 MeV; α to $^{14}\text{N}^*(0, 3.95, 4.92, 5.11, 5.69, 5.83, 6.20, 6.44)$). At that energy the reaction proceeds primarily by

direct interaction ([1973WH03](#)); multistep mechanisms may be very important ([1975WH03](#)). For τ_m measurements see Table [14.13](#). For angular correlation measurements see ([1974NO1G](#)). See also ([1971LA1B](#)) and ([1970OG1A](#), [1973OG1A](#)).



See ([1969RO1G](#), [1970LK1A](#); theor.).




At $E(^{11}\text{B}) = 114$ MeV the spectrum is dominated by groups to the 5^+ state at $E_x = 8.96$ MeV and to one or more of the states at 12.8 MeV, presumably the 4^+ one ([1974AN36](#), [1975PO10](#)). See also ([1972SC1L](#), [1973SC1J](#)). The angular distribution of the group to $^{14}\text{N}^*(8.96)$ is a smoothly varying, exponential function of angle ([1975PO10](#)): see the analyses by ([1965SA07](#), [1975PO10](#)). See also ([1970AN1D](#), [1970LK1A](#); theor.).



This reaction has been studied at $E(^{12}\text{C}) = 114$ MeV: the spectrum is dominated by $^{14}\text{N}^*(8.96)$ [$J^\pi = 5^+$] but there is substantial population also of $^{14}\text{N}^*(5.83)$ [3^-] and of a state at $E_x = 11.2$ MeV ([1974AN36](#)). See also ([1971SC1F](#), [1972SC1L](#), [1972SC21](#)).



See ([1968RO1D](#); theor.).



Observed resonances are displayed in Table [14.20](#). The decay schemes of various levels of ^{14}N , as derived from the γ -spectra in this and other reactions, are exhibited in Table [14.12](#) [see ([1970AJ04](#)) for the earlier references and see ([1972RE10](#))]. For the τ_m of bound ^{14}N states see Table [14.13](#) ([1971BI03](#), [1972RE10](#)).

Table 14.20: Levels of ^{14}N from $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$ and $^{13}\text{C}(\text{p}, \text{p})^{13}\text{C}$ ^a

E_{p} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	l_{p}	$\omega\Gamma_{\gamma}$ (eV)	$J^{\pi}; T$	$^{14}\text{N}^*$ (MeV)	Refs.
0.4485 ± 0.5	< 0.37	2	0.022	2^-	7.9669	A
0.551 ± 1	30 ± 1	0	9.2	$1^-; 1$	8.062	A, (1972RE10)
1.012 ± 2	≤ 0.2	4	≈ 0.01	$(4^-); 0$	8.490	A
1.150 ± 2	7 ± 1	1	1.3	$0^+; 1$	8.618	A, (1971BI03)
1.34 ± 50	≈ 460	0	12.8	$0^-; 1$	8.79	A
1.462 ± 3	16 ± 2	2	0.72	$3^-; 1$	8.907	A
1.523 ± 2	< 1		≈ 0.003	$5^+; 0$	8.964	A
1.540 ± 3	8 ± 2	1, (3)	0.13	2^+	8.980	A
1.701 ± 1	< 1	2	≈ 0.03	$(2^-; 0)^j$	9.129	A
1.7476 ± 0.9	0.07 ± 0.05		14.8	$2^+; 1$	9.1725	A
1.980 ± 3	13 ± 3	2	^c	$3^-, 2^-$	9.388	A, (1975NO1F)
2.110 ± 3	41 ± 2	2	6.2	$2^-; 1$	9.509	A
2.319 ± 4	15 ± 3	1		1^+	9.703	A
2.743^b	5	1	^d	$1^+, (2^+)$	10.096	A
2.885 ± 10^b	80 ± 15	0, 2		$1(^-); 0$	10.228	A
3.105 ± 7^b	33 ± 3	1	17	$2^+; 1$	10.432	A, (1971RI13)
3.20^b	140	0, 2		1^-	10.52	(1961KA04)
3.515 ± 6			^e		10.812	(1975NO1F)
3.72 ± 30^f	165 ± 30				11.00	(1971RI13)
3.771 ± 5	≤ 2		^k	$(2, 3)^+$	11.050	(1971RI13)
3.79	100			1^+	11.07	A
3.94 ± 30^g	220 ± 30				11.21	(1971RI13)
3.98^b	11	2		3^-	11.24	(1961KA04)
4.04^b	175	2		2^-	11.30	A
4.14^b	28	1		1^+	11.39	A
4.525 ± 15^h	115 ± 10		^l	1^+	11.750	A, (1971RI13)
5.325 ± 10	48 ± 7		^m		12.492	(1971RI13) ^p
5.88 ± 20^f	120 ± 30				13.01	(1971RI13)

Table 14.20: Levels of ^{14}N from $^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$ and $^{13}\text{C}(\text{p}, \text{p})^{13}\text{C}$ ^a (continued)

E_{p} (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	l_{p}	$\omega\Gamma_{\gamma}$ (eV)	$J^{\pi}; T$	$^{14}\text{N}^*$ (MeV)	Refs.
6.20 ± 100 ⁱ	1000 ± 150		ⁿ	(2 ⁻); 1	13.30	(1971RI13)
6.62 ± 20 ^f					13.69	(1971RI13)
o						
16.1				2 ⁻ ; 1	22.5	(1971RI13)
16.7				2 ⁻ ; 1	23.0	(1971RI13)
					^q	

A: See references for this state quoted in Table 14.16 in (1970AJ04).

^a See also Table 14.12.

^b Reduced width for proton emission is of the order of 1% of the Wigner limit (1961KA04).

^c Gamma decays predominantly to $^{14}\text{N}^*(5.11, 5.83, 8.9) (\approx 5 \text{ meV each})$ (1975NO1F).

^d $(2J+1)\Gamma_{\gamma} = 0.5 \pm 0.2 \text{ eV}$, $\Gamma = 12 \pm 3 \text{ eV}$ (1960RO23).

^e Observed transitions $10.81 \rightarrow 6.44 \rightarrow \text{g.s.}$: $\Gamma_{\gamma} = 15.8 \pm 5.9 \text{ meV}$ (1975NO1F). See also Tables 14.12 and 14.24.

^f Weak resonance.

^g See also Table 14.16 in (1970AJ04).

^h In the $\gamma_{3.09}$ channel the leak occurs 55 keV higher (1971RI13); interference effects may be present.

ⁱ Part of the giant dipole resonance.

^j See, however, (1975FO01) in reaction 6.

^k $(2J+1)\Gamma_{\gamma_0} = (1.2 \pm 0.3) \Gamma/\Gamma_{\text{p}} \text{ eV}$ (1971RI13).

^l $(2J+1)\Gamma_{\gamma} = (18.5 \pm 4.2) \Gamma/\Gamma_{\text{p}} \text{ eV}$; if $J = 1$, $\Gamma_{\gamma} \geq 6 \text{ eV}$ (1971RI13).

^m $(2J+1)\Gamma_{\gamma_0} = 2.3 \Gamma/\Gamma_{\text{p}} \text{ eV}$; if $\Gamma = 38 \text{ eV}$ is assumed (1971RI13).

ⁿ $(2J+1)\Gamma_{\gamma_0} \geq 200 \text{ eV}$ (1971RI13): thus the transition is dipole and $T = 1$. The resonance is asymmetric and it is suggested that two states are involved, one with $J^{\pi} = 1^-$ at $E_x = 12.7$ and the other one with 2^- at $E_x = 13.3 \text{ MeV}$.

^o Some broad structure is evident in the γ_0 , $\gamma_{3.68}$ and $\gamma_{3.85}$ yields (1971RI13).

^p See also (1974GM01).

^q Two $T = 2$ states reported by (1971RI13) are not confirmed by (1975PA18).

The low-energy capture cross-section yields an extrapolated S -factor at $E_{\text{p}} = 25 \text{ keV}$ (c.m.), $S_0 = 6.0 \pm 0.8 \text{ keV}\cdot\text{b}$ (1960HE14). See also (1970AJ04) and (1971BA1A, 1972CA1N, 1973CL1E, 1973TR1E; astrophys. considerations). The capture cross section rises from $(7.7 \pm 1.8) \times 10^{-10} \text{ b}$ at $E_{\text{p}} = 100 \text{ keV}$ to $(9.8 \pm 1.2) \times 10^{-9} \text{ b}$ at $E_{\text{p}} = 140 \text{ keV}$ (1961HE02).

Following is a summary of the reasons for the assignments of $J^{\pi}; T$ to some of the lower resonances displayed in Table 14.20: for a fuller discussion and complete references see (1970AJ04) and see Table 14.12. $^{14}\text{N}^*(7.97)$: angular distribution of the γ -rays is consistent with $J^{\pi} = 2^-$.


$^{14}N^*(8.06)$: width of resonance, isotropy of γ -rays show $l_p = 0$: $J^\pi = 1^-$ from $^{13}\text{C}(\text{p}, \text{p})$; E1 transition to g.s. is uninhibited, e.g., $T = 1$ [but 1.4% $8.06 \rightarrow 2.3$ transition [$E_x = 2312.6 \pm 0.3$ keV ([1971BI03](#))] shows $T = 0$ admixture: $\alpha^2 = 0.046$ ([1972RE10](#))]. The strong transition $8.06 \rightarrow 5.69$ [3.5% ([1972RE10](#))] permits either E1 or M1, $\Delta T = 1$. Since $5.69 \rightarrow 2.31$ is seen $^{14}N^*(5.69)$ cannot have $J^\pi = 0^+$, and 2^+ is excluded by the strength of the $8.62 \rightarrow 5.69$ transition. It is then $J^\pi = 1^-$; $T = 0$ [the isospin mixing $\alpha^2 = 0.09$ ([1972RE10](#))]; $E_x = 5690.5 \pm 1.5$ keV ([1971BI03](#)). $^{14}N^*(8.49, 8.96, 9.13)$ correspond to anomalies in the cross section. The nature of their γ -decays [see Table [14.12](#)] and the angular distribution leads to $J^\pi = 4^-, 5^+$ and 2^- ; $T = 0, 0$ and 0 , respectively ([1965DE19](#)). However, work by ([1975FO01](#)) in $^{10}\text{B}(^{6}\text{Li}, \text{d})$ shows that $^{14}N^*(9.13)$ is a doublet: the state populated in reaction 6 has $J^\pi = 3^+$. The structure at $E_p = 1.70$ MeV will therefore have to be reanalyzed.

$^{14}N^*(8.62)$ [$J^\pi = 0^+$ from $^{13}\text{C}(\text{p}, \text{p})$] shows strong transitions to $^{14}N^*(0, 3.95, 5.69)$: $T = 1$ ([1959WA16](#)). The strength of the $8.62 \rightarrow 3.95$ decay shows it is dipole and therefore $J = 1$ for $^{14}N^*(3.95)$ ([1959WA04](#)) [$E_x = 3947.6 \pm 0.4$ keV ([1971BI03](#))]. The strength of the transition $8.62 \rightarrow 6.21$ and the angular correlation $8.62 \rightarrow 6.21 \rightarrow$ g.s. is consistent with $J^\pi = 1^+$; $T = 0$ for $^{14}N^*(6.20)$ [$E_x = 6203.7 \pm 0.6$ keV ([1971BI03](#))]. $^{14}N^*(8.79)$ [$J^\pi = 0^-$ from $^{13}\text{C}(\text{p}, \text{p})$] has a large Γ_γ consistent with E1 and $T = 1$. $^{14}N^*(9.17)$: angular correlation and angular distribution measurements indicate $J^\pi = 2^+$ for that state, 3^- for $^{14}N^*(6.44)$ [see, however, Table [14.11](#)] and $J = 2$ for $^{14}N^*(7.03)$.


The angular distribution of the γ -rays from $10.23 \rightarrow 2.31$ is consistent with $J^\pi = 1^+$ for $^{14}N^*(10.23)$: $T = 0$ from $M^2(\text{M1})$ ([1963RO17](#)) [see, however, Table [14.11](#)]. The γ_0 angular distribution is consistent with $J = 2$ for $^{14}N^*(10.43)$: the similar decay characteristics of this state and of $^{14}N^*(9.17)$ suggest that they are both $J^\pi = 2^+$; $T = 1$ ([1964RO03](#)).

Below $E_p = 5.5$ MeV only γ_0 can be observed in the capture radiation. ([1971RI13](#)) have observed a number of resonances in the γ_0 yield and in the yield of the ground state γ -rays from $^{13}\text{C}^*(3.09, 3.68, 3.85)$: these are shown in Table [14.20](#) in the range $E_p = 3.7$ to 6.6 MeV [see reaction 25 in ([1970AJ04](#)) for the earlier work]. Angular distributions and measurements of Γ_{γ_0} lead to the J^π values shown. Above $E_p = 7$ MeV the γ_0 yield shows broad structure and the giant dipole resonance at $E_x = 22.5$ and 23.0 MeV ([1971RI13](#)). Measurements by ([1975PA18](#)) of the γ_0 and γ_1 90° yields for $E_x = 23$ to 33 MeV show that the $J^\pi = 1^-$; $T = 0$ giant dipole resonance is concentrated between $E_x = 15$ and 23 MeV, that the $T = 2$ resonances reported by ([1971RI13](#)) at $E_x = 23.7$ and 24.2 MeV do not exist, and that there is no evidence for the $T = 2$ GDR between $E_x = 25$ and 29 MeV ([1975PA18](#)). The 90° yields of γ -rays to $T = 0$ states ($4.9 < E_x < 5.9$ MeV) and to $T = 1$ states ($8.0 < E_x < 9.5$ MeV) have been measured for $E_x = 23$ and 26 MeV, respectively, to $E_x = 33$ MeV; the former is quite constant at $\approx 4 \mu\text{b}/\text{sr}$, the latter slowly rises to a value of $\approx 2 \mu\text{b}/\text{sr}$: by comparison the γ_1 yield is $\approx 1 \mu\text{b}/\text{sr}$ ([1975PA18](#)). See also ([1970SI1E](#), [1973BE1R](#)), ([1973SU1E](#)) and ([1975VE01](#); theor.).


The elastic scattering has been studied for $E_p = 0.14$ to 11 MeV (see (1970AJ04) for a listing of the older work) and at $E_p = 4.6$ to 5.3 MeV (1974GM01: also $p_1 \rightarrow p_3$). For observed resonances see Table 14.20; for angular distributions see ^{13}C . Polarization measurements for the p_0 group have been carried out at 7 , 14.5 and 32.9 MeV (see (1970AJ04)) and at $E_p = 30.4$ MeV (1972GR02; also for p_1). For spallation measurements see (1973RA37).




The yield of neutrons has been measured from threshold to $E_p = 13.7$ MeV: see (1970AJ04). Observed resonances are displayed in Table 14.21 (1959GI47, 1961DA09). The ratio of the reaction cross section at $E_p = 22.8$ MeV to the n_0 yield is 1.06 ± 0.07 : thus there is little competition of γ -rays from excited states of ^{13}N with neutron emission making this a convenient fast neutron calibration source (1975LI11). Polarization measurements are reported for $E_p = 6.9$ to 12.3 MeV (1965WA02: n_0) and 7 to 15 MeV (1974LI1J). See also ^{13}N and (1971GE12; theor.).



A polarization measurement has been reported at $E_p = 7$ MeV by (1969GU02). See also (1970AJ04) and ^{12}C in (1975AJ02).



At $E_p = 49.6$ MeV polarization measurements have been carried out for the tritons and ^3He ions to the mirror groups $^{11}\text{B}^*(0, 2.12, 4.45, 5.02, 6.74, 12.91)$ and $^{11}\text{C}^*(0, 2.00, 4.32, 4.80, 6.48, 12.50)$ (1974MA12). Integrated cross sections for populating the first four states in ^{11}B and ^{11}C have been measured at several energies in the range $E_p = 26.8$ to 43.1 MeV (1975MI01). See also ^{11}B and ^{11}C in (1975AJ02).



Excitation functions have been measured from $E_p = 5.5$ (α_0), 6.0 (α_1), 7.0 (α_2), 8.0 (α_3), 10 (α_4), 11 (α to $^{10}\text{B}^*(5.11)$) to 18 MeV. Total cross sections have also been obtained for the production of ^6Li , ^9Be and ^{10}B : the latter shows a great deal of structure. The consequences for astrophysical problems are discussed by (1975OB01).

Table 14.21: Resonances in $^{13}\text{C}(\text{p}, \text{n})^{13}\text{N}$
^a(1961DA09)

E_{p} (MeV)	Γ (keV)	$^{14}\text{N}^*$ (MeV)
3.76 ± 0.05	100	11.04
3.98 ± 0.02	30	11.24
4.05		11.31
4.15 ± 0.02	40	11.40
4.5 ± 0.1	100	11.7
4.7 ± 0.1	150	11.9
5.03 ^b		12.22
(5.44 ± 0.03)	(60)	(12.60)
5.53 ± 0.03	50	12.68
5.72 ± 0.03	60	12.86
6.20 ± 0.04	70	13.30
6.67 ± 0.13	250	13.74
7.0 ± 0.1	150	14.0
7.3		14.3
7.85 ± 0.08	150	14.83
7.93 ± 0.03	50	14.91
8.03 ± 0.03	50	15.00
8.7 ± 0.2	350	15.6
9.3 ± 0.1	150	16.2
10.2 ± 0.2	400	17.0
11.4 ± 0.3	600	18.1

^a See also Table 14.9 in (1959AJ76).

^b (1959GI47).

Table 14.22: ^{14}N levels from $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$ and $^{13}\text{C}({}^3\text{He}, \text{d})^{14}\text{N}$

$^{14}\text{N}^*(\text{MeV})$	$J^\pi; T$	l_p^{a}	l_p^{b}
g.s.	$1^+; 0$	1	1
2.31	$0^+; 1$	1	1
3.95	$1^+; 0$	1	1
4.92	$0^-; 0$	0	0
5.11	$2^-; 0$	2	2
5.69	$1^-; 0$	0	0
5.83	$3^-; 0$	2	2
6.20	$1^+; 0$	isotropic ^c	1
6.44	$3^+; 0$	1^{c}	1
7.03	$2^+; 0$	1	1
7.97	$2^-; 0$	d	$1, 2^{\text{g,h}}$
8.06	$1^-; 1$	0^{e}	
8.49	$4^-; 0$	$(3, 4)^1$	4^{i}
8.62	$0^+; 1$	$0^{\text{f}}, 1^1$	1^{j}
8.91	$3^-; 1$	$2^{\text{c,l}}$	$2^{\text{g,k}}$
8.98	$2^+; (0)$	$(1, 2, 3)$	
9.13	$2^-; 0$	2^1	
9.17	$2^+; 1$	$(1, 3)^1$	
9.39	$2^-, 3^-; 0$	2^1	
9.51	$2^-; 1$	2^1	
9.70	$1^+; 0$	1^1	

^a $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$: $E_{\text{d}} = 5.5$ and 6 MeV ([1966FU10](#)), $4.5, 5.0$ and 5.5 MeV ([1973BO10](#)), 6.5 MeV ([1975BO35](#)).

^b $^{13}\text{C}({}^3\text{He}, \text{d})^{14}\text{N}$: $E({}^3\text{He}) = 15$ MeV ([1966HO15](#), [1971FO05](#)).

^c ([1973BO10](#)).

^d Angular distributions not complete because groups partly masked by contaminant.

^e ([1973BO10](#)) report $l = 1$ in their Table 1: this is a typographical error (see p. 367).

^f Expected $l = 1$ ([1973BO10](#)).

^g See also ([1969HO23](#)).

^h The width obtained for this state in $^{13}\text{C}(\text{p}, \gamma)$: $(2J + 1)\Gamma_{\text{p}} = 12.6 \pm 3.6$ eV implies $l_{\text{p}} = 2$ and therefore odd parity: Γ_{p} is then 2.5 ± 0.07 eV, based on $J = 2$ ([1972BA56](#)).

ⁱ $\Gamma_{\text{p}} < 9.9 \times 10^{-2}$ eV ([1971FO05](#)).

^j $\Gamma_{\text{p}} < 18$ keV ([1971FO05](#)).

^k $\Gamma_{\text{p}} = 12.1$ keV ([1971FO05](#)).

^l ([1975BO35](#)).

Table 14.23: Relative spectroscopic factors for ^{14}N states ^a

$^{14}\text{N}^*$ (MeV)	$J^\pi; T$	$^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$				$^{13}\text{C}({}^3\text{He}, \text{d})^{14}\text{N}$			Theory ^j
		E_{d} (MeV)	4.5 – 5.5 ^b	6.0 ^c	6.5 ^d	7.0 – 12.0 ^e	13, 17 ^g	15 ^h	
g.s.	$1^+; 0$	1.00	1.00			1.00	1.00	1.00	1.00
2.31	$0^+; 1$	0.97	1.15			0.92 ± 0.05	1.77	1.53	1.24
3.95	$1^+; 0$	0.38	0.57			0.55 ± 0.03	0.51	0.39	0.29
4.92	$0^-; 0$	0.70	1.05			1.38 ^f		0.93	0.71
5.11	$2^-; 0$	0.67	1.26			1.10		0.78	0.68
5.69	$1^-; 0$	0.79	1.04			0.88		0.85	0.58
5.83	$3^-; 0$	0.87	1.03			0.94		0.51	0.59
6.20	$1^+; 0$					0.17		≈ 0.03	
6.44	$3^+; 0$							≈ 0.04	
7.03	$2^+; 0$	0.23	0.07			0.17		0.12	0.09
7.97	$2^-; 0$	0.12							0.04 ^k
8.06	$1^-; 1$	0.74							
8.49	$4^-; 0$	0.18							< 0.07
8.62	$0^+; 1$	0.10		0.03					< 0.22
8.91	$3^-; 1$	0.60		0.03					0.67
8.98	$2^+; (0)$			0.04, 0.02 ^l					
9.13	$2^-; 0$			0.04 ^m					
9.17	$2^+; 1$			0.013, 0.01 ^l					
9.39	$2^-, 3^-; 0$			0.09 ^m					
9.51	$2^-; 1$			0.12 ^m					
9.70	$1^+; 0$			0.02 ⁿ					

^a See review in (1973BO10). See also Table 14.19 in (1970AJ04).

^b (1973BO10).

^c (1966FU10).

^d (1975BO35).

^e (1968CO24).

^f This value and the ones below are at $E_d = 7$ MeV (1968CO24): see (1973BO10).

^g 13 and 17 MeV, energy averaged (1966SI02): see (1973BO10).

^h (1966HO15).

ⁱ (1971FO05): see also (1973BO10).

^j Negative parity states (1967CO32), positive parity states (1963SE19): see (1973BO10).

^k Based on $l_p = 2$: see Table 14.22 (1972BA56).

^l If $l = 1$ or 3, respectively (1975BO35).

^m Derived with $d_{\frac{5}{2}}$ form factors (1975BO35).

ⁿ Derived with a $p_{\frac{1}{2}}$ form factor (1975BO35).



Observed neutron groups are displayed in Table 14.22. Angular distributions have been reported at many energies up to $E_{\text{d}} = 12$ MeV: see (1970AJ04) for the earlier references and (1973BO10): $E_{\text{d}} = 4.5, 5.0, 5.5$ MeV; to states shown in Table 14.22), (1975BO35): $E_{\text{d}} = 6.5$ MeV; to states in Table 14.22). Comparison of relative spectroscopic factors obtained in this reaction and in reaction 32 are shown in Table 14.23: it appears that S_{rel} for $^{14}\text{N}^*(2.31)$ [$T = 1$] is smaller in this reaction than in the (^3He , d) reaction although simple DWBA calculations would suggest that the factors would be the same in both proton pickup reactions. The $\tau \cdot \mathbf{T}$ terms appears to be energy dependent: see Table 14.23 (1968CO24).

Observed γ -rays attributed to transitions in ^{14}N are shown in Table 14.20 of (1970AJ04): see Table 14.12 here for a general review of radiative decays in ^{14}N . The angular correlation of internal pairs conclusively establish the parities of $^{14}\text{N}^*(4.92, 5.11, 5.69)$ as odd (1964WA05). For τ_m measurements see Table 14.13 (1974RU1B). See also (1972RE1C) and ^{15}N .



Angular distributions have been studied at $E(^3\text{He}) = 13$ and 17 MeV (1966SI02: d_0, d_1, d_2) and 15 MeV (1966HO15, 1969HO23, 1971FO05: see Table 14.22). Relative spectroscopic factors are displayed in Table 14.23 and compared with those from $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$. See also (1967SP09) and (1970BO1K, 1971DZ07; theor.).



Angular distributions have been measured at $E_\alpha = 27$ MeV for the α -groups to $^{14}\text{N}^*(0, 2.31, 3.95, 4.92, 5.11)$: relative spectroscopic factors are 1.0, 0.2, 0.3, 4.4, 5.5 (1974KE06). See also (1974DM01; 23.2, 23.7, 25.3 MeV; t_0). [The earlier work at $E_\alpha = 46$ MeV (see (1970AJ04)) has not been published.]



At $E(^7\text{Li}) = 34$ MeV, angular distributions have been obtained for the transitions to $^{14}\text{N}^*(0, 2.31, 3.95)$ (1973SC26).



This reaction has been studied at $E(^{11}\text{B}) = 113.5 \text{ MeV}$ ([1967PO13](#)) and 114 MeV ([1974AN36](#)). The relative population of the observed ^{14}N states is dependent upon whether $^{10}\text{Be}_{\text{g.s.}}$ or $^{10}\text{Be}_{3.37}^*$ is involved: in all cases the cross section is greater if the 2^+ state in ^{10}Be is populated. Difficulties are encountered in separating the two sets of ^{14}N states: see ([1974AN36](#)).



See ^{14}C .



$$E_{\text{thresh.}} = 670.90 \pm 0.09 \text{ keV} \quad (\textcolor{red}{1974HI1C}).$$

Neutron thresholds have been observed at $E_p = 671.5 \pm 0.5$ and $3149.6 \pm 1.1 \text{ keV}$ ([1956SA06](#)) and at $E_p = 4910 \pm 8 \text{ keV}$ ([1960BA34](#)) corresponding to the ground state of ^{14}N and to excited states at $2311.9 \pm 1.2 \text{ keV}$ and $3.952 \pm 0.008 \text{ MeV}$. Angular distributions and polarizations of the neutrons corresponding to $^{14}\text{N}^*(0, 2.31, 3.95)$ have been obtained at $E_p = 6$ to 14 MeV ([1967WO05](#)) and 7.2 to 13.3 MeV ([1971WO03](#)), respectively. The angular distributions for the ground state transition require a tensor interaction: the results are consistent with a tensor strength which is not energy dependent. The polarization measurements for the 1^+ states do not clearly favor a tensor component in the effective two-body force ([1967WO05](#), [1971WO03](#)). See also ^{15}N and ([1969SC1H](#), [1972AU1A](#); theor.).



At $E(^3\text{He}) = 44.8 \text{ MeV}$, triton groups are observed corresponding to all the known levels of ^{14}N with $E_x < 7.1 \text{ MeV}$. Triton groups were also seen to unresolved states with $E_x = 8.0 - 9.5 \text{ MeV}$, to $^{14}\text{N}^*(10.43)$ and to excited states with $E_x = 12.49 \pm 0.04$, 12.83 ± 0.05 and $13.70 \pm 0.04 \text{ MeV}$. Angular distributions were obtained for nine of the triton groups and analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture. Dominant $L = 0$ transitions are found to $^{14}\text{N}^*(2.31, 3.95, 13.7)$, $L = 1$ to $^{14}\text{N}^*(5.11)$, $L = 2$ to $^{14}\text{N}^*(0, 7.03, 10.43)$ and $L = 3$ to $^{14}\text{N}^*(5.83)$ ([1969BA06](#)). See also reaction 47 and ([1969SC1H](#); theor.). Angular distributions have been studied for reaction (b) for the transitions to $^{14}\text{N}^*(0, 3.95)$ [$L = 2$ and 0 , respectively]: $^{14}\text{N}^*(3.95)$ carries at least 95% of the GT strength ([1976GOZX](#)).

40. (a) $^{14}\text{N}(\gamma, \text{n})^{13}\text{N}$	$Q_m = -10.554$
(b) $^{14}\text{N}(\gamma, \text{p})^{13}\text{C}$	$Q_m = -7.5507$
(c) $^{14}\text{N}(\gamma, \text{d})^{12}\text{C}$	$Q_m = -10.2725$
(d) $^{14}\text{N}(\gamma, \text{pn})^{12}\text{C}$	$Q_m = -12.4971$
(e) $^{14}\text{N}(\gamma, \alpha)^{10}\text{B}$	$Q_m = -11.6134$
(f) $^{14}\text{N}(\gamma, \text{n}\alpha)^9\text{B}$	$Q_m = -20.049$
(g) $^{14}\text{N}(\gamma, \text{p}\alpha)^9\text{Be}$	$Q_m = -18.199$

The total absorption over the range $E_\gamma = 9$ to 31 MeV is dominated by a single peak at 22.5 MeV [estimated $\sigma \approx 29$ mb, $\Gamma \approx 2 - 3$ MeV] and appreciable strength extending beyond 30 MeV. The cross section cannot be accounted for solely by the (γ, n) and (γ, p_0) processes: particle unstable excited states of ^{13}C , ^{13}N are involved ([1969BE92](#)). The combined (γ, n) and (γ, pn) cross section begins to rise rapidly above 18 MeV, reaches its maximum value of 15 mb at 23.3 MeV and exhibits structure at about 19, 20.5 and 26 MeV. The main peak [$\Gamma \approx 3.5$ MeV: see ([1970AJ04](#))] at 23.3 MeV appears to be split into two absorption levels ([1970BE54](#), [1975BE60](#), [1975BE1F](#): monoenergetic photons). Maxima reported in other experiments and “breaks” in the (γ, n) activation curve are listed in ([1970AJ04](#)). See also ([1968KA38](#), [1971FR11](#), [1971KA70](#)).

The (γ, p_0) and (γ, p_2) cross sections and angular distributions have been measured in the giant resonance region by ([1972CA34](#), [1974BA37](#)). The authors infer that the giant dipole states $[(\text{p}_{3/2})^{-1}(2\text{s}1\text{d})]$ which decay by p_0 emission to $^{13}\text{C}^*(3.68)$ carry $\approx 90\%$ of the E1 strength and do not contribute substantially to the (γ, p_0) process which is populated by $(\text{p}_{1/2})^{-1}(2\text{s}1\text{d})$ giant dipole states. Above $E_\gamma = 22$ MeV d-wave emission from 2^- states appears to dominate the (γ, p_0) cross section ([1972CA34](#), [1974BA37](#)). For reaction (d) see ([1970SH06](#), [1972GE11](#)) and ^{13}C . For reactions (d, e, f, g) see ([1970AJ04](#)). See also ([1972BU1J](#), [1973DI1C](#), [1975BR1F](#)) and ([1969ER1A](#), [1971AN08](#), [1972GO23](#), [1973KI05](#); theor.).

41. $^{14}\text{N}(\gamma, \gamma)^{14}\text{N}$

Absorption measurements have led to the determination of τ_m (see Table 14.13) and of Γ_γ (see Tables 14.12 and 14.24): see the references in ([1970AJ04](#)) and the newer work by ([1974RAZD](#), [1975RA22](#)). ([1966SW01](#)) find $E_x = 7029 \pm 6$ keV for $^{14}\text{N}^*(7.03)$: the angular distribution of the γ -rays is consistent with $J = 2$.

42. $^{14}\text{N}(\text{e}, \text{e}')^{14}\text{N}^*$

The r.m.s. radius of ^{14}N is given as 2.540 ± 0.020 fm ([1973FE13](#), [1975SC18](#)). See also ([1971BE25](#): 2.493 ± 0.033 fm), ([1970DA20](#): 2.64 fm), ([1970AJ04](#)) and ([1969BE21](#), [1970BR1C](#),

Table 14.24: Gamma widths ^a of unbound levels from $^{14}\text{N}(\gamma, \gamma)^{14}\text{N}$ and $^{14}\text{N}(e, e)^{14}\text{N}$

E_x (MeV)	$J^\pi; T$	Γ_γ to g.s. (eV)	Multipol.	Reaction	Refs.
8.06	$1^-; 1$	10.5 ± 6	E1	(γ, γ)	(1956GR17, 1958GR97)
8.91	$3^-; 1$	$(6.6 \pm 2.2) \times 10^{-3}$	M2	(e, e)	(1968CL05)
9.17 ^{b,c}	$2^+; 1$	7.7 ± 0.9	M1	(e, e)	(1968CL05)
		8.7 ± 1.5		(γ, γ)	(1959HA11)
		9 ± 3		(e, e)	(1962ED02)
10.43 ^b	$2^+; 1$	12.1 ± 1.5	M1	(e, e)	(1968CL05)
		20 ± 6		(e, e)	(1962ED02)

^a See also Tables 14.12, 14.13 and 14.20.

^b See also (1963BA19).

^c $\Gamma = 72 \pm 10$ eV (1965LU05), 77 ± 12 eV (1959HA11).

1973TH1B). Form factors have been determined at $E_e = 60.7$ to 122.0 MeV for $^{14}\text{N}^*(2.31, 3.95, 4.92, 5.11, 5.69, 5.83)$: the reduced transition probabilities for these states, in single-particle units, are, respectively, 0.065 ± 0.020 , 1.70 ± 0.14 , $(1.1 \pm 0.5) \times 10^{-7}$, 4.1 ± 1.0 , $(3.8 \pm 2.1) \times 10^{-8}$ and 6.1 ± 1.3 (1974EN01). Inelastic scattering (at $\theta = 180^\circ$) gives evidence for the excitation of $^{14}\text{N}^*(8.91, 9.17, 10.43)$: the Γ_{γ_0} are given in Table 14.24 (1962ED02, 1963BA19, 1968CL05). In addition (1968CL05) report the excitation of a state with $E_x = 11.01 \pm 0.07$ MeV. Partial Γ_γ for cascade transitions of $^{14}\text{N}^*(9.17, 10.43)$ are shown in Table 14.12 (1968CL05). See also (1970AJ04), (1970BR1C, 1972THZF, 1974DE1E) and (1971BA2T, 1973GA19, 1974PE08; theor.).

43. $^{14}\text{N}(n, n')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered neutrons are displayed in Table 14.23 of (1970AJ04). See also (1974HO1E) and (1971MI12; theor.). Observed γ -rays are shown in Table 14.25 (1969DI1B, 1971NY03, 1972NY02). See also (1970DI1A, 1972NI05).

44. (a) $^{14}\text{N}(p, p')^{14}\text{N}^*$

$$(b) ^{14}\text{N}(p, 2p)^{13}\text{C} \quad Q_m = -7.5507$$

$$(c) ^{14}\text{N}(p, pd)^{12}\text{C} \quad Q_m = -10.2725$$

Angular distributions of elastically and inelastically scattered protons have been measured and analyzed at a number of energies up to $E_p = 155$ MeV: see Tables 14.26 here and 14.23 in (1970AJ04).

Table 14.25: Gamma rays from $^{14}\text{N}(\text{n}, \text{n}')^{14}\text{N}^*$

E_γ^a (keV)	E_γ^b (keV)	Transition ^e
2310 ± 3	2312 ± 3 ^c	$2.31 \rightarrow \text{g.s.}$
$\equiv 3945$		$3.95 \rightarrow \text{g.s.}$
1632 ± 3	1634 ± 5 ^c	$3.95 \rightarrow 2.31$
4913 ± 3	4919 ± 10 ^d	$4.92 \rightarrow \text{g.s.}$
$\equiv 5105$	5106 ± 3	$5.11 \rightarrow \text{g.s.}$
2789 ± 2	2792 ± 3	$5.11 \rightarrow 2.31$
5686 ± 3		$5.69 \rightarrow \text{g.s.}$
3378 ± 3		$5.69 \rightarrow 2.31$
5839 ± 2	5835 ± 5 ^c	$5.83 \rightarrow \text{g.s.}$
729 ± 2	727 ± 3 ^c	$5.83 \rightarrow 5.11$
	6426 ± 10 ^d	$6.44 \rightarrow \text{g.s.}$
	7021 ± 10 ^d	$7.03 \rightarrow \text{g.s.}$

^a ([1972NY02](#)): $E_n = 6.9$ MeV.

^b ([1971NY03](#)): $E_n = 15$ MeV.

^c E_x (3.95) is then 3947 ± 5 keV and E_x (5.83) = 5833 ± 3 keV.

^d These γ -ray energies have been corrected for Doppler shift.

^e ([\(1969DI1B\)](#): $E_n = 8.6$ MeV) also reports observation of the $6.44 \rightarrow 3.95$ and $6.20 \rightarrow 2.31$ transitions.

Compound nucleus effects are appreciable below $E_p = 15$ MeV ([1973HA54](#)). The inclusion of a tensor term is necessary in the analysis of the distribution of the p₁ group (to $^{14}\text{N}^*(2.31)$ [$J^\pi = 0^+$; $T = 1$]): see ([1970CR03](#), [1973AU1E](#), [1973HA54](#)). See also ^{15}O and ([1971ES1A](#)). Observed inelastic proton groups are shown in Table 14.27. See also ([1969SC1H](#), [1970SA06](#), [1971MC15](#), [1972AU1A](#), [1973KA04](#), [1974OD01](#); theor.).

Reaction (b) at $E_p = 19$ MeV proceeds at least in part through an intermediate state in ^{14}N at $E_x \approx 11.2$ MeV ([1965DE21](#)). See also ([1970WE09](#), [1970WE1J](#)) in ^{13}C , ([1971LO25](#)) and ([1972JA1C](#); theor.). For reaction (c) see ([1970AJ04](#)).

45. $^{14}\text{N}(\text{d}, \text{d}')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered deuterons have been obtained for E_d up to 52 MeV: see Table 14.23 in ([1970AJ04](#)) and ([1967FL10](#): $E_d = 2.01$ to 5.50 MeV; d_0), ([1970DU04](#): $E_d = 5.47$ (d_1) and 6.37 (d_2) to 10.16 MeV), ([1974JA25](#): $E_d = 10.02$, 12.02 MeV):

Table 14.26: $^{14}\text{N}(\text{p}, \text{p})$ angular distribution studies ^a

E_{p} (MeV)	Angular distribution of proton groups	Refs.
7.53 – 10.54	p_0, p_1, p_2	(1970ME30)
8.6	$p_0 \rightarrow p_6$	(1973HA54)
9.54, 10.54	$p_3 \rightarrow p_9$	(1970ME30)
10.6, 12.6, 14.6	$p_0 \rightarrow p_9$	(1973HA54)
11.90 – 14.62	$p_0 \rightarrow p_{10}, p_{12}$ at $E_{\text{p}} = 14.6$ MeV, p to $^{14}\text{N}^*(9.17)$	(1971OD01)
12.737 – 19.450	p_0 ^b	(1974JA25)
14.5	p corresponding to $E_x < 10.1$ MeV: Table 14.27	(1971CU01)
17	p to $E_x < 8.5$ MeV	(1971RO1H, 1971RO1J)
18 – 49.4	p_0	(1974PI05)
24.8	p_1	(1970CR03)
29.8, 36.6, 40.0	p_0, p_1, p_2	(1971FO1K, 1973AU1E)
49.4	p_0	(1971RU04)
100	p_0	(1973CH06)

^a See also Table 14.23 in (1970AJ04) for a listing of the earlier work to $E_{\text{p}} = 155$ MeV.

^b Accurate differential cross sections, few angles.

d_0 , very accurate, at a few angles) and (1974BU06: $E_d = 15$ MeV; d_0). Inelastic deuteron groups are discussed in Table 14.27. The deuteron group to the 0^+ ; $T = 1$ state $^{14}\text{N}^*(2.31)$ is isospin forbidden: its intensity is 1 – 3% of the deuteron group to $^{14}\text{N}^*(3.95)$ for $E_d = 7.1$ to 10.2 MeV (1970DU04). The deuteron group to $^{14}\text{N}^*(8.06)$ [$T = 1$] is not observed: see Table 14.27. See also (1970AJ04), ^{16}O in (1971AJ02, 1977AJ02), (1970PI1A) and (1970ME25, 1971SI24, 1972DM01, 1972FO07, 1973FA1J, 1975GU10; theor.).

46. $^{14}\text{N}(\text{t}, \text{t})^{14}\text{N}$

Elastic differential cross sections have been measured at four angles at $E_t = 20.03$ MeV (1974JA25). See also (1973FA1J; theor.).

Table 14.27: $^{14}\text{N}^*$ levels from $^{14}\text{N}(\text{p}, \text{p}')$, (d, d') , $(^3\text{He}, ^3\text{He}')$ and (α, α')

$^{14}\text{N}^*(\text{MeV} \pm \text{keV})$				L^k	Dominant config. ⁿ	$J^\pi; T$
(p, p')	(d, d')	(^3He , $^3\text{He}'$)	(α , α')			
$2.313 \pm 5^{\text{a}}$	see text	h	see text		$(p_{\frac{1}{2}})^2$	$0^+; 1$
$3.945 \pm 5^{\text{a}}$	f	h	j	2	$(p_{\frac{1}{2}})^2 + \text{c.e.}^{\text{o}}$	$1^+; 0$
$4.910 \pm 10^{\text{a}}$	f	h	j	1, 3	$p_{\frac{1}{2}} s_{\frac{1}{2}}$	$0^-; 0$
$5.104 \pm 10^{\text{a}}$	f	h	j	1, 3	$p_{\frac{1}{2}} d_{\frac{5}{2}}$	$2^-; 0$
$5.69 \pm 30^{\text{b}}$	f	h	j	1, 3	$p_{\frac{1}{2}} s_{\frac{1}{2}}$	$1^-; 0$
$5.83 \pm 30^{\text{b}}$	f	h	j	1, 3	$p_{\frac{1}{2}} d_{\frac{5}{2}}$	$3^-; 0$
$6.23 \pm 20^{\text{b}}$	f	h	j,l		$(s_{\frac{1}{2}})^2$	$1^+; 0$
$6.46 \pm 20^{\text{b}}$	f		j,l		$s_{\frac{1}{2}} d_{\frac{5}{2}}$	$3^+; 0$
$7.03 \pm 20^{\text{b}}$	f	h	j	2	c.e. ^o	$2^+; 0$
7.97 ^c			m		$p_{\frac{1}{2}} d_{\frac{3}{2}}$	$2^-; 0$
8.06 ^c		$8.0 \rightarrow 11.0^{\text{i}}$			$p_{\frac{1}{2}} s_{\frac{1}{2}}$	$1^-; 1$
8.49 ⁿ						$(4^-); 1$
8.62 ⁿ					$(s_{\frac{1}{2}})^2$	$0^+; 1$
8.91 ⁿ					$p_{\frac{1}{2}} d_{\frac{5}{2}}$	$3^-; 0$
9.17 ^{d,n}					c.e. ^o	$2^+; 1$
9.39 ⁿ						$2^-, 3^-; 0$
9.51 ⁿ					$p_{\frac{1}{2}} d_{\frac{5}{2}}$	$1^+; 0$
9.70 ⁿ					$(p_{\frac{1}{2}})^2$	$1^+; 0$
10.1 ⁿ					$s_{\frac{1}{2}} d_{\frac{5}{2}}$	$2^+, 1^+; 0$
$11.2 \pm 200^{\text{e}}$	f,g	$11.22 \pm 50^{\text{h}}$				
$12.8 \pm 400^{\text{c}}$	f,g	$12.77 \pm 50^{\text{h}}$				
17 ^d						
21.5 ^d						

^a (1953BO70).

^b (1956BU16). See also text, (1964EA04, 1971CU01), and Table 14.26.

^c (1964EA04, 1971CU01).

^d (1958TY46).

^e (1965DE21).

^f Observed: see (1970AJ04).

^g $E_x = 11.3$ and 13 MeV (1968HI14).

^h (1969BA06).

ⁱ Unresolved structure (1969BA06).

^j (1966HA19).

^k From (α, α') : (1966HA19).

^l Relatively low cross section due to two-nucleon transition (1966HA19).

^m See also (1956MI17).

ⁿ (1971CU01): p, p'.

^o c.e. = compound elastic.

47. $^{14}\text{N}(\text{He}, \text{He}')^{14}\text{N}^*$

Angular distributions of elastically and inelastically scattered ^3He ions have been measured at $E(^3\text{He})$ up to 44.6 MeV: see Table 14.23 in (1970AJ04) and (1970KN01: 4.5, 7.0 MeV; elastic).

At $E(^3\text{He}) = 44.6$ MeV, twelve ^3He groups are reported corresponding to states in ^{14}N : see Table 14.27 (1969BA06). The angular distributions were analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture. A comparison of the cross sections of the reactions $^{14}\text{N}(^3\text{He}, t)^{14}\text{O}_{\text{g.s.}}$, $^{14}\text{N}(^3\text{He}, ^3\text{He}')^{14}\text{N}^*(2.31)$ and $^{14}\text{C}(^3\text{He}, t)^{14}\text{N}(0)$ [which all correspond to transitions between identical initial and final states] shows that they are roughly equal, as would be expected from charge independence, once detailed-balance, isospin coupling and phase-space corrections have been applied (1969BA06). See also (1973FA1J; theor.).

48. (a) $^{14}\text{N}(\alpha, \alpha')^{14}\text{N}^*$

$$(\text{b}) \quad ^{14}\text{N}(\alpha, \alpha p)^{13}\text{C} \qquad Q_m = -7.5507$$

$$(\text{c}) \quad ^{14}\text{N}(\alpha, \alpha d)^{12}\text{C} \qquad Q_m = -10.2725$$

Angular distributions of elastically and inelastically scattered α -particles have been measured for $E_\alpha = 11$ to 104 MeV: see Table 14.23 in (1970AJ04) for the earlier references and (1969FE10, 1970AG08: $E_\alpha = 19.8 - 23.1$ MeV; α_0), (1970GA25, 1971GA44: 21.7 MeV; $\alpha_0, \alpha_2, \alpha_{3+4}, \alpha_{5+6}, \alpha_{7+8}, \alpha_9$), (1972KU19: 26.6 MeV; α_0), (1972LO08: 29.98 MeV; α_0) and (1970HA1G: 104 MeV; α_0). See also (1972OE01, 1974YOZO). See Table 14.27 for the observed α -groups. Generally the

intensity of the α_1 group is weak, typically a few percent of the α_0 or α_2 groups: see (1959AJ76, 1966HA19). See also (1970CH1D).


Reduced transition probabilities are reported by (1966HA19): $B(E2)\downarrow/e^2 = 6.5$ and 3.3 fm^4 , respectively for $^{14}\text{N}^*(3.95, 7.03)$; $B(E3)\downarrow/e^2 = 40$ and 60 fm^6 , respectively for $^{14}\text{N}^*(5.11, 5.83)$. See also (1970AJ04) and ^{18}F in (1972AJ02, 1978AJ03). See also (1974HA1C) and (1971MU1H, 1972DM01, 1972RA34, 1973FA1J; theor.).



The angular distribution involving $^{6}\text{Li}_{\text{g.s.}} + ^{14}\text{N}_{\text{g.s.}}$ has been measured at $E(^{6}\text{Li}) = 32 \text{ MeV}$ by (1971GR44, 1971ZE1C). See also (1974FL1A).



See ^{9}Be in (1974AJ01).





The elastic angular distribution has been measured at $E(^{10}\text{B}) = 100 \text{ MeV}$ (1975NA15). For reaction (a) see also (1970IS1A). The elastic scattering angular distributions (reaction (b)) have been studied at $E(^{14}\text{N}) = 41, 77$ and 113 MeV (1971LI11).



Elastic angular distributions have been measured at $E(^{14}\text{N}) = 21.3 \text{ MeV}$ (1971VO01), 22.5 MeV (1969HE06), $65, 84$ and 88 MeV (1971KO11, 1974KO38, 1975KO06), 78 MeV (1970VO02: and $^{12}\text{C}^*(4.4)$), 118 MeV (1974AN36: and $^{12}\text{C}^*(4.4)$), and 155 MeV (1974BI1E, 1975NA11, 1975NA15: and various ^{12}C states). See also (1975ZE1C) and (1975MO23, 1975RA33, 1975VO1B; theor.). At $E(^{14}\text{N}) = 155 \text{ MeV}$ the selective population of certain ^{14}N states is observed and angular distributions are reported for the transitions to $^{14}\text{N}^*(0, 8.96, 12.7)$ (1975NA11).

For yield measurements see (1974JO1J, 1974ST1N, 1975ZE1C), (1972SC1L), (1972MA74, 1974GA1L, 1975DE09; theor.) and ^{12}C in (1975AJ02).



The elastic angular distribution has been measured at $E(^{14}\text{N}) = 19.3$ MeV ([1971VO01](#)). See also ([1973MC1J](#), [1973VO04](#), [1975DE09](#); theor.). For reaction (b) see ([1971VO1D](#), [1975DE09](#), [1975VO1B](#); theor.).



Elastic angular distributions have been studied for $E(^{14}\text{N}) = 4.99$ to 20.22 MeV ([1969JA15](#)). For yield measurements see ([1969JA15](#), [1973GO01](#), [1973RE13](#), [1973ST1L](#), [1974ST1N](#)). See also ([1973ST1A](#)) and ([1972PA31](#), [1973KA46](#), [1973PA1L](#), [1973PA1M](#), [1973SC1K](#), [1974VE05](#); theor.).



Elastic angular distributions have been measured for $E(^{14}\text{N}) = 8.08$ to 18.05 MeV ([1969JA15](#)), 25 MeV ([1971BO1U](#)), 79 MeV ([1976MO03](#)) and 155 MeV ([1975NA15](#)). For yield measurements see ([1969JA15](#), [1970SI09](#), [1973GO01](#), [1974ST1N](#)). See also ([1973KA46](#), [1974BA40](#), [1975MO23](#), [1975VO1B](#); theor.).



The decay proceeds almost entirely to the $J^\pi = 0^+$; $T = 1$ state of ^{14}N at 2.31 MeV: see ^{14}O . Measurement of the γ -ray energy from the decay of $^{14}\text{N}^*(2.31)$ leads to $E_x = 2312.87 \pm 0.10$ keV ([1968FR08](#)), 2312.89 ± 0.10 keV ([1967CH19](#)). See also reaction 1 in ^{14}O .



Angular distributions have been obtained for the deuterons corresponding to $^{14}\text{N}^*(0, 2.31, 3.95)$ ([1961BE12](#): $E_p = 18.6$ MeV) and to $^{14}\text{N}^*(0 - 8.06, 8.62, 8.91, 8.96 + 8.98, 9.17 - 10.43, 10.81, 11.05, 11.24 + 11.29, 11.36 - 11.66, 11.75, 11.95, 12.20, 12.50, 12.61, 12.79 + 12.82, 13.17 + 13.24, 13.71 + 13.72)$ ([1969SN04](#), [1969SN1B](#): $E_p = 39.8$ MeV). Spectroscopic factors were extracted by DWBA analysis of the $l_n = 1$ pickup angular distributions. $\Gamma = 210 \pm 30$ keV for $^{14}\text{N}^*(13.71)$. Weak deuteron groups to ^{14}N states at $E_x = 6.70, 7.40$ and 7.60 MeV are reported [see, however, Table 14.11] ([1969SN04](#)). See also ([1972PA1A](#)).



Not reported.



Observed states in ^{14}N are displayed in Table 14.28 together with the derived spectroscopic factors ([1966BA13](#), [1970BO21](#)). See also ([1962CL12](#), [1969HO23](#), [1974HO1M](#)).



At $E(^{15}\text{N}) = 30, 32$ and 45 MeV the angular distributions involving $^{14}\text{N}^*(0, 2.31)$ have been studied: they are symmetric about 90° for the transition to the $T = 1$ analog state $^{14}\text{N}^*(2.31)$ ([1975GA17](#)).




At $E_{\pi^-} = 230$ MeV, γ -rays from the population of $^{14}\text{N}^*(2.31, 3.95, 5.11, 5.83)$ are observed ([1974LI15](#)). See also ([1971KO23](#)).



Not reported.



See ([1972BO17](#)) and ([1974GO1G](#); theor.).



Angular distributions have been measured at $E_p = 27$ MeV ([1973IN05](#): to $^{14}\text{N}^*(2.31)$), $32.2, 36.6, 43.5$ MeV ([1975MI01](#): to $^{14}\text{N}^*(0, 2.31, 3.95)$), 39.8 MeV ([1973HO10](#): to $^{14}\text{N}^*(0, 2.31, 3.95)$); $L = 0 + 2, 0, 0 + 2$, respectively), 43.7 and 54.1 MeV ([1964CE02](#), [1970HA23](#), [1971FL04](#), [1974MA12](#): to $^{14}\text{N}^*(0, 2.31, 3.95, 5.11, 7.03, 9.17, 11.05)$; some with pol. p [also observed $^{14}\text{N}^*(10.43, 12.50)$] and 49.5 MeV ([1970NE1B](#), [1970NE17](#): pol. p to $^{14}\text{N}^*(0, 2.31)$]). A number of comparisons have been made of the ratio of ($\text{p}, ^{3}\text{He}$) to the $T = 1$ state at 2.31 MeV and of (p, t) to the analog ground state of ^{14}O : see reaction 10 in ^{14}O . See also ([1969SC1F](#), [1972HA1X](#), [1974AD1B](#)), ([1974ST16](#); theor.) and ^{17}F in ([1977AJ02](#)).

Table 14.28: States of ^{14}N from $^{15}\text{N}(^3\text{He}, \alpha)^{14}\text{N}$ ^a

$^{14}\text{N}^*$ (MeV \pm keV)	$J^\pi; T$	l ^a	S ^b	S ^c
0	$1^+; 0$	1	1.20	1.50
2.31	$0^+; 1$	1	1.50	2.00
3.95	$1^+; 0$	1	0.60	0.73
4.92 ^d	$0^-; 0$			
5.11	$2^-; 0$	2	0.06	
5.691 ± 8 ^e	$1^-; 0$	2	0.02	
5.83	$3^-; 0$	2	0.03	
6.20	$1^+; 0$	1	0.05	
6.44	$3^+; 0$	3	0.01	
7.032 ± 10 ^e	$2^+; 0$	1	1.03	1.00
7.97 ^d	$2^-; 0$			
8.06 ^d	$1^-; 1$			
8.62 ^d	$0^+; 1$			
8.91	$3^-; 1$	(4)	(5.5×10^{-2})	
9.17	$2^+; 1$	1	1.90	1.36
9.39 ^d	$2^-, 3^-; 0$			
9.51 ^d	$2^-, 1$			
9.70 ^d	$1^+, 1$			
10.43	$2^+; 1$	1	1.04	1.10
13.75 ^f	$1^+; 1$	1	(2.4 ± 1.0)	1.4

^a See also (1962CL12, 1974HO1M).

^b (1970BO21): $E(^3\text{He}) = 11.0$ MeV; see also (1969HO23).

^c (1966BA13): $E(^3\text{He}) = 39.8$ MeV.

^d (1969HO23).

^e (1962CL12).

^f 13.72 ± 0.04 MeV (1966BA13).



Angular distributions of α -particles have been measured at many energies up to $E_d = 40$ MeV: see Table 14.25 in (1970AJ04) for the earlier measurements and (1971GR2B: 0.9 – 2.0 MeV; α_0), (1970CA1C: 1.0 – 2.0 MeV; α_0), (1973JO13: 2 to 14 MeV; α_0, α_1), (1971TH03: 4.405, 4.578, 4.710 MeV; α_0), (1969AL13: 10.6 to 12.6 MeV; α_0), (1971JA04: 14.6, 15.0, 15.8 MeV; α_1) and (1974VAZK, 1975VA1H: 40 MeV; α_0, α_2). Alpha groups have been seen corresponding to most known states of ^{14}N with $E_x \leq 11.5$ MeV (1968JO07: $E_d = 5$ to 9 MeV) and (1974VAZK) observe $^{14}\text{N}^*(11.05)$.

The yield of the isospin forbidden α_1 group [to $^{14}\text{N}^*(2.31)$] has been studied for $E_d = 2$ to 15 MeV by (1969JO09, 1973JO13): the intensity of the isospin group is strongly dependent on E_d and on the angle of observation. The α_1 reaction appears to proceed almost exclusively by a compound nuclear process and its study leads to the determination of a large number of ^{18}F states: the average isospin impurity in ^{18}F for $10 \leq E_x \leq 20$ MeV is 3 – 10% (1973JO13). For further discussions see (1969DE29, 1971JA04) and (1970AJ04).

Measurements on the absolute cross sections of this reaction [$E_d = 3.6$ to 5.3 MeV] and its inverse [$^{14}\text{N}(\alpha, \text{d})^{16}\text{O}$] are consistent with the principle of detailed balance. An upper limit of 0.2% is assigned to the time-reversal non-invariant part of the reaction amplitudes (1971TH03).

See also (1967SP09), (1970FO1B; theor.) and ^{18}F in (1978AJ03).



At $E_\alpha = 42$ MeV the transitions involving ($^{14}\text{N}_{\text{g.s.}}$ and ${}^6\text{Li}^*(0, 3.56)$), ($^{14}\text{N}^*(2.31) + {}^6\text{Li}_{\text{g.s.}}$) and ($^{14}\text{N}^*(3.95) + {}^6\text{Li}_{\text{g.s.}}$) have been studied by (1972RU03).



See (1974MI1F).



See (1965SH10, 1965SH14).



See (1972KU1H; theor.).



See ([1974RO04](#)).



See ([1973SE02](#)) and ([1975RO20](#); astrophys. questions) in ^{18}F ([1978AJ03](#)).



See ([1967DE03](#)).

¹⁴O
(Figs. 8 and 9)

GENERAL: (See also (1970AJ04).)

Nuclear models: (1973SA30, 1974KU1F).

Special reactions involving ¹⁴O: (1971AR02, 1973BA81, 1974BA89, 1975HU14).

Reactions involving pions: (1973CH20, 1973DA37, 1975HU1D, 1975RE01).


Other topics: (1970FO1B, 1972AN05, 1972CA37, 1972KU1C, 1973GO1H, 1973KA1H, 1973PA1F, 1973RO1R, 1973SP1A, 1974BO22, 1974KU1F, 1974SE1B, 1974VA24, 1975BU1M).

Ground state: (1975BE31).



The adopted half-life of ¹⁴O is 70.599 ± 0.022 sec (1975HA45). Recent measured values are $\tau_{1/2} = 70.588 \pm 0.028$ sec (1971CL07, 1973CL12), 70.48 ± 0.15 sec (1972AL01), 70.32 ± 0.12 sec (1972SI50), 70.43 ± 0.18 sec (1974AZ01). ¹⁴O [$J^\pi = 0^+$; $T = 1$] decays predominantly to the analog state ¹⁴N*(2.31): the branching ratio to that state is $99.328 \pm 0.012\%$. This value is obtained by adopting $0.61 \pm 0.01\%$ (1966SI05) and $0.062 \pm 0.007\%$ (1969KA1B) for the branching ratios to ¹⁴N*(0, 3.95) [both 1^+ ; 0 states]. The ft value for the $0^+ \rightarrow 0^+$ transition (to ¹⁴N*(2.31)), based on the 99.328% branching ratio, the adopted $\tau_{1/2}$, a Q for the transition of 2831.83 ± 0.36 keV, is 3084.3 ± 3.6 sec (1975HA45). See also (1973CL12, 1973HA2E).

The question of how ft values for $0^+ \rightarrow 0^+$ transitions should be calculated has been studied by many other groups: see, e.g., (1972JA19, 1972WI05, 1973TO04, 1974RO21, 1974WI02; theor.). The importance of the question relates to the determination of the fundamental coupling constant G_V ; see the discussion in (1975HA45). See also (1970AJ04). The ft values for the transitions to ¹⁴N*(0, 3.95) are, respectively, $(1.9 \pm 0.2) \times 10^7$ (1969KA1B) and 1200 ± 150 (1969KA1B). For a summary of earlier values see Table 14.27 in (1970AJ04). See also (1975RA37), (1969BL1D, 1969LY1A, 1970HU08, 1970JA02, 1970LA1L, 1970PA23, 1971FA17, 1971HU10, 1971JA20, 1972AR20, 1972BE04, 1972WI28, 1972WI1C, 1973HA49, 1973WI1J, 1974AS07, 1974BO18, 1974KR09, 1975WI1K, 1975WI1E, 1976LO01; theor.).



[See also Table 14.28 in (1970AJ04).]

Table 14.29: Energy levels of ^{14}O

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ (sec) or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$0^+; 1$	$\tau_{1/2} = 70.599 \pm 0.022$ sec	β^+	1, 2, 4, 6, 7, 8, 9, 10, 11, 12
5.173 ± 10	$1^-; 1$			2, 5, 8, 10
5.920 ± 10	$0^+; 1$	$\Gamma \leq 50$ keV	p	2, 5, 8, 10
6.272 ± 10	$3^-; 1$	103 ± 6	p	2, 3, 4, 5, 8, 10
6.590 ± 10	$2^+; 1$	≤ 60	p	2, 8, 10
(6.79 ± 30)	$\pi = -$			8
7.768 ± 10	$2^+; 1$	76 ± 10	p	2, 8, 10
(8.72 ± 40)				5, 8, 10
9.715 ± 20	$(2^+); 1$			2, 5, 8, 10
9.915 ± 20	$4^+; 1$	100 ± 50		2, 3, 4, 5
10.89 ± 50				8
11.24 ± 50				8
11.97^{a}				8
12.84 ± 50				8
13.01 ± 50				8
14.15 ± 40				8
14.64 ± 60				8
17.40 ± 60				8

^a Possibly more than one level.

Observed neutron groups are displayed in Table 14.30. Angular distributions have been studied at many energies: among the recent measurements are those by (1970AD01, 1970AD02: $E(^3\text{He}) = 10$ to 15 MeV), (1972BR60: 13.3 and 18.2 MeV) and (1971GRYL, 1972GR39: 25.4 MeV). For earlier measurements see (1970AJ04). A study of n-p coincidences ($^{14}\text{O}^* \rightarrow ^{13}\text{N} + p$) leads to J and Γ assignments for $^{14}\text{O}^*(5.92, 6.27, 6.59, 7.77)$ (1973PR08): see Table 14.30. See also (1971KL1C, 1973JU1B, 1973RH1A, 1974AD11, 1974GO23), (1974GR39; theor.) and ^{15}O .



At $E(^{11}\text{B}) = 114$ MeV, $^{14}\text{O}^*(6.27, 9.9)$ [$J^\pi = 3^-$ and 4^+ , respectively] are populated: see (1973SC1J, 1974AN36).

Table 14.30: Levels of ^{14}O from $^{12}\text{C}(^{3}\text{He}, \text{n})^{14}\text{O}$

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)		L^{c}	J^π ^c
(1961TO03, 1968TO09)	(1972GR39)	(1961TO03)	(1973PR08)	
0	0			0
5.17	5.173 ± 10			1
5.905 ± 12	5.930 ± 15	≤ 60	≤ 47	0
6.30 ± 30	6.272 ± 10	120	103 ± 6	3
$6.586 \pm 12^{\text{a}}$	6.596 ± 10	≤ 60	≤ 56	(2)
	7.768 ± 10		76 ± 10	2
	9.705 ± 25			(2)
	$9.915 \pm 20^{\text{b}}$	$100 \pm 50^{\text{b}}$		4
				$(2^+)^{\text{d}}$
				(2^+)
				4^+

^a (1970AD01) report $E_x = 6.585 \pm 0.005$ MeV.

^b (1972BR60) report $E_x = 9.95 \pm 0.043$ MeV; $J^\pi = 3^-$.

^c See (1968TO09, 1970AD01, 1970AD02, 1972BR60, 1972GR39).

^d $J = 2$ follows from the n-p coincidence study of (1973PR08). The J shown for $^{14}\text{O}^*(5.91, 6.29, 7.78)$ are in accord with this work.

$$4. \ ^{12}\text{C}(^{12}\text{C}, ^{10}\text{Be})^{14}\text{O} \quad Q_m = -20.617$$

At $E(^{12}\text{C}) = 114$ MeV, the population of $^{14}\text{O}^*(0, 6.27, 9.9)$ is reported: it is suggested that $^{14}\text{O}^*(9.9)$ has a $(\text{d}_{5/2})_{4+,1}^2$ configuration (1971SC1F, 1972SC21, 1974AN36). $^{14}\text{O}_{\text{g.s.}}$ is weakly populated (1974AN36).

$$5. \ ^{12}\text{C}(^{14}\text{N}, ^{12}\text{B})^{14}\text{O} \quad Q_m = -18.515$$

At $E(^{14}\text{N}) = 118$ MeV, the population of $^{14}\text{O}^*(6.27, 9.9)$ is reported (1974AN36). At $E(^{14}\text{N}) = 155$ MeV there is some evidence for the excitation of $^{14}\text{O}^*(5.17, 5.92, 6.27, 8.74, 9.72)$ (1975NA11).

$$6. \ ^{13}\text{C}(\text{p}, \pi^-)^{14}\text{O} \quad Q_m = -137.163$$

At $E_p = 185$ MeV, an angular distribution has been obtained for the transition to $^{14}\text{O}_{\text{g.s.}}$: in the range $35^\circ - 130^\circ$ it shows no peaking (1973DA37).




See (1970AJ04) for the earlier work. See also (1970WI1B, 1971BE46, 1971TH1D) and (1970LO1B, theor.). [The earlier reported angular distributions have not been published.]



Triton groups have been observed at $E(^3\text{He}) = 44.6$ MeV to the first six states shown in Table 14.29 and to levels with $E_x = 6.79 \pm 0.03, 8.74 \pm 0.06, 9.74 \pm 0.03, 10.89 \pm 0.05, 11.24 \pm 0.05, 11.97$ (unresolved), $12.84 \pm 0.05, 13.01 \pm 0.05, 14.15 \pm 0.04, 14.64 \pm 0.06$ and 17.40 ± 0.06 MeV (1967BA13, 1969BA06). See also (1968BA1E) and reaction 47 in ^{14}N . [The states at 6.79 and 8.74 MeV reported in this reaction are relatively weakly excited and are not observed in reaction 2.]



See (1970AJ04).



Angular distributions of ground state tritons have been studied at $E_p = 27$ MeV (1973IN05), 32.2, 36.6 and 43.5 MeV (1975MI01), 43.7 and 54.1 MeV (1964CE02, 1971FL04), 43.8 MeV (1970HA23: pol. p) and 49.5 MeV (1970NE17, 1970NE1B: pol. p). Angular distributions have also been studied with polarized protons at $E_p = 43.8$ MeV to $^{14}\text{O}^*(0, 5.17, 6.27, 6.59, 7.77, 9.72)$ (1974MA12). The ratio of (p, ^3He) [to the analog state in ^{14}N at 2.31 MeV] to (p, t) [to $^{14}\text{O}_{\text{g.s.}}$] is correctly predicted by DWBA calculations: see, e.g., (1974CR01).

Triton groups have also been observed to states with $E_x = 5.21 \pm 0.04, 5.92 \pm 0.06, 6.28 \pm 0.05, 6.59, 7.78, 8.69 \pm 0.06$ [weak, not observed in reaction 2], and 9.65 ± 0.06 MeV (1971FL04) and to $^{14}\text{O}^*(6.59, 7.77, 9.71)$ (1970HA23). The J^π assignments derived from these two papers and from (1974MA12, 1975KU22) are in agreement with the values shown in Table 14.29. See also (1969SC1F, 1972HA1X, 1974AD1B), (1971KA04, 1974ST16; theor.) and ^{17}F in (1977AJ02).



At $E_\alpha = 58$ MeV the differential cross section for populating $^{14}\text{O}_{\text{g.s.}} \lesssim 0.4 \mu\text{b}$ at $\theta_{\text{lab}} = 21^\circ$ (1975VA01).

12. $^{18}\text{O}(\alpha, {}^8\text{He})^{14}\text{O}$

$$Q_m = -38.016$$

At $E_\alpha = 58$ MeV the differential cross section is 40 ± 15 nb/sr at $\theta_{\text{lab}} = 8^\circ$ ([1975JA10](#)).

^{14}F
(Not illustrated)

^{14}F has not been observed: its atomic mass excess is predicted to be 33.38 MeV which would make it unstable with respect to decay into ${}^{13}\text{O} + \text{p}$ by ≈ 3 MeV ([1973BA34](#)).

References

(Closed 31 January 1976)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1953BO70 C.K. Bockelman, C.P. Browne, W.W. Buechner and A. Sperduto, Phys. Rev. 92 (1953) 665
- 1953MA42 G. Manning and B. Singh, Proc. Phys. Soc. A66 (1953) 842
- 1953SH64 E.S. Shire, J.R. Wormald, G. Lindsay-Jones, A. Lundan and A.G. Stanley, Phil. Mag. 44 (1953) 1197
- 1954MO84 A. Moljk and S.C. Curran, Phys. Rev. 96 (1954) 395
- 1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, Phys. Rev. 96 (1954) 1316
- 1955AL16 H.R. Allan and N. Sarma, Proc. Phys. Soc. A68 (1955) 535
- 1955MC75 J.N. McGruer, E.K. Warburton and R.S. Bender, Phys. Rev. 100 (1955) 235
- 1955WI43 D.H. Wilkinson, Phys. Rev. 100 (1955) 32
- 1956BO61 T.W. Bonner, A.A. Kraus, Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956BU16 E.J. Burge and D.J. Prowse, Phil. Mag. 1 (1956) 912
- 1956GR17 G.M. Griffiths, Can. J. Phys. 34 (1956) 339
- 1956LE28 P. Lehmann, A. Leveque and R. Pick, Compt. Rend. 243 (1956) 743
- 1956MI17 D.W. Miller, B.M. Carmichael, U.C. Gupta, V.K. Rasmussen and M.B. Sampson, Phys. Rev. 101 (1956) 740
- 1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434
- 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1957WI27 D.H. Wilkinson and S.D. Bloom, Phil. Mag. 2 (1957) 63
- 1958GR97 G.M. Griffiths, Proc. Phys. Soc. A72 (1958) 337
- 1958TY46 H. Tyren and T.A.J. Maris, Nucl. Phys. 6 (1958) 82
- 1958WA02 E.K. Warburton and H.J. Rose, Phys. Rev. 109 (1958) 1199
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AR1A Armstrong, Moore and Blair, Bull. Amer. Phys. Soc. 4 (1959) 17

- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959HA11 S.S. Hanna and L. Meyer-Schutzmeister, Phys. Rev. 115 (1959) 986
- 1959HO01 H.D. Holmgren, E.A. Wolicki and R.L. Johnston, Phys. Rev. 114 (1959) 1281
- 1959RO54 H.J. Rose, W. Trost and F. Riess, Nucl. Phys. 12 (1959) 510
- 1959WA04 E.K. Warburton, H.J. Rose and E.N. Hatch, Phys. Rev. 114 (1959) 214
- 1959WA16 E.K. Warburton, Phys. Rev. 113 (1959) 595
- 1960BA34 J.K. Bair, R.D. Edge and H.B. Willard, Phys. Rev. 119 (1960) 1948
- 1960HE14 D.F. Hebbard and J.L. Vogl, Nucl. Phys. 21 (1960) 652
- 1960JA17 A.A. Jaffe, F. de S. Barros, P.D. Forsyth, J. Muto, I.J. Taylor and S. Ramavataram, Prog. Phys. Soc. 76 (1960) 914
- 1960RO13 H.J. Rose, Nucl. Phys. 19 (1960) 113
- 1960RO23 H.J. Rose, F. Riess and W. Trost, Nucl. Phys. 21 (1960) 367
- 1960WA12 E.K. Warburton and W.T. Pinkston, Phys. Rev. 118 (1960) 733
- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. 77 (1961) 853
- 1961BE12 E.F. Bennett, Phys. Rev. 122 (1961) 595
- 1961CO05 H.O. Cohn, J.K. Bair and H.B. Willard, Phys. Rev. 122 (1961) 534
- 1961DA09 P. Dagley, W. Haeberli and J.X. Saladin, Nucl. Phys. 24 (1961) 353
- 1961HE02 R.E. Hester and W.A.S. Lamb, Phys. Rev. 121 (1961) 584
- 1961KA04 E. Kashy, R.R. Perry, R.L. Steele and J.R. Risser, Phys. Rev. 122 (1961) 884
- 1961MA1L W.B. Mann, W.F. Marlow and E.E. Hughes, National Bureau of Standards (U.S.) Tech. News Bull. 45 (1961) 21
- 1961MA32 W.B. Mann, W.F. Marlow and E.E. Hughes, Int. J. Appl. Rad. Isotopes 11 (1961) 57
- 1961TO03 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. 77 (1961) 399
- 1961WA16 D.E. Watt, D. Ramsden and H.W. Wilson, Int. J. Appl. Rad. Isotopes 11 (1961) 68; Nucl. Sci. Abs. 15, 4221, Abs. 32740 (1961)
- 1962CL12 D.D. Clayton, Phys. Rev. 128 (1962) 2254
- 1962ED02 R.D. Edge and G.A. Peterson, Phys. Rev. 128 (1962) 2750
- 1962HA40 B.G. Harvey, J. Cerny, R.H. Pehl and E. Rivet, Nucl. Phys. 39 (1962) 160
- 1962LA15 N.O. Lassen, Nucl. Phys. 38 (1962) 442
- 1962OL04 I.U. Olsson, I. Karlen, A.H. Turnbull and N.J.D. Prosser, Ark. Fys. 22 (1962) 237
- 1962RO21 H.J. Rose, F. Uihlein, F. Riess and W. Trost, Nucl. Phys. 36 (1962) 583
- 1963AL21 D.E. Alburger and E.K. Warburton, Phys. Rev. 132 (1963) 790

- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963JE02 J.M.F. Jeronymo, G.S. Mani, F. Picard and A. Sadeghi, Nucl. Phys. 43 (1963) 417
- 1963MA28 G.S. Mani, P.D. Forsyth and R.R. Perry, Nucl. Phys. 44 (1963) 625
- 1963OH02 G.G. Ohlsen and R.E. Shamu, Nucl. Phys. 45 (1963) 523
- 1963PR03 F.W. Prosser, Jr., R.W. Krone and J.J. Singh, Phys. Rev. 129 (1963) 1716
- 1963RO17 H.J. Rose, W. Trost and F. Riess, Nucl. Phys. 44 (1963) 287
- 1963SE19 T. Sebe, Prog. Theor. Phys. 30 (1963) 290
- 1964BA11 M.R. Baker, C.H. Anderson and N.F. Ramsey, Phys. Rev. 133 (1964) A1533
- 1964BE12 J.A. Becker and E.K. Warburton, Phys. Rev. 134 (1964) B349
- 1964BI09 G.R. Bishop, M. Bernheim and P. Kossanyi-Demay, Nucl. Phys. 54 (1964) 353
- 1964CE02 J. Cerny and R.H. Pehl, Phys. Rev. Lett. 12 (1964) 619
- 1964DZ1A Dzubay and Blair, Phys. Rev. 134 (1964) B586
- 1964EA04 L.G. Earwaker and D.F. Hebbard, Nucl. Phys. 53 (1964) 252
- 1964HU09 E.E. Hughes and W.B. Mann, Int. J. App. Rad. Isotopes 15 (1964) 97; Nucl. Sci. Abs. 18, 2271, Abs.16961 (1964)
- 1964MI05 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 63
- 1964RO03 H.J. Rose, F. Riess and W. Trost, Nucl. Phys. 52 (1964) 481
- 1964WA05 E.K. Warburton, D.E. Alburger, A. Gallmann, P. Wagner and L.F. Chase, Jr., Phys. Rev. 133 (1964) B42
- 1964WA09 E.K. Warburton, J.W. Olness, D.E. Alburger, D.J. Bredin and L.F. Chase, Jr., Phys. Rev. 134 (1964) B338
- 1965BA06 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klubes, B. Kuhn and G. Stiller, Nucl. Phys. 64 (1965) 55
- 1965BR42 O.D. Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965CA06 R.R. Carlson and D.W. Heikkinen, Phys. Lett. 17 (1965) 305
- 1965DE19 R.W. Detenbeck, J.C. Armstrong, A.S. Figuera and J.B. Marion, Nucl. Phys. 72 (1965) 552
- 1965DE21 R.W. Detenbeck, Nucl. Phys. 74 (1965) 184
- 1965FO06 P.D. Forsyth, I.J. Taylor and R.R. Perry, Nucl. Phys. 66 (1965) 376
- 1965LA09 J.M. Lacambra, D.R. Tilley, N.R. Roberson and R.M. Williamson, Nucl. Phys. 68 (1965) 273
- 1965LO07 J.A. Lonergan and D.J. Donahue, Nucl. Phys. 74 (1965) 318
- 1965LU05 A. Luukko, Soc. Sci. Fennica, Comment. Phys. Math. 31:6 (1965)

- 1965NE06 J.O. Newton, R.S. Blake, D.J. Jacobs and J.P. Schapira, Nucl. Phys. 71 (1965) 113
- 1965PE03 R.H. Pehl, E. Rivet, J. Cerny and B.G. Harvey, Phys. Rev. 137 (1965) B114
- 1965SA07 M.W. Sachs, C. Chasman and D.A. Bromley, Phys. Rev. 139 (1965) B92
- 1965SH10 S.N. Shumilov, A.P. Klyucharev and N.Y. Rutkevich, Pisma Zh. Eksp. Teor. Fiz. 2 (1965) 213; JETP Lett. 2 (1965) 135
- 1965SH14 S.N. Shumilov, A.P. Klyucharev and N.Y. Rutkevich, Zh. Eksp. Teor. Fiz. 49 (1965) 1754; Sov. Phys. JETP 22 (1966) 1198
- 1965VA13 P.I. Vatset, L.Y. Kolesnikov and S.G. Tonapetyan, Yad. Fiz. 1 (1965) 809; Sov. J. Nucl. Phys. 1 (1965) 579
- 1965WA02 B.D. Walker, C. Wong, J.D. Anderson, J.W. McClure and R.W. Bauer, Phys. Rev. 137 (1964) B347
- 1965WA06 E.K. Warburton, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and M.F. Thomas, Phys. Rev. 138 (1965) B104
- 1966AL10 D.E. Alburger, A. Gallmann, J.B. Nelson, J.T. Sample and E.K. Warburton, Phys. Rev. 148 (1966) 1050
- 1966BA13 G.C. Ball and J. Cerny, Phys. Lett. 21 (1966) 551
- 1966CA07 R.R. Carlson, Phys. Rev. 148 (1966) 991
- 1966DI04 G.U. Din and J.L. Weil, Nucl. Phys. 86 (1966) 509
- 1966FU10 H. Fuchs, K. Grabisch, P. Kraaz and G. Roschert, Phys. Lett. 23 (1966) 363
- 1966GO15 S. Gorodetzky, R.M. Freeman, A. Gallman and F. Hass, Phys. Rev. 149 (1966) 801
- 1966HA19 B.G. Harvey, J.R. Meriwether, J. Mahoney, A. Bussiere de Nercy and D.J. Horen, Phys. Rev. 146 (1966) 712
- 1966HO15 C.H. Holbrow, R. Middleton and B. Rosner, Phys. Rev. 152 (1966) 970
- 1966LI07 K.P. Lieb, Nucl. Phys. 85 (1966) 461
- 1966LO15 J.A. Lonergan and D.J. Donahue, Nucl. Phys. 88 (1966) 465
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966RI02 F. Riess and W. Trost, Nucl. Phys. 78 (1966) 385
- 1966RI04 E. Rivet, R.H. Pehl, J. Cerny and B.G. Harvey, Phys. Rev. 141 (1966) 1021
- 1966SI02 R.H. Siemssen, G.C. Morrison, B. Zeidman and H. Fuchs, Phys. Rev. Lett. 16 (1966) 1050
- 1966SI05 G.S. Sidhu and J.B. Gerhart, Phys. Rev. 148 (1966) 1024
- 1966SW01 C.P. Swann, Phys. Rev. 148 (1966) 1119
- 1966WA1C Warburton and Alburger, Nucl. Spin-Parity Assign.; Ed., Gove (1966) 114

- 1967AL16 A.U. Aldzhauakhiri, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, *Yad. Fiz.* 6 (1967) 248; *Sov. J. Nucl. Phys.* 6 (1968) 180
- 1967BA13 G.C. Ball and J. Cerny, *Phys. Rev.* 155 (1967) 1170
- 1967BL22 R.S. Blake, E.B. Paul, C.H. Sinex and S.T. Emerson, *Nucl. Phys.* A102 (1967) 305
- 1967CH19 C. Chasman, K.W. Jones, R.A. Ristinen and D.E. Alburger, *Phys. Rev.* 159 (1967) 830
- 1967CO32 S. Cohen and D. Kurath, *Nucl. Phys.* 101 (1967) 1
- 1967DE03 L.J. Denes and W.W. Daehnick, *Phys. Rev.* 154 (1967) 928
- 1967FE06 P. Fessenden and D.R. Maxson, *Phys. Rev.* 158 (1967) 948
- 1967FL10 J.L. Flinner, J.C. Harris and B. Mulligan, *Phys. Rev.* 161 (1967) 1082
- 1967GA12 A. Gallmann, F. Haas and B. Heusch, *Phys. Rev.* 164 (1967) 1257
- 1967GO27 Y.V. Gofman, N.I. Zaika, A.V. Mokhnach, O.F. Nemets, P.L. Shmarin and A.M. Yasnogorodskii, *Yad. Fiz.* 5 (1967) 718; *Sov. J. Nucl. Soc.* 5 (1967) 510
- 1967HA20 R.L. Hahn and E. Ricci, *Nucl. Phys.* A101 (1967) 353
- 1967NE06 J.B. Nelson and W.R. Smith, *Nucl. Phys.* A96 (1967) 671
- 1967OL02 J.W. Olness, A.R. Poletti and E.K. Warburton, *Phys. Rev.* 154 (1967) 971
- 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, *Phys. Rev.* 164 (1967) 1295
- 1967SP09 A. Sperduto, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 657
- 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, *Nucl. Instrum. Meth.* 56 (1967) 325
- 1967WO05 C. Wong, J.D. Anderson, J. McClure, B. Pohl, V.A. Madsen and F. Schmittroth, *Phys. Rev.* 160 (1967) 769
- 1967YO02 F.C. Young, P.D. Forsyth and J.B. Marion, *Nucl. Phys.* A91 (1967) 209
- 1967ZA01 C.D. Zafiratos, J.S. Lilley and F.W. Slee, *Phys. Rev.* 154 (1967) 887
- 1968AL12 K.W. Allen, T.K. Alexander and D.C. Healey, *Can. J. Phys.* 46 (1968) 1575
- 1968BA1E Ball, UCRL-18263 (1968)
- 1968BE30 R.A.I. Bell, R.D. Gill, B.C. Robertson, J.S. Lopes and H.J. Rose, *Nucl. Phys.* A118 (1968) 481
- 1968BE47 F. Bella, M. Alessio and P. Fratelli, *Nuovo Cim.* B58 (1968) 232
- 1968CL05 H.-G. Clerc and E. Kuphal, *Z. Phys.* 211 (1968) 452
- 1968CO24 J.A. Cookson, *Phys. Lett.* B27 (1968) 619
- 1968FR08 J.M. Freeman, J.G. Jenkin, D.C. Robinson, G. Murray and W.E. Burcham, *Phys. Lett.* B27 (1968) 156
- 1968HI14 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, G.J. Wagner and P. Turek, *Nucl. Phys.* A111 (1968) 265

- 1968HO23 K. Hosono, J. Phys. Soc. Jpn. 25 (1968) 36
- 1968JA09 J. Janecke, T.F. Yang, R.M. Polichar and W.S. Gray, Phys. Rev. 175 (1968) 1301
- 1968JO07 J.E. Jobst, Phys. Rev. 168 (1968) 1156
- 1968JO1C Joshi, Nucl. Phys. and Solid State Phys. Symp. (1968)
- 1968KA38 N.N. Kaushal, E.J. Winhold, P.F. Yergin, H.A. Medicus and R.H. Augustson, Phys. Rev. 175 (1968) 1330
- 1968KL06 R. Klubes, F. Baldeweg and G. Stiller, Nucl. Phys. A121 (1968) 113; Erratum Nucl. Phys. A127 (1969) 693
- 1968MA29 N.F. Mangelson, B.G. Harvey and N.K. Glendenning, Nucl. Phys. A117 (1968) 161
- 1968ME10 K. Meier-Ewert, K. Bethge and K.-O. Pfeiffer, Nucl. Phys. A110 (1968) 142
- 1968OK1C O'konski and Ha, J. Chem. Phys. 49 (1968) 5354
- 1968RO1C Rose, Hausser and Warburton, Rev. Mod. Phys. 40 (1968) 591
- 1968RO1D I. Rotter, Nucl. Phys. A122 (1968) 567
- 1968TO09 J.H. Towle and G.J. Wall, Nucl. Phys. A118 (1968) 500
- 1969AL13 M.S. Alvesh, O.I. Vasileva, V.G. Nadtochii, A.V. Spasskii, I.B. Teplov and L.N. Fataeva, Yad. Fiz. 10 (1969) 481; Sov. J. Nucl. Phys. 10 (1970) 276
- 1969AR1B Armstrong, Beery, Keaton and Veeser, LA 4177 (1969)
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969BA1J Balashov, Bochum Conf. STI/PUB/232 IAEA (1969) 59
- 1969BA1L Batusov et al., Yad. Fiz. 10 (1969) 354
- 1969BE21 H.A. Bentz, Z. Naturforsch. A24 (1969) 858
- 1969BE92 N. Bezic, D. Brajnik, D. Jamnik and G. Kernel, Nucl. Phys. A128 (1969) 426
- 1969BL1D Blin-Stoyle, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 115
- 1969BO1D Bonaccorsi, Scrocco and Tomasi, J. Chem. Phys. 50 (1969) 2940
- 1969BO32 I. Borbely, T. Dolinszky, J. Ero and G. Hrehuss, Acta Phys. Acad. Sci. Hung. 26 (1969) 269
- 1969CA1A R.R. Carlson and H.W. Wyborny, Phys. Rev. 178 (1969) 1529
- 1969CO02 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A127 (1969) 95
- 1969CU10 H.-H. Cuno, G. Clausnitzer and R. Fleischmann, Nucl. Phys. A139 (1969) 657
- 1969DE29 A.E. Denisov, R.P. Kolalis, Y.A. Nemilov, V.S. Sadkovskii and E.U. Teterin, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 1376; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1271
- 1969DI1B Dickens and Perey, Nucl. Sci. Eng. 36 (1969) 280

- 1969EL1A Elliott, Cargese Lecture in Phys.; Ed., M. Jean, Vol. 3 (1969) 337
- 1969ER1A Erdas, Mosconi, Pompei and Quarati, Nuovo Cim. B64 (1969) 406
- 1969FE10 J.L. Ferrero, A. Garcia, J. Milio and F. Senent, An. Fis. 65 (1969) 307
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA01 A. Gallmann, F. Hibou and P. Fintz, Nucl. Phys. A123 (1969) 27
- 1969GA18 J. Galin, B. Gatty, M. Lefort, J. Peter, X. Tarrago and R. Basile, Phys. Rev. 182 (1969) 1267
- 1969GO1C Golovanova and Zelenskaya, Sov. J. Nucl. Phys. 8 (1969) 158
- 1969GU02 H. Guratzsch, G. Hofmann, H. Muller and G. Stiller, Nucl. Phys. A129 (1969) 405
- 1969GU1C Guratzsch and Stiller, Zentral. Kernf. Rossendorf Dresden. Rept. No. Zfk 189 (1969)
- 1969HA1F Hanna, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 591
- 1969HE06 H.-D. Helb, H. Voit, G. Ischenko and W. Reichardt, Phys. Rev. Lett. 23 (1969) 176
- 1969HO23 C.H. Holbrow, R. Middleton and W. Focht, Phys. Rev. 183 (1969) 880
- 1969IW1A Iwao, Prog. Theor. Phys. 42 (1969) 1304
- 1969JA06 R.J. Jaszczak, R.L. Macklin and J.H. Gibbons, Phys. Rev. 181 (1969) 1428
- 1969JA15 L.A. Jacobson, Phys. Rev. 188 (1969) 1509
- 1969JO09 J. Jobst, S. Messelt and H.T. Richards, Phys. Rev. 178 (1969) 1663
- 1969KA1B R.W. Kavanagh, Nucl. Phys. A129 (1969) 172
- 1969KA1C Kaschlik and Koch, Wiss. Z. Friedrich. Schiller U. Jena. Math. Naturwiss. Reihe (Germany) 18 (1969) 37
- 1969KO1A Koch and Kaschlik, Wiss. Z. Friedrich. Schiller U. Jena. Math. Naturwiss. Reihe (Germany) 18 (1969) 43
- 1969KO1B Kouloumdjian, Thesis, Univ. of Lyon (1969)
- 1969KR21 N.N. Krasnov, P.P. Dmitriev, Z.P. Dmitrieva, I.O. Konstantinov and G.A. Molin, Atomn. Energ. (USSR) 27 (1969) 125
- 1969KU1B Kurtz, Braun and Cuer, Bochum Conf. STI/PUB/232 IAEA (1969) 257
- 1969LE1B Leonardi, Cargese Lecture in Phys.; Ed., M. Jean, Vol. 3 (1969) 97
- 1969LY1A Lyutostanskii, Yad. Fiz. 10 (1969) 198
- 1969MA1C Massue et al., Bochum Conf. STI/PUB/232 IAEA (1969) 258
- 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
- 1969SC1F Schiffer, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 665
- 1969SC1G Schirmer, Kreivopustov, Oehler and Sizov, Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR, Rept. P15 4801 (1969)

- 1969SC1H Schmitroth, Thesis, Oregon State Univ. (1969)
- 1969SN02 F.D. Snyder and M.A. Waggoner, Phys. Rev. 186 (1969) 999
- 1969SN04 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1259
- 1969SN1B Snelgrove, Thesis, Michigan State Univ. (1969)
- 1969SP02 A.V. Spasskii, K.G. Tarov, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 9 (1969) 936; Sov. J. Nucl. Phys. 9 (1969) 548
- 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
- 1969WA1C Warburton and Weneser, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 173
- 1970AD01 E.G. Adelberger, A.V. Nero and A.B. McDonald, Nucl. Phys. A143 (1970) 97
- 1970AD02 E.G. Adelberger and A.B. McDonald, Nucl. Phys. A145 (1970) 497
- 1970AG08 J. Aguilar, C.F. Pineda, C.R. Bauza, J.L. Ferrero, A. Garcia and F. Senent, An. Fis. 66 (1970) 279
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1970AL26 M. Allab, M. Amiel, A. Amokrane, H. Beaumevielle, A. Henni and M. Lambert, J. Phys. (Paris) 31 (1970) 919
- 1970AN1D Anni and Taffara, Rivista Nuovo Cim. 2 (1970) 1
- 1970AN1E Andrade et al., Bull. Amer. Phys. Soc. 15 (1970) 629
- 1970AR1D Artukh et al., Nucl. Reactions Included by Heavy Ions, Heidelberg, 1969 (1970) 140
- 1970AR27 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Phys. Lett. B33 (1970) 407
- 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
- 1970BA1J Barker et al., Nucl. Reactions included by Heavy Ions, Heidelberg, 1969 (1970) 152
- 1970BA44 Y.A. Batusov, B.Z. Zalikhanov, V.M. Sidorov, D. Tuvdendorzh and V.A. Yarba, Yad. Fiz. 11 (1970) 736; Sov. J. Nucl. Phys. 11 (1970) 413
- 1970BA63 N.P. Babenko and V.D. Domkin, Yad. Fiz. 12 (1970) 692; Sov. J. Nucl. Phys. 12 (1971) 375
- 1970BE54 B.L. Berman, S.C. Fultz, J.T. Caldwell, M.A. Kelly and S.S. Dietrich, Phys. Rev. C2 (1970) 2318
- 1970BL10 J.L. Black, W.J. Caelli, W.F. Davidson and R.B. Watson, Nucl. Phys. A153 (1970) 233
- 1970BO1K Borbei and Dolinski, Sov. J. Nucl. Phys. 10 (1970) 173
- 1970BO1M Bohm et al., Nucl. Phys. B24 (1970) 248
- 1970BO20 I.I. Bondouk, F. Machali and Z.A. Saleh, Z. Phys. 236 (1970) 170

- 1970BO21 W. Bohne, H. Homeyer, H. Lettau, H. Morgenstern, J. Scheer and F. Sichelschmidt, Nucl. Phys. A154 (1970) 105
- 1970BR1C Brix, Private Communication (1970)
- 1970BU08 J.-P. Burq, G. Hadinger, J. Kouloumdjian and J. Meyer, Nucl. Phys. A149 (1970) 488
- 1970BU1D Bukhvostov and Popov, Nucl. Phys. A147 (1970) 385
- 1970CA1C Cavallaro, Cunsolo, Potenza and Rubbino, Prgm. of 56th Natl. Congress of Italian Phys. Soc., Venice, 1970 (1970) 36
- 1970CA28 J. Catala, A. Garcia, V. Martinez and F. Senent, An. Fis. 66 (1970) 297
- 1970CH1D Chesterfield and Parker, Bull. Amer. Phys. Soc. 15 (1970) 1654
- 1970CL1C Clayton, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki (1970) 35
- 1970CL1D Cloud, Thesis, Univ. of Oregon (1970)
- 1970CO1H Cohen and Kurath, Nucl. Phys. A141 (1970) 145
- 1970CR03 G.M. Crawley, S.M. Austin, W. Benenson, V.A. Madsen, F.A. Schmittroth and M.J. Stomp, Phys. Lett. B32 (1970) 92
- 1970DA20 E.B. Dally, M.G. Croissiaux and B. Schweitz, Phys. Rev. C2 (1970) 2057
- 1970DI1A Dickens and Perey, Nucl. Sci. Eng. 40 (1970) 346
- 1970DM01 T.A. Dmitrieva, Z. Oziewicz and A. Pikulski, Nucl. Phys. A155 (1970) 205
- 1970DO04 J.J. Domingo, B.W. Allardyce, C.H.Q. Ingram, S. Rohlin, N.W. Tanner, J. Rohlin, E.M. Rimmer, G. Jones and J.P. Girardeau-Montaut, Phys. Lett. B32 (1970) 309
- 1970DU04 J.R. Duray and C.P. Browne, Phys. Rev. C1 (1970) 776
- 1970DU07 J.L. Duggan, J.Y. Park, S.D. Danielopoulos, P.D. Miller, J. Lin, M.M. Duncan and R.L. Dangle, Nucl. Phys. A151 (1970) 107
- 1970FO1B Fonda, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 283
- 1970FR13 N. Freed and P. Ostrander, Lett. Nuovo Cim. 3 (1970) 673
- 1970FR1C Frahn, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 297
- 1970GA07 S. Gangadharan and R.L. Wolke, Phys. Rev. C1 (1970) 1333
- 1970GA09 A. Gallman, F. Haas, N. Balaux, B. Heusch and M. Toulemonde, Can. J. Phys. 48 (1970) 1595
- 1970GA25 A. Garcia, J. Milio and F. Senent, An. Fis. 66 (1970) 69
- 1970GU13 M.R. Gunye and C.S. Warke, Phys. Rev. C2 (1970) 2106
- 1970HA1G Habs et al., Kernforsch. Karlsruhe, Ext. Bericht 18-70 2 (1970)
- 1970HA23 J.C. Hardy, A.D. Bacher, G.R. Plattner, J.A. Macdonald and R.G. Sextro, Phys. Rev. Lett. 25 (1970) 298
- 1970HS02 S.T. Hsieh and H. Horie, Nucl. Phys. A151 (1970) 243

- 1970HU08 J.N. Huffaker and C.E. Laird, Nucl. Phys. A150 (1970) 61
- 1970IS1A Ischenko, Reichardt, Voit and Held, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 105
- 1970JA02 W. Jaus and G. Rasche, Nucl. Phys. A143 (1970) 202
- 1970JO09 D.J. Johnson and M.A. Waggoner, Phys. Rev. C2 (1970) 41
- 1970JO1D Johnson, Thesis, Univ. of Iowa (1970)
- 1970KH01 S.B. Khadkikar and M.R. Gunye, Nucl. Phys. A144 (1970) 289
- 1970KN01 A.R. Knudson and F.C. Young, Nucl. Phys. A149 (1970) 323
- 1970KO26 T.I. Kopaleishvili and I.Z. Machabeli, Yad. Fiz. 12 (1970) 525; Sov. J. Nucl. Phys. 12 (1970) 286
- 1970LA1L Laird, Thesis, Univ. of Alabama (1970)
- 1970LE20 J.W. Leonard and D.O. Wells, Nucl. Phys. A153 (1970) 657
- 1970LI1E Liebenauer, Silverstein, Kibler and Koral, Bull. Amer. Phys. Soc. 15 (1970) 521
- 1970LK1A Lkhagva and Rotter, Sov. J. Nucl. Phys. 11 (1970) 576
- 1970LO1B W.G. Love, L.J. Parish and A. Richter, Phys. Lett. B31 (1970) 167
- 1970ME25 M.M. Meier, R.L. Walter, T.R. Donoghue, R.G. Seyler and R.M. Drisko, Nucl. Phys. A159 (1970) 273
- 1970ME30 S. Messelt, Phys. Norvegica 4 (1970) 191
- 1970NE17 J.M. Nelson, N.S. Chant and P.S. Fisher, Nucl. Phys. A156 (1970) 406
- 1970NE1B J.M. Nelson, N.S. Chant and P.S. Fisher, Phys. Lett. B31 (1970) 445
- 1970NE1F Neudatchin and Wildermuth, Proc. Int. Conf. Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 327
- 1970OG1A Ogloblin, Proc. Int. Conf. Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 231
- 1970PA23 H. Paul, Nucl. Phys. A154 (1970) 160
- 1970PE1B Pearson, Rickel and Zissermann, Nucl. Phys. A148 (1970) 273
- 1970PI1A Pickles and Learn, Bull. Amer. Phys. Soc. 15 (1970) 652
- 1970RO07 M.L. Roush, L.A. West and J.B. Marion, Nucl. Phys. A147 (1970) 235
- 1970SA06 H.S. Sandhu, Nucl. Phys. A146 (1970) 163
- 1970SH06 N.K. Sherman, R.W. Gellie, K.H. Lokan and R.G. Johnson, Phys. Rev. Lett. 25 (1970) 114
- 1970SI09 R.H. Siemssen, H.T. Fortune, R. Malmin, A. Richter, J.W. Tippie and P.P. Singh, Phys. Rev. Lett. 25 (1970) 536
- 1970SI1E Sievers and Prosser, Bull. Amer. Phys. Soc. 15 (1970) 1654

- 1970SO08 C. Sonntag, H. Rebel, B. Ribbat, S.K. Thio and W.R. Gramm, Lett. Nuovo Cim. 4 (1970) 717
- 1970SO11 L.S. Sokolov and N.E. Kabakova, Izv. Akad. Nauk SSSR Ser. Fiz. 34 (1970) 2159; Bull. Acad. Sci. USSR Phys. Ser. 34 (1971) 1926
- 1970SU1B Sukhoruchkin, Sov. J. Nucl. Phys. 10 (1970) 144
- 1970UL01 N. Ullah, S.S.M. Wong and L.E.H. Trainor, Can. J. Phys. 48 (1970) 842
- 1970VA24 V.A. Vartanyan, T.A. Dmitrieva, H.-U. Jager, H.R. Kissener and R.A. Eramzhyan, Yad. Fiz. 11 (1970) 528; Sov. J. Nucl. Phys. 11 (1970) 295
- 1970VE06 A.N. Vereshchagin, I.N. Korostova and I.P. Chernov, Izv. Vys. Ucheb. Zav. Fiz. 8 (1970) 105; Sov. Phys. J. 13 (1973) 1071
- 1970VO02 W. Von Oertzen, M. Liu, C. Caverzasio, J.C. Jacmart, F. Pougheon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A143 (1970) 34
- 1970WE03 H.R. Weller, Phys. Rev. C2 (1970) 321
- 1970WE09 L.C. Welch, C.C. Chang, H.H. Forster, C.C. Kim, D.W. Devins and P.A. Deutchman, Nucl. Phys. A158 (1970) 644
- 1970WE1J L.C. Welch, Thesis, Univ. of Southern California (1970)
- 1970WI1B Williams et al., Bull. Amer. Phys. Soc. 15 (1970) 62
- 1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1
- 1971AL09 B.J. Allen and R.L. Macklin, Phys. Rev. C3 (1971) 1737
- 1971AL1D Alder and Trautmann, Nucl. Phys. A178 (1971) 60
- 1971AM1A Amato et al., Bull. Amer. Phys. Soc. 16 (1971) 511
- 1971AN08 G. Andersson, B. Forkman and B. Friberg, Nucl. Phys. A171 (1971) 529
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AR1P Artukh et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 193
- 1971BA1A Barnes, Advances in Nucl. Phys. 4 (1971) 133
- 1971BA2T Banerji, Indian J. Phys. 44 (1971) 319
- 1971BA2Y Baus-Baghidian, Univ. libre Bruxelles, Bull. No. 45 (1971)
- 1971BA68 P.H. Barker, P.M. Cockburn, A. Huber, H. Knoth, U. Matter, H.-P. Seiler and P. Marmier, Ann. Phys. 66 (1971) 705
- 1971BA70 J. Banaigs, J. Berger, J. Duflo, L. Goldzahl, O. Harff, M. Cottreau, F. Lefebvres, H. Quechon and P. Tardy-Joubert, Nucl. Instrum. Meth. 95 (1971) 307
- 1971BA96 Y.A. Batusov, S.A. Bunyatov, V.A. Vartanyan, G.R. Gulkanyan, N. Kurts, V.M. Sidorov, K.M. Chernev, R.A. Eramzhyan and V.A. Yarba, Yad. Fiz. 14 (1971) 1206; Sov. J. Nucl. Phys. 14 (1972) 673

- 1971BAWQ G.C. Ball, W.G. Davies, J.S. Forster and J.C. Hardy, Bull. Amer. Phys. Soc. 16 (1971) 536, DF11
- 1971BE25 H.A. Bentz, Z. Phys. 243 (1971) 138
- 1971BE46 R.F. Bentley, J.D. Carlson, D.A. Lind, R.B. Perkins and C.D. Zafiratos, Phys. Rev. Lett. 27 (1971) 1081
- 1971BI03 M. Bister, A. Anttila, M. Piiparinen and M. Viitasalo, Phys. Rev. C3 (1971) 1972
- 1971BO1F Bolsterli, LA-DC 12417 (1971)
- 1971BO1U Bohlen et al., J. Phys. (Paris) 32 (1971) C6-141
- 1971BO39 V.I. Borovlev, R.P. Slabospitskii, A.P. Klyucharev, G.B. Andreev and I.D. Lopatko, Yad. Fiz. 14 (1971) 47; Sov. J. Nucl. Phys. 14 (1972) 27
- 1971BO44 V.I. Borovlev, A.P. Klyucharev, R.P. Slabospitskii, G.B. Andreyev and I.D. Lopatko, Izv. Akad. Nauk. SSSR Ser. Fiz. 35 (1971) 1670; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 1521
- 1971BR44 R.C. Brown, A.A. Debenham, G.W. Greenlees, J.A.R. Griffith, O. Karban, D.C. Kocher and S. Roman, Phys. Rev. Lett. 27 (1971) 1446
- 1971BU03 A. Budzanowski, L. Freindl, W. Karcz, B. Lazarska and W. Zipper, Nucl. Phys. A161 (1971) 610
- 1971BU1K Burymov et al., Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 159
- 1971CL07 G.J. Clark, J.M. Freeman, D.C. Robinson, J.S. Ryder, W.E. Burcham and G.T.A. Squier, Phys. Lett. B35 (1971) 503
- 1971CO14 W.R. Coker, P.D. Miller, J.L. Duggan, M.M. Duncan, R.L. Dangle and J. Lin, Nucl. Phys. A168 (1971) 307
- 1971CO32 J. Coudurier, F. Merchez and C. Perrin, Lett. Nuovo Cim. 2 (1971) 1117
- 1971CU01 T.H. Curtis, H.F. Lutz, D.W. Heikkinen and W. Bartolini, Nucl. Phys. A165 (1971) 19
- 1971DE42 V.P. Denisov, L.A. Kulchitskii and I.Y. Chubukov, Yad. Fiz. 14 (1971) 889; Sov. J. Nucl. Phys. 14 (1972) 497
- 1971DU03 J.R. Duray and C.P. Browne, Phys. Rev. C3 (1971) 1867
- 1971DU1B A.K. Dutta, Phys. Rev. C3 (1971) 2095
- 1971DZ07 P.O. Dzhambalov and E.I. Dolinskii, Yad. Fiz. 14 (1971) 753; Sov. J. Nucl. Phys. 14 (1972) 423
- 1971ES1A Escudie, Tarrats and Raynal, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 705
- 1971FA09 J. Favier, T. Bressani, G. Charpak, L. Massonnet, W.E. Meyerhof and C. Zupancic, Nucl. Phys. A169 (1971) 540
- 1971FA17 S.A. Fayans, Phys. Lett. B37 (1971) 155

- 1971FE1D Fetscher et al., Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 772
- 1971FL04 D.G. Fleming, J.C. Hardy and J. Cerny, Nucl. Phys. A162 (1971) 225
- 1971FO05 H.T. Fortune, R. Middleton, J.D. Garrett and C.H. Holbrow, Phys. Rev. C3 (1971) 1441
- 1971FO1K Fox, Austin and Larson, Bull. Amer. Phys. Soc. 16 (1971) 1163
- 1971FR11 B. Friberg, G. Andersson and B. Forkman, Nucl. Phys. A171 (1971) 551
- 1971GA05 R.M. Gaedke, W. Tobocman and K.S. Toth, Phys. Rev. C3 (1971) 1444
- 1971GA44 A. Garcia, J. Milio and F. Senent, An. Fis. 67 (1971) 175
- 1971GE12 H.V. Geramb, Phys. Lett. B37 (1971) 351
- 1971GO1E Goldfarb, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 205
- 1971GR20 J.A.R. Griffith, M. Irshad, O. Karban, S.W. Oh and S. Roman, Nucl. Phys. A167 (1971) 87
- 1971GR2B Grambole et al., Zentral. Kernf. Rossendorf Dresden, Rept. No. Zfk 212 (1971)
- 1971GR44 K.O. Groeneveld, A. Richter, U. Strohbusch and B. Zeidman, Phys. Rev. Lett. 27 (1971) 1806
- 1971GRYL M.B. Greenfield, Thesis, Univ. of Tennessee (1971); Diss. Abst. Int. B32 (1971) 4793
- 1971HA1R Haeberli, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 235
- 1971HE24 H.H. Heckman, D.E. Greiner, P.J. Lindstrom and F.S. Bieser, Science 174 (1971) 1130
- 1971HI09 D. Hilscher, J.C. Davis and P.A. Quin, Nucl. Phys. A174 (1971) 417
- 1971HS05 S.T. Hsieh and S.M. Hsieh, Chin. J. Phys. (Taiwan) 9 (1971) 92
- 1971HU10 J.N. Huffaker, Nucl. Phys. A171 (1971) 87
- 1971JA02 J.D. Jafar, H.B. van der Raay, D.G. Ryan, J.A. Stiegelmair and R.K. Tandon, Nucl. Phys. A161 (1971) 105
- 1971JA04 J. Janecke, T.F. Yang, W.S. Gray and R.M. Polichar, Phys. Rev. C3 (1971) 79
- 1971JA13 H.U. Jager, H.R. Kissener and R.A. Eramzhian, Nucl. Phys. A171 (1971) 16
- 1971JA14 H.U. Jager, H.R. Kissener and R.A. Eramzhian, Nucl. Phys. A171 (1971) 584
- 1971JA17 S. Jaccard, J.-F. Germond, J. Piffaretti and J. Weber, Helv. Phys. Acta 44 (1971) 502
- 1971JA20 W. Jaus, Nucl. Phys. A177 (1971) 70
- 1971KA04 S. Kahana and D. Kurath, Phys. Rev. C3 (1971) 543
- 1971KA18 S. Kardonsky, H.L. Finston and E.T. Williams, Phys. Rev. C4 (1971) 846

- 1971KA35 G. Kaschl, G. Mairle, H. Mackh, D. Hartwig and U. Schwinn, Nucl. Phys. A178 (1971) 275
- 1971KA70 N.N. Kaushal, E.J. Winhold, R.H. Augustson, P.F. Yergin and H.A. Medicus, Nucl. Energy 25 (1971) 91
- 1971KE01 D.P. Kerr and K.T. Bainbridge, Can. J. Phys. 49 (1971) 1950
- 1971KL1C Klein, Geller, Joblin and Hollandsworth, Bull. Amer. Phys. Soc. 16 (1971) 599
- 1971KO11 I. Kohno, S. Nakajima, T. Tonuma and M. Odera, J. Phys. Soc. Jpn. 30 (1971) 910
- 1971KO23 W.J. Kossler, H.O. Funsten, B.A. MacDonald and W.F. Lankford, Phys. Rev. C4 (1971) 1551
- 1971LA1B Lanford and Fulbright, Bull. Amer. Phys. Soc. 16 (1971) 35
- 1971LI11 M. Liu, W. von Oertzen, J.C. Jacmart, F. Pougeon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A165 (1971) 118
- 1971LI1K Liebenauer, Thesis, Case Western Reserve Univ. (1970)
- 1971LO25 I.I. Loshchakov, V.I. Ostroumov and A.E. Sadilin, Yad. Fiz. 14 (1971) 676; Sov. J. Nucl. Phys. 14 (1972) 379
- 1971MA1T Massue et al., Conf. on Certain Microscopic Aspects of Nucl. Interactions, La Tourinei, 1971 (1971)
- 1971MA2A Massue, FRNC TH 110 (1971)
- 1971MA38 A.P. Maclin and J.M. Eisenberg, Can. J. Phys. 49 (1971) 1826
- 1971MA46 K. Maluszynska, M. Przytula and I.V. Sizov, Acta Phys. Pol. B2 (1971) 183
- 1971MC15 G.M. McAllen, W.T. Pinkston and G.R. Satchler, Part. Nucl. 1 (1971) 412
- 1971ME18 H.O. Meyer, P. Huber and E. Baumgartner, Helv. Phys. Acta 44 (1971) 830
- 1971MI12 D. Miljanic and V. Valkovic, Nucl. Phys. A176 (1971) 110
- 1971MU18 G.S. Mutchler, D. Rendic, D.E. Velkley, W.E. Sweeney, Jr. and G.C. Phillips, Nucl. Phys. A172 (1971) 469
- 1971MU1H Mullensiefen, Nucl. Phys. B28 (1971) 368
- 1971NO02 J.L. Norton and P. Goldhammer, Nucl. Phys. A165 (1971) 33
- 1971NY03 K. Nyberg-Ponnert, B. Jonsson and I. Bergqvist, Phys. Scr. 4 (1971) 165
- 1971OD01 Y. Oda, H. Nakamura, A. Yoshida, M. Kusuhara and M. Murakami, J. Phys. Sov. Jpn. 30 (1971) 1217
- 1971PU01 G.D. Putt, Nucl. Phys. A161 (1971) 547
- 1971RI13 F. Riess, W.J. O'Connell and P. Paul, Nucl. Phys. A175 (1971) 462
- 1971RI15 H.T. Richards and H.V. Smith, Jr., Phys. Rev. Lett. 27 (1971) 1735
- 1971RO1G Rotter, J. Phys. (Paris) 32 (1971) C6-113

- 1971RO1H Rogers, Fessenden and Madsen, Bull. Amer. Phys. Soc. 16 (1971) 830
- 1971RO1J Rogers, Thesis, Oregon State Univ. (1971)
- 1971RU04 A.A. Rush, E.J. Burge and D.A. Smith, Nucl. Phys. A166 (1971) 378
- 1971SC1F Scott et al., J. Phys. (Paris) 32 (1971) C6-275
- 1971SC29 W. Schutz, H. Theissen, K.H. Schmidt and H. Frank, Z. Naturforsch. A26 (1971) 2070
- 1971SE06 E. Seibt and C. Weddigen, Phys. Lett. B35 (1971) 509
- 1971SH26 V.S. Shirley, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1255
- 1971SI24 M.H. Simbel and A.Y. Abul-Magd, Nucl. Phys. A177 (1971) 322
- 1971SM01 L.G. Smith, Phys. Rev. C4 (1971) 22
- 1971ST40 A.I. Steshenko and G.F. Filippov, Yad. Fiz. 14 (1971) 715; Sov. J. Nucl. Phys. 14 (1972) 403
- 1971TA1A Talmi, Proc. Int. Conf. Hyperfins Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1133
- 1971TH03 S.T. Thornton, C.M. Jones, J.K. Bair, M.D. Mancusi and H.B. Willard, Phys. Rev. C3 (1971) 1065
- 1971TH1D Thirion, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971TH1E Thornton et al., Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 871
- 1971VO01 W. Von Oertzen, G. Bohlen, B. Gebauer, E.E. Habib and N. Marquardt, Phys. Lett. B34 (1971) 51
- 1971VO04 D. von Ehrenstein, L. Meyer-Schutzmeister, J.E. Monahan, A. Richter and J.C. Stoltzfus, Phys. Rev. Lett. 27 (1971) 107
- 1971VO1D Von Oertzen et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 350
- 1971WA1D Walter, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 317
- 1971WI02 H. Wilsch and G. Clausnitzer, Nucl. Phys. A160 (1971) 609
- 1971WO03 C. Wong, J.D. Anderson, V.A. Madsen, F.A. Schmittroth and M.J. Stomp, Phys. Rev. C3 (1971) 1904
- 1971WY01 H.W. Wyborny and R.R. Carlson, Phys. Rev. C3 (1971) 2185
- 1971ZA04 N.I. Zaika, Y.V. Kibkalo, E.B. Levshin, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, P.L. Shmarin, A.M. Shchur and A.M. Yasnogorodskii, Yad. Fiz. 13 (1971) 928; Sov. J. Nucl. Phys. 13 (1971) 533

- 1971ZE1C Zeidman, Strohbusch, Ritcher and Grdeneveld, J. Phys. (Paris) 32 (1971) C6-295
- 1972AG01 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, G.P. Gulkanyan, V.M. Sidorov and V.A. Yarba, Yad. Fiz. 15 (1972) 18; Sov. J. Nucl. Phys. 15 (1972) 10
- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- 1972AL01 D.E. Alburger, Phys. Rev. C5 (1972) 274
- 1972AN05 R.K. Anderson, M.R. Wilson and P. Goldhammer, Phys. Rev. C6 (1972) 136
- 1972AR20 L. Armstrong and Jr., C.W. Kim, Phys. Lett. B41 (1972) 39
- 1972AU1A Austin, MSUCL-49 (1972)
- 1972BA56 D.P. Balamuth and J.W. Noe, Phys. Rev. C6 (1972) 30
- 1972BE04 H. Behrens and W. Buhring, Nucl. Phys. A179 (1972) 297
- 1972BE1E Bertsch and Mekjian, Ann. Rev. Nucl. Sci. (1972) 25
- 1972BL04 R.A. Blue, K.J. Stout and G. Marr, Nucl. Phys. A184 (1972) 377
- 1972BO17 V.V. Boikov, B.V. Velberg, I.I. Loshchakov and V.I. Ostroumov, Yad. Fiz. 15 (1972) 425; Sov. J. Nucl. Phys. 15 (1972) 239
- 1972BO56 V.I. Borovlev, I.D. Lopatko, G.B. Andreev and R.P. Slabospitsky, Ukr. Fiz. Zh. 17 (1972) 966
- 1972BO68 H.G. Bohlen, M. Feil, A. Gamp, B. Kohlmeyer, N. Marquardt and W. Von Oertzen, Phys. Lett. B41 (1972) 425
- 1972BR60 O.D. Brill, A.D. Vongai, K.A. Gridnev and A.R. Faiziev, Yad. Fiz. 16 (1972) 640; Sov. J. Nucl. Phys. 16 (1973) 359
- 1972BU1J Burlow and Forkman, LUNP 7208 (1972)
- 1972BU29 A.P. Bukhvostov, A.M. Chatrchyan, G.E. Dogotar, R.A. Eramzhyan, N.P. Popov and V.A. Vartanjan, Acta Phys. Pol. B3 (1972) 375
- 1972CA1N Caughlan and Fowler, Nature Phys. Sci. 238 (1972) 23
- 1972CA34 R.W. Carr, J.E.E. Baglin and E.J. Bentz, Jr., Phys. Rev. C6 (1972) 2032
- 1972CA37 P. Camiz, E. Olivier, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CI01 A. Ciocanel, V. Corcalciuc, E. Duduleanu, I. Minzatu, C. Pencea and O. Salagean, Rev. Roum. Phys. 17 (1972) 107
- 1972CL1A Clayton, Encyclopedia of the Twentieth Century (1972)
- 1972CO01 J.A. Cookson, Nucl. Phys. A180 (1972) 89
- 1972CRZN H. Crannell, P.L. Hallowell, J.T. O'Brien, J.M. Finn, F.J. Kline, S. Penner, J.W. Lightbody, Jr. and S.P. Fivozinsky, Proc. Int. Conf. on Nucl. Struct. Studies Using Electron Scattering and Photoreaction, Sendai, Japan (1972) 375

- 1972DA02 J.R. Davis and G.U. Din, Nucl. Phys. A179 (1972) 101; Erratum Nucl. Phys. A211 (1973) 619
- 1972DA04 P. David, J. Debrus, H. Mommsen and A. Riccato, Nucl. Phys. A182 (1972) 234
- 1972DI1B Diener and Young, Bull. Amer. Phys. Soc. 17 (1972) 465
- 1972DM01 T.A. Dmitrieva and I.B. Teplov, Yad. Fiz. 15 (1972) 478; Sov. J. Nucl. Phys. 15 (1972) 266
- 1972EM01 J.F. Emery, S.A. Reynolds, E.I. Wyatt and G.I. Gleason, Nucl. Sci. Eng. 48 (1972) 319
- 1972FA14 G. Falldt and H. Pilkuhn, Phys. Lett. B40 (1972) 613
- 1972FO07 R.P. Fogel and S.T. Thornton, Nucl. Phys. A187 (1972) 624
- 1972GE11 R.W. Gellie, K.H. Lokan, N.K. Sherman, R.G. Johnson and J.I. Lodge, Can. J. Phys. 50 (1972) 1689
- 1972GO23 N.G. Goncharova, Yad. Fiz. 15 (1972) 242; Sov. J. Nucl. Phys. 15 (1972) 137
- 1972GO27 N.G. Goncharova and G.S. Sagijan, Vestnik Mos. Univ. Fiz. Astron. (USSR) 13 (1972) 121
- 1972GR02 P.D. Greaves, V. Hnizdo, J. Lowe and O. Karban, Nucl. Phys. A179 (1972) 1
- 1972GR39 M.B. Greenfield, C.R. Bingham, E. Newman and M.J. Saltmarsh, Phys. Rev. C6 (1972) 1756
- 1972HA1X Hardy, Symp. on Two-Nucleon Transfer and Pairing Excitations, Argonne (1972) 203; CONF-720309
- 1972HU13 M. Huez, M. Quadlia and G. Weber, Nucl. Instrum. Meth. 105 (1972) 197
- 1972HU1A W.T. Huang, C.A. Levinson and M.K. Banerjee, Phys. Rev. C5 (1972) 651
- 1972JA19 W. Jaus, Phys. Lett. B40 (1972) 616
- 1972JA1C Janiszewski and McCarthy, Nucl. Phys. A181 (1972) 97
- 1972KE02 P.W. Keaton, Jr., D.D. Armstrong, L.R. Veeser, H.T. Fortune and N.R. Roberson, Nucl. Phys. A179 (1972) 561
- 1972KO30 V.M. Kolybasov and V.A. Khangulyan, Yad. Fiz. 15 (1972) 934; Sov. J. Nucl. Phys. 15 (1972) 520
- 1972KO31 T.I. Kopaleishvili, I.Z. Machabeli and M.S. Chachkhunashvili, Yad. Fiz. 15 (1972) 1137; Sov. J. Nucl. Phys. 15 (1972) 629
- 1972KU19 B.I. Kuznetsov, P.E. Ovsyannikova and I.P. Chernov, Yad. Fiz. 15 (1972) 673; Sov. J. Nucl. Phys. 15 (1972) 377
- 1972KU1C Kuo, 1970 4th Symp. on the Struct. of Low-Medium Mass Nuclei (1972) 75
- 1972KU1H Kurath, Comments Nucl. Part. Phys. 5 (1972) 55
- 1972LA12 C.D. Latorre and J.M. Irvine, Nucl. Phys. A188 (1972) 524

- 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353
- 1972LI06 S. Lie, Nucl. Phys. A181 (1972) 517
- 1972LO08 J. Lowe and A.R. Barnett, Nucl. Phys. A187 (1972) 323
- 1972LO1D Lobov, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 881
- 1972LU1B Lucas and Root, J. Appl. Phys. 43 (1972) 3886
- 1972MA74 H. Matsuzaki, Prog. Theor. Phys. 48 (1972) 1534
- 1972MA77 Y.G. Mashkarov, R.P. Slabospitskii and V.I. Borovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 2623; Bull. Acad. Sci. USSR Phys. Ser. 36 (1973) 2274
- 1972MO1E Morrison, J. Phys. (Paris) 33 (1972) C5-111
- 1972NI05 L. Nichol and T.J. Kennett, Can. J. Phys. 50 (1972) 553.
- 1972NO08 J.W. Noe, D.P. Balamuth and R.W. Zurmuhle, Phys. Rev. C6 (1972) 780
- 1972NY02 G. Nystrom, H. Conde, B. Lundberg and L.G. Stromberg, Phys. Scr. 5 (1972) 175
- 1972OE01 H. Oeschler, H. Schroter, H. Fuchs, L. Baum, G. Gaul, H. Ludecke, R. Santo and R. Stock, Phys. Rev. Lett. 28 (1972) 694
- 1972PA1A Parkinson, At. Masses Fund. Constants, Teddington, 1971 (1972) 38
- 1972PA31 J.Y. Park, W. Scheid and W. Greiner, Phys. Rev. C6 (1972) 1565; Erratum Phys. Rev. C7 (1973) 1726
- 1972PE09 G. Perrin, Nguyen Van Sen, A. Fiore and J. Arvieux, Nucl. Instrum. Meth. 101 (1972) 603
- 1972PE15 G. Perrin, Nguyen Van Sen, J. Arvieux, A. Fiore, J.L. Durand, R. Darves-Blanc, J.C. Gondrand, F. Merchez and C. Perrin, Nucl. Phys. A193 (1972) 215
- 1972RA34 G.H. Rawitscher, Phys. Rev. C6 (1972) 1212
- 1972RE10 M.J. Renan, J.P.F. Sellschop, R.J. Keddy and D.W. Mingay, Nucl. Phys. A193 (1972) 470
- 1972RE1C Rendic, Conf. Nucl. Struct. Study with Neutrons, Budapest, 1972 (1972) 28
- 1972RO35 A. Roberts, Nucl. Phys. A196 (1972) 465
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430.
- 1972SC1L Scott et al., Communications, Proc. Aix-en-Provence Conf. 2 (1972) 75
- 1972SC21 D.K. Scott, P.N. Hudson, P.S. Fisher, C.U. Cardinal, N. Anyas-Weiss, A.D. Panagiotou and P.J. Ellis, Phys. Rev. Lett. 28 (1972) 1659
- 1972SE09 F. Seiler, Nucl. Phys. A187 (1972) 379
- 1972SI50 J. Singh, Indian J. Pure Appl. Phys. 10 (1972) 289
- 1972SM07 H.V. Smith,Jr., Phys. Rev. C6 (1972) 441

- 1972SW1A Swannack, Thesis, Carnegie-Mellon Univ. (1972)
- 1972TA21 V.K. Tartakovskii and A.V. Fursaev, *Yad. Fiz.* 15 (1972) 51; *Sov. J. Nucl. Phys.* 15 (1972) 31
- 1972TH13 C. Thibault and R. Klapisch, *Phys. Rev. C* 6 (1972) 1509
- 1972THZF H. Theissen, Springer Tracts in Mod. Phys.; Ed., Hohler, Vol. 65 (1972) 1
- 1972VA36 J.F.A. Van Hienen and P.W.M. Glaudemans, *Phys. Lett. B* 42 (1972) 301
- 1972VO06 A.A. Vorobyov, D.M. Seliverstov, V.T. Grachev, I.A. Kondurov, A.M. Nikitin, N.N. Smirnov and Y.K. Zalite, *Phys. Lett. B* 40 (1972) 102
- 1972WA1F Waheed, Juric and Adamovic, *Fizika* 4 (1972) 87
- 1972WA1G Wapstra, At. Masses Fund. Constants, Teddington, 1971 (1972) 283
- 1972WI05 D.H. Wilkinson, *Nucl. Phys. A* 179 (1972) 289
- 1972WI1C Wilkinson, Few Particle Problems, UCLA, 1972 (1972) 191
- 1972WI28 D.H. Wilkinson, *Proc. Roy. Soc. Edinburgh A* 70 (1972) 307
- 1972YO1C Youssef, *Bull. Amer. Phys. Soc.* 17 (1972) 917
- 1973AB1E Abuzeid et al., *Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR, Rept. P*15 (1973) 7218
- 1973AG05 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, V.I. Genchev, N.F. Golovanova, G.R. Gulkanyan, V.M. Sidorov, K.M. Chernev and V.A. Yarba, *Yad. Fiz.* 18 (1973) 256; *Sov. J. Nucl. Phys.* 18 (1974) 131
- 1973AG06 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, V.I. Genchev, N.F. Golovanova, G.R. Gulkanyan, V.M. Sidorov, K.M. Chernev and V.A. Yarba, *Yad. Fiz.* 18 (1973) 264; *Sov. J. Nucl. Phys.* 18 (1974) 135
- 1973AJ01 F. Ajzenberg-Selove, H.G. Bingham and J.D. Garrett, *Nucl. Phys. A* 202 (1973) 152
- 1973AL1B Aleshin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 1775
- 1973AR1B Arthur, Thesis LA 5230 T (1973)
- 1973AR1E Arnett, *Ann. Rev. Astron. Astrophys.* 11 (1973) 73
- 1973AU1B Audouze, Truran and Zimmerman, OAP-315 (1973)
- 1973AU1C Audouze, Explosive Nucleosynthesis (1973) 47
- 1973AU1D Audouze and Fricke, *Astrophys. J.* 186 (1973) 239
- 1973AU1E Austin and Fox, in Munich, 1 (1973) 388
- 1973BA2G Baer and Crowe, in Asilomar (1973) 583
- 1973BA34 G.C. Ball, G.J. Costa, W.G. Davies, J.S. Forster, J.C. Hardy and A.B. McDonald, *Phys. Rev. Lett.* 31 (1973) 395

- 1973BA68 Y.A. Batusov, S.A. Bunyatov, L. Vizireva, G.R. Gulkanyan, F. Mirsalikhova and V.M. Sidorov, *Yad. Fiz.* 18 (1973) 962; *Sov. J. Nucl. Phys.* 18 (1974) 496
- 1973BA81 L.K. Batist, E.E. Berlovich, Y.S. Blinnikov, Y.V. Elkin, Y.N. Novikov, B.M. Ovchinnikov and V.K. Tarasov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 1944; *Bull. Acad. Sci. USSR Phys. Ser.* 37 (1974) 124
- 1973BE16 Y.I. Bely, R.A. Eramzhyan, L. Majling, J. Rizek and V.A. Vartanyan, *Nucl. Phys. A204* (1973) 357
- 1973BE1R Beukens and Drake, *Bull. Amer. Phys. Soc.* 18 (1973) 775
- 1973BE25 M. Berrada and L. Marquez, *Part. Nucl.* 5 (1973) 89
- 1973BEVW Z. Berant, G. Goldring, M. Hass and Y.S. Horowitz, *J. Phys. Soc. Jpn. Suppl.* 34 (1973) 185
- 1973BI1B Biro, Sudar, Csikai and Dezho, *Neitronnaya Fiz., Pt. III*, 1973 (1974) 85
- 1973BI1G Bizard et al., *Nucl. Instrum. Meth.* 111 (1973) 445
- 1973BO10 J.R. Bobbitt, III, M.P. Etten and G.H. Lenz, *Nucl. Phys. A203* (1973) 353
- 1973BO1R Bochsler and Geiss, *Solar Phys.* 32 (1973) 3
- 1973BO30 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, *Phys. Rev. Lett.* 31 (1973) 614
- 1973BU1B J.E. Butt and D.T. King, *Phys. Rev. C8* (1973) 359
- 1973CA1B Cameron, *Explosive Nucleosynthesis* (1973) 3
- 1973CE1B Cerny et al., in *Munich*, 1 (1973) 465
- 1973CH06 C.C. Chao and J.K.P. Lee, *Can. J. Phys.* 51 (1973) 592
- 1973CH20 Il-Tong Cheon, *Phys. Rev. C8* (1973) 534
- 1973CH38 C.v. Charzewski, R.M. Dreizler and H.T. Fortune, *Nucl. Phys. A213* (1973) 429
- 1973CI02 A. Ciocanel, V. Corcalciuc and I. Minzatu, *Acta Phys. Aust.* 38 (1973) 41
- 1973CL09 C.F. Clement, *Nucl. Phys. A213* (1973) 469
- 1973CL12 G.J. Clark, J.M. Freeman, D.C. Robinson, J.S. Ryder, W.E. Burcham and G.T.A. Squier, *Nucl. Phys. A215* (1973) 429
- 1973CL1E Clayton and Woosley, in *Munich*, 2 (1973) 718
- 1973CO16 S. Cotanch and D. Robson, *Nucl. Phys. A209* (1973) 301
- 1973CO1B Colgate, *Explosive Nucleosynthesis* (1973) 248
- 1973CR1B Crannell et al., *Bull. Amer. Phys. Soc.* 18 (1973) 669
- 1973DA17 S.E. Darden, S. Sen, H.R. Hiddleston, J.A. Aymar and W.A. Yoh, *Nucl. Phys. A208* (1973) 77
- 1973DA37 S. Dahlgren, P. Grafstrom, B. Hoistad and A. Asberg, *Phys. Lett. B47* (1973) 439

- 1973DE30 V.P. Denisov and I.Y. Chubukov, *Yad. Fiz.* 17 (1973) 682; *Sov. J. Nucl. Phys.* 17 (1974) 354
- 1973DI1C Dixon, in *Asilomar* (1973) 727
- 1973EI01 J.M. Eisenberg, R. Guy, J.V. Noble and H.J. Weber, *Phys. Lett.* B43 (1973) 20; *Erratum Phys. Lett.* B45 (1973) 93
- 1973FA1J Falldt, Pilkuhn and Schlaile, *Contrib., Uppsala* (1973) 92
- 1973FA1K Fagg, in *Asilomar* (1973) 663
- 1973FE03 M. Feil, W. von Oertzen, H.G. Bohlen, A. Gamp, R.L. Walter and B. Kohlmeyer, *Z. Phys.* 260 (1973) 271
- 1973FE13 G. Fey, H. Frank, W. Schutz and H. Theissen, *Z. Phys.* 265 (1973) 401
- 1973GA19 S. Gamba, G. Ricco and G. Rottigni, *Nucl. Phys.* A213 (1973) 383
- 1973GO01 A. Gobbi, R. Wieland, L. Chua, D. Shapira and D.A. Bromley, *Phys. Rev. C7* (1973) 30
- 1973GO1H J.M.G. Gomez and J. Sesma, *Phys. Lett.* B44 (1973) 231
- 1973GR1F Grishin, Eliseev, Inogamov and Kohli, *Yad. Fiz.* 17 (1973) 1017
- 1973HA2E Hardy, Rept. AECL-4447, At. Energy Canada Ltd., Chalk River (1973)
- 1973HA40 F. Haas, R.M. Freeman, B. Heusch, S. Kohmoto and A. Gallmann, *Nucl. Phys.* A211 (1973) 289
- 1973HA49 P.S. Hauge and S. Maripuu, *Phys. Rev. C8* (1973) 1609
- 1973HA54 L.F. Hansen, S.M. Grimes, J.L. Kammerdiener and V.A. Madsen, *Phys. Rev. C8* (1973) 2072
- 1973HE09 B. Heusch and A. Gallmann, *Phys. Rev. C7* (1973) 1810
- 1973HO10 C.G. Hoot, D.K. Olsen, R.E. Brown, J.R. Maxwell and A. Scott, *Nucl. Phys.* A203 (1973) 339
- 1973HO32 V. Horsfjord, *Phys. Lett.* B45 (1973) 455
- 1973IB1B Iben, *Explosive Nucleosynthesis* (1973) 115
- 1973IG02 M. Igarashi, M. Kawai and K. Yazaki, *Prog. Theor. Phys.* 49 (1973) 825
- 1973IN05 P.D. Ingalls, *Phys. Rev. Lett.* 31 (1973) 605
- 1973JO10 R.C. Johnson, F.D. Santos, R.C. Brown, A.A. Debenham, G.W. Greenlees, J.A.R. Griffith, O. Karban, D.C. Kocher and S. Roman, *Nucl. Phys.* A208 (1973) 221
- 1973JO13 P.L. Jolivette, *Phys. Rev. C8* (1973) 1230
- 1973JU1B Julien, Polichar and Parkinson, *Bull. Amer. Phys. Soc.* 18 (1973) 617
- 1973KA04 V.V. Karapetyan, V.N. Mileev and N.N. Titarenko, *Nucl. Phys.* A203 (1973) 561

- 1973KA19 P.J. Karol, A.A. Caretto, Jr., R.L. Klobuchar, D.M. Montgomery, R.A. Williams and M.V. Yester, Phys. Lett. B44 (1973) 459
- 1973KA1D Karapetyan, Korenman and Popov, Contrib., Uppsala (1973) 154
- 1973KA1H Kao, Ann. Phys. 81 (1973) 113
- 1973KA46 T. Kaneko, Prog. Theor. Phys. 50 (1973) 1765
- 1973KE1C A.K. Kerman and M.S. Weiss, Phys. Rev. C8 (1973) 408
- 1973KI05 H.R. Kissener, R.A. Eramzhian and H.U. Jager, Nucl. Phys. A207 (1973) 78
- 1973KI12 H.R. Kissener, A. Aswad, H.U. Jager and R.A. Eramzhian, Nucl. Phys. A215 (1973) 424
- 1973KL12 F.J. Kline, H. Crannell, J.T. O'Brien, J. McCarthy and R.R. Whitney, Nucl. Phys. A209 (1973) 381
- 1973KO1D Kovar, Symp. on Heavy Ion Transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 59
- 1973KU03 D. Kurath, Phys. Rev. C7 (1973) 1390
- 1973KU1J Kubodera and Rho, Contrib., Uppsala (1973) 19
- 1973LE26 D. Lemann and H.J. Treffkorn, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1870; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 65
- 1973LI15 C.L. Lin, S. Yamaji and H. Yoshida, Nucl. Phys. A209 (1973) 135
- 1973MA1W Maute, Blatt and D'Amato, Asilomar (1973) Paper 8B6
- 1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
- 1973MC1J McMahan and Tobocman, Nucl. Phys. A202 (1973) 561
- 1973ME22 H.O. Meyer and J.A. Thomson, Phys. Rev. C8 (1973) 1215
- 1973MI1D Mitals, J. Roy. Astron. Soc. Canada 67 (1973) 197
- 1973MO15 T. Mo and H.R. Weller, Phys. Rev. C8 (1973) 972
- 1973MU05 N.C. Mukhopadhyay, Phys. Lett. B44 (1973) 33
- 1973MU11 N.C. Mukhopadhyay, Phys. Lett. B45 (1973) 309
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973MU1B Mukhopadhyay, Lett. Nuovo Cim. 7 (1973) 460
- 1973NY04 E.M. Nyman, Nucl. Phys. A215 (1973) 397
- 1973OE01 H. Oeschler, H. Fuchs and H. Schroter, Nucl. Phys. A202 (1973) 513
- 1973OG1A Ogloblin, Sov. J. Part. Nucl. 3 (1973) 467
- 1973PA1F Parker, Baglin and Skwiersky, Bull. Amer. Phys. Soc. 18 (1973) 550
- 1973PA1L Park, Danielopoulos, Scheid and Greiner, Bull. Amer. Phys. Soc. 18 (1973) 601
- 1973PA1M Park, in Munich, 1 (1973) 365

- 1973PO1D Pougheon and Roussel, in Munich, 1 (1973) 459
- 1973PR08 J.G. Pronko, R.G. Hirko and D.C. Slater, Phys. Rev. C7 (1973) 1382
- 1973RA1D Ramaty, Borner and Cohen, Astrophys. J. 181 (1973) 891
- 1973RA37 G.M. Raisbeck, J. Lestringuez and F. Yiou, Nature 244 (1973) 28
- 1973RE13 W. Reilly, R. Wieland, A. Gobbi, M.W. Sachs and D.A. Bromley, Nuovo Cim. A13 (1973) 897
- 1973RE1E Rekstad et al., in Munich, 1 (1973) 279
- 1973RE1F Rekstad et al., Phys. Norvegica 7 (1973) 103
- 1973RH1A Rhea et al., Bull. Amer. Phys. Soc. 18 (1973) 698
- 1973RO1Q Rook, in Munich, 2 (1973) 798
- 1973RO1R Robson, Nucl. Phys. A204 (1973) 523
- 1973SA1J Sackmann, Smith and Despain, OAP-324 (1973)
- 1973SA30 R. Saayman, P.R. de Kock and J.H. van der Merwe, Z. Phys. 265 (1973) 69
- 1973SC1J Scott, Symp. on Heavy Ion transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 97
- 1973SC1K Scheid, Fink and Muller, Proc. Europhys. 1972 Plitvice Lakes Conf. (1973) 144
- 1973SC1T J.M. Scalo and R.K. Ulrich, Astrophys. J. 183 (1973) 151
- 1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, Nucl. Phys. A212 (1973) 573
- 1973SE02 J.C. Sens, F. Rietsch, A. Pape and R. Armbruster, Nucl. Phys. A199 (1973) 232
- 1973SH14 R.W. Sharp and L. Zamick, Nucl. Phys. A208 (1973) 130
- 1973SI21 M.H. Simbel, Phys. Rev. C8 (1973) 75
- 1973SM1A Smith, Sackmann and Despain, Explosive Nucleosynthesis (1973) 168
- 1973SO04 J.P. Sokol, R.M. Prior, A.A. Rollefson and C.P. Browne, Phys. Rev. C7 (1973) 2154
- 1973SP1A Sprung, Nucl. Many-Body Problem, 1972, Vol. 2 (1973) 123
- 1973ST1A Stokstad, Proc. Europhys. 1972 Plitvice Lakes Conf. (1973) 179
- 1973ST1K Stronach, Kossler and Funsten, Bull. Amer. Phys. Soc. 18 (1973) 691
- 1973ST1L Stokstad, Switkowski and Wieland, Bull. Amer. Phys. Soc. 18 (1973) 1596
- 1973SU1B Sugimoto, J. Phys. Soc. Jpn. Suppl. 34 (1973) 197
- 1973SU1E Suffert, in Asilomar (1973) 741
- 1973TA1D Talbot, Explosive Nucleosynthesis (1973) 47
- 1973TA1E Talbot, Explosive Nucleosynthesis (1973) 34
- 1973TA30 K. Takahashi, M. Yamada and T. Kondoh, At. Data Nucl. Data Tables 12 (1973) 101

- 1973TH1B Theissen, in Asilomar (1973) 871
- 1973TO04 I.S. Towner and J.C. Hardy, Nucl. Phys. A205 (1973) 33
- 1973TO05 W. Tobocman, R. Ryan, A.J. Baltz and S.H. Kahana, Nucl. Phys. A205 (1973) 193
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973TR02 S. Tryti, T. Holtebekk and J. Rekstad, Nucl. Phys. A201 (1973) 135
- 1973TR1B Truran, Cosmochemistry; Ed., Cameron (1973) 23
- 1973TR1E Trimble and Reines, Rev. Mod. Phys. 45 (1973) 1
- 1973VA25 L. Van der Zwan and K.W. Geiger, Nucl. Phys. A216 (1973) 188
- 1973VO04 W. von Oertzen and W. Norenberg, Nucl. Phys. A207 (1973) 113
- 1973VO1D Volkov, in Munich, 2 (1973) 280
- 1973VO1G Vorobyov et al., in Munich, 1 (1973) 716
- 1973WE19 K.A. Weaver, J.D. Anderson, H.H. Barschall and J.C. Davis, Nucl. Sci. Eng. 52 (1973) 35
- 1973WH03 R.L. White, K.W. Kemper, L.A. Charlton and W.J. Courtney, Phys. Rev. Lett. 31 (1973) 254
- 1973WI15 J. Wilczynski, Phys. Lett. B47 (1973) 124
- 1973WI1J Wilkinson, Private Communication (1973)
- 1974AD11 A. Adam and J. Cabe, Nucl. Instrum. Meth. 121 (1974) 339
- 1974AD1B Adelberger, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 641
- 1974AD1C S.L. Adler, S. Nussinov and E.A. Paschos, Phys. Rev. D9 (1974) 2125
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974AL07 T.K. Alexander, G.J. Costa, J.S. Forster, O. Hausser, A.B. McDonald, A. Olin and W. Witthuhn, Phys. Rev. C9 (1974) 1748
- 1974AL11 D.E. Alburger and D.R. Goosman, Phys. Rev. C10 (1974) 912
- 1974AN36 N. Anyas-Weiss, J.C. Cornell, P.S. Fisher, P.N. Hudson, A. Menchaca-Rocha, D.J. Millener, A.D. Panagiotou, D.K. Scott, D. Strottman, D.M. Brink et al., Phys. Rept. 12 (1974) 201
- 1974AR16 J. Arvieux, J. Birchall, H.E. Conzett, W. Dahme, W. Haeberli and R.M. Larimer, Nucl. Instrum. Meth. 116 (1974) 605
- 1974AS07 J. Asai and H. Ogata, Nucl. Phys. A233 (1974) 55
- 1974AZ01 G. Azuelos, J.E. Crawford and J.E. Kitching, Phys. Rev. C9 (1974) 1213
- 1974BA37 J.E.E. Baglin, E.J. Bentz and R.W. Carr, Phys. Rev. C10 (1974) 24
- 1974BA40 S. Barshay, C.B. Dover and J.P. Vary, Phys. Lett. B51 (1974) 5

- 1974BA65 J. Bang, T. Elin and H. Schulz, *Yad. Fiz.* 20 (1974) 346; *Sov. J. Nucl. Phys.* 20 (1975) 184
- 1974BA89 L.K. Batist, E.E. Berlovich, L.A. Vaishnene, V.G. Konev, K.A. Mezilev and Y.N. Novikov, *Yad. Fiz.* 20 (1974) 850; *Sov. J. Nucl. Phys.* 20 (1975) 453
- 1974BE16 F.D. Becchetti, D.G. Kovar, B.G. Harvey, D.L. Hendrie, H. Homeyer, J. Mahoney, W. von Oertzen and N.K. Glendenning, *Phys. Rev. C9* (1974) 1543
- 1974BE1R Bertojo, Chui and Townes, *Science* 104 (1974) 619
- 1974BI1E Bindal et al., *Proc. Conf. Reactions between Complex Nuclei*, Vol. 1 (1974) 5
- 1974BO05 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, *Phys. Rev. C9* (1974) 836
- 1974BO18 A. Bottino, G. Ciocchetti and C.W. Kim, *Phys. Rev. C9* (1974) 2052
- 1974BO22 A. Bouyssy and N. Vinh Mau, *Nucl. Phys. A224* (1974) 331
- 1974BO51 I.I. Bondouk, *Atomkernenergie* 24 (1974) 135
- 1974BO52 I.I. Bondouk, *Atomkernenergie* 24 (1974) 199
- 1974BO53 I.I. Bondouk, *Atomkernenergie* 24 (1974) 201
- 1974BU06 C.E. Busch, T.B. Clegg, S.K. Datta and E.J. Ludwig, *Nucl. Phys. A223* (1974) 183
- 1974CA1H Cantwell et al., *Nucl. Phys. A236* (1974) 445
- 1974CH1Q Chechik, Stocker, Eyal and Fraenkel, *Proc. Conf. Reactions between Complex Nuclei*, Vol. 1 (1974) 6
- 1974CR01 D.J. Crozier and H.T. Fortune, *Phys. Rev. C10* (1974) 377
- 1974DA06 N.E. Davison, P. Fintz and A. Gallmann, *Nucl. Phys. A220* (1974) 166
- 1974DE1E De Jager, De Vries and De Vries, *At. Data Nucl. Data Tables* 14 (1974) 479
- 1974DE25 C. Detraz, F. Pougheon, M. Bernas, M. Langevin, P. Roussel and J. Vernotte, *Nucl. Phys. A228* (1974) 39
- 1974DM01 T.A. Dmitrieva, V.M. Lebedev, V.G. Markman, A.V. Spasskii and I.B. Teplov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2567; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 91
- 1974DO03 E.I. Dolinskii, V.V. Turovtsev and R. Yarmukhamedov, *Yad. Fiz.* 19 (1974) 1003; *Sov. J. Nucl. Phys.* 19 (1975) 514
- 1974DU02 T. Dubler, L. Schellenberg, H. Schneuwly, R. Engfer, J.L. Vuilleumier, H.K. Walter, A. Zehnder and B. Fricke, *Nucl. Phys. A219* (1974) 29
- 1974DU11 S. Dubnicka and O.V. Dumbrajs, *Nucl. Phys. A235* (1974) 417
- 1974DZ03 R.I. Dzhibuti and H.M. Sallam, *Yad. Fiz.* 19 (1974) 75; *Sov. J. Nucl. Phys.* 19 (1974) 40

- 1974EN01 N. Ensslin, W. Bertozi, S. Kowalski, C.P. Sargent, W. Turchinetz, C.F. Williamson, S.P. Fivozinsky, J.W. Lightbody, Jr. and S. Penner, Phys. Rev. C9 (1974) 1705
- 1974EN10 R. Engfer, H. Schneuwly, J.L. Vuilleumier, H.K. Walter and A. Zehnder, At. Data Nucl. Data Tables 14 (1974) 509
- 1974FA1A Faldt, Pilkhun and Schlails, Ann. Phys. 82 (1974) 326
- 1974FL1A Fleury and Alexander, Ann. Rev. Nucl. Sci. 24 (1974) 279
- 1974FU1B Fulcher and Mukhopadhyay, Bull. Amer. Phys. Soc. 19 (1974) 44
- 1974GA1L Galster, Ischenko, Siller and Voit, Proc. Conf. Reactions between Complex Nucl., Vol. 1 (1974) 37
- 1974GM01 K. Gmur and H.H. Muller, Helv. Phys. Acta 46 (1974) 807
- 1974GO1G Golovanova, Il'in, Neudachin and Smirnov, JETP Lett. 20 (1974) 310
- 1974GO1L Gobbi, in Nashville, Vol. 2 (1974) 211
- 1974GO1T Goosman and Gorodetzky, Bull. Amer. Phys. Soc. 19 (1974) 432
- 1974GO23 D.R. Goosman, D.F. Geesaman, F.E. Cecil, R.L. McGrath and P. Paul, Phys. Rev. C10 (1974) 1525
- 1974GR39 K.A. Gridnev and A.R. Faiziev, Yad. Fiz. 20 (1974) 911; Sov. J. Nucl. Phys. 20 (1975) 485
- 1974HA1C Hanna, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 249
- 1974HI1C Hinderling and Staub, Helv. Phys. Acta 47 (1974) 93
- 1974HO1E Hollandsworth, Bucher and Youngblood, Bull. Amer. Phys. Soc. 19 (1974) 574
- 1974HO1M Holbrow, Garrett and Fortune, Bull. Amer. Phys. Soc. 19 (1974) 16
- 1974HU14 J. Hufner, L. Tauscher and C. Wilkin, Nucl. Phys. A231 (1974) 455
- 1974IN07 A. Ingemarsson and G. Tibell, Phys. Scr. 10 (1974) 159
- 1974JA25 N. Jarmie, J.H. Jett and R.J. Semper, Phys. Rev. C10 (1974) 1748
- 1974JE1A Jolley, Wozniak, Nagarajan and Cerny, Bull. Amer. Phys. Soc. 19 (1974) 431
- 1974JO01 P.L. Jolivette, Phys. Rev. C9 (1974) 16
- 1974JO1J Johnson, Eok and Ellidtt, Bull. Amer. Phys. Soc. 19 (1974) 549
- 1974KE06 K.W. Kemper, S. Cotanch, G.E. Moore, A.W. Obst, R.J. Puigh and R.L. White, Nucl. Phys. A222 (1974) 173
- 1974KI1B Kirson, Ann. Phys. 82 (1974) 345
- 1974KL12 F.J. Kline, H. Crannell, J.M. Finn, P.L. Hallowell, J.T. O'Brien, C.W. Werntz, S.P. Fivozinsky, J.W. Lightbody and Jr., S. Penner, Nuovo Cim. A23 (1974) 137

- 1974KO07 V.F. Kosmach, A.G. Martirosova and V.I. Ostroumov, *Yad. Fiz.* 19 (1974) 241; *Sov. J. Nucl. Phys.* 19 (1974) 117
- 1974KO1G Kohler and Bethge, *Proc. Conf. Reactions between Complex Nuclei*, Vol. 1 (1974) 35
- 1974KO1P Kohlmeyer et al., *Proc. Conf. Reactions between Complex Nuclei*, Vol. 1 (1974) 7
- 1974KO25 D.G. Kovar, W. Henning, B. Zeidman, Y. Eisen and H.T. Fortune, *Phys. Rev. Lett.* 33 (1974) 1611
- 1974KO38 I. Kohno, *Sci. Pap. Inst. Phys. Chem. Res.* 68 (1974) 38
- 1974KR09 A. Kropf and H. Paul, *Z. Phys.* 267 (1974) 129
- 1974KU1F Kuo, *Ann. Rev. Nucl. Sci.* 24 (1974) 101
- 1974LE25 T.-S.H. Lee and J.D. Vergados, *Phys. Lett.* B53 (1974) 125
- 1974LI15 B.J. Lieb and H.O. Funsten, *Phys. Rev.* C10 (1974) 1753
- 1974LI1J Lisowski, Walter, Clegg and Busch, *Bull. Amer. Phys. Soc.* 19 (1974) 477
- 1974LI1K Lisowski et al., *Bull. Amer. Phys. Soc.* 19 (1974) 1076
- 1974MA12 J.A. Macdonald, J. Cerny, J.C. Hardy, H.L. Harney, A.D. Bacher and G.R. Plattner, *Phys. Rev.* C9 (1974) 1694
- 1974MI1F Miljanic, Zabel and Phillips, *Bull. Amer. Phys. Soc.* 19 (1974) 547
- 1974MU13 N.C. Mukhopadhyay and F. Cannata, *Phys. Lett.* B51 (1974) 225
- 1974NE18 V.G. Neudachin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2010; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 1
- 1974NI1A N.N. Nikolaev and M.Zh. Shmatikov, *Phys. Lett.* B52 (1974) 293
- 1974NO01 J.W. Noe, D.P. Balamuth and R.W. Zumuhle, *Phys. Rev.* C9 (1974) 132
- 1974NO1G Noe, Thesis, Univ. of Pennsylvania (1974)
- 1974OD01 Y. Oda and M. Murakami, *J. Phys. Soc. Jpn.* 37 (1974) 299
- 1974PE08 D.C. Peaslee and C.F. Williamson, *Phys. Rev.* C10 (1974) 1229
- 1974PI05 M. Pignanelli, S. Micheletti, I. Iori, P. Guazzoni, F.G. Resmini and J.L. Escudie, *Phys. Rev.* C10 (1974) 445
- 1974PI1C Pintar, Thesis, Univ. of Kansas (1974)
- 1974RAZD V.K. Rasmussen, *Bull. Amer. Phys. Soc.* 19 (1974) 999, BB12
- 1974RO04 H.H. Rossner, G. Hinderer, A. Weidinger and K.A. Eberhard, *Nucl. Phys.* A218 (1974) 606
- 1974RO21 D.C. Robinson and P.H. Barker, *Nucl. Phys.* A225 (1974) 109
- 1974RU1B Rutz, Sink and Krone, *Bull. Amer. Phys. Soc.* 19 (1974) 432
- 1974SA11 F.D. Santos, *Phys. Lett.* B51 (1974) 14

- 1974SA20 F.D. Santos, Nucl. Phys. A236 (1974) 90
- 1974SE1B Serov, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 84
- 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)
- 1974SI04 B. Singh, Nucl. Phys. A219 (1974) 621
- 1974SN1B Sneden, Astrophys. J. 189 (1974) 493
- 1974ST05 A.P. Stamp, Nucl. Phys. A220 (1974) 137
- 1974ST16 G.L. Strobel, Phys. Rev. Lett. 33 (1974) 1392
- 1974ST1G G.L. Strobel, Nucl. Phys. A232 (1974) 502
- 1974ST1N Stokstad, Switkowski and Wieland, Bull. Amer. Phys. Soc. 19 (1974) 549
- 1974TA18 L. Tauscher and W. Schneider, Z. Phys. 271 (1974) 409
- 1974TH01 C. Thibault and R. Klapisch, Phys. Rev. C9 (1974) 793
- 1974VA24 J.H. van der Merwe and C.M. Villet, Phys. Rev. C10 (1974) 1498
- 1974VAZK A. van der Woude, R.J. de Meijer, W.J. Ockels, E. Brandenburg and L.W. Put, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 1 (1974) 190
- 1974VAZO N. Van Sen, H. Muller, J.C. Gondrand, A. Fiore and G. Perrin, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 1 (1974) 180
- 1974VE05 G.A. Vershinin and P.A. Cherdantsev, Yad. Fiz. 19 (1974) 1019; Sov. J. Nucl. Phys. 19 (1974) 522
- 1974VE10 J.D. Vergados, Phys. Lett. B53 (1974) 151
- 1974WI02 D.H. Wilkinson and B.E.F. Macefield, Nucl. Phys. A232 (1974) 58
- 1974WI1F Williams and Edmunds, Sci. Prog. 61 (1974) 323
- 1974YOZO D.H. Youngblood, J.M. Moss, C.M. Rozsa, J.D. Bronson and A.D. Bacher, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974, Vol. 2 (1974) 294
- 1974ZA05 M. Zaider, J. Alster, D. Ashery, S. Cochavi, M.A. Moinester and A.I. Yavin, Phys. Rev. C10 (1974) 938
- 1974ZA1G L. Zamick, Phys. Lett. B51 (1974) 113
- 1974ZU1A Zuker, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974 (1974) 115
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975AL19 J.M. Alexander, H. Delagrange and A. Fleury, Phys. Rev. C12 (1975) 149
- 1975AR02 E.D. Arthur, W.C. Lam, J. Amato, D. Axen, R.L. Burman, P. Fessenden, R. Macek, J. Oostens, W. Shlaer, S. Sobottka et al., Phys. Rev. C11 (1975) 332
- 1975AR14 A.G. Artyukh, V.V. Volkov, G.F. Gridnev and V.L. Mikheev, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2; Bull. Acad. Sci. USSR Phys. Ser. 39 (1975) 1
- 1975AU1D Audouze, Lequeux and Vigroux, Astron. Astrophys. 43 (1975) 71

- 1975BA52 H.W. Baer, J.A. Bistirlich, N. de Botton, S. Cooper, K.M. Crowe, P. Truol and J.D. Vergados, Phys. Rev. C12 (1975) 921
- 1975BE01 F.D. Becchetti, L.T. Chua, J. Janecke and A.M. Vander Molen, Phys. Rev. Lett. 34 (1975) 225
- 1975BE1F Berman and Fultz, Rev. Mod. Phys. 47 (1975) 713
- 1975BE1U Beranger, David, Garcia and Lucas, Rev. Phys. Appl. 10 (1975) 87
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BE60 B.L. Berman, At. Data Nucl. Data Tables 15 (1975) 319
- 1975BO1W Bondouk, Ann. Phys. (Leipzig) 32 (1975) 261
- 1975BO35 J. Bommer, M. Ekpo, H. Fuchs, K. Grabisch and H. Kluge, Nucl. Phys. A251 (1975) 257
- 1975BR1F Brown, At. Data Nucl. Data Tables 15 (1975) 111
- 1975BU1M Bueno de Mesquita, Bijl and Linddner, Int. J. Appl. Rad. Isotopes 26 (1975) 551
- 1975CA1H Cannata, Lett. Nuovo Cim. 13 (1975) 319
- 1975CA1N P.J. Camillo, P.M. Fishbane and J.S. Trefil, Phys. Rev. Lett. 34 (1975) 622
- 1975CE04 F.E. Cecil, J.R. Shepard, R.E. Anderson, R.J. Peterson and P. Kaczkowski, Nucl. Phys. A255 (1975) 243
- 1975CH1V Chwieroth and Philpott, Bull. Amer. Phys. Soc. 20 (1975) 667
- 1975CH22 P. Christillin, A. Dellafiore and M. Rosa-Clot, Phys. Rev. C12 (1975) 691
- 1975CO1Q Comte, Astron. Astrophys. 39 (1975) 197
- 1975DE09 G. Delic, Phys. Rev. Lett. 34 (1975) 1468
- 1975DE1D Deutsch et al., AIP Conf. Proc. 26 (1975) 445
- 1975DI04 H. Dirim, J.P. Elliott and J.A. Evans, Nucl. Phys. A244 (1975) 301
- 1975DW1A Dwyer and Meyer, Bull. Amer. Phys. Soc. 20 (1975) 710
- 1975EI1B Eisenberg, AIP Conf. Proc. 26 (1975) 17
- 1975EN1A Endal, Astrophys. J. 195 (1975) 187
- 1975FA1A Fagg, Rev. Mod. Phys. 47 (1975) 683
- 1975FO01 H.T. Fortune, H.G. Bingham, D.J. Crozier and J.N. Bishop, Phys. Rev. C11 (1975) 302
- 1975FU01 C.B. Fulmer and D.A. Goldberg, Phys. Rev. C11 (1975) 50
- 1975GA17 A. Gamp, P. Braun-Munzinger, C.K. Gelbke, H.L. Harney, H.G. Bohlen, W. Bohne, K.D. Hildenbrand, J. Kuzminski, W. Von Oertzen and I. Tserruya, Nucl. Phys. A250 (1975) 341
- 1975GO1Y Gottlieb et al., Astrophys. J. 200 (1975) L147

- 1975GR12 K.A. Gridnev and E.F. Hefter, *Z. Phys.* A273 (1975) 99
- 1975GU10 K. Gupta, A.R. Chowdhury and T. Roy, *J. Phys. (London)* G1 (1975) 646
- 1975HA45 J.C. Hardy and I.S. Towner, *Nucl. Phys.* A254 (1975) 221
- 1975HE02 H.T. Heaton, II, J.L. Menke, R.A. Schrack and R.B. Schwartz, *Nucl. Sci. Eng.* 56 (1975) 27
- 1975HU14 J. Hufner, K. Schafer and B. Schurmann, *Phys. Rev.* C12 (1975) 1888
- 1975HU1D Hufner, *Phys. Rept.* C21 (1975) 1
- 1975IB1A Ibragimov and Kocharov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 287
- 1975JA10 J. Janecke, F.D. Becchetti, L.T. Chua and A.M. VanderMolen, *Phys. Rev.* C11 (1975) 2114
- 1975JA1A Jaros et al., *Bull. Amer. Phys. Soc.* 20 (1975) 666
- 1975JA1F Jaschek and Jaschek, *Astron. Astrophys.* 36 (1975) 401
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, *Phys. Rev.* C11 (1975) 2049
- 1975KA07 P.J. Karol, *Phys. Rev.* C11 (1975) 1203
- 1975KA1L Kaminisi, Arai and Yoshinaga, *Prog. Theor. Phys.* 53 (1975) 1855
- 1975KA26 K. Katori and K. Furuno, *Nucl. Instrum. Meth.* 127 (1975) 381
- 1975KE1A Kerridge, *Science* 188 (1975) 162
- 1975KO06 I. Kohno, *J. Phys. Soc. Jpn.* 38 (1975) 945
- 1975KU01 D. Kurath and D.J. Millener, *Nucl. Phys.* A238 (1975) 269
- 1975KU22 S. Kubono, S. Kato, M. Yasue, K. Koyama and H. Ohnuma, *J. Phys. Soc. Jpn.* 39 (1975) 1407
- 1975LE1K Lefevre, Davis and Anderson, *Bull. Amer. Phys. Soc.* 20 (1975) 584
- 1975LI11 D.A. Lind, R.J. Peterson and H.W. Fielding, *Phys. Rev.* C11 (1975) 2099
- 1975ME1F Medina et al., *Nuovo Cim.* A28 (1975) 297
- 1975MI01 S. Micheletti, M. Pignanelli and P. Guazzoni, *Phys. Rev.* C11 (1975) 64
- 1975MI12 D.J. Millener and D. Kurath, *Nucl. Phys.* A255 (1975) 315
- 1975MO1L Morinigo and Scott, *Bull. Amer. Phys. Soc.* 20 (1975) 1160
- 1975MO1M Mondal, Basak, Kasim and Husain, *Nuovo Cim.* A28 (1975) 42
- 1975MO23 P.J. Moffa, J.P. Vary, C.B. Dover, C.W. Towsley, R.G. Hanus and K. Nagatani, *Phys. Rev. Lett.* 35 (1975) 992
- 1975MU07 J.J. Murphy II, Y.M. Shin and D.M. Skopik, *Nucl. Phys.* A246 (1975) 221
- 1975NA11 K.G. Nair, C.W. Towsley, R. Hanus, M. Hamm and K. Nagatani, *Phys. Rev. Lett.* 35 (1975) 838

- 1975NA15 K.G. Nair, H. Voit, C.W. Towsley, M. Hamm, J.D. Bronson and K. Nagatani, Phys. Rev. C12 (1975) 1575
- 1975NO1D Norgaard and Arnould, Astron. Astrophys. 40 (1975) 331
- 1975NO1F Noe, Gordon and Harakeh, Bull. Amer. Phys. Soc. 20 (1975) 628
- 1975OB01 D.L. Oberg, D. Bodansky, D. Chamberlin and W.W. Jacobs, Phys. Rev. C11 (1975) 410
- 1975OL1A Olivier, McMillan and Pierce, Nucl. Instrum. Meth. 124 (1975) 289
- 1975PA18 P. Paul, H.S. Kuan and E.K. Warburton, Nucl. Phys. A254 (1975) 1
- 1975PE1E Peimbert, Ann. Rev. Astron. Astrophys. 13 (1975) 113
- 1975PO10 J.E. Poth, M.W. Sachs and D.A. Bromley, Nuovo Cim. A28 (1975) 215
- 1975PR1B Preszler, Kish, lezniak and Webber, Bull. Amer. Phys. Soc. 20 (1975) 711
- 1975RA1M Ramaty, Bull. Amer. Phys. Soc. 20 (1975) 680
- 1975RA22 V.K. Rasmussen and F.R. Metzger, Phys. Rev. C12 (1975) 706
- 1975RA33 L. Ray, G. Westfall, S.A.A. Zaidi and W.R. Coker, Phys. Lett. B59 (1975) 427
- 1975RA37 S. Raman, T.A. Walkiewicz and H. Behrens, At. Data Nucl. Data Tables 16 (1975) 451
- 1975RE01 A. Reitan, Nucl. Phys. A237 (1975) 465
- 1975RE08 L.P. Remsberg and D.G. Perry, Phys. Rev. Lett. 35 (1975) 361
- 1975RO20 C. Rolfs and W.S. Rodney, Nucl. Phys. A250 (1975) 295
- 1975RY1A Ryter, Cesarsky and Ajdouze, Astrophys. J. 198 (1975) 103
- 1975SC18 W. Schutz, Z. Phys. A273 (1975) 69
- 1975SC1H J.M. Scalo, K.H. Despain and R.K. Ulrich, Astrophys. J. 196 (1975) 805
- 1975SE03 H.P. Seiler, R. Kulessa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 151
- 1975SE04 H.P. Seiler, R. Kulessa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 159
- 1975SE07 F. Seiler, Nucl. Phys. A244 (1975) 236
- 1975SH1N Shapiro, Silberberg and Tsao, Bull. Amer. Phys. Soc. 20 (1975) 710
- 1975SM02 L.G. Smith and A.H. Wapstra, Phys. Rev. C11 (1975) 1392
- 1975SN1A Sneden and Lambert, Monthly Notices Royal Astron. Soc. 170 (1975) 533
- 1975SP04 J. Spuller, M.D. Hasinoff, S.T. Lim, D.F. Measday and T.J. Mulligan, Nucl. Phys. A248 (1975) 276
- 1975TA02 T. Takemasa, Phys. Lett. B55 (1975) 28

- 1975TE1A Tenhaken and Quin, Bull. Amer. Phys. Soc. 20 (1975) 694
- 1975TR07 S. Tryti, T. Holtebekk and F. Ugletveit, Nucl. Phys. A251 (1975) 206
- 1975TR1A Trimble, Rev. Mod. Phys. 47 (1975) 877
- 1975UD01 T. Udagawa, T. Tamura and K.S. Low, Phys. Rev. Lett. 34 (1975) 30
- 1975UL1A Ulrich, Science 190 (1975) 619
- 1975VA01 A. Vander Molen, F.D. Becchetti, J. Janecke and L. Chua, Phys. Rev. C11 (1975) 734
- 1975VA06 L. van der Zwan and K.W. Geiger, Nucl. Phys. A246 (1975) 93
- 1975VA1H Van der Woude, de Meijer and Brandenburg, Bull. Amer. Phys. Soc. 20 (1975) 597
- 1975VE01 J.D. Vergados, Nucl. Phys. A239 (1975) 271
- 1975VE05 J.D. Vergadosk, Phys. Rev. C12 (1975) 1278
- 1975VO1B Von Oertzen and Bohlen, Phys. Rept. 19 (1975) 1
- 1975WH03 R.L. White, L.A. Charlton and K.W. Kemper, Phys. Rev. C12 (1975) 1918
- 1975WI04 S.J. Wilson, Phys. Rev. C11 (1975) 1071
- 1975WI1E Wilkinson, Proc. Int. Symp. on Interaction Studies in Nucl., Germany, 1975 (1975) 147
- 1975WI1K Wilkinson, Nature 257 (1975) 189
- 1975ZA06 N.I. Zaika and P.L. Shmarin, Ukr. Fiz. Zh. 20 (1975) 238
- 1975ZE1C Zeller et al., Bull. Amer. Phys. Soc. 20 (1975) 718
- 1976DA1U Dayras, Switkowski and Tombrello, LAP-147 (1976)
- 1976GOZX C.D. Goodman, W.R. Wharton and D.C. Hensley, Bull. Amer. Phys. Soc. 21 (1976) 82, HE2
- 1976JA04 N.P. Jacob, Jr. and S.S. Markowitz, Phys. Rev. C13 (1976) 754
- 1976LO01 B.A. Logan and A. Ljubicic, Phys. Rev. C13 (1976) 361
- 1976MA04 F. Malaguti and P.E. Hodgson, Nucl. Phys. A257 (1976) 37
- 1976MO03 T. Motobayashi, I. Kohno, K. Katori, M. Yoshie, T. Ohi and H. Kamitsubo, Phys. Rev. Lett. 36 (1976) 390
- 1976PA03 K.-H. Passler, Nucl. Phys. A257 (1976) 253
- 1976YOZZ G.R. Young, E.B. Blum, E.R. Cosman, A.J. Lazzarini, S.G. Steadman and N. Tsoupas, Bull. Amer. Phys. Soc. 21 (1976) 30, BE10
- 1977AJ02 F. Ajzenberg-Selove, Nucl. Phys. A281 (1977) 1
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- HA70P Unknown Source

