

Energy Levels of Light Nuclei $A = 18$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18$ – 20 was published in *Nuclear Physics A475* (1987), p. 1. This version of $A = 18$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed June 1, 1987)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 18$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below. The introductory [Table 2](#) is also available via the link.

A. Nuclides: [\$^{18}\text{He}\$](#) , [\$^{18}\text{Be}\$](#) , [\$^{18}\text{B}\$](#) , [\$^{18}\text{C}\$](#) , [\$^{18}\text{N}\$](#) , [\$^{18}\text{O}\$](#) , [\$^{18}\text{F}\$](#) , [\$^{18}\text{Ne}\$](#) , [\$^{18}\text{Na}\$](#) , [\$^{18}\text{Mg}\$](#) , etc.

B. Tables of Recommended Level Energies:

[Table 18.1](#): Energy levels of ^{18}C

[Table 18.2](#): Energy levels of ^{18}N

[Table 18.3](#): Energy levels of ^{18}O

[Table 18.13](#): Energy levels of ^{18}F

[Table 18.22](#): Energy levels of ^{18}Ne

C. [References](#)

D. Figures: [\$^{18}\text{O}\$](#) , [\$^{18}\text{F}\$](#) , [\$^{18}\text{Ne}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{18}He

(Not illustrated)

Not observed: see (1982AV1A, 1983ANZQ; theor.).

 ^{18}Be

(Not illustrated)

^{18}Be has not been observed. It is predicted to have a mass excess of 78.43 MeV: see (1978AJ03). ^{18}Be is then unstable with respect to breakup into $^{16}\text{Be} + 2n$, $^{15}\text{Be} + 3n$, $^{14}\text{Be} + 4n$, $^{13}\text{Be} + 5n$, $^{12}\text{Be} + 6n$, $^{11}\text{Be} + 7n$ and $^{10}\text{Be} + 8n$ by, respectively, 3.1, 3.0, 6.0, 3.1, 4.9, 1.8 and 1.3 MeV, using the masses for the residual nuclei adopted by (1985WA02, 1986AJ01, 1986AJ04). See also (1983ANZQ; theor.).

 ^{18}B

(Not illustrated)

^{18}B has not been observed in the bombardment of Ta by 44 MeV/A Ar ions (1985DE60, 1985LA03, 1986PO13) or in the bombardment of Be by 12 MeV/A ^{56}Fe ions (1984MU27). ^{18}B has been predicted to have a mass excess of 53.85 MeV. It would then be unstable with respect to $^{17}\text{B} + n$ by 1.8 MeV: see (1978AJ03, 1985WA02). ^{18}B is calculated to have $J^\pi = 4^-$ and to have excited states at 0.62, 0.86 and 1.59 MeV with $J^\pi = 1^-, 2^-$ and 2^- (1985PO10). See also (1985AN1B) and (1983ANZQ; theor.).

 ^{18}C

(Fig. 4)

GENERAL: (See also (1978AJ03, 1983AJ01).)

See (1983ANZQ, 1983FR1A, 1983WI1A, 1984SA37, 1986AN07, 1986AV1B, 1986GU1D, 1987SA15).

Mass of ^{18}C : The atomic mass excess of ^{18}C is 24.923 ± 0.030 MeV, based on the Q -value of the $^{48}\text{Ca}(^{18}\text{O}, ^{18}\text{C})^{48}\text{Ti}$ reaction. ^{18}C is then bound by 4.187 MeV with respect to breakup into $^{17}\text{C} + n$ (1982FI10). See also (1986PI09).

$$1. \ ^{18}\text{C}(\beta^-)^{18}\text{N} \quad Q_m = 11.806$$

The half-life of ^{18}C , $\tau_{1/2} = 66_{-15}^{+25}$ msec, $P_n \approx 25\%$ (1987MU1J; prelim.). See also (1984KL06; theor.).

Table 18.1: Energy levels of ^{18}C

| E_x in ^{18}C (MeV \pm keV) | $J^\pi; T$ | Decay | Reactions |
|---|------------|-------------|-----------|
| 0 | $(0^+); 3$ | (β^-) | 2, 3 |
| 1.62 ± 20 | $(2^+); 3$ | | 2, 3 |

2. $^{18}\text{O}(\pi^-, \pi^+)^{18}\text{C}$ $Q_m = -25.705$

The angular distribution of the π^+ to the ground state of ^{18}C has been measured at $E_{\pi^-} = 164$ MeV by (1984GI10) [see also for excitation function at $\theta = 5^\circ$ for $E_{\pi^-} \approx 140$ to 240 MeV]. There is also some indication of the population of an excited state at $E_x = 1.55$ MeV (1984GI10). See also (1983AJ01).

3. $^{48}\text{Ca}(^{18}\text{O}, ^{18}\text{C})^{48}\text{Ti}$ $Q_m = -21.434$
 $Q_0 = -21434 \pm 30$ keV (1982FI10).

At $E(^{18}\text{O}) = 112$ MeV the ground state and an excited state at 1.62 ± 0.02 MeV are observed by (1982FI10). See also (1983AJ01).

^{18}N
(Figs. 1 and 4)

GENERAL: (See also (1983AJ01).)

See (1981NAZQ, 1983ANZQ, 1983FR1A, 1983SH44, 1983WI1A, 1984AS1D, 1984HI1A, 1986AN07, 1986BI1A, 1986HA1B, 1986MA48, 1986ME1F, 1987RI03).

Mass of ^{18}N : The atomic mass excess derived from the Q -value of the $^{18}\text{O}(^7\text{Li}, ^7\text{Be})^{18}\text{N}$ reaction is 13.117 ± 0.020 MeV (1983PU01). ^{18}N is then stable with respect to breakup into $^{17}\text{N} + n$ by 2.825 MeV. See (1983AJ01) for the earlier work.

1. $^{18}\text{N}(\beta^-)^{18}\text{O}$ $Q_m = 13.899$

The half-life of ^{18}N is 0.624 ± 0.012 sec (1982OL01). The decay branches are displayed in Table 18.9. The nature of the decay leads to $J^\pi = 1^-$ for the ^{18}N ground state (1982OL01). See also (1983SN03).

Table 18.2: Energy levels of ^{18}N

| E_x (MeV \pm keV) | $J^\pi; T$ | $\tau_{1/2}$ (msec) | Decay | Reactions |
|-----------------------|------------|---------------------|-----------|---------------|
| 0 | $1^-; 2$ | 624 ± 12 | β^- | 1, 3, 4, 5, 6 |
| 0.121 ± 10 | 2^-^a | | | 3, 4, 6 |
| 0.575 ± 25 | 2^-^a | | | 3, 6, 7 |
| 0.747 ± 10 | 3^-^a | | | 6 |
| b | | | | |
| 2.21 | | | | 6 |
| 2.42 | | | | 6 |

^a Suggested by (1984BA24). See also (1982OL01).

^b See (1984BA24) for a calculation suggesting additional states.

2. $^{14}\text{C}(^7\text{Li}, ^3\text{He})^{18}\text{N}$ $Q_m = -10.122$

The work described in (1983AJ01) has not been published.

3. $^{14}\text{C}(^{18}\text{O}, ^{14}\text{N})^{18}\text{N}$ $Q_m = -13.743$

At $E(^{18}\text{O}) = 92.2$ MeV groups are observed to the ground state of ^{18}N (unresolved) and to an excited state at $E_x = 575 \pm 25$ keV (1980NA14).

4. $^{18}\text{O}(\pi^-, \pi^0)^{18}\text{N}$ $Q_m = -9.295$

See (1983AS01, 1984AS05).

5. $^{18}\text{O}(t, ^3\text{He})^{18}\text{N}$ $Q_m = -13.881$

See (1983AJ01).

6. $^{18}\text{O}(^7\text{Li}, ^7\text{Be})^{18}\text{N}$ $Q_m = -14.761$
 $Q_0 = -14761 \pm 20$ keV (1983PU01).

At $E(^7\text{Li}) = 52$ MeV, ^7Be groups are observed corresponding to the excitation of the states displayed in Table 18.2 (1983PU01).

7. $^{18}\text{O}(^{11}\text{B}, ^{11}\text{C})^{18}\text{N}$ $Q_m = -15.881$

See (1983PU01).

¹⁸O

(Figs. 1 and 4)

GENERAL: (See also (1983AJ01).)

Shell model: (1978WI1B, 1982ZH01, 1983BR29, 1983WA17, 1983KI13, 1984AS07, 1984BA24, 1984CH1V, 1985LE1K, 1985SA29, 1986HAZY, 1986YU1B, 1987WA1H).

Cluster, collective and deformed models: (1983BE1W, 1983CL1B, 1983ME12, 1984AS04, 1984BA55, 1984GA1T, 1984SA37, 1985BA1A, 1985DE24, 1985RO1G, 1985SU10, 1986CA1T, 1986KA1D, 1986YU1B).

Special states: (1978WI1B, 1982BA62, 1982BE1Z, 1982ZH01, 1983BE1W, 1983BR29, 1983LI10, 1983ME12, 1983WA17, 1983KI13, 1984AS07, 1984BA24, 1984BA55, 1984CH1V, 1984GA1T, 1984GE1A, 1984HA14, 1984SA37, 1984ST1E, 1984WI17, 1985BR20, 1985CA1Q, 1985DE24, 1985LE1K, 1985LE1L, 1985MA1Q, 1985MA1R, 1985RO1G, 1985SA29, 1985SO12, 1985SU10, 1985YU1B, 1986AN10, 1986AN07, 1986CA1T, 1986CA27, 1986GU1A, 1986KA1D, 1986YU1B, 1987WA1H, 1987YA03).

Electromagnetic transitions and giant resonances: (1982AL30, 1982BA62, 1982BR24, 1982RI04, 1983BR29, 1983IS1F, 1983KA28, 1983KI13, 1984AS04, 1984AS07, 1984BA55, 1984BR1L, 1984GA1T, 1984HA14, 1984ST02, 1985AL21, 1985DE24, 1985LE1L, 1985SU10, 1985WI17, 1986AN10, 1986CA27, 1986ER1A, 1986HAZY).

Astrophysical questions: (1981WA1Q, 1982BU1A, 1982CA1A, 1982RO1A, 1982WI1B, 1982WO1A, 1983CO1K, 1984BL1J, 1984HA1R, 1984HA1Z, 1984LA1J, 1985HA1Z, 1985HA1R, 1986CO2C, 1986DO1L, 1986LA1C, 1986WI1L, 1987HA1C, 1987HA1D, 1987PR1A, 1987WI11).

Applications: (1982MA1Q, 1982PI1H, 1983AM1A, 1983AM1D, 1983GR1L, 1983KI1D, 1983KU1C, 1983LI1T, 1983TA1P, 1984LE1G, 1984SH2A, 1985WA1R, 1985YO1B, 1985ZA1A, 1986CO2B, 1986DR1E, 1986DU1Q, 1986EN1A, 1986HO1Q, 1986HO1L, 1986NI1C, 1986XU1C).

Complex reactions involving ¹⁸O: (1981IC02, 1982MO1K, 1982SA1Q, 1983DE26, 1983EN04, 1983FR17, 1983FR1A, 1983GR10, 1983GR28, 1983LI10, 1983OL1A, 1983SA06, 1983VA23, 1983WI1A, 1984GR08, 1984HI1A, 1984HO23, 1984NA1D, 1985DE60, 1985GAZT, 1985HO05, 1985KAZQ, 1985MA1Q, 1985PO11, 1986AN1F, 1986HA1B, 1986IR01, 1986MA19, 1986ME06, 1986PO06, 1986SA30, 1986SC28, 1986SC29, 1986SO10, 1986VAZZ, 1987LI04, 1987RI03).

Muon capture: (1983GM1A, 1987SU06).

Pion capture and reactions (See also reaction 23): (1981SEZR, 1982BI08, 1982LI15, 1982LI1M, 1982LI1N, 1983AM1C, 1983AS01, 1983BE1W, 1983GE12, 1983HO02, 1983JO06, 1983LI08, 1983LI1U, 1983MA16, 1983ME1H, 1983OS1E, 1983RO07, 1983SE16, 1983ZE1C, 1984AL20, 1984AS05, 1984BL17, 1984GR27, 1984HU1C, 1984JO01, 1984KA26, 1984SC09, 1984TR15, 1985AL15, 1985BI01, 1985CA1Q, 1985CH23, 1985GI01, 1985GM01, 1985KE1A, 1985SE08, 1985SE21, 1986BA1C, 1986FO06, 1986GE06, 1986GM01, 1986RO03, 1986SCZX, 1986SI11, 1987TE01).

Table 18.3: Energy levels of ^{18}O ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m^b or $\Gamma_{c.m.}$ (keV) | Decay | Reactions |
|--------------------------|------------|---|----------|--|
| 0 | $0^+; 1$ | — | stable | 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 |
| 1.98207 ± 0.09 | 2^+ | $\tau_m = 2.80 \pm 0.07$ psec $g = -0.287 \pm 0.015$ | γ | 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 15, 17, 18, 19, 22, 23, 24, 25, 26, 27, 29, 30, 36, 37, 39, 41, 42, 44, 45, 46, 47, 48 |
| 3.55484 ± 0.40 | 4^+ | $\tau_m = 24.8 \pm 1.2$ psec $g = -0.62 \pm 0.10$ | γ | 2, 3, 6, 8, 9, 13, 14, 15, 17, 18, 19, 22, 25, 30, 36, 37, 47, 48 |
| 3.63376 ± 0.11 | 0^+ | $\tau_m = 1.38 \pm 0.16$ psec | γ | 2, 3, 6, 8, 9, 13, 17, 19, 22, 25, 30, 36, 37, 46, 47, 48 |
| 3.92044 ± 0.14 | 2^+ | 26.5 ± 2.9 fsec | γ | 2, 3, 6, 8, 9, 13, 17, 19, 22, 25, 30, 36, 47 |
| 4.45554 ± 0.10 | 1^- | 65 ± 15 fsec | γ | 2, 3, 6, 8, 9, 13, 17, 19, 25, 30, 36, 37, 46, 47 |
| 5.09778 ± 0.54 | 3^- | 62 ± 25 fsec | γ | 2, 3, 6, 8, 9, 13, 17, 19, 22, 23, 24, 25, 30, 36, 37, 42, 47, 48 |
| 5.2604 ± 1.2 | 2^+ | 10.1 ± 0.5 fsec | γ | 2, 3, 6, 8, 9, 13, 15, 17, 22, 25, 30, 46, 47 |
| 5.3364 ± 0.6 | 0^+ | 200 ± 40 fsec | γ | 2, 3, 8, 13, 17, 22, 30, 47 |
| 5.3778 ± 1.2 | 3^+ | < 30 fsec | γ | 2, 3, 13, 17, 47 |
| 5.53024 ± 0.29 | 2^- | < 25 fsec | γ | 2, 3, 13, 19, 25, 30, 47 |
| 6.19822 ± 0.40 | 1^- | 3.7 ± 0.6 fsec | γ | 2, 3, 8, 13, 17, 19, 21, 30, 47 |
| 6.3513 ± 0.6 | (2^-) | < 35 fsec | γ | 2, 3, 13, 17, 19, 30, 47, 48 |
| 6.4044 ± 1.2 | 3^- | 30 ± 15 fsec | γ | 2, 3, 13, 30, 47 |
| 6.88045 ± 0.27 | 0^- | < 25 fsec | γ | 2, 3, 13, 19, 30, 46, 47 |

Table 18.3: Energy levels of ^{18}O ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m^b or $\Gamma_{c.m.}$ (keV) | Decay | Reactions |
|--------------------------|--------------|--|---------------------|---|
| 7.1169 ± 1.2 | 4^+ | < 25 fsec | γ, α | 2, 3, 6, 8, 9, 13, 15, 17, 22, 25, 30, 34, 36, 37, 47 |
| 7.619 ± 3 | 1^- | < 2.5 | | 2, 3, 6, 8, 13, 30, 34, 36, 37, 47 |
| 7.77107 ± 0.50 | 2^- | | γ | 2, 3, 13, 19, 47 |
| 7.864 ± 5 | 5^- | | γ | 2, 3, 6, 8, 9, 13, 17, 22, 30, 34, 36, 37, 47, 48 |
| 7.977 ± 4 | $(3^+, 4^-)$ | | γ | 2, 3, 13, 17, 47 |
| 8.039 ± 2 | 1^- | < 2.5 | γ, α | 2, 3, 6, 7, 13, 14, 15, 34, 36, 37, 47 |
| 8.125 ± 2 | 5^- | | γ, α | 2, 3, 6, 8, 9, 13, 47 |
| 8.213 ± 4 | 2^+ | 1.0 ± 0.8 | γ, n, α | 2, 3, 6, 7, 13, 25, 30, 34, 36, 37, 47 |
| 8.282 ± 3 | 3^- | 8 ± 1 | γ, n, α | 2, 3, 6, 7, 8, 9, 13, 30, 47 |
| 8.410 ± 8 | | 8 ± 6 | n, α | 7, 13, 47 |
| 8.521 ± 6 | | | | 13, 47 |
| 8.660 ± 6 | | | | 13, 47 |
| 8.817 ± 12 | (1^+) | 70 ± 12 | n, α | 7, 25, 30 |
| 8.955 ± 4 | | 43 ± 3 | n, α | 7, 13, 30 |
| 9.03 | | | | 13, 17, 30 |
| (9.10) | | | | 30 |
| 9.361 ± 6 | (3^-) | 27 ± 15 | γ, n, α | 7, 9, 13, 22, 30, 34, 36, 37 |
| 9.414 ± 18 | | ≈ 120 | n, α | 7, 9, 13, 30 |
| 9.48 ± 24 | | ≈ 65 | n, α | 7, 13 |
| 9.672 ± 7 | (3^-) | 60 ± 30 | n, α | 7, 13, 30, 34, 36, 37 |
| 9.713 ± 7 | | | | 13, 30 |
| 9.890 ± 11 | | ≈ 150 | n, α | 7, 13, 30 |
| 10.118 ± 10 | 3^- | 16 ± 4 | n, α | 7, 8, 13, 30 |
| 10.295 ± 14 | 4^+ | | n, α | 7, 8, 9, 13, 14, 30, 34, 36, 37 |
| 10.396 ± 9 | 3^- | | n, α | 7, 13, 30 |

Table 18.3: Energy levels of ^{18}O ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m^b or $\Gamma_{c.m.}$ (keV) | Decay | Reactions |
|--------------------------|------------------------|--|------------------------|-----------------------------|
| 10.595 \pm 15 | | | n, α | 7, 13 |
| 10.82 \pm 20 | | | n, α | 7 |
| 10.91 \pm 20 | | | n, α | 7, 9 |
| 10.99 \pm 20 | | | n, α | 7 |
| 11.13 \pm 20 | | | n, α | 7, 9, 46 |
| 11.39 \pm 20 | (2 ⁺) | | n, α | 7, 8 |
| 11.41 \pm 20 | (4 ⁺) | | n, α | 7, 8 |
| 11.62 \pm 20 | 5 ⁻ | 76 \pm 8 | γ , n, α | 7, 8, 9, 22, 30, 34, 36, 37 |
| 11.69 \pm 20 | 6 ⁺ | | n, α | 7, 8, 9, 30 |
| 11.82 \pm 20 | (3 ⁻) | | n, α | 7 |
| 12.04 \pm 20 | (2 ⁺) | 28 \pm 6 | γ , n, α | 7, 8, 22 |
| 12.25 \pm 20 | (1 ⁻) | | n, α | 7, 8 |
| 12.33 \pm 20 | 5 ⁻ | | n, α | 7, 8, 9 |
| 12.50 \pm 20 | 4 ⁺ | | n, α | 7, 34, 36, 37 |
| 12.53 \pm 20 | 6 ⁺ | | n, α | 7, 8, 9, 34, 36, 37 |
| 13.1 ^c | 1 ⁻ | 700 | γ , n | 20 |
| 13.8 | 1 ⁻ | 600 | γ , n | 20 |
| 14.7 | 1 ⁻ | 800 | γ , n | 20 |
| 15.8 | 1 ⁻ | 700 | γ , n | 20 |
| 16.210 \pm 10 | 1 ⁽⁻⁾ | | γ | 22 |
| 16.315 \pm 10 | (3, 2) ⁻ | | γ | 22 |
| 16.399 \pm 5 | 2 ⁻ ; 2 | < 20 | γ | 22, 25 |
| 16.948 \pm 10 | (3, 2) ⁻ | | γ | 22 |
| 17.025 \pm 10 | (> 2); 2 | 20 \pm 6 | γ | 22 |
| 17.05 | (7 ⁻) | \approx 350 | | 8 |
| 17.398 \pm 10 | 1 ⁻ ; (2) | 600 | γ , n, p | 20, 22 |
| 17.450 \pm 10 | (2, 1, 3) ⁻ | | γ | 22 |
| 17.5 | | \approx 150 | γ | 22 |
| 17.502 \pm 10 | (1, 2, 3) ⁻ | | γ | 22 |
| (17.6 \pm 200) | (8 ⁺) | | | 8 |

Table 18.3: Energy levels of ^{18}O ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m ^b or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|--------------------------|---|--|----------------|-----------|
| 17.635 \pm 10 | | | γ | 22 |
| 18.049 \pm 10 | | | γ | 22 |
| 18.2 | | ≈ 150 | γ | 22 |
| 18.5 | | ≈ 4300 | γ | 22 |
| 18.70 \pm 20 | (4 ⁻); 2 | < 20 | γ | 22 |
| 18.871 \pm 5 | 1 ⁺ ; 2 | | γ | 22 |
| 18.927 \pm 10 | (1, 2 ⁺) | | γ | 22 |
| 18.95 | (7 ⁻) | ≈ 350 | | 8 |
| 19.027 \pm 10 | (1, 3) ⁻ | | γ | 22 |
| 19.150 \pm 10 | (1 ⁻ , 2 ⁺ , 3 ⁻) | | γ | 22 |
| 19.24 \pm 20 | (> 2); 2 | < 20 | γ | 22 |
| 19.4 | 1 ⁻ ; (2) | 900 | γ, p | 20 |
| 19.7 | | ≈ 200 | γ | 22 |
| 20.2 | | ≈ 180 | γ | 22 |
| 20.36 \pm 20 | (4 ⁻); 2 | < 20 | γ | 22 |
| 21.0 | 1 ⁻ ; (1) | ≈ 150 | γ, n, p | 20, 22 |
| 22.39 \pm 40 | (4 ⁻) | 74 \pm 7 | γ | 22 |
| 22.7 | 1 ⁻ | | γ, n, p | 20 |
| 23.8 | 1 ⁻ ; (1) | ≈ 1500 | γ, n, p | 20, 22 |
| 27 | 1 ⁻ ; (2) | | γ, n, p | 20 |
| 30 | | | γ, n | 20 |
| 36 | | | γ | 20 |

^a See also Tables 18.4 and 18.11 here and 18.2 in (1983AJ01).

^b See Table 18.4 in (1978AJ03) for a display of τ_m measurements.

^c For additional states with $12.9 \leq E_x \leq 23.1$ MeV see (1983CU03) [reaction 8].

K-mesons and other meson interactions: (1983MA1V, 1983TO21, 1986MO1Q, 1986MO1A).

Anti-proton interactions: (1983GR11, 1984GI11, 1984SU07, 1985DU05, 1985LI16, 1986DU10, 1986FR10, 1986RO23, 1987BA18, 1987SP05).

Hypernuclei: (1984AS1D, 1984MO1N, 1985YA1C, 1986MO1Q, 1986MO1A).

Other topics: (1978WIIB, 1981CL05, 1982CA12, 1982VE02, 1983AR1J, 1983BR29, 1983CL1B, 1983JO1C, 1983MA35, 1983SH32, 1983UE01, 1984WI17, 1985AL21, 1985AN28, 1985CH23, 1985MA56, 1985YU1B, 1986GU1A).

Ground and 1.98 MeV states of ^{18}O : (1978WIIB, 1982BA62, 1982CA12, 1982LO13, 1983ANZQ, 1983AR1J, 1984ANZW, 1984BA55, 1984FR13, 1984HA14, 1984ST1E, 1984WE04, 1985AN28, 1985BA1A, 1985DE24, 1986CA27, 1986RO03, 1987AB03, 1987SA15).

$$\text{Isotopic abundance} = (0.200 \pm 0.012)\% \text{ (1984DE53).}$$

$$\langle r^2 \rangle^{1/2} = 2.784 \pm 0.020 \text{ fm: see reaction 22.}$$

$^{18}\text{O}^*(1.98)$

$$g = -0.287 \pm 0.015 \text{ [see (1983AJ01)]}$$

$$Q = -4.2 \pm 0.8 e \cdot \text{fm}^2 \text{ [weighted mean of } -3.6 \pm 0.9 \text{ and } -5.8 \pm 1.5 e \cdot \text{fm}^2 \text{: see (1983GR28); see also (1983AJ01)].}$$

$$B(E2; 0^+ \rightarrow 2^+) = 39.0 \pm 1.8 e^2 \cdot \text{fm}^4 \text{ [(1979FE06, 1983GR10); see also (1983AJ01)];}$$

$$= 44.8 \pm 1.3 e^2 \cdot \text{fm}^4 \text{ (1982NO04);}$$

$$= 47.6 \pm 1.0 e^2 \cdot \text{fm}^4 \text{ (1982BA06); see also (1987RA01).}$$

For a discussion of the hexadecapole deformation see (1983GR10). See also (1987RA01).

- | | |
|--|-----------------|
| 1. (a) $^{10}\text{B}(^9\text{Be}, \text{p})^{18}\text{O}$ | $Q_m = 16.8917$ |
| (b) $^{11}\text{B}(^9\text{Be}, \text{d})^{18}\text{O}$ | $Q_m = 7.6621$ |

See (1986CU02) for production cross sections of 1.98 MeV γ -rays.

- | | |
|--|---------------|
| 2. $^{12}\text{C}(^7\text{Li}, \text{p})^{18}\text{O}$ | $Q_m = 8.400$ |
|--|---------------|

Observed proton groups are displayed in Table 18.5. See also (1983AJ01, 1986SM01).

- | | |
|--|----------------|
| 3. $^{13}\text{C}(^6\text{Li}, \text{p})^{18}\text{O}$ | $Q_m = 10.704$ |
|--|----------------|

Angular distributions of proton groups have been measured to the states shown in Table 18.5 at $E(^6\text{Li}) = 28 \text{ MeV}$ (1986SM01). It is found that when σ_{tot} to a particular state in ^{18}O is large in this reaction, it is also large in the $^{12}\text{C}(^7\text{Li}, \text{p})$ reaction. The relationship between the cross sections in this reaction and in reaction 1 depends on the parity of the final states (1986SM01).

Table 18.4: Radiative decays in ^{18}O ^a

| E_i (MeV) | J_i^π | E_f (MeV) | Branch (%) | δ |
|-----------------------|-----------|-------------|-----------------------------|-----------------|
| 1.98 | 2^+ | 0 | 100 | |
| 3.55 | 4^+ | 1.98 | 100 | |
| 3.63 | 0^+ | 0 | 0.30 ± 0.06 | |
| | | 1.98 | 99.70 ± 0.06 | |
| 3.92 ^{b,c,d} | 2^+ | 0 | 11.4 ± 0.5 ^e | |
| | | 1.98 | 88.6 ± 0.5 | f |
| 4.46 ^{b,c,d} | 1^- | 1.98 | 28.6 ± 1.1 | f |
| | | 3.63 | 68.9 ± 0.7 | |
| | | 3.92 | 2.5 ± 0.6 | |
| 5.10 ^{b,c,d} | 3^- | 1.98 | 76.3 ± 0.6 | f |
| | | 3.55 | 5.8 ± 0.6 | f |
| | | 3.92 | 17.9 ± 0.4 | f |
| 5.26 ^{b,d} | 2^+ | 0 | 30.3 ± 0.9 | f |
| | | 1.98 | 56.0 ± 1.0 | 0.15 ± 0.04 |
| | | 3.55 | 1.1 ± 0.6 | |
| | | 3.63 | 0.9 ± 0.6 | |
| | | 3.92 | 8.7 ± 0.4 | |
| | | 4.46 | 3.0 ± 0.3 | |
| 5.34 | 0^+ | 0 | ^g | |
| | | 1.98 | 60 ± 2 | |
| | | 4.46 | 40 ± 2 | f |
| 5.38 | 3^+ | 1.98 | 86.5 ± 2.2 | f |
| | | 3.92 | 13.5 ± 2.2 | f |
| 5.53 | 2^- | 1.98 | 49 ± 2 | f |
| | | 3.92 | 24 ± 2 | |
| | | 4.46 | 27 ± 2 | f |
| 6.20 ^{b,d} | 1^- | 0 | 88.7 ± 0.9 | |
| | | 3.63 | 2.5 ± 0.3 | |
| | | 4.46 | 4.1 ± 0.4 | |
| | | 5.26 | 3.6 ± 0.4 | |
| | | 5.34 | 1.1 ± 0.3 | |

Table 18.4: Radiative decays in ^{18}O ^a (continued)

| E_i (MeV) | J_i^π | E_f (MeV) | Branch (%) | δ |
|-----------------------|--------------|-------------|-----------------------------|----------------------|
| 6.35 | (2^-) | 1.98 | 32 ± 2 | f |
| | | 3.92 | 55 ± 2 | f |
| | | 4.46 | 12 ± 2 | f |
| 6.40 ^{b,d} | 3^- | 1.98 | 68.1 ± 1.8 | f |
| | | 3.55 | 7.4 ± 1.2 | |
| | | 3.92 | 6.3 ± 1.0 | f |
| | | 4.46 | 2.8 ± 1.0 | |
| | | 5.10 | 9.8 ± 0.9 | |
| | | 5.26 | 5.6 ± 0.9 | |
| | | 4.46 | 100 | f |
| 6.88 | 0^- | 4.46 | 100 | f |
| 7.12 ^{b,d,i} | 4^+ | 1.98 | 27.0 ± 0.5 | $-(0.052 \pm 0.035)$ |
| | | 3.55 | 70.0 ± 6.0 ^h | |
| | | 3.92 | 1.8 ± 0.4 | |
| | | 5.10 | 1.2 ± 0.3 | |
| 7.62 ⁱ | 1^- | 0 | 23 ± 2 | |
| | | 1.98 | 62 ± 3 ^j | $-(0.027 \pm 0.008)$ |
| | | 4.46 | 8 ± 1 | $-(0.21 \pm 0.03)$ |
| | | 5.34 | 6 ± 1 | |
| | | 6.20 | 1 ± 1 | |
| 7.77 | 2^- | 1.98 | 53 ± 3 | |
| | | 4.46 | 11 ± 2 | |
| | | 5.10 | 36 ± 3 | |
| 7.86 ⁱ | 5^- | 3.55 | > 75 | |
| 7.98 | $(3^+, 4^-)$ | 3.55 | 67 ± 2 | |
| | | 5.10 | 12 ± 2 | |
| | | 5.38 | 21 ± 2 | |
| 8.04 ⁱ | 1^- | 0 | 16 ± 1 | |
| | | 1.98 | 70 ± 2 ^k | |
| | | 3.63 | 10 ± 1 | |
| | | 5.26 | 4 ± 1 | |
| 8.13 ⁱ | 5^- | 3.55 | 99 ± 1 ^l | |

Table 18.4: Radiative decays in ^{18}O ^a (continued)

| E_i (MeV) | J_i^π | E_f (MeV) | Branch (%) | δ |
|-------------------|----------------|-------------|------------|----------|
| 8.21 ⁱ | 2 ⁺ | 5.10 | 1 ± 1 | |
| | | 0 | 19 ± 4 | |
| | | 1.98 | 29 ± 3 | |
| | | 3.55 | 3 ± 1 | |
| | | 3.92 | 3 ± 1 | |
| | | 4.46 | 29 ± 3 | |
| 8.28 ⁱ | 3 ⁻ | 5.10 | 17 ± 1 | |
| | | 3.55 | 61 ± 3 | |
| | | 4.46 | 3 ± 3 | |
| | | 5.26 | 36 ± 3 | |

^a For references and additional information see Tables 18.3 in (1978AJ03, 1983AJ01).

Upper limits for other transitions are not shown.

^b M. Ruscev and M. Gai, private communication.

^c Weighted means of results in Table 18.3 and in ref. ^b.

^d See also Table 3 in (1978AJ03).

^e $\Gamma_\pi/\Gamma = (3.0 \pm 0.6) \times 10^{-3}$.

^f δ is consistent with 0.

^g $\Gamma_\pi/\Gamma \leq 2.3 \times 10^{-3}$.

^h $\Gamma_\gamma/\Gamma_\alpha = 0.9 \pm 0.1$.

ⁱ Adopted by (1987GA15): see for upper limits for the branching ratio to other states.

^j $\Gamma_\alpha\Gamma_\gamma/\Gamma = 0.34$ eV.

^k $\Gamma_\alpha\Gamma_\gamma/\Gamma = 0.89$ eV.

^l $\Gamma_\alpha\Gamma_\gamma/\Gamma = 0.22$ eV.

4. $^{13}\text{C}(^9\text{Be}, \alpha)^{18}\text{O}$ $Q_m = 12.8300$

See (1983AJ01, 1983VO1A).

5. $^{13}\text{C}(^{17}\text{O}, ^{12}\text{C})^{18}\text{O}$ $Q_m = 3.098$

See (1983AJ01) and (1984AB1A; theor.).

Table 18.5: States of ^{18}O from $^{12}\text{C}(^7\text{Li}, \text{p})$ and $^{13}\text{C}(^6\text{Li}, \text{p})$

| E_x (keV) ^a | σ_{tot} ^b (μb) | σ_{tot} ^c (μb) |
|-------------------------------|--|--|
| 0 | 41.7 ± 1.7 | 6.1 ± 0.3 |
| 1982.4 ± 4.2 | 243 ± 4 | 39 ± 1 |
| $\equiv 3555.0$ | 429 ± 5 | 56 ± 1 |
| 3636.1 ± 3.5 | 58.5 ± 1.0 | 13 ± 1 |
| 3922.4 ± 3.1 | 209 ± 3 | 36 ± 1 |
| 4457.8 ± 3.0 | 164 ± 2 | 46 ± 1 |
| 5099.3 ± 3.8 | 387 ± 5 | 74 ± 1 |
| 5252.1 ± 4.9 | 353 ± 4 | 44 ± 1 |
| 5331.6 ± 3.8 | 217 ± 2 | 35 ± 1 |
| 5377.6 ± 3.6 | | |
| 5532.2 ± 3.7 | 218 ± 3 | 45 ± 1 |
| 6194.9 ± 3.8 | 103 ± 2 | 37 ± 1 |
| 6351.1 ± 4.5 ^d | 521 ± 4 | 131 ± 2 |
| 6402.1 ± 4.5 | | |
| 6888.1 ± 6.1 | 33.8 ± 0.7 | 5.3 ± 0.4 |
| 7119.8 ± 5.1 ^e | 892 ± 10 | 208 ± 2 |
| 7619.4 ± 5.0 | 124 ± 2 | 33 ± 1 |
| 7779.3 ± 4.8 ^f | 262 ± 3 | 37 ± 1 |
| 7858.7 ± 4.9 ^g | 487 ± 6 | 101 ± 1 |
| 7972.7 ± 4.8 | 380 ± 4 | 84 ± 1 |
| 8041.5 ± 5.0 | 80 ± 1 | 19 ± 1 |
| 8133.9 ± 5.3 | 500 ± 6 | 140 ± 2 |
| 8218.3 ± 5.2 | 508 ± 6 | 48 ± 1 |
| 8291.7 ± 5.6 | 368 ± 4 | 103 ± 2 |

^a (1978FO29): $^{12}\text{C}(^7\text{Li}, \text{p})^{18}\text{O}$. See also Table 18.4 in (1983AJ01).

^b σ_{tot} ($0-180^\circ$) from $^{12}\text{C}(^7\text{Li}, \text{p})$ at $E(^7\text{Li}) = 18.0$ MeV (1978FO29). Angular distributions were also measured at 16.0 MeV. The total cross sections generally agree with a $(2J + 1)$ relationship (1978FO29). For discussions of the deviations from this relationship see (1978FO29, 1984FO15).

^c $^{13}\text{C}(^6\text{Li}, \text{p})^{18}\text{O}$ at $E(^6\text{Li}) = 28.0$ MeV (1986SM01).

^d 2^- is suggested.

^e May be an unresolved doublet.

^f $J = (2, 3)$.

^g $J^\pi = (4^+, 5^-)$.

Table 18.6: Resonances in $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$ and $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$ ^a

| E_α (MeV \pm) | Γ_{lab} (keV) | Particles out | $^{18}\text{O}^*$ (MeV) | J^π |
|--------------------------|-----------------------------|-----------------------------------|-------------------------|--------------|
| $1.140 \pm 2^{\text{b}}$ | | γ | 7.114 | 4^+ |
| $1.790 \pm 2^{\text{b}}$ | < 3 | γ | 7.619 | 1^- |
| 2.10^{b} | | γ | 7.86 | 5^- |
| $2.330 \pm 2^{\text{b}}$ | < 3 | γ, α_0 | 8.039 | 1^- |
| $2.440 \pm 2^{\text{b}}$ | | γ | 8.125 | 5^- |
| $2.554 \pm 4^{\text{b}}$ | 1.3 ± 1 | $\gamma, \text{n}, \alpha_0$ | 8.213 | 2^+ |
| $2.643 \pm 3^{\text{b}}$ | 10 ± 1 | $\gamma, \text{n}, \alpha_0$ | 8.282 | 3^- |
| 2.800 ± 7 | 10 ± 7 | n | 8.404 | |
| 3.330 ± 12 | 90 ± 15 | n, α_0 | 8.817 | |
| 3.508 ± 4 | 55 ± 3 | n, α_0 | 8.955 | |
| 4.030 ± 15 | 35 ± 20 | n, (α_0) | 9.361 | |
| 4.07 ± 40 | ≈ 150 | n, (α_0) | 9.39 | |
| 4.17 ± 40 | ≈ 70 | n, (α_0) | 9.47 | |
| 4.434 ± 10 | 80 ± 40 | n, (α_0) | 9.675 | |
| 4.70 ± 40 | ≈ 200 | n, (α_0) | 9.88 | |
| 5.004 ± 10 | 21 ± 5 | n, α_0 | 10.118 | 3^- |
| 5.23^{c} | d | n, α_0 | 10.29 | 4^+ |
| 5.34 | d | n, α_0 | 10.38 | 3^- |
| 5.60 | e | n, α_0 | 10.58 | |
| 5.90 | f | n, α_0 | 10.82 | |
| 6.02 | f | n, α_0 | 10.91 | |
| 6.13 | f | n, α_0 | 10.99 | |
| 6.30 | e | n, α_0 | 11.13 | |
| 6.64 | d | n, α_0 | 11.39 | (2^+) |
| 6.67 | d | n, α_0 | 11.41 | (4^+) |
| 6.93 | d | n, α_0 | 11.62 | 5^- |
| 7.03 | d | n, α_0 | 11.69 | 6^+ |
| 7.19 | f | n, α_0 | 11.82 | (3^-) |
| 7.47 | f | n, α_0 | 12.04 | (2^+) |
| 7.75 | g | n, α_0 | 12.25 | $(0^+, 1^-)$ |
| 7.85 | d | n, α_0 | 12.33 | 5^- |

Table 18.6: Resonances in $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$ and $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$ ^a (continued)

| E_α (MeV \pm keV) | Γ_{lab} (keV) | Particles out | $^{18}\text{O}^*$ (MeV) | J^π |
|----------------------------|-----------------------------|---------------|-------------------------|---------|
| 8.06 | ^d | n, α_0 | 12.50 | 4^+ |
| 8.10 | ^d | n, α_0 | 12.53 | 6^+ |

^a See also Table 18.4. For references see Table 18.5 in (1978AJ03). I am grateful to E.K. Warburton for his comments.

^b (1987GA15): $\Gamma_\gamma = 0.095 \pm 0.020, 0.41 \pm 0.08, 0.043 \pm 0.009, 1.07 \pm 0.22, 0.27 \pm 0.05, 0.41 \pm 0.09$ and 0.49 ± 0.13 eV, respectively for $^{18}\text{O}^*(7.11, 7.62, 7.86, 8.04, 8.13, 8.21, 8.28)$.

^c ± 10 – 20 keV for this and all higher resonances (G.E. Mitchell, private communication).

^d Γ_α , large; Γ_n , large.

^e Γ_α , small; Γ_n , small.

^f Γ_α , small; Γ_n , large.

^g Γ_α , large; Γ_n , small.

6. $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$ $Q_m = 6.2270$

Resonances in the yield of capture γ -rays are observed at $E_\alpha = 1.14, 1.79, 2.09, 2.33, 2.44, 2.55$ and 2.64 MeV: see Tables 18.6 here and 18.5 in (1978AJ03). Gamma-ray angular distribution and correlation measurements lead to $J^\pi = 4^+, 1^-, 1^-$ and 5^- for $^{18}\text{O}^*(7.11, 7.62, 8.04, 8.13)$, as well as to J^π assignments for lower states involved in the cascade decay. See also (1982AL30, 1983GA02, 1985GA1J) and (1986HA2D; astrophysics).

7. (a) $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$ $E_b = 6.2270$
 (b) $^{14}\text{C}(\alpha, n)^{17}\text{O}$ $Q_m = -1.8172$

Observed anomalies in the scattering [reaction (a)] for $E_\alpha = 2$ to 8.2 MeV and the resonances in the relative neutron yield [reaction (b)] for $E_\alpha = 2.3$ to 8.5 MeV are displayed in Table 18.6. See also (1978AJ03) and (1985DE24, 1988UM1A; theor.).

8. (a) $^{14}\text{C}(^6\text{Li}, d)^{18}\text{O}$ $Q_m = 4.752$
 (b) $^{14}\text{C}(^6\text{Li}, d\alpha)^{14}\text{C}$ $Q_m = -1.4751$

At $E(^6\text{Li}) = 34$ MeV angular distributions have been measured for the deuteron groups to many states of ^{18}O (1981CU07) [see also (1983AJ01)] including $^{18}\text{O}^*(17.6 \pm 0.2)$ (1982CU01). $J^\pi = 4^+, 2^+, 2^+, (4^+)$ and (4^+) are suggested for $^{18}\text{O}^*(7.86, 8.9, 12.04, 14.6, 17.0)$ (1981CU07). The $2^+, 4^+, 6^+$ and 8^+ members of the $K^\pi = \text{O}_2^+$ rotational band based on $^{18}\text{O}^*(3.62)$ are $^{18}\text{O}^*(5.26, 7.12, 11.69, 17.6)$ (1981CU07, 1982CU01).

Angular correlations have been measured at $E(^6\text{Li}) = 34$ MeV; these lead to the assignment of $J^\pi = 8^+$ to $^{18}\text{O}^*(17.6)$ (1982CU01) and to the assignment of $J^\pi = 4^+, 5^-, 6^+, 7^-$ and 8^+ to sixteen states in ^{18}O with $11.4 \leq E_x \leq 23.1$ MeV (1983CU03) [see also for assignment of ^{18}O states to bands]. At $E(^6\text{Li}) = 32$ MeV (1983AR11) find that the strongest groups are those to (unresolved) structures at $E_x = 17.05$ and 18.95 MeV [each $\Gamma \approx 0.35$ MeV] dominated by $J^\pi = 7^-$. $^{18}\text{O}^*(11.6, 12.6)$ with $J^\pi = (6^+, 5^-)$ and 5^- are also observed (1983AR11). [See, however, the density of states]. See also (1985KO1B; theor.).

$$9. \ ^{14}\text{C}(^7\text{Li}, t)^{18}\text{O} \quad Q_m = 3.759$$

At $E(^7\text{Li}) = 20.4$ MeV, triton groups are observed corresponding to a number of states of ^{18}O with $E_x < 12.6$ MeV. Angular distributions were obtained for some of these, including $^{18}\text{O}^*(0, 1.98, 7.12, 11.69)$ with $J^\pi = 0^+, 2^+, 4^+, 6^+$. The latter two are the most strongly populated in this reaction: they appear to be part of the ground-state rotational band: see (1972AJ02). See also (1983GA02, 1984RU1A).

$$10. \ ^{14}\text{C}(^{14}\text{C}, ^{10}\text{Be})^{18}\text{O} \quad Q_m = -5.785$$

See (1985KO04).

$$11. \ ^{14}\text{C}(^{16}\text{O}, ^{12}\text{C})^{18}\text{O} \quad Q_m = -0.9349$$

See (1978AJ03).

$$12. \ ^{15}\text{N}(^{13}\text{C}, ^{10}\text{B})^{18}\text{O} \quad Q_m = -8.0421$$

See (1983AJ01).

$$13. \ ^{16}\text{O}(t, p)^{18}\text{O} \quad Q_m = 3.7061$$

Table 18.7: States in ^{18}O from $^{16}\text{O}(t, p)$ ^a

| E_x (MeV \pm keV) | L | J^π | E_x (MeV \pm keV) | E_x (MeV \pm keV) |
|-----------------------|------|----------------|-----------------------|-----------------------|
| 0 | 0 | 0^+ | 7623 ± 18 | 9713 ± 7 |
| 1986 ± 4 | 2 | 2^+ | 7782 ± 6 | 9890 ± 11 |
| 3556 ± 2 | 4 | 4^+ | 7871 ± 2^d | 10120 ± 40 |
| 3634^b | 0 | 0^+ | 7983 ± 3^d | 10300 ± 20 |
| 3915 ± 2 | 2 | 2^+ | 8046 ± 7 | 10400 ± 10 |
| 4458 ± 3 | 1 | 1^- | 8140 ± 10 | 10610 ± 20 |
| 5105 ± 2 | 3 | 3^- | 8233 ± 9 | |
| 5258 ± 6 | 2 | 2^+ | 8294 ± 5^d | |
| 5340 ± 4 | 0 | 0^+ | 8430 ± 12 | |
| 5382 ± 4 | | | 8521 ± 3^d | |
| 5530 ± 4 | | | 8660 ± 6 | |
| 6197 ± 3 | 1 | 1^- | 9030 ± 15^e | |
| 6356 ± 7 | 1, 2 | $(1^-, 2^+)^c$ | 9362 ± 5^d | |
| 6399 ± 3 | 3 | 3^- | 9420 ± 20 | |
| 6885 ± 9 | | | 9480 ± 30 | |
| 7123 ± 7 | 4 | 4^+ | 9671 ± 8 | |

^a (1981CO13): $E_t = 15$ MeV; DWBA analysis. See also Table 18.6 in (1978AJ03).

^b Nominal energy.

^c See, however, Table 18.9.

^d Comparisons of E_x shown here with those displayed in Table 18.3 for $^{18}\text{O}^*(3.92, 5.10, 6.40, 7.77)$ suggest that the uncertainty shown may be low: ± 6 keV was arbitrarily used in calculating the best value for E_x for this state in Table 18.3.

^e This is the “average” of several unresolved levels. (1985FO11) state that the main components are at 8.96 and 9.03 MeV. [Comment: It is not clear whether these states are actually resolved.]

Proton groups corresponding to states of ^{18}O are displayed in Table 18.7 (1981CO13). See (1976LA13) for a general discussion of the properties of the states of ^{18}O . Lifetime measurements are reported in Table 18.4 of (1978AJ03). See also reaction 17, (1982AN12, 1985AN17) and (1985BA1A; theor.).

$$14. \ ^{16}\text{O}(\alpha, 2p)^{18}\text{O} \quad Q_m = -16.1080$$

At $E_\alpha = 65$ MeV, the angular distribution to $^{18}\text{O}^*(3.55)$ [$J^\pi = 4^+$] has been studied. In addition $^{18}\text{O}^*(8.04, 9.15, 10.3)$ are also populated: see (1983AJ01).

$$15. \text{ (a) } ^{16}\text{O}(^{10}\text{B}, ^8\text{B})^{18}\text{O} \quad Q_m = -14.824$$

$$\text{ (b) } ^{16}\text{O}(^{13}\text{C}, ^{11}\text{C})^{18}\text{O} \quad Q_m = -11.480$$

At $E(^{10}\text{B}) = 100$ MeV, $^{18}\text{O}^*(3.55)$ [first $(d_{5/2})_{4+}^2$ state] is preferentially populated. $^{18}\text{O}^*(1.98, 5.26, 7.12, 8.0, 8.3, 9.1)$ are also observed. At $E(^{13}\text{C}) = 105$ MeV the angular distribution to $^{18}\text{O}^*(3.55)$ has been measured. See (1983AJ01) and (1983OS07; theor.).

$$16. \text{ (a) } ^{17}\text{O}(\text{n}, \gamma)^{18}\text{O} \quad Q_m = 8.0443$$

$$\text{ (b) } ^{17}\text{O}(\text{n}, \text{n})^{17}\text{O} \quad E_b = 8.0443$$

$$\text{ (c) } ^{17}\text{O}(\text{n}, \alpha)^{14}\text{C} \quad Q_m = 1.8172$$

For reaction (a) see (1983AJ01). [The work reported there has not been published.] The scattering amplitude (bound) $a = 5.62 \pm 0.45$ fm, $\sigma_{\text{free}} = 3.55 \pm 0.25$ b. The thermal cross section for reaction (c) is 235 ± 10 mb. See (1983AJ01) for references.

$$17. \ ^{17}\text{O}(\text{d}, \text{p})^{18}\text{O} \quad Q_m = 5.8197$$

Observed proton groups are displayed in Table 18.8. At $E_d = 12$ MeV a strong, asymmetric peak is observed corresponding to $E_x = 9.0$ MeV. On the basis of this work and the measurement of the cross section to a peak at about the same energy observed in the $^{16}\text{O}(\text{t}, \text{p})$ reaction, (1985FO11) assign $J^\pi = 4^+$ and a $(1d_{5/2})(1d_{3/2})$ configuration to $^{18}\text{O}^*(9.0)$. Proton- γ coincidence measurements are shown in Table 18.4.

$$18. \ ^{17}\text{O}(^{12}\text{C}, ^{11}\text{C})^{18}\text{O} \quad Q_m = -10.677$$

Table 18.8: States of ^{18}O from $^{17}\text{O}(\text{d}, \text{p})$ ^a

| E_x (MeV \pm keV) ^b | l_n ^b | $J\pi$ ^b | S ^b |
|------------------------------------|--------------------|---------------------|------------------|
| 0 | 2 | 0^+ | 1.22 |
| 1.982 ± 10 | $0 + 2$ | 2^+ | $0.21 + 0.83$ |
| 3.552 ± 10 | 2 | 4^+ | 1.57 |
| 3.63 | 2 | 0^+ | 0.28 |
| 3.92 | $0 + 2$ | 2^+ | $0.35 + 0.66$ |
| 4.46 | 1 | 1^- | 0.03 |
| 5.10 | 3 | 3^- | 0.03 |
| 5.255 ± 10 | 0 | 2^+ | 0.35 |
| 5.34 | 2 | 0^+ | 0.16 |
| 5.375 ± 10 | 0 | 3^+ | 1.01 |
| 6.20 | 1 | 1^- | 0.03 |
| 6.35 | 1 | $\leq 3^{(-)}$ | $0.03 - 0.04$ |
| 7.110 ± 15 | 2 | 4^+ | |
| 7.855 ± 20 | | | |
| 7.962 ± 20 | | | |
| 9.0 ^c | | 4^+ | |

^a See references in Tables 18.7 in (1972AJ02) and in (1978AJ03).

^b E_x values without uncertainties are nominal. J are consistent with l_n and are used to calculate S .

^c (1985FO11). See text.

Table 18.9: Branching in $^{18}\text{N}(\beta^-)^{18}\text{O}$ ^a

| Decay to $^{18}\text{O}^*$ ^b (keV) | J^π | Branch (%) | $\log ft$ ^c |
|---|--------------------|----------------|------------------------|
| 1982.05 ± 0.09 ^d | 2^+ | 3.9 ± 1.5 | 6.73 ± 0.17 |
| 3554.13 ± 0.80 | 4^+ | < 0.6 | > 7.22 |
| 3633.70 ± 0.11 | 0^+ | < 0.35 | > 7.43 |
| 3920.42 ± 0.14 | 2^+ | < 0.43 | > 7.29 |
| 4455.52 ± 0.10 | 1^- | 54.6 ± 1.1 | 5.107 ± 0.033 |
| 5097.60 ± 0.60 | 3^- | < 0.43 | > 7.03 |
| 5530.17 ± 0.32 | 2^- | 3.1 ± 0.4 | 6.10 ± 0.06 |
| 6198.22 ± 0.40 | 1^- | 1.4 ± 0.2 | 6.28 ± 0.07 |
| 6349.76 ± 1.0 | (2^-) | 2.2 ± 0.3 | 6.04 ± 0.07 |
| 6880.45 ± 0.27 | 0^- ^e | 14.8 ± 0.8 | 5.066 ± 0.040 |
| 7771.07 ± 0.50 | 2^- ^e | 5.0 ± 0.5 | 5.26 ± 0.06 |

^a (1982OL01).

^b For γ -ray branchings see Table 18.4.

^c It is estimated that $(15 \pm 6)\%$ of the transitions are to states of ^{18}O which do not γ -decay. The uncertainty in this number is reflected in the $\log ft$ values but not in the branching ratios. The $\log ft$ values were calculated using 13899 ± 20 keV for the $^{18}\text{N}(\beta^-)^{18}\text{O}$ Q -value (see also “mass of ^{18}N ” in the “General” section of ^{18}N) and the $\tau_{1/2}$ value given in reaction 1 of ^{18}N , 624 ± 12 msec (1982OL01). I am indebted to Dr. E.K. Warburton for his calculations of the $\log ft$ values.

^d $E_\gamma = 1981.933 \pm 0.09$ keV is adopted by (1982OL01).

^e See (1982OL01).

Angular distributions involving $^{18}\text{O}^*(0, 1.98, 3.55)$ have been studied at $E(^{12}\text{C}) = 115$ MeV: see (1983AJ01).

19. $^{18}\text{N}(\beta^-)^{18}\text{O}$ $Q_m = 13.899$

The transitions observed in the β^- decay are displayed in Table 18.9 (1982OL01).

20. (a) $^{18}\text{O}(\gamma, n)^{17}\text{O}$ $Q_m = -8.0443$
 (b) $^{18}\text{O}(\gamma, 2n)^{16}\text{O}$ $Q_m = -12.1879$

| | |
|--|---------------------------|
| (c) $^{18}\text{O}(\gamma, \text{p})^{17}\text{N}$ | $Q_{\text{m}} = -15.942$ |
| (d) $^{18}\text{O}(\gamma, \text{t})^{15}\text{N}$ | $Q_{\text{m}} = -15.8336$ |
| (e) $^{18}\text{O}(\gamma, \alpha)^{14}\text{C}$ | $Q_{\text{m}} = -6.2270$ |

Table 18.10: Resonances in $^{18}\text{O} + \gamma$

| E_{x} (MeV) ^a | | | | σ (mb) | Γ (MeV) |
|-----------------------------------|----------------------|-----------------------|----------------------|--------------------------------------|----------------|
| (γ, tot) | (γ, n) | $(\gamma, 2\text{n})$ | (γ, p) | | |
| 9.1 | 9.1 | | | 1.1 ^b | 0.6 |
| 10.3 | 10.3 | | | 5.3 ^b | 0.9 |
| 11.4 | 11.4 | | | 9.0 ^b | 0.7 |
| 13.1 | 13.1 | 13.2 | | 8.6 ^b | 0.7 |
| 13.8 | 13.8 | 13.9 | | 6.9 ^b | 0.6 |
| 14.7 | 14.7 | 14.8 | | 13.1 ^b | 0.8 |
| 15.8 | 15.7 | 15.8 | | 10.9 ^b | 0.7 |
| 17.3 ^c | 17.1 | | 17.5 | 10.1 ^b , 1.2 ^e | 0.6 |
| 19.4 ^c | | (19.1) | 19.4 | 10.0 ^b , 1.8 ^e | 0.9 |
| 21.1 ^d | | 21.1 | 21.0 | 9.7 ^b , 1.2 ^e | |
| 22.6 | (22.6) | 22.7 | 22.7 | | |
| 23.7 ^d | 23.7 | 23.5 | 23.7 | 17.7 ^b , 6.1 ^e | 1.6 |
| 27 ^c | 27 | | 27–28 | | |
| 30 ^f | 30 | | | | |
| 36 ^f | | | | | |

^a (1979WO04). See also Table 18.9 in (1983AJ01) and (1987JU07).

^b $\sigma(\gamma, \text{n}) + 2\sigma(\gamma, 2\text{n})$.

^c $T = 2$: see (1979WO04).

^d $T = 1$: see (1979WO04).

^e $\sigma(\gamma, \text{p})$.

^f Weak and broad resonances: may indicate the presence of particle-hole states at these high energies.

The cross sections for the (γ, p) , (γ, n) , $(\gamma, 2\text{n})$ and (γ, tot) [tot = total absorption] have been measured with monoenergetic photons to 42 MeV: observed resonances are displayed in Table 18.10. All three of the partial cross sections have substantial strength in the giant resonance region; the $(\gamma, 2\text{n})$ cross section is a significant fraction of $\sigma(\gamma, \text{tot})$ and is even larger than $\sigma(\gamma, \text{p})$.

Above the GDR the partial cross sections decrease but the integrated $\sigma(\gamma, \text{tot})$ between 29 and 42 MeV is about one-third of the value integrated from threshold to 42 MeV. The relative strengths of partial cross sections leads to the T assignments shown in Table 18.10. The $T_<$ and $T_>$ components of the ^{18}O photo absorption cross section are also derived (1979WO04). Structures in the (γ, α_0) cross section are reported at $E_x = 18.2, 20.9, 22.1$ and 24.2 MeV (1982BA03; $E_{\text{brems.}}$). The decay of the GDR to ^{14}C , ^{15}N , ^{16}O , ^{17}N and ^{17}O states has been studied: see (1983AJ01). Less than 20% of the decays of states with $14.5 < E_x < 20$ MeV goes via the n_0 channel (1987JU07). See (1978AJ03) for the earlier work, (1985SH1P), (1985PY01) and (1983KA28, 1984VA1G, 1985PY01, 1985VA1C; theor.).

21. $^{18}\text{O}(\gamma, \gamma)^{18}\text{O}$

The Γ_{γ_0} of $^{18}\text{O}^*(6.20)$ is 0.18 ± 0.03 eV, assuming $\Gamma_{\gamma_0}/\Gamma = 0.88$; $E_x = 6202.7 \pm 0.8$ keV: see (1978AJ03).

22. $^{18}\text{O}(e, e)^{18}\text{O}$

The ^{18}O charge radius, $\langle r^2 \rangle^{1/2} = 2.784 \pm 0.020$ fm, based on studies of the elastic charge form factors for $E_e = 70$ to 370 MeV, the resulting determinations of the difference in the ^{18}O and ^{16}O radii, and the rms radius of ^{16}O : see (1983AJ01).

Inelastic scattering has been reported to many states of ^{18}O : see (1983AJ01) and Table 18.11 here (1982NO04, 1983BE36, 1986MA48). See also (1987DE43) and (1986GU05, 1986ST15, 1987MI1E; theor.).

Table 18.11: Some states of ^{18}O from $^{18}\text{O}(e, e')$ ^a

| E_x (MeV \pm keV) | Γ (keV) | $J^\pi; T$ | Mult. | Transition probability (in $e^2 \cdot \text{fm}^{2\lambda}$) |
|--------------------------------|----------------|------------|-------|--|
| 1.98 ^b | | $2^+; 1$ | C2 | 44.8 ± 1.3 |
| 3.55 ^b | | $4^+; 1$ | C4 | $(9.04 \pm 0.90) \times 10^2$ |
| 3.92 ^b | | $2^+; 1$ | C2 | 22.2 ± 1.0 |
| 5.26 ^b | | $2^+; 1$ | C2 | 28.3 ± 1.5 |
| 7.12 ^b | | $4^+; 1$ | C4 | $(1.31 \pm 0.06) \times 10^4$ |
| 9.36 ± 20 ^c | 10 ± 7 | $T = 1$ | | |
| 11.65 ± 20 ^c | 76 ± 8 | $T = 1$ | | |
| 12.06 ± 60 ^c | 28 ± 6 | $T = 1$ | | |
| 16.210 ± 10 ^{d,e} | | $1^{(-)}$ | | |

Table 18.11: Some states of ^{18}O from $^{18}\text{O}(e, e')$ ^a (continued)

| | Γ (keV) | $J^\pi; T$ | Mult. | Transition probability (in $e^2 \cdot \text{fm}^{2\lambda}$) |
|----------------------------------|----------------|-------------------|-----------|--|
| 16.315 ± 10 ^{d,e} | | $(3, 2)^-$ | | |
| 16.399 ± 5 ^{d,f} | < 20 | $2^-; 2^g$ | M2 | $(64 \pm 8) \times 10^{-2}$ |
| 16.948 ± 10 ^{d,e} | | $(3, 2)^-$ | | |
| 17.025 ± 10 ^{d,e,f} | 20 ± 6 | $(> 2); 2$ | | |
| (17.35 ± 60) ^f | 680 ± 250 | | | |
| 17.398 ± 10 ^{d,e} | | $(2, 1, 3)^-$ | | |
| 17.450 ± 10 ^{d,e} | | $(2, 1, 3)^-$ | | |
| 17.5 ^d | ≈ 150 | | | |
| 17.502 ± 10 ^{d,e} | | $(1, 2, 3)^-$ | | |
| 17.635 ± 10 ^{d,e} | | | | |
| 18.049 ± 10 ^{d,e} | | ^d | | |
| 18.2 ^d | ≈ 150 | | | |
| (18.48 ± 20) ^f | 90 ± 34 | | | |
| 18.5 ^d | ≈ 4300 | | | |
| 18.70 ± 20 ^f | < 20 | $(4^-); 2$ | M4 | 63 ± 8 |
| 18.871 ± 5 ^d | | $1^+; 2$ | M1 | $(3.1 \pm 0.4) \times 10^{-2}$ |
| 18.927 ^{d,e} | | $1; (2^+)$ | | |
| 19.027 ± 10 ^{d,e} | | $(1, 3)^-$ | | |
| 19.150 ± 10 ^{d,e} | | $1^-; (2^+, 3^-)$ | | |
| 19.24 ± 20 ^f | < 20 | $(> 2); 2$ | | |
| 19.7 ^d | ≈ 200 | | | |
| 20.2 ^d | ≈ 180 | | | |
| 20.36 ± 20 ^f | < 20 | $(4^-); 2$ | M4 | 66 ± 6 |
| 21.0 ^d | ≈ 150 | | | |
| (21.43 ± 20) ^f | 49 ± 37 | | | |
| 22.39 ± 40 ^{d,f} | 74 ± 7 | (4^-) | M4 | 400 ± 32 |
| 23.8 ^d | ≈ 1300 | | | |

^a Additional states have been excited: see reaction 28 in (1983AJ01). For ground state see text here.

^b (1982NO04).

^c Mark Manley, private communication.

^d (1983BE36).

^e Weakly excited.

^f (1986MA48).

^g See Fig. 4 (isobar diagram for $A = 18$) for missing $T = 2$ strength.

23. (a) $^{18}\text{O}(\pi^\pm, \pi^\pm)^{18}\text{O}$
(b) $^{18}\text{O}(\pi^\pm, \pi^\pm\text{p})^{17}\text{N}$ $Q_m = -15.942$
(c) $^{18}\text{O}(\pi^-, \pi^-\text{n})^{17}\text{O}$ $Q_m = -8.0443$

Angular distributions for the scattering to $^{18}\text{O}^*(0, 1.98, 5.10)$ have been reported at $E_{\pi^\pm} = 29.2$ to 230 MeV [see (1983AJ01)] and at 50 MeV (1984TA1A; $^{18}\text{O}^*(0, 1.98)$) at 140, 180 and 220 MeV (1984SE1A; $^{18}\text{O}^*(1.98)$), at 164 MeV (1987CH14; $^{18}\text{O}^*(0, 1.98, 4.46, 5.10)$) and at 180 MeV (1987MO07; $^{18}\text{O}^*(1.98)$). $\langle r_n^2 \rangle^{1/2} = 2.81 \pm 0.03$ fm, $\langle r_n^2 \rangle^{1/2} - \langle r_p^2 \rangle^{1/2} = 0.03 \pm 0.03$ fm: see (1983AJ01). For a discussion of the proton matter distribution in ^{18}O see (1985BA27). For total reaction cross sections at $E_\pi = 50$ and 65 MeV see (1986ME1C; prelim.). At $E_\pi = 165$ MeV, the cross section for reaction (c) is larger for ^{18}O than for ^{16}O while reaction (b) has a lower cross section (1982PI06). For the $(\pi^+, 2\text{p})$, (π^+, pn) and (π^-, pn) reactions at $E_\pi = 165$ MeV see (1984AL20, 1986AL22). See also the “General” section here.

24. $^{18}\text{O}(\text{n}, \text{n})^{18}\text{O}$

Angular distributions have been measured for $E_n = 2.9$ to 24 MeV [see (1972AJ02, 1983AJ01)] and at $E_n = 5.0$ to 7.5 MeV (1986KO10; n_0, n_1).

25. $^{18}\text{O}(\text{p}, \text{p})^{18}\text{O}$

Angular distributions have been measured for $E_p = 0.84$ to 135 MeV [see (1978AJ03, 1983AJ01)], at $E_p = 135$ MeV (1986KE05; p_1) and at $E_{\bar{p}} = 800$ MeV (1982GL08; \bar{p} to $^{18}\text{O}^*(0, 1.98, 7.12)$).

At $E_p = 24.5$ MeV (1974ES02) have studied the angular distributions of the proton groups to $^{18}\text{O}^*(1.98, 3.55, 3.63, 3.92, 4.46, 5.10, 5.26, 5.53, 7.12)$: a modified DWBA analysis leads to $J^\pi = 2^+, 4^+, 0^+, 2^+, 1^-, 3^-, 2^+, 2^-$ and 4^+ for these states. A coupled-channels calculation suggests $\beta_2 = 0.37 \pm 0.03, 0.56 \pm 0.06$ and 0.18 ± 0.04 for $^{18}\text{O}^*(1.98, 5.10, 7.12)$. Such calculations also support evidence for a rotational band involving $^{18}\text{O}^*(0, 1.98, 7.12)$. The 3^- state at 5.10 MeV

is strongly excited and collective in nature: $B(E3) = 1120 e^2 \cdot \text{fm}^6$. For $^{18}\text{O}^*(1.98, 3.92, 5.26)$, $B(E2) = 45, 8.3$ and $24 e^2 \cdot \text{fm}^4$ (1974ES02). The 800-MeV data indicates that $^{18}\text{O}^*(7.12)$ can be described only if a large hexadecapole deformation is assumed (1982GL08). At $E_p = 201$ MeV forward angle $\sigma(\theta)$ have been measured to $^{18}\text{O}^*(8.21, 8.82, 16.40)$: it is proposed that $^{18}\text{O}^*(8.82)$ has $J^\pi = 1^+$ and that additional 1^+ strength is located in a group centered at $E_x \approx 10.1$ MeV as well as in the region $E_x = 12.4$ to 15 MeV. The $1^+; T = 2$ state $^{18}\text{O}^*(18.87)$, reported in (e, e'), is not observed (1987DJ01).

$^{18}\text{O}^*(1.98)$ has $|g| = 0.287 \pm 0.015$ [$\tau_m = 2.99 \pm 0.12$ psec]. $^{18}\text{O}^*(3.55)$ has $|g| = 0.62 \pm 0.10$ suggesting a mainly $(d_{5/2})^2$ configuration for this state: see (1983AJ01). See also ^{19}F , (1983CE1A) and (1982KU04, 1982NA13, 1984PH02, 1986DE1G, 1986PE1E; theor.).

26. $^{18}\text{O}(\bar{p}, \bar{p})^{18}\text{O}$

Angular distributions are reported with 178.4 MeV antiprotons to $^{18}\text{O}^*(0, 1.98)$ (1986BR04, 1986LE13). For atomic effects see (1986KO22). See also (1984PO1A), (1985LI16; theor.) and the “General” section here.

27. $^{18}\text{O}(d, d)^{18}\text{O}$

Angular distributions have been reported at $E_d = 7.0$ to 15.0 MeV: see (1972AJ02, 1983AJ01). See also ^{20}F .

28. $^{18}\text{O}(t, t)^{18}\text{O}$

See (1972AJ02).

29. $^{18}\text{O}(^3\text{He}, ^3\text{He})^{18}\text{O}$

The elastic scattering has been studied at $E(^3\text{He}) = 11.0$ to 41 MeV [see (1972AJ02, 1983AJ01)] and at 14 MeV (1982AB04), at 25 MeV (1982VE13) [the matter radius, $\langle r_m^2 \rangle^{1/2} = 2.59 \pm 0.12$ fm] and at $E(^3\text{He}) = 33$ MeV (1983LE03; also A_y ; and also to $^{18}\text{O}^*(1.98)$). See also (1985HA11, 1987CO07; theor.).

30. $^{18}\text{O}(\alpha, \alpha)^{18}\text{O}$

Angular distributions of many α -groups have been measured in the range $E_\alpha = 21$ to 40.5 MeV [see (1978AJ03)], at 23.5 MeV (1984SA28; to $^{18}\text{O}^*(1.98, 3.56 + 3.63, 3.92, 4.45, 5.1 - 5.53)$) and at 54.1 MeV (1987AB03; g.s.). The transitions to $^{18}\text{O}^*(4.46, 5.10)$ are $L = 1$ and 3, respectively, fixing $J^\pi = 1^-$ and 3^- for these states. Measurements of α -groups near 180° for $E_\alpha = 20.0$ to 23.4 MeV confirm assignments of natural parity for $^{18}\text{O}^*(1.98, 3.55, 3.63, 3.92, 4.46, 5.10, 5.26, 5.34, 6.20, 6.40, 7.12, 7.62, 7.86, 8.22, 8.29, 8.82, 8.96, 9.03, 9.10, 9.36, 9.41, 9.67, 9.72 \pm 0.03, 9.88, 10.12, 10.30, 10.40, 11.62, 11.69)$. [See, however, Table 18.3.] Levels at $E_x = 5.38, 8.48$ and 8.64 MeV were not observed, and those at 5.53, 6.35 and 6.88 MeV were populated weakly indicating unnatural parity: $J^\pi = 3^+$ and 2^- respectively for $^{18}\text{O}^*(5.38, 5.53)$.

Alpha-gamma correlation measurements involving ^{18}O states below $E_x = 6.4$ MeV [see Table 18.4] lead to $J^\pi = 1^-$ and 3^- for $^{18}\text{O}^*(6.20, 6.40)$. Other J^π values agree with previous assignments. The transitions $3.92 \rightarrow 1.98$ and $5.26 \rightarrow 1.98$ are almost pure M1. For τ_m measurements see Table 18.4 in (1978AJ03). For references see (1983AJ01). See also (1983WI12, 1984BR1P, 1984LA01, 1987AS01; theor.).

31. (a) $^{18}\text{O}(^6\text{Li}, ^6\text{Li})^{18}\text{O}$
 (b) $^{18}\text{O}(^7\text{Li}, ^7\text{Li})^{18}\text{O}$

See (1972AJ02, 1983AJ01).

32. $^{18}\text{O}(^9\text{Be}, ^9\text{Be})^{18}\text{O}$

See (1972AJ02, 1983BI13, 1985BE1A) and (1984HA43; theor.).

33. (a) $^{18}\text{O}(^{10}\text{B}, ^{10}\text{B})^{18}\text{O}$
 (b) $^{18}\text{O}(^{11}\text{B}, ^{11}\text{B})^{18}\text{O}$

An elastic angular distribution has been reported at $E(^{11}\text{B}) = 115$ MeV: see (1983AJ01). For reaction (a) see (1974AJ01).

34. (a) $^{18}\text{O}(^{12}\text{C}, ^{12}\text{C})^{18}\text{O}$
 (b) $^{18}\text{O}(^{13}\text{C}, ^{13}\text{C})^{18}\text{O}$
 (c) $^{18}\text{O}(^{14}\text{C}, ^{14}\text{C})^{18}\text{O}$
 (d) $^{18}\text{O}(^{12}\text{C}, \alpha)^{12}\text{C}^{14}\text{C}$ $Q_m = -6.2270$

Elastic angular distributions have been studied at $E(^{18}\text{O}) = 32.3$ to 57.5 MeV for reaction (a) [as well as at $E(^{18}\text{O}) = 70, 100$ and 140 MeV (1982HE07)] and at $E(^{18}\text{O}) = 31$ MeV for reaction (b). Yields and fusion cross sections are reported by (1982BA49, 1982HE07, 1985BE40, 1985CA01, 1986GA13). See also (1983VO1B). For the earlier work see (1983AJ01). For reaction (c) see (1986STZY).

Angular correlations (reaction (d)) have been studied at $E(^{18}\text{O}) = 82$ MeV. $^{18}\text{O}^*(7.10, 7.62, 7.86, 8.04, 8.22, 10.30, 11.59, 12.55)$ are observed: the first seven of these have $J^\pi = 4^+, 1^-, 5^-, 1^-, 2^+, 4^+, 5^-$ (1984BH01, 1984RA07). In addition $^{18}\text{O}^*(9.33, 9.65)$ are also populated [$\Gamma \approx 0.3$ MeV]: a possible interpretation of the data is that these two are 3^- states and that there is in addition a very wide (> 1 MeV) 2^+ state at ≈ 9.5 MeV (1984RA17). See also (1983CA1N), (1983BI13, 1984HA53) and (1982GI1C, 1982LO13, 1983CI08, 1983MA29, 1984FR1A, 1985CH1R, 1986HA13; theor.).

35. $^{18}\text{O}(^{15}\text{N}, ^{15}\text{N})^{18}\text{O}$

See (1983DU13).

36. $^{18}\text{O}(^{16}\text{O}, ^{16}\text{O})^{18}\text{O}$

Angular distributions have been measured at many energies for $E(^{16}\text{O}) = 24$ to 54.5 MeV and $E(^{18}\text{O}) = 25$ to 52 MeV, involving besides $^{18}\text{O}_{\text{g.s.}}, ^{18}\text{O}^*(1.98, 3.55 + 3.63, 3.92, 4.46, 5.10, 7.12)$. At $E(^{18}\text{O}) = 126$ MeV $^{18}\text{O}^*(9.0)$ is relatively strongly populated. See (1983AJ01). For yields and fusion cross sections, including the effect of $^{18}\text{O}^*(1.98)$, see (1985TH03, 1985WU03, 1986GA13, 1986TH01). See also (1983DU13), (1986BA69) and (1983SH04, 1985MA1T, 1986DAZS, 1986VI08, 1987LO01; theor.).

37. (a) $^{18}\text{O}(^{17}\text{O}, ^{17}\text{O})^{18}\text{O}$

(b) $^{18}\text{O}(^{18}\text{O}, ^{18}\text{O})^{18}\text{O}$

Angular distributions involving $^{18}\text{O}^*(0, 1.98)$ are reported at $E(^{17}\text{O}) = 36$ MeV. Angular distributions [reaction (b)] have been studied at $E(^{18}\text{O}) = 20$ to 52 MeV. $^{18}\text{O}^*(3.55 + 3.63, 4.46, 5.10, 7.12)$ are also populated; see (1978AJ03, 1983AJ01). See also (1986GA13) and (1982CI1C, 1983DU13).

38. $^{18}\text{O}(^{19}\text{F}, ^{19}\text{F})^{18}\text{O}$

The elastic scattering has been studied at $E(^{19}\text{F}) = 27, 30$ and 33 MeV: see (1983AJ01). See also (1986GA13) and (1982OH05; theor.).

39. (a) $^{18}\text{O}(^{24}\text{Mg}, ^{24}\text{Mg})^{18}\text{O}$
(b) $^{18}\text{O}(^{26}\text{Mg}, ^{26}\text{Mg})^{18}\text{O}$

Angular distributions are reported for reaction (a) at $E(^{18}\text{O}) = 29$ and 35 MeV to $^{18}\text{O}^*(0, 1.98)$. See (1983AJ01, 1983BI13) and (1982LO13, 1983CI08, 1984FR1A, 1985HU04, 1986HA13, 1986OS05; theor.).

40. $^{18}\text{O}(^{27}\text{Al}, ^{27}\text{Al})^{18}\text{O}$

The elastic angular distribution has been studied at $E(^{18}\text{O}) = 100$ MeV (1981ME13). See also (1985DEZZ), (1983AJ01, 1983BI13) and (1982LO13, 1983CI08, 1984FR1A, 1985HU04; theor.).

41. $^{18}\text{O}(^{28}\text{Si}, ^{28}\text{Si})^{18}\text{O}$

Elastic angular distributions are reported at $E(^{18}\text{O}) = 36$ to 56 MeV [see (1983AJ01)] and at 351.7 MeV (1984BUZX; also to $^{18}\text{O}^*(1.98)$). See also (1986BL08, 1986GA13), (1983BI13) and (1982LO13, 1983CI08, 1984FR1A, 1985HU04, 1986HOZF; theor.).

42. (a) $^{18}\text{O}(^{40}\text{Ca}, ^{40}\text{Ca})^{18}\text{O}$
(b) $^{18}\text{O}(^{44}\text{Ca}, ^{44}\text{Ca})^{18}\text{O}$
(c) $^{18}\text{O}(^{48}\text{Ca}, ^{48}\text{Ca})^{18}\text{O}$

Angular distributions have been measured at $E(^{18}\text{O}) = 62.1$ MeV [reaction (a)] for the transitions to $^{18}\text{O}^*(0, 1.98, 5.10)$ (1982RE14). For fusion study [reaction (b)] see (1984DE38). See also (1985DE38, 1986OS05, 1986RO12; theor.).

43. $^{18}\text{F}(\beta^+)^{18}\text{O} \quad Q_m = 1.6554$

See ^{18}F .

Table 18.12: ^{18}O states from $^{19}\text{F}(t, \alpha\gamma)$ ^a

| E_x (keV) | J^π | E_x (keV) | J^π |
|--------------------|---------|------------------|-------------------|
| 1982.16 ± 0.20 | | 5530.5 ± 0.6 | 1, 2 |
| 3555.07 ± 0.45 | | 6196.3 ± 1.2 | 1 |
| 3634.50 ± 0.40 | | 6351.3 ± 0.6 | 1, 2 |
| 3920.6 ± 0.4 | | 6404.4 ± 1.2 | |
| 4456.1 ± 0.5 | | 6881.6 ± 1.2 | 0, (1) |
| 5098.5 ± 1.2 | | 7116.9 ± 1.2 | |
| 5260.4 ± 1.2 | | 7.75 | 1 \rightarrow 4 |
| 5336.4 ± 0.6 | | 7.98 | 1 \rightarrow 5 |
| 5377.8 ± 1.2 | | b | |

^a (1973OL02): see Table 18.4 for branching ratios and Table 18.3 for τ_m . See also Table 18.10 in (1983AJ01).

^b Alpha groups are also reported to ^{18}O states with $E_x = 7.60, 7.75, 7.84, 7.96, 8.02, 8.11, 8.19, 8.26, 8.39, 8.48, 8.64$ MeV (± 20 keV).

44. $^{19}\text{F}(\gamma, p)^{18}\text{O}$ $Q_m = -7.9942$

(1985KE03) have measured the yields of proton groups to $^{18}\text{O}^*(0, 1.98)$ [and to unresolved states] for E_{bs} in the GDR range. See also (1978AJ03) and ^{19}F .

45. $^{19}\text{F}(n, d)^{18}\text{O}$

Angular distributions have been measured at $E_n = 14$ to 14.4 MeV: see (1972AJ02). See also (1978AJ03) and ^{20}F .

46. $^{19}\text{F}(d, ^3\text{He})^{18}\text{O}$ $Q_m = -2.5007$

Many states of ^{18}O ($E_x < 14.6$ MeV) have been populated in this reaction: see Table 18.8 in (1978AJ03). [Comment: Note, however, density of states.] Analyzing powers for the ground-state transition are reported at $E_d = 12.4$ MeV (1983EN02). See also (1983KI13; theor.).

$$47. \text{}^{19}\text{F}(t, \alpha)\text{}^{18}\text{O} \quad Q_m = 11.8198$$

See Table 18.12.

$$48. \text{}^{22}\text{Ne}(d, \text{}^6\text{Li})\text{}^{18}\text{O} \quad Q_m = -8.194$$

At $E_d = 80$ MeV angular distributions have been measured for the ${}^6\text{Li}$ groups to the ground state of ${}^{18}\text{O}$ and to excited states at 1.98, 3.57, 5.10, 6.30, 7.8, 9.4 [± 0.04] MeV (1984OE02) [see also for S_{rel}]. For earlier work see (1983AJ01).

¹⁸F
(Figs. 2 and 4)

GENERAL: (See also (1983AJ01).)

Shell model: (1978WI1B, 1982ZH01, 1983BR29, 1983KI13, 1984MI1H, 1984MI17, 1985LE1K, 1986YU1B).

Cluster, collective and deformed models: (1983ME12, 1984QU1A, 1985BA1A, 1987ER05).

Special states: (1978WI1B, 1982ZH01, 1983BI1C, 1983BR29, 1983ME12, 1983KI13, 1984AD1E, 1984HA14, 1984HO1H, 1984MI1H, 1984MI17, 1985AD1A, 1985HA18, 1985LE1K, 1985MI10, 1985SO12, 1985YU1B, 1986AN07, 1986CA27, 1986HA1E, 1986HA1Q, 1986SI1K, 1986TO1D, 1986WI1P, 1986YU1B, 1987ER05).

Electromagnetic transitions: (1982RI04, 1983BR29, 1983KI13, 1984BI10, 1984HA14, 1984MI1H, 1984QU1A, 1985AD1A, 1986CA27, 1986SI1K, 1987ER05, 1987KI03).

Astrophysical questions: (1982WI1B).

Applications: (1982HI1H, 1982YA1C, 1984HI1D, 1984NI1C, 1985HU02, 1986HI1B, 1987HI1B).

Complex reactions involving ¹⁸F: (1983DE26, 1983JA05, 1983OL1A, 1983WI1A, 1984GR08, 1984HI1A, 1984HO23, 1985BE40, 1985CU1C, 1985DR07, 1985GR1K, 1985PO11, 1986AV07, 1986BA3G, 1986GR1A, 1986HA1B, 1986HO27, 1986PO06, 1986SC28, 1986SC29, 1986VA23, 1987RI03, 1987RO10).

Pion capture and reactions (See also reaction 32.): (1982LI15, 1983HO02, 1986GE06, 1986NA1K).

Hypernuclei: (1984AS1D).

Other topics: (1978WI1B, 1983AD1B, 1983AG1C, 1983AR1J, 1983BI1C, 1983BR29, 1983SH32, 1984BI10, 1984HO1H, 1984MI1H, 1985AD1A, 1985AN28, 1985MI10, 1985YU1B, 1986LA29, 1986NA1K, 1986TO1D, 1987KI03).

Ground and 1.12 MeV states of ¹⁸F: (1978WI1B, 1983ANZQ, 1983AR1J, 1984HA14, 1985AN28, 1985HA18, 1986CA27).

$$\mu_{1.12} = +2.86 \pm 0.03 \text{ nm [see (1983AJ01)]}$$

$$Q_{1.12} = 0.13 \pm 0.03 \text{ b [see (1983AJ01)].}$$

1. ¹⁸F(β^+)¹⁸O $Q_m = 1.6554$

The positron decay is entirely to the ground state of ¹⁸O [$J^\pi = 0^+$, $T = 1$]; the half-life is 109.77 ± 0.05 min [see Table 18.11 in (1972AJ02)]; $\log ft = 3.554$. The fact that the β^+ transition to ¹⁸O_{g.s.} is allowed indicates $J^\pi = 1^+$ for ¹⁸F_{g.s.}. The ratio $\epsilon_K/\beta^+ = 0.030 \pm 0.002$:

see (1978AJ03). See also (1985BR29), (1982KA1C; applications), (1985AR1A, 1985KL1A, 1986HE1C, 1987GOZX; astrophysics) and (1983CA03; theor.) [see for discussions of weak magnetism form factor and of search for heavy neutrinos].

Table 18.13: Energy levels of ^{18}F ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | K^π | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|------------|---------|--|-----------|---|
| 0 | $1^+; 0$ | 0^+ | $\tau_{1/2} = 109.77 \pm 0.05$ min | β^+ | 1, 4, 5, 6, 9, 10, 12, 13, 15, 21, 23, 24, 25, 29, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 |
| 0.93720 ± 0.06 | $3^+; 0$ | 0^+ | $\tau_m = 67.6 \pm 2.5$ psec $g = +0.56 \pm 0.05$ | γ | 2, 6, 9, 10, 13, 21, 23, 25, 30, 34, 35, 37, 38, 39, 40, 42 |
| 1.04155 ± 0.08 | $0^+; 1$ | | 2.55 ± 0.45 fsec | γ | 6, 9, 21, 25, 30, 33, 34, 36, 37, 38, 40, 41 |
| 1.08054 ± 0.12 | $0^-; 0$ | 0^- | 27.5 ± 1.9 fsec | γ | 6, 9, 10, 21, 25, 34, 36, 37, 38, 39, 40, 42 |
| 1.12136 ± 0.15 | $5^+; 0$ | 0^+ | 234 ± 10 nsec $\mu = +2.86 \pm 0.03$ nm $Q = 0.13 \pm 0.036$ b | γ | 5, 6, 9, 10, 13, 14, 21, 22, 25, 30, 31, 34, 36, 38, 40, 42 |
| 1.70081 ± 0.18 | $1^+; 0$ | 1^+ | 955 ± 27 fsec | γ | 6, 10, 21, 25, 33, 34, 38, 40, 41, 42 |
| 2.10061 ± 0.10 | $2^-; 0$ | 0^- | 5.1 ± 0.5 psec | γ | 6, 10, 13, 21, 23, 25, 34, 38, 40, 42 |
| 2.52335 ± 0.18 | $2^+; 0$ | 1^+ | 590 ± 24 fsec | γ | 6, 10, 21, 25, 30, 38, 40 |
| 3.06184 ± 0.18 | $2^+; 1$ | | < 1.2 fsec | γ | 6, 21, 25, 30, 34, 37, 38, 40, 41 |
| 3.13387 ± 0.15 | $1^-; 0$ | 1^- | 0.39 ± 0.02 psec | γ | 6, 10, 21, 25, 34, 37, 38, 40 |
| 3.3582 ± 1.0 | $3^+; 0$ | 1^+ | 0.44 ± 0.03 psec | γ | 6, 10, 21, 34, 38, 40, 42 |
| 3.72419 ± 0.22 | $1^+; 0$ | | $2.7^{+4.1}_{-2.7}$ fsec | γ | 6, 10, 21, 23, 25, 33, 34, 38, 40, 42 |
| 3.79149 ± 0.22 | $3^-; 0$ | 1^- | 1.91 ± 0.13 psec | γ | 5, 10, 21, 23, 25, 34, 38, 40, 42 |

Table 18.13: Energy levels of ^{18}F ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | K^π | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|--------------|---------|--|---------------------|---------------------------------------|
| 3.83917 ± 0.22 | $2^+; 0$ | | 19.0 ± 2.7 fsec | γ | 6, 10, 21, 23, 25, 30, 34, 38, 40, 42 |
| 4.11590 ± 0.25 | $3^+; 0$ | | 91 ± 22 fsec | γ | 6, 10, 21, 23, 25, 30, 34, 38, 40, 42 |
| 4.2258 ± 0.7 | $2^-; 0$ | (1^-) | 110 ± 15 fsec | γ | 6, 10, 21, 23, 34, 38, 40, 42 |
| 4.36015 ± 0.26 | $1^+; 0$ | | 27 ± 10 fsec | γ | 10, 21, 25, 33, 34, 38, 40, 42 |
| 4.3981 ± 0.7 | $4^-; 0$ | 0^- | 58 ± 12 fsec | γ | 6, 10, 13, 14, 21, 34, 38, 40, 42 |
| 4.652 ± 2 | $4^+; 1$ | | < 10 fsec | γ | 6, 21, 24, 30, 34, 38, 40 |
| 4.753 ± 3 | $0^+; 1$ | | | γ | 21, 34, 37, 38, 40, 42 |
| 4.8483 ± 0.5 | $5^-; 0$ | 1^- | 5.2 ± 0.9 psec | γ | 5, 23 |
| 4.860 ± 2 | $1^-; 0$ | | 66 ± 18 fsec | γ, α | 6, 21, 38, 40, 42 |
| 4.9636 ± 0.8 | $2^+; 1$ | | < 4 fsec | γ | 6, 21, 30, 38, 40 |
| 5.2976 ± 1.5 | $4^+; 0$ | 1^+ | 30 ± 5 fsec | γ, α | 6, 9, 10, 11, 21, 38, 40 |
| 5.502 ± 2 | $3^{(-)}; 0$ | | 63 ± 25 fsec | γ, α | 6, 10, 21, 38, 40 |
| 5.60338 ± 0.27 | 1^+ | | 15 ± 10 fsec | γ, α | 6, 8, 25, 38, 40, 42 |
| 5.60486 ± 0.28 | $1^-; 0 + 1$ | | $\Gamma < 1.2$ keV | γ, α | 6, 8, 10, 21, 25, 38, 40, 42 |
| 5.673 ± 2 | $1^-; 0 + 1$ | | < 0.8 | γ, α | 6, 8, 10, 21, 25, 38, 40, 42 |
| 5.786 ± 2.4 | $2^-; 0$ | | $\tau_m = 15 \pm 10$ fsec | γ, α | 6, 21, 38, 40, 42 |
| 6.0964 ± 1.1 | $4^-; 0$ | 1^- | $\Gamma = 0.24 \pm 0.03$ | γ, p, α | 6, 10, 21, 25, 29, 38, 40, 42 |
| 6.108 ± 3 | $(1^+); 0$ | | 0.034 ± 0.003 | γ, p, α | 6, 8, 21, 23, 29, 40, 42 |
| 6.13647 ± 0.33 | $0^+; 1$ | | ≤ 1 | γ, p | 21, 25, 27, 40, 42 |
| 6.1632 ± 0.9 | $3^+; 1$ | | 14 ± 0.5 | γ, p, α | 21, 25, 27, 29, 42 |
| 6.2404 ± 0.8 | $3^-; 0 + 1$ | | 0.19 ± 0.03 | γ, p, α | 6, 21, 25, 27, 29, 40 |
| 6.242 ± 3 | $3^-; 0 + 1$ | | 0.18 ± 0.04 | γ, p, α | 6, 8, 21, 25, 29, 40 |
| 6.262 ± 2.5 | $1^+; 0$ | | 0.60 ± 0.12 | γ, p, α | 6, 8, 10, 21, 29, 33, 40 |

Table 18.13: Energy levels of ^{18}F ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | K^π | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|------------------|---------|--|---------------------|-----------------------------|
| 6.2832 ± 0.9 | $2^+; 1$ | | 10.0 ± 0.5 | γ, p, α | 21, 25, 27, 29 |
| 6.3105 ± 0.8 | $3^+; 0$ | | 0.95 ± 0.14 | γ, p, α | 6, 21, 25, 27, 29, 42 |
| 6.3855 ± 1.7 | $2^+; 0 + 1$ | | 0.49 ± 0.09 | γ, p, α | 6, 21, 25, 29, 40 |
| 6.4849 ± 1.5 | $3^+; 0$ | | 0.40 ± 0.10 | γ, p, α | 6, 21, 25, 29, 40, 42 |
| 6.5670 ± 1.5 | $5^+; 0$ | 1^+ | 0.56 ± 0.13 | γ, p, α | 6, 8, 9, 10, 11, 21, 29, 40 |
| 6.633 ± 10 | 1 | | 80 ± 2 | p, α | 29, 40 |
| 6.6437 ± 0.8 | $2^-; 1$ | | 0.60 ± 0.07 | γ, p, α | 6, 7, 21, 25, 29 |
| 6.647 ± 4 | 1^- | | 91 ± 4 | p, α | 8, 10, 29 |
| 6.777 ± 1.4 | $4^+; 0$ | | 9.2 ± 1.0 | γ, p, α | 21, 25, 27, 29, 40 |
| 6.8031 ± 1.5 | $1^+, 2, 3^+; 0$ | | ≤ 2 | γ, p | 10, 21, 25, 27, 40 |
| 6.809 ± 5 | 2^- | | 88 ± 2 | p, α | 7, 8, 29 |
| 6.811 | (2^+) | | 3.0 ± 0.5 | p, α | 29 |
| 6.857 ± 10 | (3^-) | | 5.0 ± 1.0 | p, α | 29, 40 |
| 6.8774 ± 1.7 | $3, 4^-; 0$ | | ≤ 2 | γ, p, α | 21, 25, 29 |
| 7.201 ± 2 | $(4^+); 0$ | | 6.5 | p, α | 8, 29, 40 |
| 7.247 ± 2 | $(1^+); 0$ | | 46.5 | p, α | 8, 29 |
| 7.291 ± 2 | 3^- | | 38 | p, α | 7, 8, 27, 29 |
| 7.315 ± 4 | $(3^-; 0)$ | | 52 | p, α | 29, 40 |
| 7.336 ± 2 | $1^-; 1$ | | 16 ± 2 | γ, p | 25, 27 |
| 7.406 ± 2 | 1^+ | | 14.6 ± 1.4 | p | 27 |
| 7.447 ± 10 | | | 140 | p, α | 29 |
| 7.454 ± 2 | 1^- | | 6 | p | 27 |
| 7.478 ± 2 | (2) | | 12 ± 3 | γ, p, α | 25, 27, 29 |
| (7.485 \pm 2) | (1^-) | | 32 | p | 27 |
| 7.506 ± 2 | 4^- | | 12 ± 2 | p, α | 27, 29 |
| 7.513 ± 2 | | | < 4 | γ, p | 25 |
| 7.528 ± 2 | $2^-; 1$ | | 16.5 ± 3.0 | γ, p, α | 25, 27, 29 |
| 7.532 ± 5 | | | 75 | p, α | 27, 29 |
| 7.555 ± 2 | (1^-) | | 30 | p | 27 |
| 7.584 ± 2 | | | 9 ± 2 | γ, p, α | 25, 27, 29 |
| 7.685 ± 2 | $3^+, 4^+$ | | 36 ± 4 | p, α | 27, 29 |
| 7.729 ± 4 | ≥ 1 | | 66 ± 5 | p, α | 27, 29 |

Table 18.13: Energy levels of ^{18}F ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | K^π | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------------|--|-------------------|--|--------------------------|---------------|
| 7.763 \pm 4 | | | 70 | p | 27 |
| 7.878 \pm 3 | ≥ 2 | | 20 | p, α | 27, 29 |
| 7.899 \pm 2 | (2 ⁻) | | 38 | p, α | 7, 8, 29 |
| 7.941 \pm 12 | (1 ⁺) | | 112 | p, α | 7, 8, 29 |
| 8.064 \pm 6 | ≥ 4 | | 60 | p, α | 27, 29 |
| 8.115 \pm 8 | | | 96 | p | 27 |
| 8.209 \pm 2 | 2 ⁻ | | 52 | p, α | 27, 29 |
| 8.238 \pm 2 | 4 ⁺ | | 20 | p | 27 |
| 9.207 \pm 15 ^b | 3, 4 ⁻ ; 0 | | | p, d, α | 16, 17, 18 |
| 9.50 | 2, 3 ⁺ ; 0 | | | n, d, α | 16, 18 |
| 9.58 \pm 20 ^c | 6 ⁺ | 1 ⁺ | | d, α | 9, 10, 11, 22 |
| 10.58 \pm 50 | | | | | 11 |
| 11.22 \pm 30 | 7 ⁺ | 1 ⁺ | | d, α | 9, 10, 11, 22 |
| 13.83 | 4 ⁻ , 5 ⁺ | | 60 | d, α | 18 |
| 14.02 | 4 ⁻ , 5 ⁺ | | 60 | d, α | 18 |
| 14.10 | 4 ⁻ , 5 ⁺ | | 60 | d, α | 18 |
| 14.18 \pm 40 | (8 ⁺) | (1 ⁺) | | d, α | 9, 10, 11 |
| 15.09 | 4 ⁻ , 5 ⁺ | | | d, α | 18 |
| 15.34 | 5 ⁺ , 6 ⁻ | | | d, α | 18 |
| 15.79 \pm 100 | | | | | 11 |
| 16.07 | 4 ⁻ , 5 ⁺ | | 220 | d, α | 18 |
| 16.72 | 4 ⁻ , 5 ⁺ | | 60 | d, α | 18 |
| 17.43 | 4 ⁻ , 5 ⁺ , 6 ⁻ | | 70 | d, α | 18 |
| 18.62 \pm 120 | | | | | 11 |
| (19.00 \pm 150) | | | (500 \pm 150) | γ , ^3He | 12 |
| 20.1 \pm 200 | (2 ⁻ ; 1) | | 1600 \pm 100 | γ , ^3He | 12 |
| 22.7 \pm 200 | (2 ⁻ ; 1) | | 1200 \pm 100 | γ , ^3He | 12 |
| (24.1 \pm 200) | | | (1400 \pm 300) | γ , ^3He | 12 |

^a See also Table 18.14 for radiative transitions and 18.15 for τ_m .

^b Uncertainty estimated by reviewer.

^c For other states with $E_x < 9.6$ MeV see footnote (e) in Table 18.17 of (1978AJ03) and Table 18.16 here. For other states with $10.0 < E_x < 19.6$ MeV see Table 18.16 here, and Tables 18.14 and 18.16 in (1978AJ03). These two tables in (1978AJ03) display the states deduced from the yields of the isospin-forbidden α_1 groups in $^{14}\text{N} + \alpha$ and $^{16}\text{O} + d$, respectively. (1976CH24) reports 151 isospin mixed natural-parity states with $10.4 < E_x < 17.5$ MeV [$^{14}\text{N}(\alpha, \alpha_1)$] and (1973JO13) reports 138 such states with $9.2 < E_x < 19.4$ MeV [$^{16}\text{O}(d, \alpha_1)$] of which 16 have $E_x > 17.5$ MeV. In the region $10.4 < E_x < 20.8$ MeV some 167 states with mixed isospin and natural parity have been reported. See also reaction 29.

$$\begin{aligned} 2. \text{ (a) } & {}^{10}\text{B}({}^9\text{Be}, n){}^{18}\text{F} & Q_m = 14.4539 \\ & \text{ (b) } & {}^{11}\text{B}({}^9\text{Be}, 2n){}^{18}\text{F} & Q_m = 2.9998 \end{aligned}$$

See (1986CU02) for production cross sections of 0.94 MeV γ -rays.

$$\begin{aligned} 3. \text{ (a) } & {}^{12}\text{C}({}^6\text{Li}, d){}^{16}\text{O} & Q_m = 5.6868 & E_b = 13.212 \\ & \text{ (b) } & {}^{12}\text{C}({}^6\text{Li}, \alpha){}^{14}\text{N} & Q_m = 8.7972 \\ & \text{ (c) } & {}^{12}\text{C}({}^6\text{Li}, {}^6\text{Li}){}^{12}\text{C} \end{aligned}$$

Cross sections for these and other charged particle channels have been measured for $E({}^6\text{Li}) = 1.9$ to 36 MeV [see (1978AJ03, 1983AJ01)].

The cross section for the isospin-forbidden α_1 group [to $^{14}\text{N}^*(2.31), 0^+, T = 1$] is 1 to 2% of the cross section to the allowed α_0 and α_2 groups for $E({}^6\text{Li}) = 3.2$ to 6 MeV while for 9 to 14 MeV it varies from 0.4 to 1.8%. At 20 MeV, the α_1 yield is 0.02% of the allowed yield. Structures are reported at $E({}^6\text{Li}) = 11.0$ and 13.0 MeV in the α_0 yield, at 11.5 and (13.0) MeV in the α_1 yield and at ≈ 11.7 and 12.8 MeV in the α_2 yield. A resonance is also reported in the α_1 yield at $E({}^6\text{Li}) = 4.2$ MeV: $E_x = 15.99 \pm 0.02$ MeV, $\Gamma_{\text{c.m.}} = 290 \pm 30$ keV, $J^\pi = 2^+$ (one-level BW fit). It is suggested that this resonance is due to 2^+ states with $T = 0$ and $T = 1$ which are unresolved. Cross sections for populating $^{16}\text{O}^*(8.87, 10.36, 11.08, 11.10)$ are reported by (1981GL02).

The excitation functions for the ${}^6\text{Li}$ ions to $^{12}\text{C}^*(0, 4.43)$ shows a single isolated structure at $E({}^6\text{Li}) = 22.8$ MeV, in the range 20–36 MeV, with $\Gamma \approx 0.8$ MeV. It is unlikely to be due to an isolated state in ^{18}F . At $E({}^6\text{Li}) = 20$ MeV analyzing power measurements are reported for many deuteron and α groups and for elastically scattered ${}^6\text{Li}$ ions. VAP measurements for elastic scattering are also reported at $E_{\vec{d}} = 9.0$ and 19.2 MeV (1983RU09) and at 150 MeV (1986TA1B, 1986KA1C).

Table 18.14: Radiative decays in ^{18}F ^a

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------|--------------|-------------|-----------------|---|
| 0.94 | $3^+; 0$ | 0 | 100 | |
| 1.04 | $0^+; 1$ | 0 | 100 | |
| 1.08 | $0^-; 0$ | 0 | 100 | |
| 1.12 | $5^+; 0$ | 0.94 | 100 | |
| 1.70 | $1^+; 0$ | 0 | 29.8 ± 1.3 | |
| | | 1.04 | 70.2 ± 1.3 | |
| 2.10 | $2^-; 0$ | 0 | 38 ± 1 | $\Gamma_\gamma = (4.6 \pm 2.2) \times 10^{-5} \text{ eV}$ |
| | | 0.94 | 31 ± 1 | $\Gamma_\gamma = (4.0 \pm 1.9) \times 10^{-5} \text{ eV}$ |
| | | 1.08 | 31 ± 1 | |
| 2.52 | $2^+; 0$ | 0 | 74.9 ± 1.8 | $\delta = 3.0 \pm 1.0$ |
| | | 0.94 | 21.5 ± 1.2 | $\delta = -(1.5 \pm 0.6)$ |
| | | 1.70 | 3.9 ± 0.6 | $\delta = 0.94 \pm 0.4$ |
| 3.06 | $2^+; 1$ | 0 | 23.2 ± 0.8 | |
| | | 0.94 | 76.7 ± 0.8 | |
| | | 1.04 | 0.11 ± 0.03 | |
| 3.13 | $1^-; 0$ | 0 | 39 ± 2 | $\delta = +(0.07 \pm 0.05)$ |
| | | | | $\Gamma_\gamma = (5.7 \pm 2) \times 10^{-4} \text{ eV}$ |
| | | 1.04 | 34 ± 2 | $\Gamma_\gamma = (7.3 \pm 2.7) \times 10^{-4} \text{ eV}$ |
| | | 1.08 | 25 ± 2 | $\Gamma_\gamma = (4.8 \pm 1.8) \times 10^{-4} \text{ eV}$ |
| | | 1.70 | 2.0 ± 0.5 | $\delta = +(0.22 \pm 0.15)$ |
| 3.36 | $3^+; 0$ | 0 | 45 ± 5 | |
| | | 0.94 | 9 ± 3 | |
| | | 1.70 | 40 ± 4 | |
| | | 2.10 | < 3 | |
| | | 2.52 | 6 ± 3 | $\delta = -0.4_{-0.5}^{+0.3}$ |
| 3.72 | $1^+; 0$ | 0 | 5 ± 2 | |
| | | 1.04 | 91 ± 2 | $\Gamma_\gamma = (1.3 \pm 0.2) \times 10^{-3} \text{ eV}^c$ |
| | | 3.06 | 4 ± 2 | |
| 3.79 | $3^-; 0$ | 2.10 | 68 ± 4 | $\delta = -(0.22 \pm 0.06)$ |
| | | 2.52 | 2.2 ± 1.1 | |
| | | 3.06 | 30 ± 3 | $\delta = -(0.09 \pm 0.09)$ |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------------|--------------|-------------|---------------|--|
| 3.84 | $2^+; 0$ | 0 | 38 ± 2 | $\delta = -(1.8 \pm 0.5)$ |
| | | 0.94 | 8.9 ± 1.4 | $\delta = -(0.3 \pm 0.3)$ |
| | | 1.70 | 3.0 ± 1.0 | |
| | | 3.06 | 50 ± 3 | $\delta = -(0.1 \pm 0.3)$ |
| 4.12 | $3^+; 0$ | 0 | 5 ± 3 | |
| | | 3.06 | 95 ± 3 | $\delta = +0.06 \pm 0.07$ |
| 4.23 | $2^-; 0$ | 0 | 23 ± 2 | $\delta = 0.15 \pm 0.15$ |
| | | 0.94 | 49 ± 3 | $\delta = 0.0 \pm 0.2$ |
| | | 1.08 | 3.2 ± 1.0 | |
| | | 1.70 | 9.3 ± 1.2 | |
| | | 2.10 | 15 ± 5 | |
| | | 3.13 | 0.9 ± 0.6 | |
| 4.36 | 1^+ | 3.06 | 100 | |
| 4.40 | $4^-; 0$ | 0.94 | 13 ± 4 | $\delta = -(0.2 \pm 0.3)$ |
| | | 1.12 | 60 ± 6 | $\delta = -(0.2 \pm 0.2)$ |
| | | 2.10 | 27 ± 3 | |
| 4.65 | $4^+; 1$ | 0.94 | 17 ± 3 | |
| | | 1.12 | 83 ± 3 | $\delta = 0.15 \pm 0.15$ |
| 4.75 | $0^+; 1$ | 0 | 92 ± 4 | |
| | | 1.70 | 8 ± 4 | |
| 4.85 ^b | $5^-; 0$ | 1.12 | 65 ± 4 | |
| | | 3.79 | 35 ± 4 | |
| 4.86 | $1^-; 0$ | 1.04 | 65 ± 11 | |
| | | 1.08 | 8 ± 6 | |
| | | 3.06 | 23 ± 7 | $\delta = -(0.4 \pm 0.4)$ |
| | | 3.13 | 4 ± 3 | |
| 4.96 | $2^+; 1$ | 0 | 100 | $\delta = 1.2 \pm 0.7$ |
| 5.30 | $4^+; 0$ | 0.94 | 9 ± 2 | $\delta = -(0.3 \pm 0.1)$ |
| | | 1.12 | 7 ± 2 | $\delta = -(1.1 \pm 0.5)$ |
| | | 2.52 | 78 ± 3 | $\Gamma_\gamma = 12 \pm 4 \text{ meV}^c$ |
| | | 3.36 | 5 ± 1 | $\delta = 2.5 \pm 0.8$ |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------|--------------|-------------|----------------|--|
| | | 4.65 | 1.3 ± 0.3 | |
| 5.50 | $3^{(-)}; 0$ | 3.06 | 100 | $\Gamma_\gamma = 2.1 \pm 0.7 \text{ meV}^c$ |
| 5.603 | 1^+ | 0 | 16.7 ± 2.3 | |
| | | 1.04 | 3.8 ± 1.2 | |
| | | 3.06 | 79.5 ± 5.9 | $\Gamma_\gamma = 0.48 \pm 0.05 \text{ eV}^c$ |
| 5.605 | $1^-; 0 + 1$ | 0 | 6.7 ± 1.2 | |
| | | 1.04 | 4.2 ± 0.8 | |
| | | 1.08 | 54.3 ± 3.1 | $\Gamma_\gamma = 0.87 \pm 0.07 \text{ eV}^c$ |
| | | 3.06 | 2.6 ± 1.4 | |
| | | 3.13 | 32.2 ± 2.5 | $\delta = -0.05 \pm 0.02$ |
| 5.67 | $1^-; 0 + 1$ | 0 | 6.2 ± 0.4 | $\delta = -0.01 \pm 0.04$ |
| | | 1.04 | 8.1 ± 0.7 | |
| | | 1.08 | 52 ± 3 | $\Gamma_\gamma = 0.46 \pm 0.06 \text{ eV}^c$ |
| | | 1.70 | 0.8 ± 0.3 | |
| | | 2.10 | 0.4 ± 0.2 | |
| | | 3.06 | 4.0 ± 0.4 | $\delta = 0.04 \pm 0.06$ |
| | | 3.13 | 28.5 ± 2.0 | $\delta = +0.10 \pm 0.03$ |
| 5.79 | $2^-; 0$ | 0.94 | 40 ± 8 | |
| | | 1.08 | 60 ± 8 | |
| 6.10 | $4^-; 0$ | 0.94 | 4.9 ± 0.9 | $\Gamma_\gamma = 51 \pm 10 \text{ meV}^c$ |
| | | 1.12 | 55 ± 3 | |
| | | 2.10 | 27 ± 2 | |
| | | 3.79 | 1.4 ± 0.3 | |
| | | 4.12 | 1.8 ± 0.3 | |
| | | 4.40 | 0.7 ± 0.3 | |
| | | 4.65 | 8.7 ± 0.7 | |
| 6.10 | $(1^+); 0$ | 0 | 24 ± 3 | |
| | | 0.94 | 11 ± 3 | |
| | | 2.10 | 20 ± 6 | |
| | | 3.06 | 45 ± 5 | |
| 6.14 | $0^+; 1$ | 0 | 50 ± 3 | $\Gamma_\gamma > 1.6 \text{ eV}$ |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------|--------------|-------------|-----------------|--|
| 6.16 | $3^+; 1$ | 1.70 | 12 ± 2 | $\Gamma_\gamma = 0.96 \pm 0.26 \text{ eV}^c$ |
| | | 3.72 | 36 ± 3 | |
| | | 4.36 | 2.1 ± 0.4 | |
| | | 5.603 | 0.19 ± 0.02 | |
| | | 0 | 0.2 ± 0.2 | |
| | | 0.94 | 51 ± 3 | |
| | | 1.12 | 1.0 ± 0.1 | |
| | | 2.52 | 5.5 ± 0.4 | |
| | | 3.06 | 1.3 ± 0.3 | |
| | | 3.79 | 11.6 ± 1.3 | |
| | | 3.84 | 25.0 ± 1.6 | |
| | | 4.12 | 1.5 ± 0.3 | |
| | | 4.23 | 0.9 ± 0.3 | |
| 6.242 | $3^-; 0 + 1$ | 4.40 | 2.0 ± 0.2 | $\Gamma_\gamma = 0.80 \pm 0.11 \text{ eV}^c$ |
| | | 0.94 | 4.6 ± 0.3 | |
| | | 2.10 | 71.5 ± 3.0 | |
| | | 3.36 | 1.1 ± 0.4 | |
| | | 3.79 | 10.6 ± 0.5 | |
| | | 3.84 | 1.0 ± 0.2 | |
| | | 4.12 | 0.5 ± 0.2 | |
| | | 4.23 | 7.8 ± 0.4 | |
| | | 4.40 | 2.9 ± 0.3 | |
| | | 0.94 | 4.1 ± 0.3 | |
| 6.241 | $3^-; 0 + 1$ | 2.10 | 71.2 ± 3.0 | $\Gamma_\gamma = 0.73 \pm 0.11 \text{ eV}^c$ |
| | | 3.36 | 0.8 ± 0.3 | |
| | | 3.79 | 11.6 ± 0.6 | |
| | | 3.84 | 0.9 ± 0.2 | |
| | | 4.12 | 1.1 ± 0.4 | |
| | | 4.23 | 8.2 ± 0.4 | |
| | | 4.40 | 2.1 ± 0.3 | |
| | | 0 | (100) | |
| 6.26 | $1^+; 0$ | 0 | (100) | |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | | |
|-------------|----------------|-------------|----------------|--|-----------------------------|
| 6.28 | $2^+; 1$ | 0 | 0.3 ± 0.1 | $\Gamma_\gamma = 1.8 \pm 0.5 \text{ eV}^c$ | |
| | | 0.94 | 67 ± 3 | | |
| | | 1.04 | 1.3 ± 0.1 | | |
| | | 1.70 | 5.7 ± 0.6 | | |
| | | 2.10 | 1.2 ± 0.3 | | |
| | | 2.52 | 0.3 ± 0.2 | | |
| | | 3.13 | 0.7 ± 0.3 | | |
| | | 3.36 | 2.3 ± 0.3 | | |
| | | 3.72 | 1.4 ± 0.5 | | |
| | | 3.84 | 15.8 ± 1.4 | | |
| | | 4.12 | 3.9 ± 0.2 | | |
| | | 4.36 | 0.5 ± 0.4 | | |
| 6.31 | $3^+; 0$ | 0 | 4.0 ± 0.7 | $\Gamma_\gamma = 0.17 \pm 0.04 \text{ eV}^c$ | |
| | | 0.94 | 10.6 ± 1.0 | | |
| | | 1.70 | 3.0 ± 0.8 | | |
| | | 2.52 | 4.0 ± 0.5 | | |
| | | 3.06 | 57 ± 3 | | $\delta = -(0.03 \pm 0.10)$ |
| | | 3.72 | 1.4 ± 0.7 | | |
| | | 3.84 | 4.6 ± 1.0 | | |
| | | 4.12 | 2.4 ± 1.7 | | |
| | | 4.96 | 13.0 ± 1.5 | | $\delta = -(0.01 \pm 0.14)$ |
| | | 6.39 | $2^+; 0 + 1$ | | 0 |
| 0.94 | 75 ± 3 | | | $\delta = -(0.25 \pm 0.10)$ | |
| 1.70 | 6.8 ± 1.7 | | | | |
| 3.84 | 14.1 ± 1.6 | | | $\delta = 0.1 \pm 0.2$ | |
| 4.12 | 2.3 ± 0.5 | | | | |
| 6.48 | $3^+; 0$ | 0 | 13 ± 2 | $\Gamma_\gamma = 74 \pm 21 \text{ meV}^c$ | |
| | | 0.94 | 33 ± 2 | | |
| | | 1.12 | 10 ± 2 | | |
| | | 1.70 | 4 ± 2 | | |
| | | 2.52 | 4 ± 2 | | |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------|--------------------|-------------|----------------|---|
| 6.57 | $5^+; 0$ | 3.06 | 21 ± 3 | $\Gamma_\gamma = 26 \pm 5 \text{ meV}^{\text{c,d}}$ |
| | | 3.79 | 4 ± 2 | |
| | | 3.84 | 9 ± 2 | |
| | | 4.96 | 2 ± 2 | |
| | | 0.94 | 15.2 ± 1.6 | |
| 6.64 | $2^-; 1$ | 3.36 | 83 ± 3 | $\Gamma_\gamma = 1.4 \pm 0.4 \text{ eV}^{\text{c}}$ |
| | | 5.30 | 2.3 ± 0.6 | |
| | | 0.94 | 8.9 ± 0.6 | |
| | | 2.10 | 58 ± 3 | |
| | | 3.13 | 22.0 ± 1.3 | |
| 6.78 | $4^+; 0$ | 3.72 | 0.9 ± 0.2 | $\Gamma_\gamma = 0.31 \pm 0.08 \text{ eV}^{\text{c}}$ $\delta = -(0.35 \pm 0.18)$ $\delta = -(1.4 \pm 1.1)$ $\delta = 0.13 \pm 0.13$ |
| | | 3.79 | 2.4 ± 0.2 | |
| | | 4.12 | 1.0 ± 0.3 | |
| | | 4.86 | 2.6 ± 0.2 | |
| | | 5.50 | 4.0 ± 0.3 | |
| 6.80 | $1^+, 2, 3^+; (0)$ | 0.94 | 12.6 ± 0.9 | |
| | | 1.12 | 25.2 ± 1.3 | |
| | | 4.65 | 62 ± 2 | |
| | | 0 | 20 ± 2 | |
| | | 0.94 | 20 ± 2 | |
| 6.88 | $3, 4^-; 0$ | 3.06 | 50 ± 3 | |
| | | 3.84 | 3.0 ± 1.6 | |
| | | 4.96 | 7.0 ± 1.7 | |
| | | 2.10 | 9 ± 2 | |
| | | 4.65 | 91 ± 2 | |
| 7.34 | $1^-; 1$ | 0 | 4 ± 0.5 | |
| | | 1.08 | 54 ± 2 | |
| | | 2.10 | 18 ± 1 | |
| | | 3.06 | 1 ± 0.5 | |
| | | 3.13 | 8 ± 0.5 | |

Table 18.14: Radiative decays in ^{18}F ^a (continued)

| E_i (MeV) | $J_i^\pi; T$ | E_f (MeV) | Branch (%) | |
|-------------|--------------|-------------|--------------|--|
| | | 4.23 | 15 ± 0.6 | |
| 7.48 | (2) | 0.94 | 100 | |
| 7.52 | | 0.94 | 5 ± 4 | |
| | | 2.10 | 7 ± 5 | |
| | | 3.79 | 33 ± 5 | |
| | | 4.40 | 55 ± 7 | |
| 7.53 | 2^- | 0 | 10 ± 3 | |
| | | 0.94 | 14 ± 6 | |
| | | 2.10 | 50 ± 9 | |
| | | 3.79 | 26 ± 7 | |
| 7.59 | | 0 | 18 ± 7 | |
| | | 0.94 | 14 ± 12 | |
| | | 1.12 | 9 ± 7 | |
| | | 4.65 | 59 ± 16 | |

^a For earlier references see Tables 18.11 in (1978AJ03) and 18.12 in (1983AJ01). See these tables also for upper limits for transitions to other states.

^b (1982FR15): see reactions 6 and 23.

^c Γ_γ = total radiative width for this state.

^d $\Gamma_\alpha = \Gamma \approx 560$ eV, $\Gamma_p < 4.5$ eV.

For fusion studies see (1982DE30, 1987PA12). For references to earlier work and for additional comments see (1978AJ03, 1983AJ01). See also (1980KR1F), (1984MU1D; theor.) and ^{12}C in (1985AJ01), ^{14}N in (1986AJ01) and ^{16}O in (1986AJ04).

4. $^{12}\text{C}(^9\text{Be}, t)^{18}\text{F}$ $Q_m = -4.4754$

Angular distributions are reported at $E(^9\text{Be}) = 12$ to 27 MeV to $^{18}\text{F}_{\text{g.s.}}$ and to the unresolved states at 1 MeV: see (1983AJ01). For excitation functions see (1982HU06, 1983JA09).

5. $^{12}\text{C}(^{11}\text{B}, \alpha n)^{18}\text{F}$ $Q_m = -2.7015$

Table 18.15: Lifetime measurements of some ^{18}F states

| $^{18}\text{F}^*$ (MeV) | $J^\pi; T$ | τ_m | Refs. |
|-------------------------|------------|--|-----------------------------|
| 0.94 | $3^+; 0$ | 67.6 ± 2.5 psec | mean ^a |
| 1.04 | $0^+; 1$ | 2.7 ± 0.4 fsec | ^b |
| | | <u>2.2 ± 0.6 fsec</u> | (1983CA21) |
| | | 2.55 ± 0.45 fsec | see (1983CA21) ^c |
| 1.08 | $0^-; 0$ | 27.5 ± 1.9 psec | mean ^a |
| 1.12 | $5^+; 0$ | 234 ± 10 nsec | mean ^b |
| 1.70 | $1^+; 0$ | 0.971 ± 0.30 psec | (1982BA40) |
| | | <u>0.897 ± 0.057 psec</u> | (1983MO16) ^d |
| | | 0.955 ± 0.027 psec | mean |
| 2.10 | $2^-; 0$ | 5.12 ± 0.56 psec | (1982BA40) |
| | | <u>4.93 ± 0.78 psec</u> | (1983MO16) |
| | | 5.06 ± 0.46 psec | mean |
| 2.52 | $2^+; 0$ | 0.605 ± 0.029 psec | (1982BA40) |
| | | <u>0.554 ± 0.045 psec</u> | (1983MO16) |
| | | 0.590 ± 0.024 psec | mean |
| 3.06 | $2^+; 1$ | < 1.2 fsec | (1982BA40) ^{a,e} |
| 3.13 | $1^-; 0$ | 0.403 ± 0.018 psec | (1982BA40) |
| | | <u>0.343 ± 0.022 psec</u> | (1983MO16) |
| | | 0.39 ± 0.02 psec ^A | |
| 3.36 | $3^+; 0$ | 0.435 ± 0.041 psec | (1982BA40) |
| | | <u>0.451 ± 0.034 psec</u> | (1983MO16) |
| | | 0.44 ± 0.03 psec ^A | |
| 3.72 | $1^+; 0$ | 4 ± 2 fsec | (1973RO04) |
| | | $2.7^{+4.1}_{-2.7}$ fsec ^A | (1982BA40) ^e |
| 3.79 | $3^-; 0$ | 1.91 ± 0.17 psec | (1982BA40) |
| | | <u>1.90 ± 0.20 psec</u> | (1983MO16) |
| | | 1.91 ± 0.13 psec | mean |
| 3.84 | $2^+; 0$ | 17.4 ± 3.6 fsec | (1982BA40) |
| | | <u>21 ± 4 fsec</u> | (1983MO16) |
| | | 19.0 ± 2.7 fsec | mean |
| 4.12 | $3^+; 0$ | 91 ± 22 fsec | (1973RO06) |

Table 18.15: Lifetime measurements of some ^{18}F states (continued)

| $^{18}\text{F}^*$ (MeV) | $J^\pi; T$ | τ_m | Refs. |
|-------------------------|--------------|--------------------|------------|
| 4.23 | $2^-; 0$ | 110 ± 15 fsec | (1973RO06) |
| 4.36 | $1^+; 0$ | 27 ± 10 fsec | (1973RO06) |
| 4.40 | $4^-; 0$ | 58 ± 12 fsec | (1973RO06) |
| 4.65 | $4^+; 1$ | < 10 fsec | (1973RO06) |
| 4.85 | $5^-; 0$ | 5.2 ± 0.9 psec | (1982KO24) |
| 4.86 | $1^-; 0$ | 66 ± 18 fsec | (1973RO06) |
| 4.96 | $2^+; 1$ | < 4 fsec | (1973RO06) |
| 5.30 | $4^+; 0$ | 30 ± 5 fsec | (1973RO05) |
| 5.50 | $3^{(-)}; 0$ | 63 ± 25 fsec | (1973RO06) |
| 5.79 | $2^-; 0$ | 15 ± 10 fsec | (1973RO06) |

A = adopted.

^a See Table 18.12 in (1978AJ03).

^b See Table 18.13 in (1983AJ01).

^c See also (1985KE1C).

^d See also (1982MO09).

^e See also (1983MO16).

$\tau_m = 5.2 \pm 0.9$ psec for $^{18}\text{F}^*(4.85)$ [$5^-; T = 0$]. The E1 strength is $(3.4 \pm 0.6) \times 10^{-6}$ W.u. for the transition to $^{18}\text{F}^*(1.12)$ [$5^+; T = 0$] and the E2 strength is 14.8 ± 2.6 W.u. for that to $^{18}\text{F}^*(3.79)$ [$3^-; 0$]. The latter strength, which is that of a highly collective transition, corresponds to a quadrupole moment $Q_0 = 395 \pm 35$ mb and suggests that $^{18}\text{F}^*(4.85)$ is the 5^- state of a (strongly decoupled) $K^\pi = 1^-$ band (1982KO24). See also Tables 18.14 and 18.15.

6. $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$ $Q_m = 4.4152$

The non-resonant S -factor for this reaction, $S \approx 0.7$ MeV · b: see (1978AJ03). A number of resonances have been observed for $E_\alpha < 3$ MeV: see Table 18.16. Studies of these, principally by the Toronto and Queen's groups [see references in (1978AJ03, 1983AJ01)] in conjunction with work on $^{14}\text{N}(\alpha, \alpha)$, $^{16}\text{O}(^3\text{He}, \text{p})$, $^{17}\text{O}(\text{p}, \gamma)$ and $^{17}\text{O}(\text{p}, \alpha)$ [see Tables 18.18, 18.19, 18.20] have led to the determination of branching ratios, mixing ratios and widths (Table 18.14), lifetimes (Table 18.15) and the E_x , J^π and K^π assignments for ^{18}F states with $E_x < 6.9$ MeV. The reader is referred to the series of papers by the Toronto group for the most complete and definitive arguments on the parameters of the low-lying states of ^{18}F .

Table 18.16: Resonances in $^{14}\text{N} + \alpha$ below $E_\alpha = 5 \text{ MeV}$ ^a

| E_α (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $(2J + 1)\Gamma_\gamma\Gamma_\alpha/\Gamma$ (eV) | $J^\pi; T$ | E_x (MeV) |
|-------------------------------|--------------------|---------------------------------|---|--|--------------------|
| | | | $< 2 \times 10^{-5}$ | | 4.657 |
| 0.559 | γ | | $(2.8 \pm 0.5) \times 10^{-4}$ | 1; 0 | 4.850 |
| 0.698 | | | $< 0.5 \times 10^{-4}$ | 2 ⁺ ; 1 | 4.958 |
| 1.136 \pm 3 | γ | | 0.084 \pm 0.004 | 4 ⁺ ; 0 | 5.299 |
| 1.398 \pm 3 | γ | | 0.022 \pm 0.003 | 3 ⁽⁻⁾ ; 0 | 5.502 |
| 1.527 | γ, α_0 | | 1.44 \pm 0.14 | 1 ⁺ | 5.603 ^e |
| 1.529 \pm 2 | γ, α_0 | < 1.2 | 2.60 \pm 0.21 | 1 ⁻ ; 0 + 1 | 5.604 ^f |
| 1.618 \pm 2 | γ, α_0 | < 0.8 | 1.4 \pm 0.2 ^b | 1 ⁻ ; 0 + 1 | 5.673 ^g |
| 1.765 \pm 4 | γ | | 0.047 \pm 0.018 | 2 ⁻ ; 0 | 5.788 |
| 2.160 \pm 4 | γ | | 0.20 \pm 0.04 | 4 ⁻ ; 0 | 6.095 |
| 2.166 \pm 7 | γ, α_0 | | 0.08 \pm 0.03 | 1, 2, 3 ⁽⁻⁾ ; 0 | 6.100 |
| | | | ^c | | |
| 2.348 \pm 3 | γ, α_0 | < 0.8 | | 3 ⁻ ; 0 + 1 | 6.241 ^h |
| 2.372 \pm 3 | γ, α_0 | < 3 | | 1 ⁺ ; (0) | 6.260 ⁱ |
| | | | ^d | | |
| 2.438 \pm 4 | γ | | 0.52 \pm 0.12 | 3 ⁺ ; 0 | 6.311 |
| 2.532 \pm 4 | γ | | 1.6 \pm 0.4 | 2 ⁺ ; 0 + 1 | 6.384 |
| | γ | | 0.16 \pm 0.06 | 3 ⁺ ; (0) | 6.480 |
| 2.767 \pm 4 | γ, α_0 | (< 0.8) | 0.29 \pm 0.06 | 5 ⁺ ; 0 | 6.567 |
| 2.870 \pm 4 | γ, p_0 | < 1.6 | 2.7 \pm 0.5 | 2 ⁻ ; 1 | 6.647 |
| 2.870 \pm 6 | α_0 | 93 \pm 5 | $\Gamma_\alpha/\Gamma = 0.85$ | 1 ⁻ | 6.647 |
| | | | 0.12 \pm 0.07 | 4 ⁺ ; 0 | 6.78 |
| | | | < 0.2 | 1 ⁺ , 2, 3 ⁺ ; (0) | 6.803 |
| 3.080 \pm 6 | p_0, α_0 | 101 \pm 5 | | 2 ⁻ | 6.810 |
| 3.576 \pm 4 | α_0 | < 4 | | (4 ⁺) | 7.196 |
| 3.67 | α_0 | 45 \pm 10 | | (1 ⁺) | 7.27 |
| 3.72 | p_0, α_0 | 53 \pm 6 | | (3 ⁻) | 7.31 |
| 4.00 | p_0, α_0 | 35 | | (3 ⁻) | 7.53 |
| 4.05 | p_0, α_0 | 60 | | | 7.57 |
| 4.11 | p_0, α_0 | 40 | | | 7.61 |

Table 18.16: Resonances in $^{14}\text{N} + \alpha$ below $E_\alpha = 5 \text{ MeV}$ ^a (continued)

| E_α (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $(2J + 1)\Gamma_\gamma\Gamma_\alpha/\Gamma$ (eV) | $J^\pi; T$ | E_x (MeV) |
|-------------------------------|------------------|---------------------------------|---|-------------------|----------------|
| 4.28 | p_0, α_0 | 120 | | | 7.74 |
| 4.50 | p_0, α_0 | 30 | | (2 ⁻) | 7.92 |
| 4.55 | p_0, α_0 | 70 | | (1 ⁺) | 7.95 |

^a References are displayed in Tables 18.13 of (1972AJ02, 1978AJ03). Higher resonances observed in $^{14}\text{N}(\alpha, \alpha_1)$ are listed in Table 18.14 of (1978AJ03).

^b $\omega\gamma = 0.45 \pm 0.02$ (1982BE29).

^c ≤ 0.07 for $^{18}\text{F}^*(6.11, 6.16)$ (1973RO03).

^d ≤ 0.03 for $^{18}\text{F}^*(6.28)$ (1973RO03).

^e $\Gamma_\alpha = 42.8 \pm 1.6 \text{ eV}$, $\Gamma_\gamma = 0.485 \pm 0.046 \text{ eV}$, $l_\alpha = 0$ (1980MA26). See also Table 18.19.

^f $\Gamma_\alpha = 32.0 \pm 2.1 \text{ eV}$, $\Gamma_\gamma = 0.891 \pm 0.074 \text{ eV}$, $l_\alpha = 1$. ΔE_x for $^{18}\text{F}^*(5.603, 5.605)$ is $1.84 \pm 0.04 \text{ keV}$ (1980MA26). See also Table 18.19.

^g $\Gamma_\alpha = 130 \pm 5 \text{ eV}$, $\Gamma_\gamma = 1.4 \pm 0.3 \text{ eV}$, $l_\alpha = 1$ (1980MA26).

^h This resonance corresponds to two states at $E_x = 6240$ and 6242 keV . The lower member of the doublet (both of which have $J^\pi = 3^-$ and mixed isospin) has $\Gamma_\alpha = 133 \pm 4 \text{ eV}$, $\Gamma_\gamma = 0.80 \pm 0.11 \text{ eV}$; the higher has $\Gamma_\alpha = 137 \pm 4 \text{ eV}$, $\Gamma_\gamma = 0.73 \pm 0.11 \text{ eV}$ (1979KI12).

ⁱ $\Gamma_\alpha = 580 \pm 12 \text{ eV}$, $\Gamma_p = 25_{-25}^{+35} \text{ eV}$ (1979KI12).

No evidence is seen for the excitation of the (forbidden) state at $E_x = 4.753 \text{ MeV}$ [$J^\pi = 0^+$; $T = 1$] (1981LE1A, 1983LE08). See also (1985GO27; applications) and (1982BA1D, 1985AR1A, 1985KL1A, 1985MA1A, 1986HE1C, 1987GOZX; astrophysics).

$$7. \ ^{14}\text{N}(\alpha, p)^{17}\text{O} \qquad Q_m = -1.1914 \qquad E_b = 4.4152$$

Observed resonances are displayed in Table 18.16. See also ^{17}O in (1986AJ04).

$$8. \text{ (a) } ^{14}\text{N}(\alpha, \alpha)^{14}\text{N} \qquad E_b = 4.4152$$

$$\text{ (b) } ^{14}\text{N}(\alpha, 2\alpha)^{10}\text{B} \qquad Q_m = -11.6123$$

$$\text{ (c) } ^{14}\text{N}(\alpha, ^6\text{Li})^{12}\text{C} \qquad Q_m = -8.7972$$

Observed anomalies in the elastic scattering [reaction (a)] are exhibited in Table 18.16. Resonances in the α_1 isospin-forbidden yield are displayed in Table 18.14 of (1978AJ03). In the α_1 study, carried out for $E_\alpha = 7.6 \rightarrow 16.9 \text{ MeV}$, a partial-wave analysis involving a method of removing ambiguities and parametrizing S -matrix elements gives the level parameters of 151 isospin

mixed, natural-parity states in ^{18}F with $10.4 < E_x < 17.5$ MeV. Many of these states have also been reported in the $^{16}\text{O}(\text{d}, \alpha_1)$ reaction [Table 18.16 of (1978AJ03)]. The agreement is best for low-lying 2^+ or 4^+ states, and is quite good for 3^- and 5^- states, while for high- J states the greater centrifugal barrier for $^{16}\text{O} + \text{d}$ at the same E_x relatively suppresses high- J states in the $^{16}\text{O} + \text{d}$ work. A study of the energy dependence of averaged intensities of the partial waves shows some indication that the lower partial waves reconserve isospin as E_x increases.

The total cross sections for formation of ^{10}B and ^6Li have been studied for $E_\alpha = 21$ to 42 MeV [see (1978AJ03)], as has the cross section for production of 1.64 and 2.31 MeV γ -rays from threshold to $E_\alpha = 26$ MeV [(1985DY05)]: see for astrophysical implications.] See also (1984RE14, 1986CO11) and (1986ALZZ; theor.).

$$\begin{aligned} 9. \text{ (a) } & ^{14}\text{N}(^6\text{Li}, \text{d})^{18}\text{F} & Q_m &= 2.940 \\ & \text{(b) } & ^{14}\text{N}(^6\text{Li}, \text{d}\alpha)^{14}\text{N} & Q_m &= -1.4751 \end{aligned}$$

At $E(^6\text{Li}) = 36$ MeV angular distributions have been measured for the deuteron groups to $^{18}\text{F}^*(5.34 [4^+], 6.56 [5^+], 9.58, 11.2, 14.1)$. Angular correlations lead to $J^\pi = 6^+$ and 8^+ for $^{18}\text{F}^*(9.58, 14.1)$ and the data are consistent with $J^\pi = 7^+$ for $^{18}\text{F}^*(11.2)$ (1983ET02). For the earlier work see (1978AJ03).

$$10. \ ^{14}\text{N}(^7\text{Li}, \text{t})^{18}\text{F} \quad Q_m = 1.947$$

At $E(^7\text{Li}) = 36$ MeV the $K^\pi = 1^+$ band appears to be selectively populated. States at $E_x = 9.58 \pm 0.02, 11.22 \pm 0.03$ and 14.18 ± 0.04 MeV are strongly populated. It is suggested that the first two are the 6^+ and 7^+ members of that band: see reaction 8. [Angular distributions are reported for $^{18}\text{F}^*(1.70, 2.10, 2.52, 3.36, 4.40, 5.30, 6.57, 9.58, 11.22, 14.18)$.] See (1978AJ03) for the earlier work, and (1986NE1A; $E(^{14}\text{N}) = 150$ MeV). See also (1986CO11; theor.).

$$\begin{aligned} 11. \text{ (a) } & ^{14}\text{N}(^{11}\text{B}, ^7\text{Li})^{18}\text{F} & Q_m &= -4.249 \\ & \text{(b) } & ^{14}\text{N}(^{13}\text{C}, ^9\text{Be})^{18}\text{F} & Q_m &= -6.2324 \end{aligned}$$

These reactions have been studied at $E(^{11}\text{B}) = 115$ MeV and $E(^{13}\text{C}) = 105$ MeV. Differential cross sections at three angles are reported for the transitions to $^{18}\text{F}^*(9.58, 10.57 \pm 0.07, 11.2)$ in reaction (a) and to $^{18}\text{F}^*(5.30, 6.57, 9.58, 10.60 \pm 0.08, 11.2)$ in reaction (b). In addition to these states $^{18}\text{F}^*(14.18)$ is strongly excited in both reactions, and transitions to $^{18}\text{F}^*(15.79 \pm 0.10, 18.62 \pm 0.12)$ are also reported: see (1983AJ01).

12. (a) $^{15}\text{N}(^3\text{He}, \gamma)^{18}\text{F}$ $Q_m = 14.1596$
 (b) $^{15}\text{N}(^3\text{He}, \alpha)^{14}\text{N}$ $Q_m = 9.7445$ $E_b = 14.1596$

Excitation functions have been measured for $E(^3\text{He}) = 2.5$ to 16 MeV for the γ_0 and $\gamma_{1\rightarrow 4}$ yields. Resonances are observed corresponding to $E_x = (19.00 \pm 0.15)$ [$\gamma_{1\rightarrow 4}$], 20.1 ± 0.2 [γ_0 , $\gamma_{1\rightarrow 4}$], 22.7 ± 0.2 [γ_0 , $\gamma_{1\rightarrow 4}$] and (24.1 ± 0.2) MeV [$\gamma_{1\rightarrow 4}$], with $\Gamma_{\text{c.m.}} = 0.5 \pm 0.15$, 1.6 ± 0.1 , 1.2 ± 0.1 and 1.4 ± 0.3 MeV, respectively. The γ_0 yield is dominated by $^{18}\text{F}^*(20.1)$ (1983WA05) [see for $(2J + 1)\Gamma_{^3\text{He}}\Gamma_\gamma$ values]. It is suggested that the structures decaying by γ_0 have $J^\pi = 2^-$ (and possibly $T = 1$) (1983WA05). For analyzing power measurements at $E(^3\text{He}) = 33$ MeV see (1986DR03).

13. $^{15}\text{N}(^6\text{Li}, \text{t})^{18}\text{F}$ $Q_m = -1.636$

At $E(^6\text{Li}) = 30$ MeV preferential excitation of odd-parity states of ^{18}F below $E_x = 5$ MeV is reported. Angular distributions of the tritons to $^{18}\text{F}^*(0, 0.94, 2.10, 4.40)$ [$J^\pi = 1^+, 3^+, 2^-, 4^-$] are all strongly forward peaked: see (1978AJ03).

14. (a) $^{15}\text{N}(^{11}\text{B}, ^8\text{Li})^{18}\text{F}$ $Q_m = -13.049$
 (b) $^{15}\text{N}(^{12}\text{C}, ^9\text{Be})^{18}\text{F}$ $Q_m = -12.1194$

These reactions have been studied with $E(^{11}\text{B}) = E(^{12}\text{C}) = 115$ MeV. Reaction (a) is dominated by the transitions to $^{18}\text{F}^*(1.12)$ [presumably the $J^\pi = 5^+$ state, although the group is unresolved] and to $^{18}\text{F}^*(7.15, 9.45)$ [$J^\pi = (7^-)$ and (6^-)]. In reaction (b) no single state is strongly preferentially populated. Differential cross sections for $^{18}\text{F}^*(4.40, 6.10, 7.15, 9.45)$ [$J^\pi = 4^-, (5^-), (7^-), (6^-)$], are fitted by FRDWBA: see (1983AJ01).

15. $^{16}\text{O}(\text{d}, \gamma)^{18}\text{F}$ $Q_m = 7.5256$

The capture cross section rises from $0.1 \mu\text{b}$ at $E_d = 0.4$ MeV to $25 \mu\text{b}$ at 3.5 MeV: Γ_γ over this range is ≈ 2 eV: see (1972AJ02).

16. (a) $^{16}\text{O}(\text{d}, \text{n})^{17}\text{F}$ $Q_m = -1.6241$ $E_b = 7.5256$
 (b) $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ $Q_m = 1.9191$

Excitation functions and polarization studies have been carried out to $E_d = 17$ MeV [see (1978AJ03, 1983AJ01)] and at $E_d \approx 5.6$ to 8.3 MeV (1985GR1B; p_0, p_3, p_4). Structures attributed to states in ^{18}F are displayed in Table 18.17. See also ^{17}O and ^{17}F in (1986AJ04), (1985JOZZ, 1986SE1B) and (1983DA31; applications).

Table 18.17: Maxima in the yields of $^{16}\text{O} + \text{d}$ ^a

| E_d (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | E_x (MeV) |
|-----------------------|--------------------------------|------------------------------|---------------|-------------|
| 0.895 | p_1, α_0 | 210 ± 25 | | (8.320) |
| 1.048 | p_1, d_0, α_0 | 88 ± 10 | 1^+ | 8.456 |
| 1.199 | α_0 | 230 ± 30 | | (8.590) |
| 1.298 | p_1, d_0, α_0 | 13 ± 3 | | (8.678) |
| 1.325 | d_0, α_0 | | | (8.702) |
| 1.482 | α_0 | 40 ± 5 | | (8.842) |
| 1.563 | d_0, α_0 | 121 ± 15 | | (8.914) |
| 1.616 | α_0 | 19 ± 15 | | (8.961) |
| 1.765 | d_0, α_0 | 141 ± 10 | | (9.093) |
| 1.885 | p_0, p_1, d_0, α_0 | 108 ± 12 | $3, 4^-; 0$ | 9.200 |
| 2.22 | n_0, α_0 | | $2, 3^+; 0$ | 9.50 |
| 2.28 | α_0 | | $2, 3^+; 0$ | (9.55) |
| 2.34 | n_0, p_1 | | | (9.60) |
| 2.55 | p_1 | | | (9.79) |
| 2.92 | n, p_0, p_1 | | | 10.12 |
| 3.05 | α_0 | | $3, 4^-; 0$ | 10.24 |
| 3.13 | $n, p_1, \alpha_0, \alpha_1$ | | $\geq 2; 0$ | 10.31 |
| 3.37 | n_0, p_0, p_1, α_1 | | | 10.52 |
| 3.47 | α_0 | | $4, 5^+; 0$ | 10.61 |
| 3.68 | n, p_0, p_1, α_1 | | 2^+ | 10.79 |
| 3.80 | p_0, α_0 | | $\geq 2^+; 0$ | 10.90 |
| 3.94 | n, p_1, α_1 | | | 11.03 |
| 3.95 | p_1, α_0 | ≈ 35 | $3, 4^-; 0$ | 11.03 |
| 4.07 | n, p_1 | | | 11.14 |
| 4.38 | p_1, α_0 | | $4, 5^+; 0$ | 11.42 |
| 4.57 | α_0 | | $5, 6^-; 0$ | 11.58 |
| 4.80 | d_0, α_0 | | $\geq 3; 0$ | 11.79 |
| 4.93 | α_0 | | $5, 6^-; 0$ | 11.90 |
| 5.05 ± 15 | α_4 | 40 | | 12.01 |
| 5.11 | $\alpha_0, \alpha_2, \alpha_4$ | 60 | $4, 5^+; 0$ | 12.06 |
| 5.17 | α_0 | 55 | $T = 0$ | 12.12 |

Table 18.17: Maxima in the yields of $^{16}\text{O} + \text{d}$ ^a (continued)

| E_d (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | E_x (MeV) |
|-----------------------|--------------------------------|------------------------------|-------------|-------------|
| 5.32 | α_0 | 70 | | 12.25 |
| 5.34 | α_0, α_2 | 170 | | 12.27 |
| 5.40 | α_0, α_4 | 130 | | 12.32 |
| 5.47 | α_4 | 80 | | 12.38 |
| 5.49 | $\alpha_2, \alpha_3, \alpha_4$ | 120 | | 12.40 |
| 5.59 | α_0, α_2 | 120 | | 12.49 |
| 5.65 | α_0, α_2 | 140 | | 12.54 |
| 5.77 | α_0 | 180 | 2^+ | 12.65 |
| 5.80 | $\alpha_0, \alpha_2, \alpha_4$ | 160 | | 12.68 |
| 5.81 | α_3, α_4 | 80 | 5^- | 12.69 |
| 5.91 | α_2 | 160 | | 12.77 |
| 6.00 | α_0 | 120 | | 12.85 |
| 6.11 | α_0, α_4 | 120 | | 12.95 |
| 6.19 | α_2, α_3 | 200 | $\geq 4; 0$ | 13.02 |
| 6.25 | α_0, α_4 | 150 | $T = 0$ | 13.08 |
| 6.30 | α_0, α_2 | 160 | | 13.12 |
| 6.34 | α_0, α_3 | 160 | $5, 6^-; 0$ | 13.16 |
| 6.38 | α_0, α_3 | 145 | $T = 0$ | 13.19 |
| 6.43 | α_0, α_2 | 120 | | 13.24 |
| 6.46 | α_0, α_4 | 100 | | 13.26 |
| 6.54 | α_0, α_2 | 135 | | 13.33 |
| 6.61 | $\alpha_2, \alpha_3, \alpha_4$ | 120 | | 13.40 |
| 6.64 | α_0, α_2 | 200 | | 13.42 |
| 6.66 | α_0 | 100 | | 13.44 |
| 6.72 | α_2 | 100 | | 13.79 |
| 6.73 | α_2 | 100 | | 13.50 |
| 6.80 | α_2, α_3 | 140 | | 13.56 |
| 6.84 | $\alpha_0, \alpha_2, \alpha_4$ | 150 | | 13.60 |
| 6.94 | α_0, α_3 | 90 | | 13.69 |
| 7.10 | α_3, α_4 | 60 | $4^-, 5^+$ | 13.83 |
| 7.27 | α_3 | 150 | | 13.98 |

Table 18.17: Maxima in the yields of $^{16}\text{O} + \text{d}$ ^a (continued)

| E_d (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | E_x (MeV) |
|-----------------------|--|------------------------------|-------------|-------------|
| 7.31 | α_2 | 60 | $4^-, 5^+$ | 14.18 |
| 7.34 | $\alpha_0, \alpha_3, \alpha_4$ | 200 | | 14.04 |
| 7.38 | α_0, α_3 | 210 | | 14.08 |
| 7.41 | α_3 | 60 | $4^-, 5^+$ | 14.02 |
| 7.49 | α_0 | 220 | | 14.18 |
| 7.58 | α_0 | 200 | $\geq 4; 0$ | 14.26 |
| 7.62 | α_4 | 85 | | 14.29 |
| 7.66 | $\alpha_0, \alpha_2, \alpha_4$ | 130 | $T = 0$ | 14.33 |
| 7.67 | $\alpha_0, \alpha_2, \alpha_3, \alpha_4$ | 250 | $T = 0$ | 14.34 |
| 7.74 | α_3 | 200 | $3^+, 4^-$ | 14.40 |
| 7.80 | α_0, α_4 | 70 | | 14.45 |
| 7.82 | α_0, α_2 | 225 | | 14.47 |
| 7.99 | α_4 | 200 | | 14.62 |
| 8.02 | α_0 | 150 | | 14.65 |
| 8.03 | α_3 | 310 | | 14.66 |
| 8.07 | α_0 | 120 | | 14.69 |
| 8.08 | α_3, α_4 | 310 | | 14.70 |
| 8.21 | α_2 | 250 | | 14.82 |
| 8.25 | α_4 | 380 | | 14.85 |
| 8.30 | $\alpha_0, \alpha_2, \alpha_3$ | 210 | | 14.90 |
| 8.34 | α_4 | 115 | | 14.93 |
| 8.37 | α_0 | 130 | | 14.96 |
| 8.37 | α_0, α_3 | 250 | | 14.96 |
| 8.40 | α_0 | 310 | | 14.99 |
| 8.43 | α_4 | 120 | | 15.01 |
| 8.52 | α_3, α_4 | 160 | $4^-, 5^+$ | 15.09 |
| 8.52 | α_2 | 150 | | 15.09 |
| 8.56 | α_2 | 220 | | 15.13 |
| 8.58 | α_4 | 180 | | 15.15 |
| 8.61 | α_0, α_3 | 200 | | 15.17 |
| 8.65 | α_0, α_2 | 135 | | 15.21 |

Table 18.17: Maxima in the yields of $^{16}\text{O} + \text{d}$ ^a (continued)

| E_d (MeV \pm keV) | Particles out | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | E_x (MeV) |
|-----------------------|--------------------------------|------------------------------|-----------------|-------------|
| 8.72 | α_2, α_4 | 120 | | 15.27 |
| 8.76 | α_2 | 160 | | 15.30 |
| 8.79 | α_0 | 200 | | 15.33 |
| 8.80 | $\alpha_0, \alpha_3, \alpha_4$ | 200 | $5^+, 6^-$ | 15.34 |
| 8.89 | α_3 | 110 | | 15.42 |
| 8.93 | α_3, α_4 | 190 | | 15.46 |
| 8.97 | α_2, α_4 | 210 | | 15.49 |
| 9.00 | α_0, α_2 | 190 | | 15.52 |
| 9.62 | α_3 | 220 | $4^-, 5^+$ | 16.07 |
| 10.35 | α_3 | 60 | $4^-, 5^+$ | 16.72 |
| 11.15 | α_3 | 70 | $4^-, 5^+, 6^-$ | 17.43 |

^a For references see Table 18.15 in (1978AJ03). This table does not include the structures in α_1 leading to mixed isospin states in ^{18}F : for the latter see Table 18.16 in (1978AJ03).

17. $^{16}\text{O}(\text{d}, \text{d})^{16}\text{O}$

$$E_b = 7.5256$$

The yield of elastically scattered deuterons and elastic polarization measurements have been reported for $E_d = 0.65$ to 56 MeV: see (1978AJ03, 1983AJ01). More recent measurements are those by (1985GR1B) [excitation functions for $E_d \approx 5.6$ to 8.3 MeV] and the polarization studies at $E_d = 20.5$ MeV (1984FR14; TAP), 56 MeV (1986MA32; VAP, TAP) and 200, 400 and 700 MeV (1987NG01; VAP, TAP). See also (1982AO06, 1983AO03, 1986NG1B, 1986YA1R, 1987MA09; theor.) and ^{16}O in (1986AJ04).

18. $^{16}\text{O}(\text{d}, \alpha)^{14}\text{N}$

$$Q_m = 3.1104$$

$$E_b = 7.5256$$

The yields of various groups of α -particles have been measured for $E_d \leq 20$ MeV: see (1978AJ03, 1983AJ01). The yield curves have been fitted in terms of a large number of states in ^{18}F : see Tables 18.17 here, and 18.16 in (1978AJ03).

A detailed study by (1973JO13) of the isospin-forbidden α_1 yield, analyzed by S -matrix theory, identifies a large number of isospin mixed states in ^{18}F , possibly as many as 138 with $9.2 < E_x < 19.4$ MeV. The reaction mechanism appears to be almost entirely compound nuclear. The isospin

impurity, averaged over 1 MeV intervals, is 3–10% for the above E_x range. The average coherence width increases from ≈ 100 keV at $E_x = 14$ MeV to ≈ 500 keV at $E_x = 20$ MeV. The level densities appear to be consistent with predictions of the Fermi-gas model (1973JO13). See also (1985JOZZ). [For mixed isospin states observed in $^{14}\text{N}(\alpha, \alpha_1)$ see Table 18.14 in (1978AJ03).] Polarization measurements are reported for $E_d = 6.8$ to 16 MeV: see (1978AJ03, 1983AJ01).

$$19. \text{}^{16}\text{O}(\text{d}, \text{}^6\text{Li})^{12}\text{C} \qquad Q_m = -5.6868 \qquad E_b = 7.5256$$

Polarization measurements for the transitions to $^{12}\text{C}^*(0, 4.4)$ are reported at $E_d = 18$ and 22 MeV (1987TA07; VAP, TAP) and 51.7 MeV (1986YA12; VAP; also to $^{12}\text{C}^*(14.1)$).

$$20. \text{}^{16}\text{O}(\text{t}, \text{n})^{18}\text{F} \qquad Q_m = 1.2690$$

See (1987BO16; applied) and (1983AJ01).

$$21. \text{}^{16}\text{O}(\text{}^3\text{He}, \text{p})^{18}\text{F} \qquad Q_m = 0.321$$

Excitation energies derived from measurements of γ -rays are displayed in Table 18.18 together with l -assignments obtained from distorted-wave analyses, and J^π ; T and K^π assignments from branching ratios, radiative widths, linear polarization, γ -ray angular distributions and τ_m measurements [see also Tables 18.14 and 18.15]. Studies of this reaction, together with the work on $^{14}\text{N}(\alpha, \gamma)$ and $^{17}\text{O}(\text{p}, \gamma)$, have defined the low-lying states of ^{18}F .

The g -factor of $^{18}\text{F}^*(0.94)$ [$J^\pi = 3^+$] is $+(0.56 \pm 0.05)$: see (1983AJ01). The circular polarization of the 1.08 MeV \rightarrow g.s. γ -ray, $P_\gamma = (-10 \pm 18) \times 10^{-4}$ (1982AH07), $(2.7 \pm 5.7) \times 10^{-4}$ (1985BI03), $(1.6 \pm 5.6) \times 10^{-4}$ (1985EV03), $(1.7 \pm 5.8) \times 10^{-4}$ (1987PA07). The weak pion-nucleon coupling constant deduced from the weighted average of all recent P_γ measurements [$(1.2 \pm 3.9) \times 10^{-4}$] is $(0.3_{-0.3}^{+1.0}) \times 10^{-7}$. Together with PNC matrix elements in other experiments this suggests that the isovector weak NN interaction may be strongly suppressed compared with the isoscalar weak NN interaction (1985EV03, 1987PA07). For a measurement of the ICC of the 0.94, 1.02, 1.04 and 1.08 MeV γ -rays see (1986KR04). See also (1978AJ03, 1983AJ01), (1984BE1A; applications), (1987ER05; theor.) and ^{19}Ne .

$$22. \text{}^{16}\text{O}(\alpha, \text{d})^{18}\text{F} \qquad Q_m = -16.3211$$

Table 18.18: States in ^{18}F from $^{16}\text{O}(^3\text{He}, p\gamma)^{18}\text{F}$ ^a

| E_x (keV) ^b | l ^a | $J\pi; T$ ^c | $K\pi$ ^c |
|--------------------------|------------------|----------------------------|---------------------|
| 0 | 0 | $1^+; 0$ | 0^+ |
| 937.1 ± 0.4 | 2 | $3^+; 0$ | 0^+ |
| 1040.9 ± 0.5 | 0 | $0^+; 1$ | |
| 1080.1 ± 0.5 | | $0^-; 0$ | 0^- |
| 1119.0 ± 0.6 | 4 | $5^+; 0$ | 0^+ |
| 1701.4 ± 0.7 | 0 | $1^+; 0$ | 1^+ |
| 2099.9 ± 0.6 | | $2^-; 0$ | 0^- |
| 2523.4 ± 0.7 | 2 | $2^+; 0$ | 1^+ |
| 3061.2 ± 0.5 | 2 | $2^+; 1$ | |
| 3132.8 ± 0.6 | | $1^-; 0$ | 1^- |
| 3358.2 ± 1.0 | | $3^+; 0$ | 1^+ |
| 3725.4 ± 0.8 | | $1^+; 0$ | |
| 3790.6 ± 0.9 | | $3^-; 0$ | 1^- |
| 3838.4 ± 0.7 | 2 | $2^+; 0$ | |
| 4114.5 ± 0.9 | | $3^+; 0$ | |
| 4225.8 ± 0.7 | | $2^{(-)}; 0$ | (1^-) |
| 4361.0 ± 0.7 | | $1^{(+)}$ | |
| 4398.1 ± 0.7 | | $3^-, 4^-; 0$ ^d | (0^-) |
| 4652 ± 2 | 4 | $4^+; 1$ | |
| 4753 ± 3 | | $(0^+; 1)$ | |
| 4860 ± 2 | | $1^{(-)}; 0$ | |
| 4963.6 ± 0.8 | | $2^+; 1$ | |
| 5297.6 ± 1.5 | | 4^+ | 1^+ |
| 5502 ± 2 | | $3^{(-)}; 0$ | |
| 5603 ± 2 | | $1^-; 0 + 1$ | |
| 5669 ± 2 | | $1^-; 0 + 1$ | |
| 5785 ± 3 | | $2^-; 0$ | |
| 6097.4 ± 1.4 | | $4^-; 0$ | 1^- |
| 6108 ± 3 | | $1, 2, 3^{(-)}; 0$ | |
| 6138.3 ± 1.0 | | $0^+; 1$ | |
| 6164.0 ± 1.0 | | $3^+; 1$ | |

Table 18.18: States in ^{18}F from $^{16}\text{O}(^3\text{He}, p\gamma)^{18}\text{F}$ ^a (continued)

| E_x (keV) ^b | l ^a | $J^\pi; T$ ^c | K^π ^c |
|---------------------------|------------------|-------------------------|----------------------|
| 6241.2 ± 1.0 | | $3^-; 1$ | |
| 6263 ± 3 | | 1^+ | |
| 6284.0 ± 1.0 | | $2^+; 0 + 1$ | |
| 6310.5 ± 0.8 | | $3^+; 0$ | |
| 6383 ± 3 | | $2^+; 0 + 1$ | |
| 6480 ± 2 | | $3^+; (0)$ | |
| 6567.0 ± 1.5 | | 5^+ | 1^+ |
| 6643.0 ± 1.5 | | $2^-; 1$ | |
| 6777 ± 2 ^c | | 4^+ | |
| 6803.0 ± 1.5 | | $1^+, 2, 3^+; (0)$ | |
| 6878 ± 2 ^c | | $3^{(-)}, 4^-; (0)$ | |

^a For earlier results derived from measurements of proton spectra and of γ -rays, see Table 18.18 in (1972AJ02). See also Tables 18.14 and 18.15 here.

^b (1973RO03): γ -ray measurements.

^c See Table 18.17 in (1978AJ03).

^d See p. 179 of (1979KI12).

Angular distributions of the deuteron groups to $^{18}\text{F}^*(1.12)$ [$J^\pi = 5^+$] have been studied at $E_\alpha = 28.0$ to 33.6 MeV: see (1983AJ01). At $E_\alpha = 65.3$ MeV a number of angular distributions are reported to ^{18}F states with $E_x \leq 11.4$ MeV: $^{18}\text{F}^*(9.49, 10.54)$ are suggested to have $J^\pi = 6^-$ and 7^+ respectively (1986KA36). See, however, reactions 9 and 10.

23. $^{16}\text{O}(^6\text{Li}, \alpha)^{18}\text{F}$ $Q_m = 6.050$

Angular distributions have been measured at $E(^6\text{Li}) = 5.5$ to 34 MeV [see (1983AJ01)] and at $E(^6\text{Li}) = 48$ MeV (1984CO05; $\alpha_0, \alpha_1, \alpha_4$). (1982FR15) report the excitation of a state at $E_x = 4848.3 \pm 0.5$ keV which decays (35 ± 4)% to $^{18}\text{F}^*(3.79)$ [$E_\gamma = 1056.8 \pm 0.4$ keV] and (65 ± 4)% to $^{18}\text{F}^*(1.12)$. Alpha- γ angular correlations are consistent with $J^\pi = 5^-$, and $T = 0$ (1982FR15). See also (1986GL02) and (1986IC01; theor.).

24. (a) $^{16}\text{O}(^{11}\text{B}, ^9\text{Be})^{18}\text{F}$ $Q_m = -8.290$
 (b) $^{16}\text{O}(^{13}\text{C}, ^{11}\text{B})^{18}\text{F}$ $Q_m = -11.153$
 (c) $^{16}\text{O}(^{14}\text{N}, ^{12}\text{C})^{18}\text{F}$ $Q_m = -2.7468$

Table 18.19: Excited states of ^{18}F from $^{17}\text{O}(\text{p}, \gamma)^{18}\text{F}$ ^a

| E_x (keV) | E_x (keV) |
|--------------------|--------------------|
| 937.18 ± 0.06 | 3724.19 ± 0.22 |
| 1041.55 ± 0.08 | 3791.49 ± 0.22 |
| 1080.54 ± 0.12 | 3839.17 ± 0.22 |
| 1121.36 ± 0.15 | 4115.90 ± 0.25 |
| 1700.81 ± 0.18 | 4360.15 ± 0.26 |
| 2100.61 ± 0.10 | 5603.38 ± 0.27 |
| 2523.35 ± 0.18 | 5604.86 ± 0.28 |
| 3061.84 ± 0.18 | 5668 ± 2 |
| 3133.87 ± 0.15 | 6136.47 ± 0.33 |

^a See also Table 18.20 here, and Table 18.17 in (1983AJ01).

See (1983AJ01).

$$25. \ ^{17}\text{O}(\text{p}, \gamma)^{18}\text{F} \quad Q_m = 5.6065$$

Gamma-ray measurements lead to the very accurate E_x determinations for ^{18}F states below 6.2 MeV: see Table 18.19. Observed resonances are displayed in Table 18.20; branching ratios, radiative widths and multipole mixing ratios are shown in Table 18.14; and τ_m in Table 18.15.

The direct capture cross section has been studied for $E_p = 0.3$ to 1.9 MeV: $^{18}\text{F}^*(5.603, 5.605, 5.668, 5.786)$ have $J^\pi = 1^+, 1^-, 1^-$ and 2^- . The 1^- states have mixed isospin. For astrophysical considerations see (1978AJ03, 1983AJ01) and (1982RO1A).

$$26. \ ^{17}\text{O}(\text{p}, \text{n})^{17}\text{F} \quad Q_m = -3.5432 \quad E_b = 5.6065$$

Observed resonances are displayed in Table 18.20. Analyzing power measurements are reported at $E_{\bar{p}} = 135$ MeV (1983PUZZ; n_0).

$$27. \ ^{17}\text{O}(\text{p}, \text{p})^{17}\text{O} \quad E_b = 5.6065$$

Table 18.20: Resonances in $^{17}\text{O} + \text{p}$ ^a

| E_p (keV) | Yield of | $\Gamma_{\text{c.m.}}$ (keV) | $(2J + 1)\Gamma_\gamma\Gamma_p/\Gamma$ (eV) | $J^\pi; T$ | E_x (MeV \pm keV) |
|---------------------------|-------------------------|------------------------------|---|--|-----------------------|
| 517.0 \pm 1.0 | γ, α_0 | 0.24 \pm 0.03 | 0.26 \pm 0.05 | 4 ⁻ ; 0 | 6.095 |
| 525 | α_0 | 0.034 \pm 0.003 | | (1 ⁺) | 6.102 |
| 561.2 \pm 1.0 | γ | ≤ 1 | 2.2 \pm 0.6 | 0 ⁺ ; 1 | 6.136 |
| 587.1 \pm 1.0 | γ, p_0, α_0 | 14 \pm 0.5 | 6.7 \pm 1.8 | 3 ⁺ ; 1 | 6.161 |
| 670.5 \pm 1.0 | γ, p_0, α_0 | 0.19 \pm 0.03 | (c) | 3 ⁻ ; 0 + 1 | 6.239 |
| 673.0 | γ, α_0 | 0.18 \pm 0.04 | (c) | 3 ⁻ ; 0 + 1 | 6.242 |
| 690 \pm 4 | α_0 | 0.60 \pm 0.12 | ≤ 0.02 | 1 ⁺ ; 0 | 6.258 |
| 714.2 \pm 1.0 | γ, p_0, α_0 | 10.0 \pm 0.5 | 9.1 \pm 2.3 | 2 ⁺ ; 1 | 6.281 |
| 741 \pm 2 | γ, p_0, α_0 | 0.95 \pm 0.14 | 0.64 \pm 0.17 | 3 ⁺ ; 0 | 6.306 |
| 826 \pm 2 | γ, α_0 | 0.40 \pm 0.09 | 0.60 \pm 0.18 | 2 ⁺ ; 0 + 1 | 6.386 |
| 926 \pm 2 | γ, α_0 | 0.40 \pm 0.10 | 0.36 \pm 0.15 | 3 ⁺ ; 0 | 6.481 |
| 1015 | α_0 | 0.56 \pm 0.13 | ≤ 0.0023 | 5 ⁺ ; 0 | 6.565 |
| 1090 | α_0 | 80 \pm 2 | | 1 | 6.635 |
| 1098.9 \pm 0.4 | γ, α | 0.60 \pm 0.07 | 4.3 \pm 1.2 | 2 ⁻ ; 1 | 6.6439 |
| 1101 \pm 4 | α_0 | 89 \pm 5 | | | 6.646 |
| 1240 \pm 2 ^b | γ, p_0, α_0 | 9.2 \pm 1.0 | 2.8 \pm 0.7 | 4 ⁺ ; 0 | 6.777 |
| 1270 | γ, p_0 | ≤ 2 | 0.54 \pm 0.20 | 1 ⁺ , 2, 3 ⁺ ; 0 | 6.8031 \pm 1.5 |
| 1274 \pm 5 | α_0 | 88 \pm 2 | | 2 ⁻ | 6.809 |
| 1276 | α_0 | 3.0 \pm 0.5 | | (2 ⁺) | 6.811 |
| 1338 | α_0 | 5.0 \pm 1.0 | | (3 ⁻) | 6.870 |
| 1345 \pm 3 | γ, α_0 | ≤ 2 | 1.0 \pm 0.4 | 3, 4 ⁻ ; 0 | 6.876 |
| 1687.5 \pm 1 | α_0 | 6.5 | 3.9 | (4 ⁺); 0 | 7.199 |
| 1738 \pm 2 | α_0 | 46.5 | 8.8 | (1 ⁺); 0 | 7.247 |
| 1784 \pm 2 | p_0, α_0 | 38 | 47 | 3 ⁻ | 7.291 |
| 1810 \pm 4 | α_0 | 52 | 8.5 | (3 ⁻ ; 0) | 7.315 |
| 1832.5 \pm 1 | γ, p_0, p_1 | 16 \pm 2 | ^d | 1 ⁻ ; 1 | 7.336 |
| 1906 \pm 2 | p_0, p_1 | 14.6 \pm 1.4 | | 1 ⁺ | 7.406 |
| 1950 \pm 10 | α_0 | 140 | 5.6 | | 7.447 |
| 1957 \pm 2 | p_0 | 6 | | 1 ⁻ | 7.454 |
| 1983 \pm 2 | γ, p_1, α_0 | 12 \pm 3 | 1.5 | (2) | 7.478 |
| (1990 \pm 2) | p_0 | 32 | | (1 ⁻) | (7.485) |
| 2012 \pm 2 | p_0, α_0 | 12 \pm 2 | 7.2 | 4 ⁻ | 7.506 |
| 2020 \pm 2 | γ | < 4 | | | 7.513 |

Table 18.20: Resonances in $^{17}\text{O} + \text{p}$ ^a (continued)

| E_p (keV) | Yield of | $\Gamma_{\text{c.m.}}$ (keV) | $(2J + 1)\Gamma_\gamma\Gamma_p/\Gamma$ (eV) | $J^\pi; T$ | E_x (MeV \pm keV) |
|-----------------|------------------------------|------------------------------|---|-------------------------|-----------------------|
| 2036 ± 2 | $\gamma, p_0, p_1, \alpha_0$ | 16.5 ± 3.0 | 5.5 ^e | $2^-; 1$ | 7.528 |
| 2040 ± 5 | p_1, α_0 | 75 | | | 7.532 |
| 2064 ± 2 | p_0 | 30 | | (1^-) | 7.555 |
| 2095 ± 2 | $\gamma, p_0, p_1, \alpha_0$ | 9 ± 2 | 3.7 ^f | ^g | 7.584 |
| 2202 ± 2 | p_0, p_1, α_0 | 36 ± 4 | 25.1 | $3^+, 4^+$ ^g | 7.685 |
| 2248 ± 4 | p_1, α_0 | 66 ± 5 | 28.2 | ≥ 1 | 7.729 |
| 2284 ± 4 | p_1 | 70 | | | 7.763 |
| 2406 ± 3 | p_1, α_0 | 20 | 24.4 | ≥ 2 | 7.878 |
| 2429 ± 2 | α_0 | 38 | 42 | (2^-) | 7.899 |
| 2473 ± 12 | α_0 | 112 | 80 | (1^+) | 7.941 |
| 2603 ± 6 | p_1, α_0 | 60 | 11 | ≥ 4 | 8.064 |
| 2657 ± 2 | p_1 | 96 | | | 8.115 |
| 2757 ± 8 | p_0, α_0 | 52 | 63 | 2^- | 8.209 |
| 2788 ± 2 | p_0 | 20 | | 4^+ | 8.238 |
| 2928 | α_0 | ≈ 50 | | | 8.370 |
| 3915 ± 20 | n | 95 | | | 9.302 |
| (4163 ± 20) | n | 19 | | | (9.536) |
| 4235 ± 10 | n | 33 | | | 9.604 |
| 4330 ± 10 | n | 33 | | | 9.694 |
| 4490 ± 20 | n | ≈ 100 | | | 9.845 |
| (4790 ± 10) | n | 28 | | | (10.128) |
| 4900 ± 20 | n | ≈ 140 | | | 10.232 |

^a For references see Tables 18.18 in (1978AJ03, 1983AJ01).

^b See footnote ^d in Table 18.18 (1978AJ03).

^c This corresponds to a doublet of 3^- , mixed isospin states, separated by 2.09 ± 0.04 keV. $\omega_{\gamma p, \gamma} = 2.04 \pm 0.45$ eV for the lower resonance and 1.16 ± 0.26 eV for the higher one.

^d $\Gamma_\gamma = 3.5 \pm 1.0$ eV.

^e $\Gamma_\gamma = 0.44 \pm 0.10$ eV.

^f $\Gamma_\gamma = 0.11 \pm 0.03$ eV.

^g Assumed to be unresolved.

The elastic scattering has been studied for $E_p = 0.5$ to 13 MeV [see (1978AJ03, 1983AJ01)]: observed anomalies are displayed in Table 18.20. Analyzing powers have been measured at $E_{\bar{p}} = 89.7$ MeV (1985VO12; p_0).

$$28. \text{}^{17}\text{O}(\text{p}, \text{t})\text{}^{15}\text{O} \quad Q_{\text{m}} = -11.3257 \quad E_{\text{b}} = 5.6065$$

Analyzing powers have been reported at $E_{\text{p}} = 89.7$ MeV for the triton groups to a number of ^{15}O states (1985VO12).

$$29. \text{}^{17}\text{O}(\text{p}, \alpha)\text{}^{14}\text{N} \quad Q_{\text{m}} = 1.1914 \quad E_{\text{b}} = 5.6065$$

The yield of α_0 shows a number of resonances for $E_{\text{p}} = 0.49$ to 3.0 MeV: see Table 18.20. The R -matrix fit of (1979KI13), obtained from data from $E_{\text{p}} = 400$ to 1400 keV, confirms the earlier result [see, e.g., reaction 31 in (1978AJ03)] that a significant quantity of ^{17}O is burned up in the (p, γ) rather than in the (p, α) reaction for a wide range of stellar temperatures (1979KI13). See also (1982RO1A, 1987RO25) and (1986BA89; theor.).

$$30. \text{}^{17}\text{O}(\text{}^3\text{He}, \text{d})\text{}^{18}\text{F} \quad Q_{\text{m}} = 0.1130$$

At $E(^3\text{He}) = 15$ MeV DWBA analysis of angular distributions of deuteron groups corresponding to states of ^{18}F with $E_{\text{x}} < 5$ MeV have led to J^{π} values and spectroscopic information: see (1972AJ02). See also (1987ER05; theor.).

$$31. \text{}^{17}\text{O}(\text{}^{12}\text{C}, \text{}^{11}\text{B})\text{}^{18}\text{F} \quad Q_{\text{m}} = -10.351$$

See (1983AJ01).

$$32. \text{}^{18}\text{O}(\pi^+, \pi^0)\text{}^{18}\text{F} \quad Q_{\text{m}} = 2.949$$

See (1983AS01, 1984AS05).

$$33. \text{}^{18}\text{O}(\text{p}, \text{n})\text{}^{18}\text{F} \quad Q_{\text{m}} = -2.4387$$

(1983AN05) have studied the distribution of Gamow-Teller (GT) strength. At $E_{\text{p}} = 135$ MeV angular distributions have been studied to the 0^+ state at 1.04 MeV and to the 1^+ states $^{18}\text{F}^*(0, 1.70, 3.72, 4.36, 6.26)$ as well as to possible 1^+ ; $T = 1$ groups at $E_{\text{x}} = 9.9, 10.9$ and 11.9 MeV. 82% of the observed strength lies in the ground state group and 5.5% in the ($T = 1$) states. The observed GT strength is $\approx \frac{2}{3}$ of that expected from the simple sum rule (1983AN05). See also (1985TAZY), (1985WA24), (1986MA1P, 1987HI1B; applied) and the earlier work in (1978AJ03).

Table 18.21: Branching in $^{18}\text{Ne}(\beta^+)^{18}\text{F}$ ^a

| Decay to $^{18}\text{F}^*$ (MeV) | $J^\pi; T$ | E_{γ_0} (keV) | Branch (%) ^b | $\log f_0 t$ ^c |
|-------------------------------------|------------|---------------------------------|----------------------------------|---------------------------|
| 0 | $1^+; 0$ | | 92.11 ± 0.21 | 3.096 ± 0.004 |
| 1.04 ^d | $0^+; 1$ | 1041.5 ± 0.3 | 7.70 ± 0.21 | 3.473 ± 0.013 |
| 1.08 ^d | $0^-; 0$ | 1080.76 ± 0.13 ^b | $(2.07 \pm 0.28) \times 10^{-3}$ | 7.012 ± 0.059 |
| 1.70 | $1^+; 0$ | 1699.9 ± 0.3 ^e | 0.188 ± 0.006 | 4.477 ± 0.015 |

^a For the earlier work see Tables 18.19 in (1983AJ01) and 18.20 in (1978AJ03).

^b (1983AD03). See also (1982HE04).

^c Based on $\tau_{1/2} = 1672 \pm 8$ msec: see (1983AD03).

^d The splitting of the 0^+ and 0^- states is 39.20 ± 0.11 keV (1983AD03).

^e And 659.2 ± 0.3 keV for the γ -ray to $^{18}\text{F}^*(1.04)$ (1982HE04).

34. $^{18}\text{O}(^3\text{He}, t)^{18}\text{F}$ $Q_m = -1.675$

At $E(^3\text{He}) = 16$ MeV, the triton spectrum is dominated by strong groups to $^{18}\text{F}^*(0, 0.94)$ and to the 0^+ and 2^+ , $T = 1$ states $^{18}\text{F}^*(1.04, 3.06)$. Angular distributions have been studied to these and many other states at this energy and at $E(^3\text{He}) = 17.3$ MeV. At $E(^3\text{He}) = 33$ MeV A_y measurements for t_0 have been reported. See (1983AJ01) for references.

35. $^{18}\text{O}(^6\text{Li}, ^6\text{He})^{18}\text{F}$ $Q_m = -5.162$

See (1978AJ03).

36. $^{18}\text{Ne}(\beta^+)^{18}\text{F}$ $Q_m = 4.446$

The half-life of ^{18}Ne is 1672 ± 8 msec [see ^{18}Ne]. The decay is to $^{18}\text{F}^*(0, 1.04, 1.08, 1.70)$: see Table 18.21.

37. $^{19}\text{F}(\gamma, n)^{18}\text{F}$ $Q_m = -10.4320$

Cross sections have been reported to 30 MeV for the transitions to $^{18}\text{F}^*(0.94, 1.04, 1.08, 3.06, 3.13, 4.75)$: see (1983AJ01).

38. $^{19}\text{F}(\text{p}, \text{d})^{18}\text{F}$ $Q_{\text{m}} = -8.2074$

Angular distributions have been reported to many states of ^{18}F with $E_{\text{x}} \lesssim 6$ MeV: see Table 18.20 in (1983AJ01). See also (1986VAZV [$E_{\text{p}} = 18.6$ MeV; d_0, d_2]), (1983BEYY) and (1981CL05, 1983KI13; theor.).

39. $^{19}\text{F}(\text{d}, \text{t})^{18}\text{F}$ $Q_{\text{m}} = -4.1747$

See (1972AJ02, 1978AJ03).

40. $^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$ $Q_{\text{m}} = 10.1458$

See (1978AJ03).

41. $^{20}\text{Ne}(\text{p}, ^3\text{He})^{18}\text{F}$ $Q_{\text{m}} = -15.562$

See (1978AJ03).

42. $^{20}\text{Ne}(\text{d}, \alpha)^{18}\text{F}$ $Q_{\text{m}} = 2.7915$

At $E_{\text{d}} = 11$ MeV α -groups are observed to many states of ^{18}F with $E_{\text{x}} < 7$ MeV. Weak or absent (each $\leq 0.3\%$ of the total yield at 30°) are the groups corresponding to $^{18}\text{F}^*(1.04, 3.06, 4.66, 4.74, 4.96)$: $T = 1$. Measurements of the TAP for $E_{\text{d}} = 10.25$ to 12.0 MeV leads to assignments of $2^-, 1^+, 0^+, 1^-, 1^+, 3^+, 3^+$ to $^{18}\text{F}^*(4.23, 4.36, 4.75, 4.86, 5.603, 6.16, 6.48)$. See (1972AJ02, 1978AJ03, 1983AJ01) for references and for other results and (1987HI1B; applications).

43. $^{21}\text{Ne}(\text{p}, \alpha)^{18}\text{F}$ $Q_{\text{m}} = -1.744$

See (1987GOZX).

44. $^{23}\text{Na}(\text{d}, ^7\text{Li})^{18}\text{F}$ $Q_{\text{m}} = -12.176$

See (1984NE1A).

^{18}Ne
(Figs. 3 and 4)

GENERAL: (See also (1983AJ01).)

Model calculations: (1982ZH01, 1983BR29, 1984SA37, 1985RO1G).

Special states: (1982ZH01, 1983BI1C, 1983BR29, 1984SA37, 1985RO1G, 1986AN10, 1986AN07).

Electromagnetic transitions: (1982BR24, 1982RI04, 1983BR29, 1985AL21, 1986AN10).

Astrophysical questions: (1982WI1B, 1987WI11).

Complex reactions involving ^{18}Ne : (1986HA1B).

Pion capture and reactions (See also reaction 4.): (1982AS1B, 1983HO02, 1983JO06, 1983LI08, 1983LI1U, 1983OS1E, 1984JO01, 1984KA26, 1985GM01, 1986BA1C, 1986FO06, 1986GE06).

Other topics: (1983BI1C, 1983BR29, 1983BR1E, 1985AL21, 1985AN28, 1986ST15).

Ground state of ^{18}Ne : (1983ANZQ, 1985AN28).

For $B(E2)$ of $^{18}\text{Ne}^*(1.89)$ and other parameters see (1987RA01) and Table 2 in the Introduction here.

1. $^{18}\text{Ne}(\beta^+)^{18}\text{F}$ $Q_m = 4.446$

The half-life of ^{18}Ne is 1672 ± 8 msec: see (1978AJ03) and (1983AD03). The decay is primarily to $^{18}\text{F}^*(0, 1.04, 1.70)$. In addition there is an extremely weak branch [$(2.07 \pm 0.28) \times 10^{-3}\%$] to $^{18}\text{F}^*(1.08)$ [$J^\pi = 0^-$; $T = 0$] (1983AD03): see Table 18.21 for the parameters of the decay. The parity mixing in the $^{18}\text{F}^*(1.04, 1.08)$ $0^+ - 0^-$ doublet has been studied by (1983AD03). See also (1982HE04). For the earlier work see, in particular, (1981AD01, 1981HA06). See also (1983AD1C, 1983AD1B, 1983AD1D, 1984AD1E, 1985AD1A, 1985BR29) and (1984HA58, 1986BR1X, 1986HA1Q, 1986TO1D, 1987KI03; theor.).

2. $^{16}\text{O}(^3\text{He}, n)^{18}\text{Ne}$ $Q_m = -3.196$

See Table 18.24. See also (1983AJ01).

3. $^{16}\text{O}(^{10}\text{B}, ^8\text{Li})^{18}\text{Ne}$ $Q_m = -18.951$

Table 18.22: Energy levels of ^{18}Ne ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|--------------|--|-----------|------------|
| 0 | $0^+; 1$ | $\tau_{1/2} = 1672 \pm 8$ msec | β^+ | 1, 2, 4, 5 |
| 1.8873 ± 0.2 | 2^+ | $\tau_m = 0.67 \pm 0.06$ psec | γ | 2, 4, 5 |
| 3.3762 ± 0.4 | 4^+ | 4.4 ± 0.6 psec | γ | 2, 3, 5 |
| 3.5763 ± 2.0 | 0^+ | 4 ± 2 psec | γ | 2, 5 |
| 3.6164 ± 0.6 | 2^+ | 63_{-20}^{+30} fsec | γ | 2, 5 |
| 4.519 ± 8 | 1^- | $\Gamma \leq 20$ | (p) | 2, 5 |
| 4.590 ± 8 | 0^+ | ≤ 20 | (p) | 2, 5 |
| 5.090 ± 8 | $(2^+, 3^-)$ | 40 ± 20 | (p) | 2, 5 |
| 5.146 ± 7 | $(2^+, 3^-)$ | 25 ± 15 | | 2, 5 |
| 5.453 ± 10 | | ≤ 50 | | 5 |
| 6.297 ± 10 | (4^+) | ≤ 60 | | 2, 5 |
| 6.353 ± 10 | | ≤ 60 | | 5 |
| 7.059 ± 10 | $(1^-, 2^+)$ | 180 ± 50 | | 2 |
| 7.713 ± 10 | | ≤ 50 | | 2, 5 |
| 7.910 ± 10 | $(1^-, 2^+)$ | ≤ 50 | | 2 |
| 7.950 ± 10 | | ≤ 60 | | 5 |
| 8.086 ± 10 | | ≤ 50 | | 2 |
| 8.500 ± 30 | | ≤ 120 | | 2 |
| 9.201 ± 9 | | ≤ 50 | | 5 |

^a See also Table 18.23.

At $E(^{10}\text{B}) = 100$ MeV the angular distribution to $^{18}\text{Ne}^*(3.38)$ [$(d_{5/2})_{4+}^2$ state] which is preferentially populated has been studied. $^{18}\text{Ne}^*(1.89)$ is also observed. See (1983AJ01). See also (1983OS07; theor.).

$$4. \ ^{18}\text{O}(\pi^+, \pi^-)^{18}\text{Ne} \quad Q_m = -6.101$$

Angular distributions have been studied at $E_{\pi^+} = 164$ and 292 MeV [see (1983AJ01)] and at 48.3 MeV (1985AL15; to $^{18}\text{Ne}_{\text{g.s.}}$) and 100 to 292 MeV (1985SE08; to $^{18}\text{Ne}_{\text{g.s.}}$). The excitation functions for production of $^{18}\text{Ne}^*(0, 1.89)$ have been measured for $E_{\pi^+} = 80$ to 292 MeV: see (1983AJ01, 1985SE08). See also (1984BA1B, 1985GIZX, 1986ANZY), (1985GI01; theor.) and the ‘‘General’’ section here.

Table 18.23: Branching ratios and lifetimes of ^{18}Ne states ^a

| E_i (MeV) | J_i^π | E_f (MeV) | Branch (%) | τ_m (psec) |
|-------------|-----------|-------------|-------------------------|---------------------------|
| 1.89 | 2^+ | 0 | 100 | 0.67 ± 0.06 |
| 3.38 | 4^+ | 1.89 | 100 ^b | 4.4 ± 0.6 |
| 3.58 | 0^+ | 1.89 | 100 ^c | 4 ± 2 |
| 3.62 | 2^+ | 0 | 9 ± 2 | |
| | | 1.89 | 91 ± 2 ^d | $0.063^{+0.030}_{-0.020}$ |

^a For references see Table 18.24 in (1978AJ03).

^b Ground state decay is $< 1\%$.

^c Ground state decay is $< 5\%$.

^d The mixing ratio, δ , is consistent with 0.

5. $^{20}\text{Ne}(p, t)^{18}\text{Ne}$ $Q_m = -20.026$

Observed triton groups are displayed in Table 18.24 as are J^π derived from DWBA analysis of angular distributions: The 0_3^+ state, identified at $E_x = 4.59$ MeV, appears to have largely $s_{1/2}^2$ configuration based on its large downward shift with respect to the analog state in ^{18}O (1981NE09).

^{18}Na
(not observed)

^{18}Na has not been observed; its atomic mass excess has been estimated to be 25.32 MeV; it is then unbound with respect to proton emission by 1.6 MeV: see (1978AJ03). See also (1986AN07) and (1983ANZQ; theor.).

^{18}Mg , etc.
(not observed)

See (1986AN07) and (1983ANZQ; theor.).

Table 18.24: States in ^{18}Ne from $^{16}\text{O}(^3\text{He}, n)$ and $^{20}\text{Ne}(p, t)$ ^a

| E_x (MeV \pm keV) | | $\Gamma_{c.m.}$ (keV) ^b | $J\pi$ ^{a,b} |
|-----------------------------|----------------|------------------------------------|-----------------------|
| A | B | | |
| 0 | | | 0^+ |
| 1.8873 ± 0.2 | 1.886 ± 10 | | 2^+ |
| 3.3762 ± 0.4 | 3.375 ± 10 | | 4^+ |
| 3.5763 ± 2.0 | 3.580 ± 10 | | 0^+ |
| 3.6164 ± 0.6 | 3.612 ± 10 | | 2^+ |
| 4.513 ± 13 | 4.522 ± 10 | ≤ 20 | 1^- |
| 4.587 ± 13 | 4.592 ± 10 | ≤ 20 | 0^+ |
| 5.075 ± 13 | 5.099 ± 10 | 40 ± 20 | $(2^+, 3^-)$ |
| 5.141 ± 10 | 5.151 ± 10 | 25 ± 15 | $(2^+, 3^-)$ |
| | 5.453 ± 10 | ≤ 50 | |
| 6.291 ± 30 ^c | 6.297 ± 10 | ≤ 60 | (4^+) |
| | 6.353 ± 10 | ≤ 60 | |
| 7.062 ± 12 ^a | | 180 ± 50 | $(1^-, 2^+)$ |
| 7.712 ± 20 | 7.713 ± 10 | ≤ 50 | |
| 7.915 ± 12 ^a | | ≤ 50 | $(1^-, 2^+)$ |
| | 7.949 ± 10 | ≤ 60 | |
| 8.100 ± 14 ^a | | ≤ 50 | |
| 8.50 ± 30 | | ≤ 120 | |
| | 9.198 ± 10 | ≤ 50 | |

A: $^{16}\text{O}(^3\text{He}, n)^{18}\text{Ne}$: for references see Table 18.23 (1978AJ03) and (1981NE09).

B: $^{20}\text{Ne}(p, t)^{18}\text{Ne}$: (1981NE09).

^a See also Table 18.23 in (1978AJ03).

^b (1981NE09).

^c $\Gamma = 180 \pm 60$ keV.

References

(Closed 01 June 1987)

- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- 1973JO13 P.L. Jolivette, Phys. Rev. C8 (1973) 1230
- 1973OL02 J.W. Olness, E.K. Warburton and J.A. Becker, Phys. Rev. C7 (1973) 2239
- 1973RO03 C. Rolfs, A.M. Charlesworth and R.E. Azuma, Nucl. Phys. A199 (1973) 257
- 1973RO04 C. Rolfs, W.E. Kieser, R.E. Azuma and A.E. Litherland, Nucl. Phys. A199 (1973) 274
- 1973RO05 C. Rolfs, H.P. Trautvetter, R.E. Azuma and A.E. Litherland, Nucl. Phys. A199 (1973) 289
- 1973RO06 C. Rolfs, I. Berka and R.E. Azuma, Nucl. Phys. A199 (1973) 306
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974ES02 J.L. Escudie, R. Lombard, M. Pignanelli, F. Resmini and A. Tarrats, Phys. Rev. C10 (1974) 1645; Erratum Phys. Rev. C11 (1975) 639
- 1976CH24 L.C. Chen, Phys. Rev. C14 (1976) 2069
- 1976LA13 R.D. Lawson, F.J.D. Serduke and H.T. Fortune, Phys. Rev. C14 (1976) 1245
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- 1978FO29 H.T. Fortune, L.R. Medsker and J.N. Bishop, Nucl. Phys. A309 (1978) 221
- 1978WI1B B.H. Wildenthal, Nukleonika 23 (1978) 459
- 1979FE06 M.P. Fewell, A.M. Baxter, D.C. Kean, R.H. Spear and T.H. Zabel, Nucl. Phys. A321 (1979) 457
- 1979KI12 W.E. Kieser, R.E. Azuma, I. Berka, K.P. Jackson, A.B. McDonald, H.B. Mak, and W. McLatchie, Nucl. Phys. A327 (1979) 172
- 1979KI13 W.E. Kieser, R.E. Azuma and K.P. Jackson, Nucl. Phys. A331 (1979) 155
- 1979WO04 J.G. Woodworth, K.G. McNeill, J.W. Jury, R.A. Alvarez, B.L. Berman, D.D. Paul and P. Meyer, Phys. Rev. C19 (1979) 1667
- 1980KR1F Kraus and Linck, Proc. Int. Conf. on Nucl. Phys., Berkeley (1980) 555
- 1980MA26 H.-B. Mak, G.T. Ewan, H.C. Evans, J.D. MacArthur, W. McLatchie and R.E. Azuma, Nucl. Phys. A343 (1980) 79
- 1980NA14 F. Naulin, C. Detraz, M. Bernas, D. Guillemaud, E. Kashy, M. Langevin, F. Pougheon, P. Roussel and M. Roy-Stephan, J. Phys. Lett. 41 (1980) L79

- 1981AD01 E.G. Adelberger, C.D. Hoyle, H.E. Swanson and R.D. von Lintig, Phys. Rev. Lett. 46 (1981) 695
- 1981CL05 C.F. Clement and S.M. Perez, J. Phys. (London) G7 (1981) L293
- 1981CO13 M.E. Cobern, L.C. Bland, H.T. Fortune, G.E. Moore, S. Mordechai and R. Middleton, Phys. Rev. C23 (1981) 2387
- 1981CU07 A. Cunsolo, A. Foti, G. Imme, G. Pappalardo, G. Raciti and N. Saunier, Phys. Rev. C24 (1981) 476
- 1981GL02 C.W. Glover and K.W. Kemper, Nucl. Phys. A366 (1981) 469
- 1981HA06 W.C. Haxton, Phys. Rev. Lett. 46 (1981) 698
- 1981IC02 A. Ichimura and M. Ichimura, Prog. Theor. Phys. 66 (1981) 913
- 1981LE1A F. Leccia, P. Mennrath, A. Morales, J. Morales, R. Nunez-Lagos and M. Plo, An. Fis. A77 (1981) 114
- 1981ME13 M.C. Mermaz, J. Barrette and H.E. Wegner, Phys. Rev. C24 (1981) 2148
- 1981NAZQ F. Naulin, C. Detraz, M. Roy-Stephan, M. Bernas, J. de Boer, D. Guillemaud, M. Langevin, F. Pougheon, P. Roussel, 4th Int. Conf. on Nucl. Far from Stability, Helsingon, Denmark (1981) 376; CERN Rept. 81-09 (1981)
- 1981NE09 A.V. Nero, E.G. Adelberger and F.S. Dietrich, Phys. Rev. C24 (1981) 1864
- 1981SEZR K.K. Seth, 4th Int. Conf. on Nucl. Far from Stability, Helsingon, Denmark (1981) 655; CERN Rept. 81-09 (1981)
- 1981WA1Q R.K. Wallace and S.E. Woosley, Astrophys. J. Suppl. 45 (1981) 389
- 1982AB04 M.S. Abdel-Wahab, L. Potvin, R. Roy, P. Bricault, R. Larue, D. Pouliot, C. Rioux and R.J. Slobodrian, Can. J. Phys. 60 (1982) 1595
- 1982AH07 G. Ahrens, W. Harfst, J.R. Kass, E.V. Mason, H. Schober, G. Steffens, H. Waeffler, P. Bock and K. Grotz, Nucl. Phys. A390 (1982) 486
- 1982AL30 Y. Alhassid, M. Gai and G.F. Bertsch, Phys. Rev. Lett. 49 (1982) 1482
- 1982AN12 M.S. ANTONY, J. Phys. G8 (1982) 1659
- 1982AO06 K. Aoki and H. Horiuchi, Prog. Theor. Phys. 68 (1982) 2028
- 1982AS1B E. Aslanides, T. Bressani, M. Caria, E. Chiavassa, S. Costa, G. Dellacasa, P. Fassnacht, M. Gallio, F. Hibou, B. Minetti et al., Proc. Int. Conf. on Nucleus-Nucleus Collisions, 26 Sept.-1 Oct. 1982, East Lansing, MI, USA (1982) 2
- 1982AV1A I.K. Averyanov, A.I. Golubev, A.A. Sadovoi, Yad. Fiz. 35 (1982) 833; Sov. J. Nucl. Phys. 35 (1982) 484
- 1982BA03 K. Bangert, U.E.P. Berg, G. Junghans, R. Stock and K. Wienhard, Nucl. Phys. A376 (1982) 15
- 1982BA06 G.C. Ball, T.K. Alexander, W.G. Davies, J.S. Forster and I.V. Mitchell, Nucl. Phys. A377 (1982) 268

- 1982BA1D C.A. Barnes, *Essays in Nucl. Astrophys.* (1982) 193
- 1982BA40 G.C. Ball, T.K. Alexander, W.G. Davies, J.S. Forster, I.V. Mitchell, J. Keinonen and H. B. Mak, *Nucl. Phys. A386* (1982) 333
- 1982BA49 Z. Basrak, R. Caplar, C. Beck, R.M. Freeman and F. Haas, *Phys. Rev. C26* (1982) 1774
- 1982BA62 F.C. Barker, *Aust. J. Phys.* 35 (1982) 377
- 1982BE1Z J.J. Bevelacqua, *Rev. Mex. Fis.* 28 (1982) 573
- 1982BE29 H.W. Becker, W.E. Kieser, C. Rolfs, H.P. Trautvetter and M. Wiescher, *Z. Phys. A305* (1982) 319
- 1982BI08 B.L. Birbrair and Yu.A. Kalashnikov, *J. Phys. (London) G8* (1982) 1531
- 1982BR24 B.A. Brown, B.H. Wildenthal, W. Chung, S.E. Massen, M. Bernas, A.M. Bernstein, R. Miskimen, V.R. Brown and V. A.Madsen, *Phys. Rev. C26* (1982) 2247
- 1982BU1A Burbidge and Burbidge, *Essays in Nucl. Astrophys.* (1982) 11
- 1982CA12 E. Caurier and A. Poves, *Nucl. Phys. A385* (1982) 407
- 1982CA1A Cameron, *Essays in Nucl. Astrophys.* (1982) 23
- 1982CI1C N. Cindro, *Nukleonika* 27 (1982) 137
- 1982CU01 A. Cunsolo, A. Foti, G. Imme, G. Pappalardo, G. Raciti and N. Saunier, *Phys. Lett. B112* (1982) 121
- 1982DE30 L.C. Dennis, K.M. Abdo, A.D. Frawley and K.W. Kemper, *Phys. Rev. C26* (1982) 981
- 1982FI10 L.K. Fifield, J.L. Durell, M.A.C. Hotchkis, J.R. Leigh, T.R. Ophel and D.C. Weisser, *Nucl. Phys. A385* (1982) 505
- 1982FR15 R.M. Freeman, P.A. DeYoung, L.J. Satkowiak, M.A. Xapsos and J.J. Kolata, *Nucl. Phys. A385* (1982) 516
- 1982GI1C R. Giordano, F. Porto, S. Sambataro and A. Scalia, in *MSU* (1982) 101
- 1982GL08 C. Glashausser, R. de Swiniarski, K. Jones, S. Nanda, F.T. Baker, M. Grimm, V. Penumetcha, A. Scott, G. Adams, G. Igo et al., *Phys. Lett. B116* (1982) 215
- 1982HE04 A.M. Hernandez and W.W. Daehnick, *Phys. Rev. C25* (1982) 2957
- 1982HE07 B. Heusch, C. Beck, J.P. Coffin, P. Engelstein, R.M. Freeman, G. Guillaume, F. Haas and P. Wagner, *Phys. Rev. C26* (1982) 542
- 1982HI1H R.R. Highfill, *Bull. Amer. Phys. Soc.* 27 (1982) 777
- 1982HU06 M. Hugi, J. Lang, R. Muller, J. Sromicki, E. Ungricht, K. Bodek, L. Jarczyk, B. Kamys, A. Strzalkowski and H. Witala, *Phys. Rev. C25* (1982) 2403
- 1982KA1C K. Kakihara, Y. Kasida and T. Ido, *Radioisotopes* 31 (1982) 357
- 1982KO24 J.J. Kolata, E.G. Funk and J.D. Hinnefeld, *Phys. Rev. C26* (1982) 1750

- 1982KU04 P.D. Kunz, J.S. Vaagen, J.M. Bang and B.S. Nilsson, Phys. Lett. B112 (1982) 5
- 1982LI15 B.J. Lieb, H.S. Plendl, C.E. Stronach, H.O. Funsten and V.G. Lind, Phys. Rev. C26 (1982) 2084
- 1982LI1M B. Liu and H.-C. Chiang, Phys. Energ. Fortis Phys. Nucl. 6 (1982) 595
- 1982LI1N B. Liu and H.-C. Chiang, Phys. Energ. Fortis Phys. Nucl. 6 (1982) 743
- 1982LO13 M. Lozano, J.I. Escudero and G. Madurga, J. Phys. (London) G8 (1982) 1259
- 1982MA1Q B. Maurel, G. Amsel and J.P. Nadai, Nucl. Instrum. Meth. Phys. Res. 197 (1982) 1
- 1982MO09 R. Moro, A. Brondi, A. D'Onofrio, V. Roca, M. Romano, F. Terrasi and B. Delaunay, Lett. Nuovo Cim. 33 (1982) 407
- 1982MO1K C.B.O. Mohr, Aust. J. Phys. 35 (1982) 1
- 1982NA13 S. Nanda and C. Glashausser, Phys. Rev. C26 (1982) 758
- 1982NO04 B.E. Norum, M.V. Hynes, H. Miska, W. Bertozzi, J. Kelly, S. Kowalski, F.N. Rad, C.P. Sargent, T. Sasanuma, W. Turchinets et al., Phys. Rev. C25 (1982) 1778
- 1982OH05 S. Ohkubo, Phys. Rev. C25 (1982) 2498
- 1982OL01 J.W. Olness, E.K. Warburton, D.E. Alburger, C.J. Lister and D.J. Millener, Nucl. Phys. A373 (1982) 13
- 1982PI06 E. Piassetzky, A. Altman, J. Lichtenstadt, A.I. Yavin, D. Ashery, W. Bertl, L. Felawka, H.K. Walter, F.W. Schleputz, R.J. Powers et al., Phys. Rev. C26 (1982) 2702
- 1982PI1H G. Pinto, Bull. Amer. Phys. Soc. 27 (1982) 792
- 1982RE14 K.E. Rehm, W. Henning, J.R. Erskine and D.G. Kovar, Phys. Rev. C26 (1982) 1010
- 1982RI04 L. Ricken, D. Bohle, G. Domogala, K. Glasner and E. Kuhlmann, Z. Phys. A306 (1982) 67
- 1982RO1A Rodney and Rolfs, Essay in Nucl. Astrophys. (1982) 171
- 1982SA1Q G.R. Satchler, Lecture Notes in Phys. 168 (1982) 25
- 1982VE02 J.J.M. Verbaarschot, G.A. Timmer and P.J. Brussaard, Nucl. Phys. A378 (1982) 280
- 1982VE13 J. Vernotte, G. Berrier-Ronsin, J. Kalifa and R. Tamisier, Nucl. Phys. A390 (1982) 285
- 1982WI1B M. Wiescher and K.-U. Kettner, Astrophys. J. 263 (1982) 891
- 1982WO1A Woosley and Weaver, Essays in Nucl. Astrophys. (1982) 377
- 1982YA1C T. Yamada, Bull. Amer. Phys. Soc. 27 (1982) 813
- 1982ZH01 M.V. Zhukov, J.M.G. Gomez and J. Bang, Phys. Scr. 25 (1982) 522
- 1983AD03 E.G. Adelberger, M.M. Hindi, C.D. Doyle, H.E. Swanson, R.D. von Lintig and W.C. Haxton, Phys. Rev. C27 (1983) 2833

- 1983AD1B E.G. Adelberger, Proc. NATO Advanced Study Institute on Symmetries in Nuclear Structure, 16-28 Aug. 1982, Dronnten, Netherlands; Eds., Abrahams, Allaart and Dieperink (1983) 55
- 1983AD1C E.G. Adelberger, Comments Nucl. Part. Phys. 11 (1983) 189
- 1983AD1D Adelberger, Proc. Int. Conf. on Nucl. Phys., August-Sept. 1983; Eds., P. Blasi and R.A. Ricci (1983) 499
- 1983AG1C Agalarov, Kerimov and Safin, in Moscow (1983) 494
- 1983AJ01 F. Ajzenberg-Selove, Nucl. Phys. A392 (1983) 1; Erratum Nucl. Phys. A413 (1984) 168
- 1983AM1A G. Amsel and J.A. Davies, Nucl. Instrum. Meth. Phys. Res. 218 (1983) 177
- 1983AM1C P. Amaudrez, Th.S. Bauer, J. Domingo, C.H.Q. Ingram, G. Kyle, D. Renker, G. Smith and R. Stamminger, SIN Newsl. 15 (1983) 48
- 1983AM1D G. Amsel and B. Maurel, Nucl. Instrum. Meth. Phys. Res. 218 (1983) 183
- 1983AN05 B.D. Anderson, A. Fazely, R.J. McCarthy, P.C. Tandy, J.W. Watson, R. Madey, W. Bertozzi, T.N. Buti, J.M. Finn, J. Kelly et al., Phys. Rev. C27 (1983) 1387
- 1983ANZQ Y. Ando, M. Uno and M. Yamada, JAERI-M-83-025 (1983)
- 1983AO03 K. Aoki and H. Horiuchi, Prog. Theor. Phys. (Kyoto) 69 (1983) 857
- 1983AR11 K.P. Artemov, V.Z. Gol'dberg, M.S. Golovkov, B.G. Novatskii, I.P. Petrov, V.P. Rudakov, Serikov and V.A. Timofeev, Yad. Fiz. 37 (1983) 1351; Sov. J. Nucl. Phys. 37 (1983) 805
- 1983AR1J Arima et al., in Florence (1983) 207
- 1983AS01 D. Ashery, D.F. Geesaman, R.J. Holt, H.E. Jackson, J.R. Specht, K.E. Stephenson, R.E. Segel, P. Zupranski, H.W. Baer, J.D. Bowman et al., Phys. Rev. Lett. 50 (1983) 482
- 1983BE1W A.M. Bernstein, V.R. Brown and V.A. Madsen, Comments Nucl. Part. Phys. 11 (1983) 1 or 203
- 1983BE36 D. Bender, A. Richter, E. Spamer, E.J. Ansaldo, C. Rangacharyulu and W. Knupfer, Nucl. Phys. A406 (1983) 504
- 1983BEYY R.B. Begzhanov, G.S. Valiev, I. Borbely, I.R. Gulamov, T. Iskhakov, A.M. Mukhamedzhanov, G.K. Ni, E.A. Romanovsky, A.V. Tsupin and R. Yarmukhamedov, in Moscow (1983) 282
- 1983BI13 J.R. Birkelund and J.R. Juizenga, Ann. Rev. Nucl. Part. Sci. 33 (1983) 265
- 1983BI1C P.G. Bizzeti, Riv. Nuovo Cim. 6 (1983) 1
- 1983BR1E Bromley, Proc. Int. Conf. on Nucl. Phys., Aug.-Sept. 1983, Florence; Eds., P. Blasi and R.A. Ricci; Vol. 2 (1983) 3
- 1983BR29 B.A. Brown and B.H. Wildenthal, Phys. Rev. C28 (1983) 2397

- 1983CA03 F.P. Calaprice and D.J. Millener, Phys. Rev. C27 (1983) 1175
- 1983CA1N Carlin et al., in Florence (1983) 590
- 1983CA21 W.N. Catford, E.F. Garman, D.M. Pringle and L.K. Fifield, Nucl. Phys. A407 (1983) 255
- 1983CE1A Cereda, de Leo, Micheletti and Pignanelli, in Florence (1983) 234
- 1983CI08 O. Civitarese, B.V. Carlson, M.S. Hussein and A. Szanto de Toledo, Phys. Lett. B125 (1983) 22
- 1983CL1B Clement et al., in Florence (1983) 126
- 1983CO1K J.J. Cowan, A.G.W. Cameron and J.W. Truran, Astrophys. J. 265 (1983) 429
- 1983CU03 A. Cunsolo, A. Foti, G. Imme, G. Pappalardo, G. Raciti and N. Saunier, Lett. Nuovo Cim. 37 (1983)193
- 1983DA31 J.A. Davies, T.E. Jackman, H. Plattner and I. Bubb, Nucl. Instrum. Meth. Phys. Res. 218 (1983) 141
- 1983DE26 P.A. De Young, J.J. Kolata, L.J. Satkowiak and M.A. Xapsos, Phys. Rev. C28 (1983) 692
- 1983DU13 G.G. Dussel, A.O. Gattone and E.E. Maqueda, Phys. Rev. Lett. 51 (1983) 2366
- 1983EN02 F. Entezami, J.D. Brown, K.S. Dhuga, O. Karban, J.M. Nelson and S. Roman, Nucl. Phys. A405 (1983) 69
- 1983EN04 Y.M. Engel and R.D. Levine, Phys. Rev. C28 (1983) 2321
- 1983ET02 M.C. Etchegoyen, D. Sinclair, A. Etchegoyen and E. Belmont Moreno, Nucl. Phys. A402 (1983) 87
- 1983FR17 R.M. Freeman, C. Beck, F. Haas, B. Heusch and J.J. Kolata, Phys. Rev. C28 (1983) 437
- 1983FR1A W.A. Friedman and W.G. Lynch, Phys. Rev. C28 (1983) 950
- 1983GA02 M. Gai, M. Ruscev, A.C. Hayes, J.F. Ennis, R. Keddy, E.C. Schloemer, S.M. Sterbenz and D.A. Bromley, Phys. Rev. Lett. 50 (1983) 239
- 1983GE12 P.M. Gensini, Lett. Nuovo Cim. 38 (1983) 469
- 1983GM1A M. Gmitro, H.R. Kissener, P. Truol and R.A. Eramzhyan, Fiz. Elem. Chastits At. Yadra 14 (1983) 773; Sov. J. Part. Nucl. 14 (1983) 323
- 1983GR10 E.E. Gross, J.R. Beene, K.A. Erb, M.P. Fewell, D. Shapira, M.J. Rhoades-Brown, G.R. Satchler and C.E. Thorn, Nucl. Phys. A401 (1983) 362
- 1983GR11 A. Green, W. Stepien-Rudzka and S. Wycech, Nucl. Phys. A399 (1983) 307
- 1983GR1L Grootes, Nature 303 (1983) 753
- 1983GR28 E.E. Gross and M.P. Fewell, Nucl. Phys. A411 (1983) 329

- 1983HO02 P. Hoodbhoy, R.A. Freedman, G.A. Miller and E.M. Henley, Phys. Rev. C27 (1983) 277
- 1983IS1F Ishkhanov and Kapitonov, in Florence (1983) 203
- 1983JA05 U. Jahnke, G. Ingold, H. Homeyer, M. Burgel, Ch. Egelhaaf, H. Fuchs and D. Hilscher, Phys. Rev. Lett. 50 (1983) 1246
- 1983JA09 L. Jarczyk, B. Kamys, Z. Rudy, A. Strzalkowski, H. Witala, M. Hugi, J. Lang, R. Muller, J. Sromicki and H.H. Wolter. Phys. Rev. C28 (1983) 700
- 1983JO06 M.B. Johnson and E.R. Siciliano, Phys. Rev. C27 (1983) 1647
- 1983JO1C M.B. Johnson and E.R. Siciliano, Phys. Rev. C27 (1983) 730
- 1983KA28 I.M. Kapitonov, Yad. Fiz. 38 (1983) 612; Sov. J. Nucl. Phys. 38 (1983) 364
- 1983KI13 M.M. King Yen, S.T. Hsieh and D.S. Chuu, J. Phys. G9 (1983) 1347
- 1983KI1D J.S. Killingley, Nature 301 (1983) 594
- 1983KU1C Kutschera, Radiocarbon 25 (1983) 677
- 1983LE03 P.M. Lewis, A.K. Basak, J.D. Brown, P.V. Drumm, O. Karban, E.C. Pollacco and S. Roman, Nucl. Phys. A395 (1983) 204
- 1983LE08 F. Leccia, Ph. Hubert, P. Mennrath, A. Morales, J. Morales, R. Nunez-Lagos and M. Plo, Nuovo Cim. A74 (1983) 28
- 1983LI08 L.C. Liu, Phys. Rev. C27 (1983) 1611
- 1983LI10 J.S. Lilley, B.R. Fulton, D. Banes, T.M. Cormier, I.J. Thompson, S. Landowne and H.H. Wolter, Phys. Lett. B128 (1983) 153
- 1983LI1T C.-D. Lien, L. Wielunski and M.-A. Nicolet, Nucl. Instrum. Meth. Phys. Res. 213 (1983) 463
- 1983LI1U X.-H. Liu and Y.-G. Li, Phys. Energ. Fortis Phys. Nucl. 7 (1983) 197
- 1983MA16 C.J. Martoff, J.A. Bistirlich, C.W. Clawson, K.M. Crowe, M. Koike, J.P. Miller, S.S. Rosenblum, W.A. Zajc, H.W. Baer, A.H. Wapstra et al., Phys. Rev. C27 (1983) 1621
- 1983MA1V Mahalanabis and Sinha Rey, in Florence (1983) 406
- 1983MA29 N. Marquardt, Phys. Rev. C28 (1983) 202
- 1983MA35 F.J. Margetan and J.P. Vary, Phys. Rev. C28 (1983) 907
- 1983ME12 A.C. Merchant, J. Phys. G9 (1983) 1169
- 1983ME1H Metsch, Klingenberg and Huber, in Florence (1983) 378
- 1983MO16 R. Moro, A. Brondi, A. D'Onofrio, V. Roca, M. Romano, F. Terrasi, B. Delaunay and H. Dumont, Lett. Nuovo Cim. 38 (1983) 7
- 1983OL1A D.L. Olson, B.L. Berman, D.E. Greiner, H.H. Heckman, P.J. Lindstrom and H.J. Crawford, Phys. Rev. C28 (1983) 1602

- 1983OS07 A. Osman and S.S. Abdel-Aziz, *Acta Phys. Hung.* 54 (1983) 9
- 1983OS1E E. Oset, D. Strottman, M.J. Vicente-Vacas and W.-H. Ma, *Nucl. Phys.* A408 (1983) 461
- 1983PU01 G.D. Putt, L.K. Fifield, M.A.C. Hotchkis, T.R. Ophel and D.C. Weisser, *Nucl. Phys.* A399 (1983) 190
- 1983PUZZ B. Pugh, W. Bertozzi, T.N. Buti, J.M. Finn, C. Hyde, J.J. Kelly, M.A. Kovash, B. Murdock, B.D. Anderson, A.R. Baldwin, A. Fazely, C. Lebo, R. Madey, J.W. Watson and C.C. Foster, *Bull. Amer. Phys. Soc.* 28 (1983) 690, DG6
- 1983RO07 R.M. Rockmore, *Phys. Rev.* C27 (1983) 2150
- 1983RU09 K. Rusek, Z. Moroz, R. Caplar, P. Egelhof, K.-H. Mobius, E. Steffens, I. Koenig, A. Weller and D. Fick, *Nucl. Phys.* A407 (1983) 208
- 1983SA06 M. Sato, M. Sasagase, Y. Nagashima, J. Schimizu, T. Nakagawa, Y. Fukuchi and T. Mikumo, *Phys. Rev.* C27 (1983) 2621
- 1983SE16 U. Sennhauser, E. Piasetzky, H.W. Baer, J.D. Bowman, M.D. Cooper, H.S. Matis, H.J. Ziock, J. Alster, A. Erell, M.A. Moinester et al., *Phys. Rev. Lett.* 51 (1983) 1324
- 1983SH04 V. Shkolnik and Y.C. Tang, *Nucl. Phys.* A397 (1983) 132
- 1983SH32 J. Shurpin, T.T.S. Kuo and D. Strottman, *Nucl. Phys.* A408 (1983) 310
- 1983SH44 R.K. Sheline, *Aust. J. Phys.* 36 (1983) 825
- 1983SN03 K.A. Snover, E.G. Adelberger, P.G. Ikossi and B.A. Brown, *Phys. Rev.* C27 (1983) 1837
- 1983TA1P D.S. Tannhauser, J.A. Kilner and B.C.H. Steele, *Nucl. Instrum. Meth. Phys. Res.* 218 (1983) 504
- 1983TO21 F.M. Toyama, *Prog. Theor. Phys.* 70 (1983) 1299
- 1983UE01 K. Ueta, H. Miyake and A. Mizukami, *Phys. Rev.* C27 (1983) 389
- 1983VA23 J. van der Plicht, H.C. Britt, M.M. Fowler, Z. Fraenkel, A. Gavron, J.B. Wilhelmy, F. Plasil, T.C. Awes and G.R. Young, *Phys. Rev.* C28 (1983) 2022
- 1983VO1A G. Vourvopoulos, *IEEE Trans. Nucl. Sci.* 30 (1983) 1119
- 1983VO1B G. Vourvopoulos et al., in *Florence* (1983) 639
- 1983WA05 C.E. Waltham, S.H. Chew, J. Lowe, J.M. Nelson and A.R. Barnett, *Nucl. Phys.* A395 (1983) 119
- 1983WA17 M. Waroquier, G. Wenes and K. Heyde, *Nucl. Phys.* A404 (1983) 298
- 1983WI12 D. Wintgen, H. Friedrich and K. Langanke, *Nucl. Phys.* A408 (1983) 239
- 1983WI1A Wilczynski, *Proc. Int. Conf. on Nucl. Phys., Aug.-Sept. 1983, Florence*; Eds., P. Blasi and R.A. Ricci; Vol. 2 (1983) 305
- 1983ZE1C B. Zeidman and D.F. Geesaman, *Nucl. Phys.* A396 (1983) 419

- 1984AB1A Abe and Park, RIFP-508 (1984)
- 1984AD1E E.G. Adelberger, AIP Conf. Proc. 123 (1984) 300
- 1984AL20 A. Altman, D. Ashery, E. Piasetzky, J. Lichtenstadt, A.I. Yavin, W. Bertl, L. Felawka, H.K. Walter, R.J. Powers, R.G. Winter et al., Phys. Lett. B144 (1984) 337
- 1984ANZW G.S. Anagnostatos, H.S. Kosmas, S.E. Massen and C.N. Panos, Bull. Amer. Phys. Soc. 29 (1984) 1029, AC10
- 1984AS04 H.J. Assenbaum, K. Langanke and A. Weiguny, Z. Phys. A318 (1984) 35
- 1984AS05 D. Ashery, D.F. Geesaman, R.J. Holt, H.E. Jackson, J.R. Specht, K.E. Stephenson, R.E. Segel, P. Zupranski, H.W. Baer, J.D. Bowman et al., Phys. Rev. C30 (1984) 946
- 1984AS07 Y.I. Assafiri and I. Morrison, Nucl. Phys. A427 (1984) 460
- 1984AS1D F. Asai, H. Bando and M. Sano, Phys. Lett. B145 (1984) 19
- 1984BA1B Bauer et al., 10th Int. Conf. on Particles and Nuclei, 30 July-3 Aug. 1984, Heidelberg (1984) F21
- 1984BA24 F.C. Barker, Aust J. Phys. 37 (1984) 17
- 1984BA55 D. Baye and P. Descouvemont, Phys. Lett. B146 (1984) 285
- 1984BE1A K. Bethge, Bull. Amer. Phys. Soc. 29 (1984) 1080
- 1984BH01 R.K. Bhowmik, W.D.M. Rae and B.R. Fulton, Phys. Lett. B136 (1984) 149
- 1984BI10 M. Bini, P.G. Bizzeti and P. Sona, Lett. Nuovo Cim. 41 (1984) 191
- 1984BL17 C.L. Blilie, D. Dehnhard, M.A. Franey, D.H. Gay, D.B. Holtkamp, S.J. Seestrom-Morris, P.J. Ellis, C.L. Morris and D.J. Millener, Phys. Rev. C30 (1984) 1989
- 1984BL1J J.H. Black and P.L. Smith, Astrophys. J. 277 (1984) 562
- 1984BR1L D.A. Bromley, Nuovo Cim. A81 (1984) 360
- 1984BR1P H.P. Brall, R. Stademann, K. Langanke and A. Weiguny, Fizika 16 (1984) 7
- 1984BUZX B.L. Burks, D.J. Horen, R.L. Auble, F.E. Bertrand, J.L. Blankenship, J.L.C. Ford, Jr., E.E. Gross, D.C. Hensley, R.O. Sayer, D. Shapira et al., Bull. Amer. Phys. Soc. 29 (1984) 1026, AB6
- 1984CH1V W.H. Chung and J.H. Lee, New Phys. (Korea) 24 (1984) 267
- 1984CO05 J. Cook, L.C. Dennis, K.W. Kemper, T.R. Ophel, A.F. Zeller, C.F. Maguire and Z. Kui, Nucl. Phys. A415 (1984) 114
- 1984DE38 D.M. De Castro Rizzo, E. Bozek, S. Cavallaro, B. Delaunay, J. Delaunay, H. Dumont, M.G. Saint-Laurent and F. Terrasi, Nucl. Phys. A427 (1984) 151
- 1984DE53 P. De Bievre, M. Gallet, N.E. Holden and I.L. Barnes, J. Phys. Chem. Ref. Data 13 (1984) 809
- 1984FO15 H.T. Fortune, Phys. Rev. C30 (1984) 848

- 1984FR13 H. Friedrich, Phys. Lett. B146 (1984) 135
- 1984FR14 R. Frick, H. Clement, G. Graw, P. Schiemenz, N. Seichert and T.-H Sun, Z. Phys. A319 (1984) 133
- 1984FR1A P. Frobrich, Phys. Rept. 116 (1984) 337
- 1984GA1T M. Gai, Proc. Int. Symp. on In-Beam Nucl. Spectrosc., 14-18 May 1984, Debrecen, Hungary (1984) 331
- 1984GE1A D.F. Geesaman, AIP Conf. Proc. 123 (1984) 150
- 1984GI10 R. Gilman, H.T. Fortune, L.C. Bland, R.R. Kiziah, C.F. Moore, P.A. Seidl, C.L. Morris and W.B. Cottingham, Phys. Rev. C30 (1984) 962
- 1984GI11 W.R. Gibbs and W.B. Kaufmann, Phys. Lett. B145 (1984) 1
- 1984GR08 R.E.L. Green, R.G. Korteling and K.P. Jackson, Phys. Rev. C29 (1984) 1806
- 1984GR27 S.J. Greene, C.J. Harvey, P.A. Seidl, R. Gilman, E.R. Siciliano and M.B. Johnson, Phys. Rev. C30 (1984) 2003
- 1984HA14 P. Halse, J.P. Elliott and J.A. Evans, Nucl. Phys. A417 (1984) 301
- 1984HA1R M.J. Harris and D.L. Lambert, Astrophys. J. 281 (1984) 739
- 1984HA1Z M.J. Harris and D.L. Lambert, Astrophys. J. 285 (1984) 674
- 1984HA43 Q. Haider and B. Cujec, Nucl. Phys. A429 (1984) 116
- 1984HA53 Q. Haider and F.B. Malik, At. Data Nucl. Data Tables 31 (1984) 185
- 1984HA58 J.C. Hardy and I.S. Towner, J. Phys. (Paris) 45 (1984) C4-417
- 1984HI1A A.S. Hirsch, A. Bujak, J.E. Finn, L.J. Gutay, R.W. Minich, N.T. Porile, R.P. Scharenberg and B.C. Stringfellow, Phys. Rev. C29 (1984) 508
- 1984HI1D R.D. Hichwa, Bull. Amer. Phys. Soc. 29 (1984) 1072
- 1984HO1H B.R. Holstein, AIP Conf. Proc. 123 (1984) 1110
- 1984HO23 H. Homeyer, M. Burgel, Ch. Egelhaaf, H. Fuchs and G. Thoma, Z. Phys. A319 (1984) 143
- 1984HU1C Hupke et al., in Panic (1984) I50
- 1984JO01 M.B. Johnson, E.R. Siciliano, H. Toki and A. Wirzba, Phys. Rev. Lett. 52 (1984) 593
- 1984KA26 T. Karapiperis, M. Kobayashi and M. Hirata, Phys. Lett. B144 (1984) 23
- 1984KL06 H.V. Klapdor, J. Metzinger and T. Oda, At. Data Nucl. Data Tables 31 (1984) 81
- 1984LA01 K. Langanke, R. Stademann and D. Frekers, Phys. Rev. C29 (1984) 40
- 1984LA1J W.D. Langer, T.E. Graedel, M.A. Frerking and P.B. Armentrout, Astrophys. J. 277 (1984) 581
- 1984LE1G Lecolle, C. R. Acad. Sci. Ser. II-Mec. Phys. 298 (1984) 211
- 1984MI17 H. Miyake, A. Mizukami and K. Ueta, Nuovo Cim. A84 (1984) 225

- 1984MI1H Millener, Drexel Univ. Symp., Unpublished (1984)
- 1984MO1N Motoba, Bando, Ikeda and Yamamoto, in Panic (1984) M8
- 1984MU1D D. Mukhopadhyay and G. Grawert, J. Phys. (Paris) 45 (1984) C6-435
- 1984MU27 J.A. Musser and J.D. Stevenson, Phys. Rev. Lett. 53 (1984) 847
- 1984NA1D S. Nagamiya, J. Randrup and T.J.M. Symons, Ann. Rev. Nucl. Part. Sci. 34 (1984) 155
- 1984NE1A Nemets, Rudchik and Chuvilski, in Alma Ata (1984) 334
- 1984NI1C R.J. Nickles, Bull. Amer. Phys. Soc. 29 (1984) 1072
- 1984OE02 W. Oelert, G. Palla, B. Rubio, M.G. Betigeri, C. Mayer-Boricke, P. Turek and H.T. Fortune, Phys. Rev. C30 (1984) 1378
- 1984PH02 D.L. Pham and R. de Swiniarski, Nuovo Cim. A83 (1984) 294
- 1984PO1A Poth et al., 10th Int. Conf. on Particles and Nuclei, 30 July-3 Aug. 1984, Heidelberg (1984) L1
- 1984QU1A Quader and Erikson, Bull. Amer. Phys. Soc. 29 (1984) 630
- 1984RA07 W.D.M. Rae and R.K. Bhowmik, Nucl. Phys. A420 (1984) 320
- 1984RA17 W.D.M. Rae and R.K. Bhowmik, Nucl. Phys. A427 (1984) 142
- 1984RE14 S.M. Read and V.E. Viola, Jr., At. Data Nucl. Data Tables 31 (1984) 359
- 1984RU1A Ruscev, Ph. D. Thesis, Yale Univ. (1984)
- 1984SA28 V.S. Sadkovsky, G.A. Feofilov, A.E. Denisov, R.P. Kolalis and L. Peres Tamaio, Izv. Akad. Nauk. SSSR Ser. Fiz. 48 (1984) 995; Bull. Acad. Sci. USSR Phys. Ser. 48 (1984) 158
- 1984SA37 M. Sakai, At. Data Nucl. Data Tables 31 (1984) 399
- 1984SC09 I. Schwanner, G. Backenstoss, W. Kowald, L. Tauscher, H.-J. Weyer, D. Gotta and H. Ullrich, Nucl. Phys. A412 (1984) 253
- 1984SE1A Seestrom-Morris et al., 10th Int. Conf. on Particles and Nuclei, 30 July-3 Aug. 1984, Heidelberg (1984) F1
- 1984SH2A N.J. Shackleton, J. Backman, H. Zimmerman, D.V. Kent, M.A. Hall, D.G. Roberts, D. Schnitker, J.G. Baldauf, A. Desprairies, R. Homrighausen et al., Nature 307 (1984) 620
- 1984ST02 M.R. Strayer, R.Y. Cusson, A.S. Umar, P.-G. Reinhard, D.A. Bromley and W. Greiner, Phys. Lett. B135 (1984) 261
- 1984ST1E Stone, Moszkowski, Mathews and Bloom, Bull. Amer. Phys. Soc. 29 (1984) 630
- 1984SU07 T. Suzuki and H. Narumi, Nucl. Phys. A426 (1984) 413
- 1984TA1A Tacik et al., 10th Int. Conf. on Particles and Nuclei, 30 July-3 Aug. 1984, Heidelberg (1984) F13

- 1984TR15 M.A. Troitsky, A.V. Tsybulnikov and N.I. Chekunaev, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 48 (1984) 370; *Bull. Acad. Sci. USSR Phys. Ser.* 48 (1984) 149
- 1984VA1G V.V. Varlamov, B.S. Ishkhanov, Yu.P. Pyt'ev, A.P. Chernyaev and D.V. Yudin, *Moscow Univ. Phys. Bull.* 39 (1984) 62
- 1984WE04 E. Wesolowski, *J. Phys. (London)* G10 (1984) 321
- 1984WI17 B.H. Wildenthal, *Prog. Part. Nucl. Phys.* 11 (1984) 5
- 1985AD1A E.G. Adelberger and W.C. Haxton, *Ann. Rev. Nucl. Part. Sci.* 35 (1985) 501
- 1985AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A433 (1985) 1; Erratum *Nucl. Phys.* A449 (1986) 155
- 1985AL15 A. Altman, R.R. Johnson, U. Wienands, N. Hessey, B.M. Barnett, B.M. Forster, N. Grion, D. Mills, F.M. Rozon, G.R. Smith et al., *Phys. Rev. Lett.* 55 (1985) 1273
- 1985AL21 T.K. Alexander, B. Castel and I.S. Towner, *Nucl. Phys.* A445 (1985) 189
- 1985AN17 M.S. Antony, J. Britz, J.B. Bueb and V.B. Ndocko-Ndongue, *Nuovo Cim.* A88 (1985) 265
- 1985AN1B Anne et al., in *Visby* (1985) 31
- 1985AN28 M.S. Antony, J. Britz, J.B. Bueb and A. Pape, *At. Data Nucl. Data Tables* 33 (1985) 447
- 1985AR1A W.D. Arnett and F.-K. Thielemann, *Astrophys. J.* 295 (1985) 589
- 1985BA1A J.M. Bang, F.G. Gareev, W.T. Pinkston and J.S. Vaagen, *Phys. Rept.* 125 (1985) 253
- 1985BA27 B.M. Barnett, W. Gyles, R.R. Johnson, R. Tacik, K.L. Erdman, H.W. Roser, D.R. Gill, E.W. Blackmore, S. Martin, C.A. Wiedner et al., *Phys. Lett.* B156 (1985) 172
- 1985BE1A M. Beckerman, *Phys. Rept.* 129 (1985) 145
- 1985BE40 C. Beck, F. Haas, R.M. Freeman, B. Heusch, J.P. Coffin, G. Guillaume, F. Rami and P. Wagner, *Nucl. Phys.* A442 (1985) 320
- 1985BI01 B.L. Birbrair, V.N. Fomenko, A.B. Gridnev and Yu.A. Kalashnikov, *J. Phys. (London)* G11 (1985) 471
- 1985BI03 M. Bini, T.F. Fazzini, G. Poggi and N. Taccetti, *Phys. Rev. Lett.* 55 (1985) 795
- 1985BR20 V.R. Brown, A.M. Bernstein and V.A. Madsen, *Phys. Lett.* B164 (1985) 217
- 1985BR29 B.A. Brown and B.H. Wildenthal, *At. Data Nucl. Data Tables* 33 (1985) 347
- 1985CA01 N. Carlin Filho, M.M. Coimbra, J.C. Acquadro, R. Liguori Neto, E.M. Szanto, E. Farrelly-Pessoa and A. Szanto de Toledo, *Phys. Rev.* C31 (1993) 152
- 1985CA1Q J.A. Carr, F. Petrovich and J.J. Kelly, *AIP Conf. Proc.* 124 (1985) 230
- 1985CH1R Cha and Park, *Bull. Amer. Phys. Soc.* 30 (1985) 1769
- 1985CH23 H.-C. Chiang and M.B. Johnson, *Phys. Rev.* C32 (1985) 531

- 1985CU1C J.B. Cumming, Phys. Rev. C32 (1985) 1445
- 1985DE24 P. Descouvemont and D. Baye, Phys. Rev. C31 (1985) 2274
- 1985DE38 D.M. de Castro Rizzo and N. Alamanos, Nucl. Phys. A443 (1985) 525
- 1985DE60 C. Detraz, J. Phys. Soc. Jpn. Suppl. 54 (1985) 27
- 1985DEZZ P. DeYoung, R. McGrath, J. Alexander, M. Gordon, R. Lefferts, D.M. De Castro-Rizzo, L. Vaz and X. Lu, Bull. Amer. Phys. Soc. 30 (1985) 708, AH10
- 1985DR07 B.J. Dropesky, G.W. Butler, G.C. Giesler, C.J. Orth and R.A. Williams, Phys. Rev. C32 (1985) 1305
- 1985DU05 O. Dumbrajs, Phys. Scr. 31 (1985) 485
- 1985DY05 P. Dyer, D. Bodansky, D.D. Leach, E.B. Norman and A.G. Seamster, Phys. Rev. C32 (1985) 1873
- 1985EV03 H.C. Evans, G.T. Ewan, S.-P. Kwan, J.R. Leslie, J.D. MacArthur, H.-B. Mak, W. McLatchie, S.A. Page, P. Skensved, S.-S. Wang et al., Phys. Rev. Lett. 55 (1985) 791
- 1985FO11 H.T. Fortune, L.C. Bland and W.D.M. Rae, J. Phys. (London) G11 (1985) 1175
- 1985GA1J M. Gai, J. Phys. Soc. Jpn. Suppl. 54 (1985) 470
- 1985GAZT S.B. Gazes, C.R. Albiston, Y. Chan, H.R. Schmidt, R.G. Stokstad and R. Kamermans, Bull. Amer. Phys. Soc. 30 (1985) 1280, EB8
- 1985GI01 R. Gilman, L.C. Bland, P.A. Seidl, C.F. Moore, C.L. Morris, S.J. Greene and H.T. Fortune, Nucl. Phys. A432 (1985) 610
- 1985GIZX R. Gilman, H.T. Fortune, K.S. Dhuga, J.D. Zumbro, C.L. Morris, S.J. Greene, J.A. Faucett, W.B. Cottingham, G.R. Burleson, P.A. Seidl et al., Bull. Amer. Phys. Soc. 30 (1985) 783, GH4
- 1985GM01 M. Gmitro, J. Kvasil and R. Mach, Phys. Rev. C31 (1985) 1349
- 1985GO27 C.R. Gossett, Nucl. Instrum. Meth. Phys. Res. B10-11 (1985) 722
- 1985GR1B C. Grama, N. Grama and Gh. Voiculescu, Rev. Roum. Phys. 30 (1985) 23
- 1985GR1K E.O. Grigoryan, A.G. Khudaverdyan, A.P. Bezverkhaya and N.A. Demekhina, Izv. Akad. Nauk. Arm. SSR Fiz. 20 (1985) 69; Sov. J. Contemp. Phys. 20 (1985) 12
- 1985HA11 J.S. Hanspal, R.J. Griffiths and N.M. Clarke, Phys. Rev. C31 (1985) 1138
- 1985HA18 S.S. Hanna and J.W. Hugg, Hyperfine Interactions 21 (1985) 59
- 1985HA1R M.J. Harris, D.L. Lambert and V.V. Smith, Astrophys. J. 299 (1985) 375
- 1985HA1Z M.J. Harris, D.L. Lambert and V.V. Smith, Astrophys. J. 292 (1985) 620
- 1985HO05 J.J. Hogan, J. Asher and D.J. Parker, Phys. Rev. C31 (1985) 477
- 1985HU02 I. Huszar and R. Weinreich, J. Radioanal. Nucl. Chem. Lett. 93 (1985) 349

- 1985HU04 M.S. Hussein, B.V. Carlson, O. Civitarese and A. Szanto De Toledo, Phys. Rev. Lett. 54 (1985) 2659
- 1985JOZZ P.L. Jolivet, M.J. Honkanen, M. Young and E. Moser, Bull. Amer. Phys. Soc. 30 (1985) 1248, AC7
- 1985KAZQ S.G. Kadenskii, S.D. Kurgalin and Yu.M. Chuvilskii, in Leningrad (1985) 438
- 1985KE03 E. Kerkhove, R. van de Vyver, H. Ferdinande, D. Ryckbosch, P. van Otten, P. Berkvens and E. Van Camp, Phys. Rev. C32 (1985) 368
- 1985KE1A Kelly, PP 85-202, Unpublished (1985)
- 1985KE1C J. Keinonen, AIP Conf. Proc. 125 (1985) 557
- 1985KL1A H.V. Klapdor, Fortschr. Phys. 33 (1985) 1
- 1985KO04 D. Konnerth, W. Trombik, K.G. Bernhardt, K.A. Eberhard, R. Singh, A. Strzalkowski and W. Trautmann, Nucl. Phys. A436 (1985) 538
- 1985KO1B Kozyr, in Leningrad (1985) 411
- 1985LA03 M. Langevin, E. Quiniou, M. Bernas, J. Galin, J.C. Jacmart, F. Naulin, F. Pougheon, R. Anne, C. Detraz, D. Guerreau et al., Phys. Lett. B150 (1985) 71
- 1985LE1K J.H. Lee and W.-H. Chang, New Phys. (Korean Phys. Soc.) 25 (1985) 101
- 1985LE1L J.H. Lee and W.-H. Chang, New Phys. (Korean Phys. Soc.) 25 (1985) 381
- 1985LI16 J. Lichtenstadt, A.I. Yavin, S. Janouin, P. Birien, G. Bruge, A. Chaumeaux, D. Drake, D. Garreta, D. Legrand, M.C. Lemaire et al., Phys. Rev. C32 (1985) 1096
- 1985MA1A G.J. Mathews and R.A. Ward, Rept. Prog. Phys. 48 (1985) 1371
- 1985MA1Q S. Marsh and W.D.M. Rae, Phys. Lett. B153 (1985) 21
- 1985MA1R V.A. Madsen and V.R. Brown, AIP Conf. Proc. 124 (1985) 171
- 1985MA1T Maglione, Vitturi and Pollaroio, in Visby (1985) 44
- 1985MA56 E. Maglione, G. Pollarolo, A. Vitturi, R.A. Broglia and A. Winther, Phys. Lett. B162 (1985) 59
- 1985MI10 R.E. Mischke, Nucl. Phys. A434 (1985) 505c
- 1985PO10 N.A.F.M. Poppelier, L.D. Wood and P.W.M. Glaudemans, Phys. Lett. B157 (1985) 120
- 1985PO11 D.N. Poenaru, M. Ivascu, A. Sandulescu and W. Greiner, Phys. Rev. C32 (1985) 572
- 1985PY01 R.E. Pywell, B.L. Berman, J.G. Woodworth, J.W. Jury, K.G. McNeill and M.N. Thompson, Phys. Rev. C32 (1985) 384
- 1985RO1G D.J. Rowe, Rept. Prog. Phys. 48 (1985) 1419
- 1985SA29 H. Sagawa, B.A. Brown and O. Scholten, Phys. Lett. B159 (1985) 228

- 1985SE08 P.A. Seidl, C.F. Moore, S. Mordechai, R. Gilman, K.S. Dhuga, H.T. Fortune, J.D. Zumbro, C.L. Morris, J.A. Faucett and G.R. Burleson, Phys. Lett. B154 (1985) 255
- 1985SE21 R. Seki, Prog. Theor. Phys. 74 (1985) 511
- 1985SH1P N.K. Sherman, W.F. Davidson, S. Raman, W. Delbianco and G. Kajrys, AIP Conf. Proc. 125 (1985) 221
- 1985SO12 H.Q. Song, Z.X. Wang, Y.H. Cai and W.Z. Huang, Chin. J. Nucl. Phys. 7 (1985) 320
- 1985SU10 Y. Suzuki, A. Yamamoto and K. Ikeda, Nucl. Phys. A444 (1985) 365
- 1985TAZY T.N. Taddeucci, C.D. Goodman, C.C. Foster, R.C. Byrd, C. Gaarde, J.S. Larsen, D.J. Horen, J. Rapaport and E. Sugarbaker, Bull. Amer. Phys. Soc. 30 (1985) 1255, BB4
- 1985TH03 J. Thomas, Y.T. Chen, S. Hinds, K. Langanke, D. Meredith, M. Olson and C.A. Barnes, Phys. Rev. C31 (1985) 1980
- 1985VA1C Varlamov et al., in Leningrad (1985) 339
- 1985VO12 K.F. von Reden, W.W. Daehnick, S.A. Dytman, R.D. Rosa, J.D. Brown, C.C. Foster, W.W. Jacobs and J.R. Comfort, Phys. Rev. C32 (1985) 1465
- 1985WA02 A.H. Wapstra and G. Audi, Nucl. Phys. A432 (1985) 1
- 1985WA1R M. Wahlen and T. Yoshinari, Nature 313 (1985) 780
- 1985WA24 J.W. Watson, W. Pairsuwan, B.D. Anderson, A.R. Baldwin, B.S. Flanders, R. Madey, R.J. McCarthy, B.A. Brown, B.H. Wildenthal and C.C. Foster, Phys. Rev. Lett. 55 (1985) 1369
- 1985WI17 B.H. Wildenthal, B.A. Brown and I. Sick, Phys. Rev. C32 (1985) 2185
- 1985WU03 J.Q. Wu, G. Bertsch and A.B. Balantekin, Phys. Rev. C32 (1985) 1432
- 1985YA1C T. Yamada, T. Motoba, K. Ikeda and H. Bando, Prog. Theor. Phys. Suppl. 81 (1985) 104
- 1985YO1B T. Yosshinari and M. Wahlen, Nature 317 (1985) 349
- 1985YU1B Z.-S. Yuan, Y.-P. Shen, Y.-S. Wang, Z.-H. Qiu and T.T.S. Kuo, Phys. Energ. Fortis Phys. Nucl. 9 (1985) 586
- 1985ZA1A R. Zahn, B. Markussen and J. Thiede, Nature 314 (1985) 433
- 1986AJ01 F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1
- 1986AJ04 F. Ajzenberg-Selove, Nucl. Phys. A460 (1986) 1
- 1986AL22 A. Altman, D. Ashery, E. Piasetzky, J. Lichtenstadt, A.I. Yavin, W. Bertl, L. Felawka, H.K. Walter, R.J. Powers, R.G. Winter et al., Phys. Rev. C34 (1986) 1757
- 1986ALZZ M.M. Alam and F.B. Malik, Bull. Amer. Phys. Soc. 31 (1986) 770, AG11
- 1986AN07 M.S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables 34 (1986) 279
- 1986AN10 M.S. Antony, J. Britz and J.B. Bueb, Nuovo Cim. A91 (1986) 283

- 1986AN1F B. Andersson, S. Garpman, G. Gustafson, H.-A. Gustafsson, B. Nilsson-Almqvist, I. Otterlund and E. Stenlund, *Phys. Scripta* 34 (1986) 451
- 1986ANZY T. Anderl, A. Altman, R.R. Johnson, B.M. Barnett, R. Rui, F.M. Rozon, G.R. Smith, M. Sevier, R.P. Trelle, U. Wienands et al., *Bull. Amer. Phys. Soc.* 31 (1986) 856, HI5
- 1986AV07 A.R. Avakyan, G.A. Vartapetyan, E.O. Grigoryan and N.A. Demekhina, *Yad. Fiz.* 44 (1986) 566; *Sov. J. Nucl. Phys.* 44 (1986) 187
- 1986AV1B Avdeichikov, in *Dubna* (1986) 122
- 1986BA1C H.W. Baer and G.A. Miller, *Comments Nucl. Part. Phys.* 15 (1986) 269
- 1986BA3G V.P. Bamblevskij, *Kernenergie* 29 (1986) 64
- 1986BA69 D. Baye, *Nucl. Phys.* A460 (1986) 581
- 1986BA89 G. Baur, *Phys. Lett.* B178 (1986) 135
- 1986BI1A R. Bimbot, S. Della-Negra, M. Manasijevic, P. Aguer, G. Bastin, R. Anne, H. Delagrangé, Y. Schutz, F. Hubert, Y. Gono et al., *J. Phys. (Paris)* 47 (1986) C4-241
- 1986BL08 K. Blatt, K. Becker, B. Heck, H. Jansch, H. Leucker, D. Fick, R. Caplar, R. Butsch, D. Kramer, K.-H. Mobius et al., *Phys. Rev. Lett.* 57 (1986) 819
- 1986BR04 G. Bruge, A. Chaumeaux, P. Birien, D.M. Drake, D. Garreta, S. Janouin, D. Legrand, M.C. Lemaire, B. Mayer, J. Pain et al., *Phys. Lett.* B169 (1986) 14
- 1986BR1X Brown, in *Harrogate* (1986) 119
- 1986CA1T F. Catara, M. Sambataro, A. Insolia and A. Vitturi, *Phys. Lett.* B180 (1986) 1
- 1986CA27 M. Carchidi, B.H. Wildenthal and B.A. Brown, *Phys. Rev.* C34 (1986) 2280
- 1986CO11 S.G. Cooper, *J. Phys. (London)* G12 (1986) 869
- 1986CO2B D.D. Cohen, A. Katsaros and D. Garton, *Nucl. Instrum. Meth. Phys. Res.* B15 (1986) 555
- 1986CO2C E.R. Comben, J.M. Brown, T.C. Steimle, K.R. Leopold and K.M. Evenson, *Astrophys. J.* 305 (1986) 513
- 1986CU02 B. Cujec, B. Dasmahapatra, Q. Haider, F. Lahlou and R.A. Dayras, *Nucl. Phys.* A453 (1986) 505
- 1986DAZS C.H. Dasso and A. Vitturi, in *Harrogate* (1986) 221
- 1986DE1G R. de Swiniarski and D.L. Pham, *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 932
- 1986DO1L J.F. Dominy, G. Wallerstein and N.B. Suntzeff, *Astrophys. J.* 300 (1986) 325
- 1986DR03 P.V. Drumm, O. Karban, A.K. Basak, P.M. Lewis, S. Roman and G.C. Morrison, *Nucl. Phys.* A448 (1986) 93
- 1986DR1E Druffel and Benavides, *Nature* 321 (1986) 58
- 1986DU10 O. Dumbrajs, H. Heiselberg, A.S. Jensen, A. Miranda, G.C. Oades and J.M. Richard, *Nucl. Phys.* A457 (1986) 491

- 1986DU1Q J.-C. Duplessy, M. Arnold, P. Maurice, E. Bard, J. Duprat and J. Moyes, *Nature* 320 (1986) 350
- 1986EN1A Engelmann and Bardy, Rept. CEA-R-5340 (1986)
- 1986ER1A R.A. Eramzhyan, B.S. Ishkhanov, I.M. Kapitonov and V.G. Neudatchin, *Phys. Rept.* 136 (1986) 229
- 1986FO06 H.T. Fortune and R. Gilman, *Phys. Rev. C* 33 (1986) 2171
- 1986FR10 E. Friedman and J. Lichtenstadt, *Nucl. Phys. A* 455 (1986) 573
- 1986GA13 E.N. Gazis, C.T. Papadopoulos, R. Vlastou and A.C. Xenoulis, *Phys. Rev. C* 34 (1986) 872
- 1986GE06 W.J. Gerace, W.J. Leonard and D.A. Sparrow, *Phys. Rev. C* 34 (1986) 353
- 1986GL02 K. Glasner, L. Ricken and E. Kuhlmann, *Nucl. Phys. A* 452 (1986) 150
- 1986GM01 M. Gmitro and A.I. Steshenko, *Yad. Fiz.* 43 (1986) 333; *Sov. J. Nucl. Phys.* 43 (1986) 213
- 1986GR1A C. Gregoire and B. Tamain, *Ann. Phys. (France)* 11 (1986) 323
- 1986GU05 I.S. Gulkarov and R.Kh. Vakil, *Yad. Fiz.* 43 (1986) 809; *Sov. J. Nucl. Phys.* 43 (1986) 515
- 1986GU1A Gulkarov and Pinkus, in *Kharkov* (1986) 172
- 1986GU1D D. Guerreau, *J. Phys. (Paris)* 47 (1986) C4-207
- 1986HA13 Q. Haider and F.B. Malik, *J. Phys. (London)* G12 (1986) 537
- 1986HA1B B.G. Harvey, *J. Phys. (Paris)* 47 (1986) C4-29
- 1986HA1E H.L. Harney, A. Richter and H.A. Weidenmuller, *Rev. Mod. Phys.* 58 (1986) 607
- 1986HA1Q Haxton, *Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., No. 68, Vol. 2* (1986) 415
- 1986HA2D M. Hashimoto, K. Nomoto, K. Arai and K. Kaminisi, *Astrophys. J.* 307 (1986) 687
- 1986HAZY A.C. Hayes, D.A. Bromley and D.J. Millener, *Bull. Amer. Phys. Soc.* 31 (1986) 815, EI3
- 1986HE1C R.B.C. Henry, *Publ. Astron. Soc. Soc. Pac.* 98 (1986) 1044
- 1986HI1B Hichwa, *Bull. Amer. Phys. Soc.* 31 (1986) 1335
- 1986HO1L N. Horvatincic, I. Krajcar-Bronic, J. Pezdic, D. Srdoc and B. Obelic, *Nucl. Instrum. Meth. Phys. Res. B* 17 (1986) 550
- 1986HO1Q S.E. Hoffman, M. Wilson and D.S. Stakes, *Nature* 321 (1986) 55
- 1986HO27 D.J. Horen, B.L. Burks, M.A.G. Fernandes, R.L. Auble, F.E. Bertrand, J.L. Blankenship, J.L.C. Ford, Jr., E.E. Gross, D.C. Hensley, R.O. Sayer et al., *Phys. Lett.* B181 (1986) 38

- 1986HOZF D.J. Horen, M.A.G. Fernandes, B.L. Burks and G.R. Satchler, in Harrogate (1986) 399
- 1986IC01 H. Ichihara and H. Yoshida, Nucl. Phys. A448 (1993) 546
- 1986IR01 M. Iriondo, D. Jerrestam and R.J. Liotta, Nucl. Phys. A454 (1986) 252
- 1986KA1C M. Kamimura, Y. Sakuragi, M. Yahiro and M. Tanifuji, J. Phys. Soc. Jpn. Suppl. 55 (1986) 205
- 1986KA1D Karban, in Dubna (1986) 83
- 1986KA36 Y. Kadota, K. Ogino, K. Obori, Y. Taniguchi, T. Tanabe, M. Yasue and J. Schimizu, Nucl. Phys. A458 (1986) 523
- 1986KE05 J. Kelly, W. Bertozzi, T.N. Buti, J.M. Finn, F.W. Hersman, M.V. Hynes, C. Hyde-Wright, B.E. Norum, A.D. Bacher, G.T. Emery et al., Phys. Lett. B169 (1986) 157
- 1986KO10 P.E. Koehler, H.D. Knox, D.A. Resler, R.O. Lane and G.F. Auchampaugh, Nucl. Phys. A453 (1986) 429
- 1986KO22 Th. Kohler, P. Blum, G. Buche, A.D. Hancock, H. Koch, A. Kreissl, H. Poth, U. Raich, D. Rohmann, G. Backenstoss et al., Phys. Lett. B176 (1986) 327
- 1986KR04 A. Krasznahorkay, T. Kibedi and Zs. Dombradi, Z. Phys. A323 (1986) 125
- 1986LA1C D.L. Lambert, B. Gustafsson, K. Eriksson and K.H. Hinkle, Astrophys. J. Suppl. 62 (1986) 373
- 1986LA29 J. Lang, Th. Maier, R. Muller, F. Nessi-Tedaldi, Th. Roser, M. Simonius, J. Sromicki and W. Haeberli, Phys. Rev. C34 (1986) 1545
- 1986LE13 M.-C. Lemaire, P. Birien, G. Bruge, D.M. Drake, D. Garreta, S. Janouin, D. Legrand, B. Mayer, J. Pain, J.C. Peng et al., Nucl. Phys. A456 (1986) 557
- 1986MA19 J.F. Mateja, A.D. Frawley, L.C. Dennis and K. Sartor, Phys. Rev. C33 (1986) 1649
- 1986MA1P Martin et al., Bull. Amer. Phys. Soc. 31 (1986) 1304
- 1986MA32 N. Matsuoka, H. Sakai, T. Saito, K. Hosono, M. Kondo, H. Ito, K. Hatanaka, T. Ichihara, A. Okihana, K. Imai et al., Nucl. Phys. A455 (1986) 413
- 1986MA48 D.M. Manley, B.L. Berman, W. Bertozzi, J.M. Finn, F.W. Hersman, C.E. Hyde-Wright, M.V. Hynes, J.J. Kelly, M.A. Kovash, S. Kowalski et al., Phys. Rev. C34 (1986) 1214
- 1986ME06 M.C. Mermaz, T. Suomijarvi, R. Lucas, B. Berthier, J. Matuszek, J.P. Coffin, G. Guillaume, B. Heusch, F. Jundt and F. Rami, Nucl. Phys. A456 (1986) 186
- 1986ME1C O. Meirav, A. Altman, E. Friedman, D.R. Gill and R.R. Johnson, in Harrogate (1986) A2
- 1986ME1F D.F. Measday, Czech. J. Phys. 36 (1986) 395
- 1986MO1A T. Motoba, Czech. J. Phys. B36 (1986) 435

- 1986MO1Q T. Motoba, Nucl. Phys. A450 (1986) 209c
- 1986NA1K G. Nardulli, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1000
- 1986NE1A Nemets et al., in Kharkov (1986) 372
- 1986NG1B Nguyen and Ye, in Harrogate (1986) C168
- 1986NI1C D. Nitz, D. Ciampa, T. Hemmick, D. Elmore, P.W. Kubik, S.L. Olsen, T. Gentile, H. Kagan, P. Haas and P.F. Smith, AIP Conf. Proc. 150 (1986) 1143
- 1986OS05 A. Osman and S.S. Abdel-Aziz Indian, J. Pure Appl. Phys. 24 (1986) 65
- 1986PE1E F. Petrovich, J.A. Carr and H. McManus, Ann. Rev. Nucl. Part. Sci. 36 (1986) 29
- 1986PI09 C. Pillai, L.W. Swenson, D.J. Vieira, G.W. Butler, J.M. Wouters, S.H. Rokni, K. Vaziri and L.P. Remsberg, Radiat. Eff. 96 (1986) 41
- 1986PO06 D.N. Poenaru, W. Greiner, K. Depta, M. Ivascu, D. Mazilu and A. Sandulescu, At. Data Nucl. Data Tables 34 (1986) 423
- 1986PO13 F. Pougheon, D. Guillemaud-Mueller, E. Quiniou, M.G. Saint Laurent, R. Anne, D. Bazin, M. Bernas, D. Guerreau, J.C. Jacmart, S.D. Hoath et al., Europhys. Lett. 2 (1986) 505
- 1986RO03 R. Rockmore and B. Saghai, Phys. Rev. C33 (1986) 576
- 1986RO12 G. Royer, J. Phys. (London) G12 (1986) 623
- 1986RO23 D. Rohmann, H. Barth, A.D. Hancock, H. Koch, Th. Kohler, A. Kreissl, H. Poth, U. Raich, A. Wolf, L. Tauscher et al., Z. Phys. A325 (1986) 261
- 1986SA30 H. Sato and Y. Okuhara, Phys. Rev. C34 (1986) 2171
- 1986SC28 C.J.S. Scholz, L. Ricken and E. Kuhlmann, Z. Phys. A325 (1986) 203
- 1986SC29 H.R. Schmidt, S.B. Gazes, Y. Chan, R. Kamermans and R.G. Stokstad, Phys. Lett. B180 (1986) 9
- 1986SCZX R.A. Schumacher, P. Amaudruz, Q. Ingram, U. Sennhauser, H. Breuer, N.S. Chant, A. Feldman, B.S. Flanders, F. Khazaie, D.J. Mack et al., Bull. Amer. Phys. Soc. 31 (1986) 1220, CC1
- 1986SE1B N. Seichert, W. Assmann, H. Clement, G. Graw, C. Hategan, H. Kader, F. Merz and P. Schiemenz, J. Phys. Soc. Jpn. Suppl. 55 (1986) 646
- 1986SI11 E.R. Siciliano, M.D. Cooper, M.B. Johnson and M.J. Leitch, Phys. Rev. C34 (1986) 267
- 1986SI1K Simonius, J. Phys. Soc. Jpn. Suppl. 55 (1986) 523
- 1986SM01 M.J. Smithson, D.L. Watson and H.T. Fortune, Phys. Rev. C33 (1986) 509
- 1986SO10 L.G. Sobotka, D.G. Sarantites, H. Puchta, F.A. Dilmanian, M. Jaaskelainen, M.L. Halbert, J.H. Barker, J.R. Beene, R.L. Ferguson, D.C. Hensley et al., Phys. Rev. C34 (1986) 917

- 1986ST15 A.I. Steshenko, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 50 (1986) 953; *Bull. Acad. Sci USSR Phys. Ser.* 50 (1986) 121
- 1986STZY S.M. Sterbenz, M. Gai, J.F. Shriner, Jr., P.D. Cottle, D.A. Bromley, M. Morando and R.A. Ricci, *Bull. Amer. Phys. Soc.* 31 (1986) 839, GJ1
- 1986TA1B Tanaka et al., *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 764
- 1986TH01 J. Thomas, Y.T. Chen, S. Hinds, D. Meredith and M. Olson, *Phys. Rev.* C33 (1986) 1679
- 1986TO1D I.S. Towner, *Ann. Rev. Nucl. Part. Sci.* 36 (1986) 115
- 1986VA23 C.P.M. van Engelen, E.A. Bakkum, R.J. Meijer and R. Kamermans, *Nucl. Phys.* A457 (1986) 375
- 1986VAZV G.S. Valiev, I.R. Gulamov, Yu.I. Denisov, T. Iskhakov, A.M. Mukhamedzhanov, G.K. Ni, E.A. Romanovsky, V.A. Stepanenko and R. Yarmukhamedov, in *Kharkov (1986)* 292
- 1986VAZZ S.P. Van Verst, J.A. Carr and K.W. Kemper, *Bull. Amer. Phys. Soc.* 31 (1986) 64, HD3
- 1986VI08 A. Vitturi and C.H. Dasso, *Nucl. Phys.* A458 (1986) 157
- 1986WI1L T.L. Wilson, E. Serabyn, C. Henkel and C.M. Walmsley, *Astron. Astrophys.* 158 (1986) L1
- 1986WI1P D. Wilkinson, *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 347
- 1986XU1C D.-Y. Xu, Z. Yan, Q.-W. Zhang, Z.-D. Shen, Y.-Y. Sun and L.-F. Ye, *Nature* 321 (1986) 854
- 1986YA12 T. Yamaya, J.I. Hirota, K. Takimoto, S. Shimoura, A. Sakaguchi, S. Kubono, M. Sugitani, S. Kato, T. Suehiro and M. Fukada, *Phys. Rev.* C34 (1986) 2369
- 1986YA1R T. Yamaya et al., *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 730, 732
- 1986YU1B Z.-S. Yuan, Y.-P. Sheng, Y.-S. Wang, Z.-H. Qui and T.T.S. Kuo, *Chin. Phys.* 6 (1986) 611
- 1987AB03 H. Abele, H.J. Hauser, A. Korber, W. Leitner, R. Neu, H. Plappert, T. Rohwer, G. Staudt, M. Strasser, S. Welte et al., *Z. Phys.* A326 (1987) 373
- 1987AS01 H.J. Assenbaum, K. Langanke and A. Weiguny, *Phys. Rev.* C35 (1987) 755
- 1987BA18 C.J. Batty, *Phys. Lett. B*189 (1987) 393
- 1987BO16 N. Bordes, G. Blondiaux, C.J. Maggiore, M. Valladon, J.L. Debrun, R. Coquille and M. Gauneau, *Nucl. Instrum. Meth. Phys. Res.* B24-25 (1987) 722
- 1987CH14 S. Chakravarti, D. Dehnhard, M.A. Franey, S.J. Seestrom-Morris, D.B. Holtkamp, C.L. Blilie, A.C. Hayes, C.L. Morris and D.J. Millener, *Phys. Rev.* C35 (1987) 2197
- 1987CO07 J. Cook, *Nucl. Phys.* A465 (1987) 207

- 1987DE43 H. De Vries, C.W. De Jager and C. De Vries, *At. Data Nucl. Data Tables* 36 (1987) 495
- 1987DJ01 C. Djalali, G.M. Crawley, B.A. Brown, V. Rotberg, G. Caskey, A. Galonsky, N. Marty, M. Morlet and A. Willis, *Phys. Rev. C* 35 (1987) 1201
- 1987ER05 T. Erikson, K.F. Quader, G.E. Brown and H.T. Fortune, *Nucl. Phys. A* 465 (1987) 123
- 1987GA15 M. Gai, R. Keddy, D.A. Bromley, J.W. Olness and E.K. Warburton, *Phys. Rev. C* 36 (1987) 1256
- 1987GOZX J. Gorres, M. Wiescher, U. Giesen, H.W. Becker, C. Rolfs and H.P. Trautvetter, *Bull. Amer. Phys. Soc.* 32 (1987) 1062, EG9
- 1987HA1C Harris, Lambert and Goldman, *Mon. Notic. Roy. Astron. Soc.* 224 (1987) 237
- 1987HA1D Harris et al., *Astrophys. J.* 316 (1987) 294
- 1987HI1B R.D. Hichwa, E.A. Hugel, J.J. Moskwa and R.R. Raylman, *Nucl. Instrum. Meth. Phys. Res. B* 24-25 (1987) 932
- 1987JU07 J.W. Jury, P.C.-K. Kuo, K.G. McNeill, C.K. Ross, H.R. Weller and S. Raman, *Phys. Rev. C* 36 (1987) 1243
- 1987KI03 M. Kirchbach, *J. Phys. (London)* G13 (1987) 135
- 1987LI04 J.S. Lilley, M.A. Nagarajan, D.W. Banes, B.R. Fulton and I.J. Thompson, *Nucl. Phys. A* 463 (1987) 710
- 1987LO01 M. Lozano and A. Vitturi, *Phys. Rev. C* 35 (1987) 367
- 1987MA09 Q. Ma and N. Austern, *Nucl. Phys. A* 463 (1987) 620
- 1987MI1E Millener, Private Communication (1987)
- 1987MO07 C.L. Morris, S.J. Seestrom-Morris, D. Dehnhard, C.L. Blilie, R. Gilman, G.P. Gilfoyle, J.D. Zumbro, M.G. Burlein, S. Mordechai, H.T. Fortune et al., *Phys. Rev. C* 35 (1987) 1388
- 1987MU1J Mueller, GANIL (1987) 87-06; XXV Int. Winter Meeting on Nucl. Phys., Bormio, Italy (1987)
- 1987NG01 Nguyen Van Sen, Ye Yanlin, J. Arvieux, G. Gaillard, B. Bonin, A. Boudard, G. Bruge, J.C. Lugol, T. Hasegawa, F. Soga et al., *Nucl. Phys. A* 464 (1987) 717
- 1987PA07 S.A. Page, H.C. Evans, G.T. Ewan, S.-P. Kwan, J.R. Leslie, J.D. MacArthur, W. McLatchie, P. Skensved, S.-S. Wang, H.-B. Mak et al., *Phys. Rev. C* 35 (1987) 1119
- 1987PA12 S.J. Padalino, K. Sartor, L.C. Dennis and K.W. Kemper, *Phys. Rev. C* 35 (1987) 1692
- 1987PR1A N. Prantzos, M. Arnould and J.-P. Arcoragi, *Astrophys. J.* 315 (1987) 209
- 1987RA01 S. Raman, C.H. Malarkey, W.T. Milner, C.W. Nestor, Jr. and P.H. Stelson, *At. Data Nucl. Data Tables* 36 (1987) 1
- 1987RI03 J. Richert and P. Wagner, *Nucl. Phys. A* 466 (1987) 132

- 1987RO10 G. Royer, Y. Raffray, A. Oubahadou and B. Remaud, Nucl. Phys. A466 (1987) 139
- 1987RO25 C. Rolfs, H.P. Trautvetter and W.S. Rodney, Rept. Prog. Phys. 50 (1987) 233
- 1987SA15 H. Sagawa and H. Toki, J. Phys. (London) G13 (1987) 453
- 1987SP05 D.A. Sparrow, Phys. Rev. C35 (1987) 1410
- 1987SU06 T. Suzuki, D.F. Measday and J.P. Roalsvig, Phys. Rev. C35 (1987) 2212
- 1987TA07 Y. Tagishi, Y. Aoki, M. Kurokawa, T. Murayama, T. Sakai, M. Takei, M. Tomizawa and K. Yagi, Phys. Rev. C35 (1987) 1153
- 1987TE01 A. Tellez-Arenas, R.J. Lombard and J.P. Maillet, J. Phys. (London) G13 (1987) 311
- 1987WA1H M. Waroquier, J. Ryckebusch, J. Moreau, K. Heyde, N. Blasi, S.Y. van der Werf and G. Wenes, Phys. Rept. 148 (1987) 249
- 1987WI11 M. Wiescher, V. Harms, J. Gorres, F.-K. Theilemann and L.J. Rybarczyk, Astrophys. J. 316 (1987) 162
- 1987YA03 B.-J. Yang and Z. Hwang, Phys. Rev. C35 (1987) 851
- 1988UM1A K. Umezawa, T. Kurio, J. Yamane, F. Shoji, K. Oura, T. Hanawa and S. Yano, Nucl. Instrum. Meth. Phys. Res. B33 (1988) 634

