

Energy Levels of Light Nuclei $A = 20$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18-20$ was published in *Nuclear Physics A190* (1972), p. 1. This version of $A = 20$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1971)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 20$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{20}\text{C}\$](#) , [\$^{20}\text{N}\$](#) , [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [\$^{20}\text{Mg}\$](#) , [\$^{20}\text{Al}\$](#)

B. Tables of Recommended Level Energies:

[Table 20.1](#): Energy levels of ^{20}O

[Table 20.4](#): Energy levels of ^{20}F

[Table 20.15](#): Energy levels of ^{20}Ne

[Table 20.35](#): Energy levels of ^{20}Na

C. [References](#)

D. Figures: [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{20}C
(Not illustrated)

^{20}C has not been observed: see (1960ZE03).

^{20}N
(Not illustrated)

^{20}N has been observed in the bombardment of ^{232}Th by 122 MeV ^{18}O ions (1969AR13, 1970AR1D) and in the 3 GeV proton bombardment of ^{197}Au (1970RA1A): it is particle stable. See also (1960ZE03, 1961BA1C, 1971BU1E).

^{20}O
(Figs. 9 and 13)

GENERAL:

Model calculations: (1959BR1E, 1960TA1C, 1962TA1B, 1963PA03, 1964CO24, 1964MO1E, 1964PA1D, 1964TR1A, 1965DE1H, 1965FE02, 1966AR10, 1966BR04, 1966TR02, 1967FE01, 1967FL13, 1967LA1H, 1967PI1B, 1968AR02, 1968BE1U, 1968CO1N, 1968FL1C, 1968GU1E, 1968HA17, 1968HA1P, 1968MO1G, 1968PA1Q, 1969FE1A, 1969KU1G, 1969SO08, 1971AR25).

Other theoretical calculations: (1961JA1E, 1966KE16, 1967ST1N, 1968SU1C, 1969SC14, 1971LA1D, 1971LE1H).

General experimental work: (1969AR13, 1971AR02, 1971AR1P).

1. $^{20}\text{O}(\beta^-)^{20}\text{F}$ $Q_m = 3.815$

^{20}O decays to $^{20}\text{F}^*(1.06)$ [$J^\pi = 1^+$] with a half-life of 13.57 ± 0.1 sec (1970MA42), 13.6 ± 1.0 sec (1960SC01). Using $E_\beta(\text{max}) = 2.76$ MeV and $\tau_{1/2} = 13.57 \pm 0.1$ sec, $\log ft = 3.73$. Upper limits for the branching to other states of ^{20}F are shown in Table 20.2 (1970MA42). See also (1959AM13) and (1957TH14, 1970AN27, 1970MC23; theor.).

2. $^{18}\text{O}(\text{t}, \text{p})^{20}\text{O}$ $Q_m = 3.079$
 $Q_0 = 3.086 \pm 0.015$ (1960JA13);
 $Q_0 = 3.076 \pm 0.010$ (1962HI06).

Table 20.1: Energy levels of ^{20}O

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ (sec)	Decay	Reactions
0	$0^+; 2$	13.57 ± 0.1	β^-	1, 2, 3
1.672 ± 5	2^+		γ	2
3.568 ± 5	4^+		(γ)	2
4.065 ± 5	2^+		(γ)	2
4.446 ± 7	0^+		(γ)	2
4.838 ± 7			(γ)	2
4.997 ± 7			(γ)	2
5.220 ± 7			(γ)	2
5.298 ± 7			(γ)	2
5.382 ± 7			(γ)	2
5.603 ± 7			(γ)	2
(5.83 ± 20)			(γ)	2

Table 20.2: Branching in $^{20}\text{O}(\beta^-)^{20}\text{F}$ (1970MA42)

Decay to $^{20}\text{F}^*$ (MeV)	J^π	Branch (%)	$\log ft$
0.98	$1^+(2^+, 3^+)$	< 0.6	> 6.0
1.06	1^+	100.0	3.73
1.31	2^+	< 0.8	> 5.7
1.84	$2, 1^+$	< 1.9	> 4.8
1.97		< 1.4	> 4.8
2.04	2^+	< 2.0	> 4.6
2.19	2^+	< 1.0	> 4.7

Table 20.3: Energy levels of ^{20}O from $^{18}\text{O}(t, p)^{20}\text{O}$

E_x (MeV \pm keV)		L	J^π
(1960JA13)	(1962HI06)		
0	0	0 ^{a,b}	0 ⁺
1.682 \pm 20	1.672 \pm 5	2 ^{a,b}	2 ⁺
	3.568 \pm 5	4 ^a	4 ⁺
4.091 \pm 25	4.065 \pm 5	2 ^{a,b}	2 ⁺
4.449 \pm 25	4.446 \pm 7	0 ^a	0 ⁺
	4.838 \pm 7		
	4.997 \pm 7		
	5.220 \pm 7		
	5.298 \pm 7		
	5.382 \pm 7		
	5.603 \pm 7		
	(5.83 \pm 20)		

^a (1964MI05): $E_t = 10.0$ MeV.

^b (1965MO19): $E_t = 5.55$ MeV.

Observed proton groups are displayed in Table 20.3: angular distributions lead to the L and J^π values shown (1960JA13, 1962HI06, 1964MI05, 1965MO19). The $J = 2$ assignment for $^{20}\text{O}^*(1.67)$ is confirmed by the angular correlation of the 1.67 MeV γ -rays (1970NI03). See also (1959JA01, 1967CH1L, 1969BA1Z) and (1964TR1A, 1966TR02, 1967DO1B; theor.).

3. $^{18}\text{O}(^{18}\text{O}, ^{16}\text{O})^{20}\text{O}$

$$Q_m = -0.628$$

See (1970MA42).

²⁰F
(Figs. 10 and 13)

GENERAL:

Model calculations: (1959BR1E, 1963KU19, 1964MO1E, 1965DE1H, 1965DE1M, 1966CH1G, 1966PI1B, 1967BO09, 1967GU05, 1967GU1D, 1968AR02, 1968CO11, 1968GU1E, 1968HA17, 1968HA1P, 1969HO32, 1970AN27, 1970BA66, 1971AR25, 1971JO01, 1971WI01).

Other theoretical calculations: (1967ST1N, 1968CE01, 1968DW1A, 1969SC14, 1971LE1H, 1971TE06).

General experimental work: (1970FA01, 1971AR02).

Ground state: $\mu = +2.0935 \pm 0.009$ nm (1967GU14; see also (1963TS01, 1969FU11)).

See also (1963KU19, 1964LI14, 1964ST1B, 1967SH14, 1968PE16, 1971AR25).

1. $^{20}\text{F}(\beta^-)^{20}\text{Ne}$ $Q_m = 7.026$

^{20}F decays principally to $^{20}\text{Ne}^*(1.63)$ with a half-life of 11.03 ± 0.06 sec [see Table 20.5]: see ^{20}Ne .

2. $^{17}\text{O}(t, p)^{19}\text{O}$ $Q_m = 3.521$ $E_b = 14.159$

See (1962MO08).

3. $^{17}\text{O}(\alpha, p)^{20}\text{F}$ $Q_m = -5.656$

Not reported.

4. $^{18}\text{O}(d, n)^{19}\text{F}$ $Q_m = 5.768$ $E_b = 12.369$

An excitation function has been measured for $E_d = 0.95$ to 3.30 MeV (1969HE1P). See also ^{19}F .

5. $^{18}\text{O}(d, p)^{19}\text{O}$ $Q_m = 1.732$ $E_b = 12.369$

The proton yield has been measured for $E_d = 0.79$ to 0.88 MeV (1956AH1A). See also ^{19}O .

Table 20.4: Energy levels of ^{20}F ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or Γ_{cm} (keV)	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 11.03 \pm 0.06$ sec	β^-	1, 8, 11, 19, 20, 21, 23, 28, 29, 30
0.65595 ± 0.15	3^+	$\tau_m = 0.40 \pm 0.05$ psec	γ	7, 8, 10, 11, 19, 29
0.8229 ± 0.2	$4^+, 2^+$	79 ± 6 psec	γ	7, 8, 10, 11, 19, 29
0.9838 ± 0.2	$(1^+, 2^+, 3^+)$	1.6 ± 0.3 psec	γ	8, 11, 19, 29
1.05693 ± 0.16	1^+	45 ± 13 fsec	γ	8, 11, 19, 22, 29
1.30922 ± 0.16	2^+	0.9 ± 0.2 psec	γ	8, 11, 19, 23
1.8244 ± 1.3	(3^-)		γ	8, 28, 29
1.8434 ± 0.3	$2, 1^+$	30 ± 20 fsec	γ	8, 11, 19, 29
1.9706 ± 0.3			γ	8, 11
2.0439 ± 0.3	2^+	37 ± 16 fsec	γ	8, 11, 19
2.1946 ± 0.5	2^+	< 12 fsec	γ	8, 11, 19
2.865 ± 1.5	$(2, 3, 4)^-$		γ	8, 11, 19
2.9662 ± 0.4	$2^+, 3^+$	60 ± 40 fsec	γ	8, 11, 19
3.1746 ± 1.2			γ	19
3.4884 ± 0.2	1^+	44 ± 11 fsec	γ	8, 11, 19, 29
3.5259 ± 0.4	0^+	30 ± 15 fsec	γ	11, 19, 29
3.5871 ± 0.3	$(1, 2, 3)^+$	30 ± 30 fsec	γ	8, 11, 19
3.6810 ± 0.4			γ	8, 11, 19
3.761 ± 2			(γ)	8, 19, 29
3.9662 ± 1.4			γ	8, 11, 19
4.0824 ± 0.4	1^+		γ	8, 11, 19
4.1989 ± 2.7			(γ)	19, 29
4.2077 ± 2.6			(γ)	19
4.2766 ± 0.5	$1^+, 2^+$		γ	11, 19
4.3115 ± 2.6	$0^+, 1^+$		(γ)	19
4.5838 ± 3.0			(γ)	19
4.5922 ± 2.9			(γ)	19
4.7302 ± 2.9			(γ)	19
4.7638 ± 2.7			(γ)	19
4.8916 ± 2.8			(γ)	19
4.8982 ± 2.8			(γ)	19
5.0402 ± 3.1	$(0, 1, 2)^-$		(γ)	11, 19

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or Γ_{cm} (keV)	Decay	Reactions
5.0655 \pm 3.1			(γ)	19
5.2240 \pm 3.1	(0, 1, 2) ⁻		(γ)	19
5.2810 \pm 3.3	(0, 1, 2) ⁻		(γ)	19
5.3171 \pm 2.7			(γ)	19
5.3445 \pm 3.3			(γ)	19
5.4131 \pm 0.6			γ	11
5.4503 \pm 3.8			(γ)	19
5.4554 \pm 3.2			(γ)	19
5.4634 \pm 3.3			(γ)	19
5.5547 \pm 0.6			γ	11
5.6203 \pm 3.3			(γ)	19
(5.713 \pm 2)			γ	11
5.7628 \pm 3.4			(γ)	19
5.8091 \pm 2.9	(0, 1, 2) ⁻		(γ)	19
5.9361 \pm 0.3	1 ⁻ , 2 ⁻		γ	11, 19
6.0173 \pm 0.3	1 ⁻ , 2 ⁻		γ	11, 19
6.0446 \pm 0.4			γ	11, 19
6.25 \pm 20			(γ)	9
6.513 \pm 33	0 ⁺ ; 2		(γ)	19, 28
6.6013 \pm 0.3	0 ⁺ , 1 ⁺		γ	11
6.616			n	12
6.627	2 ⁻	$\Gamma = 0.29$ keV	γ , n	11, 12, 19
6.632			n	12
6.634			n	12
6.637			n	12
6.648	1 ⁻	1.62	γ , n	11, 12
6.668			n	12
6.685	0 ⁻	3.80	n	12
6.692	1 ⁻	5.23	(γ), n	11, 12, 13
6.696	2 ⁻	1.05	(γ), n	11, 12
6.699	1 ⁺	2.85	(γ), n	11, 12
6.709	0 ⁻	1.14	n	12
6.717	0 ⁻	0.95	n	12

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or Γ_{cm} (keV)	Decay	Reactions
6.729	(0 ⁻)		n	12
6.732	(0 ⁻)		n	12
6.737	(0 ⁻)		n	12
6.742	(0 ⁻)		n	12
6.746	(0 ⁻)		n	12
6.791	0 ⁺	1.9	n	12
6.835	1 ⁺	1.7	n	12
6.837	1 ⁺	0.4	n	12
6.856	1 ⁺	1.3	n	12
6.858	1	19	γ, n	11, 12, 13
7.005	0 ⁽⁻⁾	24	γ, n	11, 12, 13, 19
7.076	(1 ⁺)	24	γ, n	11, 12, 13
7.171	(2 ⁺)	14	γ, n	11, 12
7.311	(1)	33	γ, n	11, 12
(7.355)	(1)	19	n	12, 13
7.410	(2 ⁺)	10	γ, n	11, 12, 13
7.489	(2)	57	n	12
7.503	(0)	85	γ, n	11, 13
7.670	(2 ⁺)	60	γ, n	11, 12
7.80	(1, 2)	100	γ, n	11, 12
8.05 \pm 100	2 ⁺ ; 2			28
8.15	(1)	190	γ, n	11, 12
8.50		140	n	12
8.74		≤ 30	n	12
8.77		76	n	12
8.99		140	n	12
9.69		140	n	12
9.85		120	n, α	12, 18
10.024 \pm 10		200	n, α	12, 18
10.10 \pm 50			n, α	18
10.228 \pm 10	0 ⁻ , 1	≈ 200	n, α	12, 18
10.480 \pm 10		≈ 10	n, α	12, 18
(10.641 \pm 10)	1, 2	≈ 60	n	12

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or Γ_{cm} (keV)	Decay	Reactions
10.807 \pm 10 (10.89)	$0^-, 1$	≈ 330	n, α	12, 18
(11.045 \pm 10)			n, α	18
(11.130 \pm 10)		≈ 30	n	12
(11.244 \pm 10)		< 25	n	12, 18
11.49 \pm 50 (11.73)		< 25	n	12, 18
12.0		n, α	18	
12.2		n, α	18	
12.39		n, α	18	
12.82		n, α	18	
13.2		n, α	18	
13.66		n, α	18	
14.0		n, α	18	

^a See also Tables 20.6, 20.9 and 20.14.

6. $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$

$$Q_m = 4.245$$

$$E_b = 12.369$$

Excitation functions have been measured for $E_d = 0.9$ to 2.0 MeV (1960AM03; $\alpha_0, \alpha_1, \alpha_2, \alpha_3$), 0.5 to 3.0 MeV (1957BO04; α) and 9.6 to 11.5 MeV (1970BO08; $\alpha_4, \alpha_5, \alpha_6, \alpha_{7+8}, \alpha_{16}, \alpha_{17}$). At the lower energies a number of sharp structures are reported: see (1957BO04, 1960AM03). At the higher energies, there is no substantial resonant structure with widths less than a few hundred keV (1970BO08).

7. $^{18}\text{O}(\text{t}, \text{n})^{20}\text{F}$

$$Q_m = 6.112$$

The lifetime of $^{20}\text{F}^*(0.82)$, $\tau_m = 79 \pm 6$ psec and $J = 2, 4$. The 167-keV γ -ray from this state to $^{20}\text{F}^*(0.63)$ is observed to have a multipole mixing ratio near zero for both of the above J values. The results are consistent with $^{20}\text{F}^*(0.82)$ having $J^\pi = 4^+$ and being the third member of the ground state rotational band (1971PR10).

Table 20.5: The half-life of ^{20}F ^a

$\tau_{1/2}$ (sec)	Ref.
12.5 ± 2	(1959VA10)
11.2 ± 0.1	(1960SC01)
11.56 ± 0.05	(1962MA38)
11.36 ± 0.07	(1963GL01)
10.81 ± 0.11	(1967FL16)
10.31 ± 0.07	(1967YU01)
11.03 ± 0.06	(1970WI05)
11.03 ± 0.06	“best” value

^a See also (1952AJ38).

8. $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$

$$Q_m = 6.876$$

$$Q_0 = 6.8752 \pm 0.0015 \text{ (1970RO06)}.$$

Proton groups have been observed to states of ^{20}F with $E_x < 4.1$ MeV (1970RO06): see Table 20.7. Directional correlation and branching ratio measurements [see Table 20.6] lead to the J^π assignments shown in Table 20.7 (1970QU04). See also (1967BI01, 1967QU01). Angular distributions of the protons corresponding to $^{20}\text{F}^*(1.06, 3.49)$ are very similar at $E(^3\text{He}) = 18$ MeV: both show an $L = 0 + 2$ pattern, and therefore both are $J^\pi = 1^+$ states. This implies that $^{20}\text{F}^*(3.53)$ which is only weakly excited has $J^\pi = 0^+$: see the discussion in (1971FO14) and in reaction 19 [see also (1971BE19)]. See also (1969BA1Z).

9. $^{18}\text{O}(\alpha, \text{d})^{20}\text{F}$

$$Q_m = -11.478$$

Not reported.

10. (a) $^{18}\text{O}(^6\text{Li}, \alpha)^{20}\text{F}$

$$Q_m = 10.895$$

(b) $^{18}\text{O}(^7\text{Li}, \alpha \text{n})^{20}\text{F}$

$$Q_m = 3.645$$

$\tau_m \approx 1$ psec for $^{20}\text{F}^*(0.66)$ and 76 ± 20 psec for $^{20}\text{F}^*(0.82)$: see Table 20.14 (1969NI09).

Table 20.6: Radiative transitions in ^{20}F

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	Refs.
0.66	3^+	0	2^+	100	(1968SP01, 1970HO05, 1970QU04)
0.82 ^a	$4^+, 2^+$	0	2^+	36 ± 10	(1970QU04)
				42 ± 4	(1969HE20)
		0.66	3^+	64 ± 10	(1970QU04)
				58 ± 6	(1969HE20)
0.98 ^b	$(1^+, 2^+, 3^+)$	0	2^+	90 ± 10	(1970QU04)
				≥ 95	(1969HE20)
		0.66	3^+	10 ± 10	(1970QU04)
				≤ 5	(1969HE20)
				< 1	(1968SP01)
		0.82	$4^+, 2^+$	< 1	(1968SP01)
1.06	1^+	0	2^+	100	(1968SP01, 1970QU04)
		0.66	3^+	< 1	(1968SP01)
		0.82	$4^+, 2^+$	< 1	(1968SP01)
1.31	2^+	0	2^+	100	(1968SP01, 1969HO20, 1970QU04)
				≥ 86	(1969HE20)
		0.66	3^+	≤ 14	(1969HE20)
		0.82	$4^+, 2^+$	< 2	(1968SP01)
		0.98	$(1^+, 2^+, 3^+)$	< 1	(1968SP01)
		1.06	1^+	< 1	(1968SP01)
1.82	(3^-)	0	2^+	< 20	(1970QU04)
		0.82	$4^+, 2^+$	> 80	(1970QU04)
1.84 ^c	$2, 1^+$	0	2^+	> 95	(1970QU04)
		0.82	$4^+, 2^+$	< 5	(1970QU04)
1.97 ^d		0	2^+	20 ± 10	(1970QU04)
		0.82	$4^+, 2^+$	50 ± 15	(1970QU04)
		1.31	2^+	30 ± 10	(1970QU04)
2.04 ^e	2^+	0	2^+	20 ± 5	(1970QU04)
		0.66	3^+	80 ± 5	(1970QU04)
2.19 ^f	2^+	0	2^+	45 ± 10	(1970QU04)

Table 20.6: Radiative transitions in ^{20}F (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	Refs.
2.87	$2^+, 3^+$	0.82	$4^+, 2^+$	55 ± 10	(1970QU04)
		0	2^+	(100)	(1970QU04)
2.97 ^c		0	2^+	24 ± 3	(1969HE20)
				45 ± 10	(1970QU04)
		0.66	3^+	14 ± 2	(1969HE20)
		0.82	$4^+, 2^+$	62 ± 6	(1969HE20)
				55 ± 10	(1970QU04)
3.17 ^g		0	2^+	< 5	(1970QU04)
		0.98	$(1^+, 2^+, 3^+)$	> 95	(1970QU04)
3.49 ^h		1^+	0	2^+	68 ± 4
	0.98		$(1^+, 2^+, 3^+)$	7 ± 1	(1969HO20)
	1.06		1^+	7 ± 1	(1969HO20)
	1.31		2^+	10 ± 2	(1969HO20)
	1.84		$2, 1^+$	8 ± 2	(1969HO20)
3.53 ^b	0^+	1.06	1^+	100	(1968SP01, 1969HO20)
		3.59	$(1, 2, 3)^+$	0	2^+
				60	(1969HO20)
	2.04	2^+		34	(1968SP01)
				40	(1969HO20)
3.68	0	2^+		33	(1968SP01)
3.97	$(1, 2, 3)^+$	0.66	3^+	67	(1968SP01)
		0	2^+	77	(1969HA04)
		0.98	$(1^+, 2^+, 3^+)$	6	(1969HA04)
		1.31	2^+	17	(1969HA04)
		4.08 ⁱ	1^+	0	2^+
1.06	2^+	65 ± 5		(1969HO20)	
4.28	$(1, 2)^+$	1.06	1^+	100	(1968SP01)
5.41		2.04	2^+	100	(1968SP01)
5.55		0	2^+	24	(1968SP01)
		1.31	2^+	76	(1968SP01)

Table 20.6: Radiative transitions in ^{20}F (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	Refs.		
5.71	$(1, 2)^-$	1.06	1^+	60	(1969HA04)		
5.94 ^j		4.28	$(1, 2)^+$	40	(1969HA04)		
		0	2^+	7	(1968SP01)		
		0.66	3^+	35	(1968SP01)		
		0.98	$(1^+, 2^+, 3^+)$	4	(1968SP01)		
		1.31	2^+	1	(1968SP01)		
		1.84	$2, 1^+$	7	(1968SP01)		
		1.97		31	(1968SP01)		
		2.19	2^+	5	(1968SP01)		
		3.49	1^+	10	(1968SP01)		
6.02 ^j	$(1, 2)^-$	0	2^+	29	(1968SP01)		
				40	(1968BL1C)		
		0.66	3^+	3	(1968SP01)		
		0.98	$(1^+, 2^+, 3^+)$	19	(1968SP01)		
				23	(1968BL1C)		
		1.06	1^+	1	(1968BL1C)		
		1.31	2^+	2	(1968BL1C)		
		1.84	$2, 1^+$	6	(1968SP01)		
				6	(1968BL1C)		
		2.19	2^+	3	(1968SP01)		
		2.97		8	(1968SP01)		
		3.49	1^+	19	(1968SP01)		
				18	(1968BL1C)		
		3.59	$(1, 2, 3)^+$	10	(1968BL1C, 1968SP01)		
		4.08	1^+	3	(1968SP01)		
		6.04	$(1, 2)^-$	1.31	2^+	46	(1968SP01)
				1.84	$2, 1^+$	54	(1968SP01)
6.60	$0^+, 1^+$	see Table 20.9					
6.63	2^-	see Table 20.9					
6.65	1^-	see Table 20.9					

^a See also (1967BE36, 1968SP01, 1969HO20, 1970HO05, 1971PR10).

^b See also (1967BE36).

^c See also (1968SP01, 1970HO05).

^d See also (1967BE36, 1968SP01).

^e See also (1967BE36, 1968SP01, 1969HA04, 1970HO05).

^f See also (1968SP01, 1969HO20).

^g See also (1969HO20).

^h See also (1967BE36, 1968SP01, 1969HA04).

ⁱ See also (1968SP01).

^j See also (1969HA04).

11. $^{19}\text{F}(n, \gamma)^{20}\text{F}$

$$Q_m = 6.601$$

$$Q_0 = 6.6013 \pm 0.0010 \text{ (1967RA24);}$$

$$Q_0 = 6.602 \pm 0.002 \text{ (1967BL04, 1968BL1C);}$$

$$Q_0 = 6.011 \pm 0.0003 \text{ (1968SP01);}$$

$$Q_0 = 6.6020 \pm 0.0006 \text{ (1969HA04).}$$

The thermal capture cross section is 9.8 ± 0.7 mb (1963GL01: see (1964ST25)). At $E_n = 30$ keV it is reported to be 4.5 ± 1.0 mb (1963MA60). A number of resonances have been observed for $E_n \leq 1.65$ MeV: see Table 20.8 (1959GA08). The primary γ -rays resulting from capture at thermal energies ($^{20}\text{F}^*(6.60)$; $J^\pi = 0^+, 1^+$) and at $E_n = 27$ and 48 keV ($^{20}\text{F}^*(6.63, 6.65)$; $J^\pi = 2^-$ and 1^- , respectively) have been studied by several groups: see Table 20.9 (1965BI03, 1967BE36, 1967VA08, 1968BL1C, 1968SP01, 1969HA04). See also (1962BI1C, 1963NA08). There is some disagreement between the results of the various groups but it appears that the decay of $^{20}\text{F}^*(6.60)$ is dominated by two intense transitions (probably E1) to $^{20}\text{F}^*(5.94, 6.02)$ [thus $J^\pi = 1^-, 2^-$]. If the ground state transition is mainly M1, these two E1 transitions are (in terms of W.u.) about 150 times stronger than the M1 transition (1968SP01). It appears also that at $^{20}\text{F}^*(6.63, 6.65)$ [$J^\pi = 2^-$ and 1^- , respectively] the E1 transitions to the ground state are very weak, even though other E1 transitions in the decay of these two states have approximately normal strengths (1965BI03, 1967BE36). Branching ratios for other ^{20}F states involved in this reaction are shown in Table 20.6 (1967BE36, 1968BL1C, 1968SP01, 1969HA04).

Table 20.10 displays excitation energies for ^{20}F states involved in cascade and in primary γ -transitions (1968SP01, 1969HA04). (1968BL1C) suggest 4^+ for $^{20}\text{F}^*(0.83)$ since neither $^{20}\text{F}^*(6.60)$ nor $^{20}\text{F}^*(6.02)$ [$J^\pi = 0^+, 1^+$, and $J^\pi = 1^-$, respectively] decays to it and 3^+ for $^{20}\text{F}^*(0.66)$ since the 6.60 MeV state decays to it and $^{20}\text{F}^*(6.02)$ does not. See also (1958GR01, 1961WA03, 1963GI1D) and (1964ST25).

12. $^{19}\text{F}(n, n)^{19}\text{F}$

$$E_b = 6.601$$

Table 20.7: States in ^{20}F from $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$

$E_x^{\text{ a}}$ (keV)	$J\pi^{\text{ c}}$
0	$2^+{}^{\text{ d}}$
657.2 ± 1.3	3^+
823.5 ± 1.5	$2^+, 4^+$
982.9 ± 1.3	$1^+, 2^+, 3^+$
1058.1 ± 1.4	1^+
1309.1 ± 1.4	$(1^+, 2^+)$
$1824.4 \pm 1.6^{\text{ b}}$	$1, 2, 3, 5$
$1843.0 \pm 1.7^{\text{ b}}$	$2, 1^+, 3^+$
1971.9 ± 1.6	
2044.0 ± 1.6	2^+
2195.5 ± 2.0	$3^+, 2^+, 1^+$
2868.2 ± 2.3	
2967.1 ± 2.0	$2^+, 3^+$
3487.8 ± 2.2	$1^+{}^{\text{ d}}$
3586.3 ± 2.2	
3681.0 ± 2.5	
3761.0 ± 3.1	
3966.9 ± 2.8	
4083.7 ± 2.9	

^a (1970RO06).

^b (1967QU01) find $E_x = 1824.4 \pm 2.1$ and 1843.0 ± 2.2 keV.

^c From (p, γ) correlation measurements (1970QU04). See also (1967BI01, 1967QU01).

^d Known from β -decay: see reaction 29 in ^{20}Ne .

Table 20.8: Resonances in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

E_n ^b (keV)	$\sigma(n, \gamma)$ ^c (mb)	Γ_γ (eV)	Γ (keV)	J^π	E_x (MeV)
27	308	1.1 ^e	0.4 ^f	2 ⁻	6.627
48	36	1.6	1.5 ^f	1 ⁻	6.647
100	4.4	2.2	12 ^f	1 ⁻	6.696
177					6.769
270 ^d	1.4	3.9	20	1	6.858
308					6.894
388					6.970
425	0.31	1.5	25	0	7.005
500	0.30	1.9	25	0, 1	7.076
600 ^d	1.8	8.1	15	1	7.171
760	0.11	2.9	60	1	7.323
865	0.11		60		7.423
950	0.06	2.8	95	0	7.503
1125	0.09	3.9	80	1	7.670
1290	0.27	8.6	75	1, 2	7.826
1635	0.07	7.5	180	1	8.154

^a (1959GA08). See also Tables 20.6 and 20.9.

^b (1963MA58) report a resonance at $E_n = 15.5$ keV which (1967BL1J) do not observe. However (1967BL1J) report a narrow resonance at $E_n = 84$ keV.

^c At resonance after correction for resolution.

^d See also (1950HE92).

^e $\Gamma_n = 0.38 \pm 0.10$ keV, $\Gamma_\gamma = 0.55 \pm 0.15$ keV (1963MA58).

^f From σ_t measurements.

Table 20.9: Primary capture transitions in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

Final state $^{20}\text{F}^*$ (MeV)	I_γ ^b from $^{20}\text{F}^*(6.60)$		I_γ ^b from $^{20}\text{F}^*(6.63)$		I_γ ^b from $^{20}\text{F}^*(6.65)$	
	(1968SP01)	(1969HA04)	(1965BI03)	(1967BE36)	(1965BI03)	(1967BE36)
0	11	10	< 1		< 1	5 ± 3
0.66	< 0.1		4	6 ± 2	5	9
0.82	< 0.1				(2)	} 13
0.98	1.5	2			5	
1.06 ^c	5	6			5	
1.31	3	3	25	32 ± 2	5	12
1.84	2	2				
1.97	0.1		22	50 ± 2	5	} 42 ± 2
2.04	6	6	25		20	
2.19			4		6	
2.87			3			
2.97	< 0.2				11	
3.49	2.5	2				
3.53	2		5	} 8 ± 1	10	} 20 ± 2
3.59	4.4	4				
3.68	1	1				
3.97		1				
4.08		1	4	5 ± 2	4	
4.28	1	1				
5.04			3			
5.41	0.5					
5.55	3					
5.94	17	13				
6.02 ^c	38	48				
6.04	2					

^a See also Tables 20.6 and 20.8 and (1968BL1C).

^b In units of photons/100 captures.

^c E_γ for the transitions (6.60 \rightarrow 0), (6.60 \rightarrow 1.06) and (6.60 \rightarrow 6.02) are, respectively, 6599.8 ± 3.0 , 5534.9 ± 2.0 and 583.6 ± 0.5 keV (1967VA08).

Table 20.10: States of ^{20}F involved in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

E_x (keV)	
(1968SP01)	(1969HA04)
0	0
656.3 ± 0.3	656 ± 1
822.9 ± 0.3	
983.8 ± 0.3	984 ± 1
1057.2 ± 0.3	1057 ± 1
1309.1 ± 0.3	1309 ± 1
1843.4 ± 0.3	1843 ± 1
1970.6 ± 0.3	
2044.2 ± 0.4	2044 ± 1
2194.5 ± 0.6	2194 ± 2
2965.8 ± 0.5	2966 ± 2
3488.3 ± 0.3	3488 ± 2
3526.0 ± 0.5	
3587.3 ± 0.3	3588 ± 2
3681.0 ± 0.4	3681 ± 2
	3967 ± 2
4082.2 ± 0.5	4085 ± 2
4276.7 ± 0.5	4275 ± 2
5413.1 ± 0.6	
5554.7 ± 6	
	5713 ± 2
5936.0 ± 0.3	5937 ± 1
6017.3 ± 0.3	6018 ± 1
6044.6 ± 0.4	
6601.1 ± 0.3	6602.0 ± 0.6

^a For measurements of I_γ , see (1968SP01, 1969HA04).

The scattering amplitude (bound) is $a = 5.74 \pm 0.09$ fm (1963BA1F). The coherent scattering cross section is 3.9 ± 0.1 b (1964ST25). See also (1961WI1A, 1969BA1P).

The total cross section has been measured for $E_n = 1$ to 300 keV (1964HI04), 0.2 to 2.2 MeV (1964EL02), 0.45 to 1.20 MeV (1966CA14), 2.5 to 15 MeV (1971FO1P, 1971FO24), 2.61 to 2.83 MeV (1965SO1A), 3.35 to 5.07 MeV (1960TS02), 7 to 14 MeV (1957PE1B), 13.70 to 14.60 MeV (1968HU17), 14.5 MeV (1970AN1F), and 17.7 to 29.1 MeV (1960PE25). See also (1966GA1K, 1969AN1E). For earlier measurements see (1959AJ76). Observed resonances in the total cross section are shown in Table 20.11 (1958WI36, 1960TS02, 1964HI04, 1964ST25, 1966CA14). The level density may be appreciably greater than that shown in Table 20.11: see (1964HI04) and (1964EL02).

Angular distributions and polarization measurements have been carried out over the range $E_n = 0.2$ to 2.2 MeV (1964EL02). The results have been interpreted in terms of doorway states: see (1967AF01, 1967MO1M, 1968LE1P). Angular distributions have also been measured at $E_n = 14.1$ MeV (1970CL03) and for $E_n = 0.66$ to 2.92 MeV (1958WI36). See also (1970GA1A). Polarization measurements are reported by (1962OT01) at $E_n = 3.5$ MeV. See also (1959MA1C, 1965WY1B, 1966EL1C, 1967EL1F) and (1964ME1B, 1967FE1F, 1971BU02, 1971GI05; theor.).

13. $^{19}\text{F}(n, n')^{19}\text{F}^*$

$$E_b = 6.601$$

The excitation function for 0.110, 0.197, 1.24 and 1.37 MeV γ -rays has been measured from threshold to 2.2 MeV by (1955FR1B). See also (1966VE1B, 1971BE1G). Resonances are reported at $E_n = (100), 270, 420, (500), 780, 830, 880$ and 950 keV, corresponding to the excitation of ^{20}F states at $E_x = (6.70), 6.86, 7.00, (7.08), 7.34, 7.39, 7.44$ and 7.50 MeV (1955FR1B). See also (1966VE1B). The cross section for excitation of $^{19}\text{F}^*(2.78)$ has been studied for $E_n = 3$ to 3.6 MeV. The very slow rise for the first 300 keV above threshold is consistent with the large spin difference between $^{19}\text{F}^*(2.78)$ and the ground state [$J^\pi = \frac{9}{2}^+$ for the former, $\frac{1}{2}^+$ for the latter]. See also (1970CL03) and (1964ST25).

14. $^{19}\text{F}(n, 2n)^{18}\text{F}$

$$Q_m = -10.431$$

$$E_b = 6.601$$

Cross sections have been measured from $E_n = 10$ to 37 MeV (1961BR1A), 12.4 to 20.9 MeV (1965PI03), 12.6 to 19.6 MeV (1965BO42), 12.7 to 19.4 MeV (1967ME22), 13.5 to 15.0 MeV (1967BO24), 13.6 to 14.6 MeV (1968VO1B), 13.9 to 14.8 MeV (1964ST1F), 14.1 MeV (1962CE1B), 14.2 MeV (1965NA1C), 14.4 MeV (1961RA06, 1963RA1A) and at 14.7 MeV (1967PA27). See (1959AJ76) for earlier reports. See also (1960MC05, 1961WI1C, 1963PI1B, 1964HE18, 1965GO1C, 1967CS02), (1964ST25, 1966JE1B) and (1969CH1N; theor.).

15. $^{19}\text{F}(n, p)^{19}\text{O}$

$$Q_m = -4.036$$

$$E_b = 6.601$$

Cross sections have been measured for $E_n = 12.6$ to 19.6 MeV (1965BO42), 14.4–14.8 MeV

Table 20.11: Resonances in $^{19}\text{F}(n, n)^{19}\text{F}$

E_n (keV)	Γ_{lab} (keV)	J^π	$^{20}\text{F}^*$ (MeV)	Refs.
15.25			6.616	(1964HI04)
27.35	0.30	2^-	6.627	(1964HI04)
31.9			6.632	(1964HI04)
34.0			6.634	(1964HI04)
38.0			6.637	(1964HI04)
49.7	1.70	1^-	6.648	(1964HI04)
70.5			6.668	(1964HI04)
87.8	4.00	0^-	6.685	(1964HI04)
96.0	5.50	1^-	6.692	(1964HI04)
100.0	1.10	2^-	6.696	(1964HI04)
102.4	3.00	1^+	6.699	(1964HI04)
113.2	1.20	0^-	6.709	(1964HI04)
121.7	1.00	0^-	6.717	(1964HI04)
134.0		(0^-)	6.729	(1964ST25)
138.0		(0^-)	6.732	(1964ST25)
143.1		(0^-)	6.737	(1964ST25)
148.0		(0^-)	6.742	(1964ST25)
152.5		(0^-)	6.746	(1964ST25)
200.2	2.0	0^+	6.791	(1964HI04)
246.0	1.8	1^+	6.835	(1964HI04)
247.8	0.4	1^+	6.837	(1964HI04)
267.7	1.4	1^+	6.856	(1964HI04)
270	20	1	6.858	(1964ST25)
425	25	0	7.005	(1964ST25)
500	25	(1^+)	7.076	(1964ST25, 1966CA14)
600	15	(2^+)	7.171	(1964ST25, 1966CA14)
747	35	(1)	7.311	(1966CA14)
794	20	(1)	(7.355)	(1966CA14)
852	11	(2^+)	7.410	(1966CA14)
935	60	(2)	7.489	(1958WI36, 1964ST25, 1966CA14)
1100	50	(2^+)	7.65	(1958WI36, 1964ST25, 1966CA14)
1250	150		7.79	(1958WI36, 1964ST25)
1620	220		8.14	(1958WI36, 1964ST25)

Table 20.11: Resonances in $^{19}\text{F}(n, n)^{19}\text{F}$ (continued)

E_n (keV)	Γ_{lab} (keV)	J^π	$^{20}\text{F}^*$ (MeV)	Refs.
2000	150		8.50	(1958WI36, 1964ST25)
2250	≤ 30		8.74	(1958WI36, 1964ST25)
2280	80		8.77	(1958WI36, 1964ST25)
2520	150		8.99	(1958WI36, 1964ST25)
3250	150		9.69	(1958WI36, 1964ST25)
3420	130		9.85	(1958WI36, 1960TS02, 1964ST25)
3460 ± 10			(9.887)	(1960TS02)
3505 ± 10			(9.929)	(1960TS02)
3560 ± 10			(9.982)	(1960TS02)
3605 ± 10	200		10.024	(1958WI36, 1960TS02, 1964ST25)
3820 ± 10	≈ 200	$0^-, 1$	10.228	(1960TS02)
4085 ± 10	≈ 10		10.480	(1960TS02)
4255 ± 10	≈ 60	1, 2	10.641	(1960TS02)
4430 ± 10	≈ 330	$0^-, 1$	10.807	(1960TS02)
4680 ± 10	≈ 30		11.045	(1960TS02)
4770 ± 10	< 25		11.130	(1960TS02)
4890 ± 10	< 25		11.244	(1960TS02)
(4935)			(11.29)	(1960TS02)

(1962KA37, 1966MI14, 1966PR14, 1967CS03, 1967PA27, 1968RE07) and for $E_n = 18.2$ to 21.0 MeV (1965PI03). See also (1959AJ76), (1961DA16, 1963PI1B), (1960BU1C, 1963LE1D, 1964ST25, 1966JE1B, 1971CU1B, 1971PR09) and ^{19}O .

16. $^{19}\text{F}(n, d)^{18}\text{O}$ $Q_m = -5.768$ $E_b = 6.601$

See (1960VE06, 1967VA12, 1968RE07, 1969LI1F) and ^{18}O .

17. $^{19}\text{F}(n, t)^{17}\text{O}$ $Q_m = -7.557$ $E_b = 6.601$

See (1967VA12, 1968RE07) and ^{17}O in (1971AJ02).

18. $^{19}\text{F}(n, \alpha)^{16}\text{N}$ $Q_m = -1.523$ $E_b = 6.601$

Table 20.12: Resonances in $^{19}\text{F}(n, \alpha)^{16}\text{N}$

E_n (MeV \pm keV)			E_x (MeV)
(1955MB01)	(1960SM03)	(1961DA16)	
3.4			9.8
3.61 ± 50			10.03
3.69 ± 50			10.10
3.77 ± 50	3.75 ± 50	3.85	10.17
4.11 ± 50	4.08 ± 50	4.1	10.49
4.42 ± 50	4.36 ± 50	4.35	10.77
	4.52 ^a		10.89
4.86 ± 50	4.79 ± 50	4.80	11.18
	5.15 ± 50		11.49
	5.40 ^a		11.73
		5.7	12.0
5.9 ± 100	5.9 ^a		12.2
		6.10	12.39
		6.55	12.82
		6.9	13.2
		7.44	13.66
		7.8	14.0

^a Not resolved.

Observed resonances are shown in Table 20.12 (1955MB01, 1960SM03, 1961DA16): see graph in (1964ST25). Excitation functions are reported from threshold to $E_n = 6$ MeV (1960SM03), $E_n = 8.2$ MeV (1961DA16) and for $E_n = 14.2$ to 21.0 MeV (1965PI03). See also (1959AJ76), (1962KA37, 1963PI1B, 1966BH05, 1966KN02, 1968RE07, 1969BR1F), (1960BU1C, 1963CH20, 1966JE1B), (1964GA1A; theor.) and ^{16}N in (1971AJ02).

19. $^{19}\text{F}(d, p)^{20}\text{F}$

$$Q_m = 4.377$$

$$Q_0 = 4.3777 \pm 0.0009 \text{ (1970RO06; see also (1967SP09))}.$$

Measurements of proton spectra and γ -rays have led to the identification of a number of states of ^{20}F : see Table 20.13 (1969HE20, 1969HO20, 1970RO06). Angular distributions have been studied at $E_d = 0.6$ MeV (1970SC14), 0.8 to 2.5 MeV (1964EL01), 1.32 to 2.03 MeV (1959ON26),

Table 20.13: States in ^{20}F from $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$

E_x (keV) ^a			l_n ^d	J^π ^e	$(2J_f + 1)S^j$
(1970RO06) ^b	(1969HO20) ^c	(1969HE20) ^c			
0			2	$1^+, 2^+, 3^+$	≤ 0.06
654.9 ± 1.0	655.9 ± 0.2	655.4 ± 0.5	2	$1^+, 2^+, 3^+$	2.59
821.6 ± 1.0	823.0 ± 0.3	822.6 ± 0.7	2	$1^+, 2^+, 3^+$	$\lesssim 0.3$
983.3 ± 1.0	983.9 ± 0.3	983.4 ± 0.7	2	$1^+, 2^+, 3^+$	$\lesssim 0.3$
1056.3 ± 1.0	1057.0 ± 0.2	1055.2 ± 0.6	0	$0^+, 1^+$	0.019
1310.8 ± 1.1	1309.3 ± 0.2	1308.0 ± 0.9	iso., 2 ^g	$(1^+, 2^+, 3^+)$	$\lesssim 0.02$
1843.4 ± 1.2	1843.5 ± 0.7				$\lesssim 0.11$
	2043.7 ± 0.5		2	$1^+, 2^+, 3^+$	2.32
2195.1 ± 1.5	2194.5 ± 0.6		2	$1^+, 2^+, 3^+$	0.50
2863.7 ± 1.6			3 ^h	$2^-, 3^-, 4^-$	$\lesssim 0.01$
2966.6 ± 1.7	2966.8 ± 0.6	2964.5 ± 2.0	1	$0^-, 1^-, 2^-$	0.36
3171.8 ± 2.2	3175.6 ± 1.3		$\lesssim 0.2$		
	3488.5 ± 0.3		0	$0^+, 1^+$	1.20
3525.5 ± 2.6	3525.9 ± 0.5		0	$0^+, 1^+$	0.28
3586.4 ± 2.7	3586.5 ± 0.6		2 ^g	$1^+, 2^+, 3^+$	0.36
3681.0 ± 2.5					$\lesssim 0.04$
3760.8 ± 2.7					
3964.5 ± 2.5					$\lesssim 0.04$
4080.9 ± 2.5	4082.5 ± 0.8		0	$0^+, 1^+$	0.18
4198.9 ± 2.7					
4207.7 ± 2.6					
4276.3 ± 2.8			2 ^g	$1^+, 2^+, 3^+$	0.07
4311.5 ± 2.6			0	$0^+, 1^+$	0.27
4583.8 ± 3.0					
4592.2 ± 2.9					
4730.2 ± 2.9					
4763.8 ± 2.7					
4891.6 ± 2.8					
4898.2 ± 2.8					
5040.2 ± 3.1			1	$0^-, 1^-, 2^-$	
5065.5 ± 3.1					

Table 20.13: States in ^{20}F from $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$ (continued)

E_x (keV) ^a			l_n ^d	J^π ^e	$(2J_f + 1)S^j$
(1970RO06) ^b	(1969HO20) ^c	(1969HE20) ^c			
5224.0 ± 3.1			1	0 ⁻ , 1 ⁻ , 2 ⁻	
5281.0 ± 3.3			1	0 ⁻ , 1 ⁻ , 2 ⁻	
5317.1 ± 2.7					
5344.5 ± 3.3					
5450.3 ± 3.8					
5455.4 ± 3.2					
5463.4 ± 3.3					
5620.3 ± 3.3					
5762.8 ± 3.4					
5809.1 ± 2.9			(1)	(0 ⁻ , 1 ⁻ , 2 ⁻)	
5933.9 ± 3.3 ^f			1	0 ⁻ , 1 ⁻ , 2 ⁻	
6015.0 ± 3.8 ^f			1 ⁱ	0 ⁻ , 1 ⁻ , 2 ⁻	
6043.3 ± 3.7					
		(1956EL1A)			
		6.25 ± 20			
		6.52 ± 20			
		6.63 ± 20	1	0 ⁻ , 1 ⁻ , 2 ⁻	
		6.81 ± 20	1	0 ⁻ , 1 ⁻ , 2 ⁻	
		6.98 ± 20	1	0 ⁻ , 1 ⁻ , 2 ⁻	
		7.20 ± 20	1	0 ⁻ , 1 ⁻ , 2 ⁻	

^a See also (1959AJ76, 1960RI05, 1964LO1A).

^b From measurements of proton groups.

^c From measurements of γ -rays.

^d (1956EL1A): see also (1963RO21, 1964EL01, 1964LO1A, 1966SC09, 1968BI09, 1970FO1J, 1970SC14, 1970ZA02, 1971LA1F, 1971RO06, 1972LA1N).

^e J^π derived from I_n measurements only. Unique assignments derived from (p- γ) correlation work are discussed in the text and displayed in Table 20.6.

^f (1971RO06) find $E_x = 5.932 \pm 0.005$ and 6.013 ± 0.005 MeV, respectively.

^g (1963RO21): $E_d = 2$ MeV.

^h (1963RO21) find $l_n = 1$.

ⁱ (1971RO06).

^j (1970FO1J) and H.T. Fortune, private communication.

1.35 to 2.00 MeV (1969BA2M), 1.5 MeV (1964LO1A), 2 MeV (1963RO21, 1968BI09), 2.6 to 4.0 MeV (1970ZA02), 3 MeV (1971LA1F, 1972LA1N), 8.9 MeV (1956EL1A, 1971RO06), 9 and 13 MeV (1966SC09) and 16 MeV (1970FO1J). See also (1959AJ76).

Branching ratio and angular correlation measurements, together with lifetime determinations [see Table 20.14] permit a unique choice of J^π in many cases, from among the J^π values stemming from direct interaction analyses of angular distributions [see Table 20.13]. These J^π values are displayed in Table 20.6 (1969HE20, 1969HO20, 1970HO05, 1970RO06). See also (1964CH21, 1965NE07).

(1970QU1A), using polarized deuterons, have looked at the protons corresponding to $^{20}\text{F}^*(0.66, 2.04)$. These two states are found to be populated by a $j = \frac{5}{2}$ neutron transfer. This result, together with (p- γ) correlation data, provides a unique $J^\pi = 3^+$ assignment for $^{20}\text{F}^*(0.66)$ and is in agreement with 2^+ for $^{20}\text{F}^*(2.04)$. Transitions to $^{20}\text{F}^*(2.19, 2.97)$ appear to be admixed $j = \frac{3}{2}, \frac{5}{2}$ transfer implying $J = 2$ for both these states (1970QU1A). See also (1972HA2H). Comparison of the results of this reaction and of the $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$ and $^{22}\text{Ne}(\text{d}, \alpha)^{20}\text{F}$ reactions leads to assignments of $J^\pi = 1^+$ and 0^+ for $^{20}\text{F}^*(3.49, 3.53)$ (1971FO14).

See also (1959GO1C, 1960RA23, 1963BO1C, 1967HE1D, 1968CA1E) and (1958SA1A, 1959BO1C, 1959HO1D, 1959SI1A, 1960NA1A, 1962DA06, 1963RO13, 1966NE1C, 1971BU02; theor.).

$$20. \text{}^{19}\text{F}(\text{t}, \text{d})^{20}\text{F} \quad Q_{\text{m}} = 0.344$$

See (1963HU1C).

$$21. \text{}^{19}\text{F}(^{19}\text{F}, ^{18}\text{F})^{20}\text{F} \quad Q_{\text{m}} = -3.833$$

See (1971GA13).

$$22. \text{}^{20}\text{O}(\beta^-)^{20}\text{F} \quad Q_{\text{m}} = 3.815$$

See ^{20}O .

$$23. \text{}^{20}\text{Ne}(\text{n}, \text{p})^{20}\text{F} \quad Q_{\text{m}} = -6.244$$

See (1971KA18).

$$24. \text{}^{20}\text{Ne}(\text{t}, ^3\text{He})^{20}\text{F} \quad Q_{\text{m}} = -7.008$$

Table 20.14: Lifetime measurements of some ^{20}F states

$^{20}\text{F}^*$ (MeV)	τ_m	Refs.
0.66 ^a	0.36 ± 0.07 psec	(1969HE20)
	0.37 ± 0.06 psec	(1970HO05)
	0.42 ± 0.05 psec	A
0.82 ^b	76 ± 20 psec	(1969NI09)
	≥ 4.4 psec	(1969HE20)
	$0.9^{+0.9}_{-0.4}$ psec	(1970HO05)
	79 ± 6 psec	(1971PR10)
0.98	$1.3^{+0.6}_{-0.4}$ psec	(1969HE20)
	1.8 ± 0.4 psec	(1970HO05)
1.06	≤ 92 fsec	(1969HE20)
	45 ± 13 fsec	(1970HO05)
1.31	$1.1^{+0.4}_{-0.3}$ psec	(1969HE20)
	0.8 ± 0.3 psec	(1970HO05)
1.84	30 ± 20 fsec	(1970HO05)
2.04	≤ 38 fsec	(1969HE20)
	37 ± 16 fsec	(1970HO05)
2.19	≤ 46 fsec	(1969HE20)
	< 12 fsec	(1970HO05)
2.97	≤ 62 fsec	(1969HE20)
	60 ± 40 fsec	(1970HO05)
3.49	≤ 47 fsec	(1969HE20)
	44 ± 11 fsec	(1970HO05)
3.53	≤ 32 fsec	(1969HE20)
	30 ± 15 fsec	(1970HO05)
3.59	30 ± 30 fsec	(1970HO05)

A: E.K. Warburton, J.W. Olness, G.A.P. Engelbertink and T.K. Alexander, private communication.

^a See also (1969NI09).

^b See also (1971PR10).

Not reported.

25. (a) $^{21}\text{Ne}(\text{n}, \text{d})^{20}\text{F}$ $Q_{\text{m}} = -10.780$
(b) $^{21}\text{Ne}(\text{d}, ^3\text{He})^{20}\text{F}$ $Q_{\text{m}} = -7.511$

Not reported.

26. $^{21}\text{Ne}(\text{t}, \alpha)^{20}\text{F}$ $Q_{\text{m}} = 6.809$

Not reported.

27. $^{22}\text{Ne}(\text{n}, \text{t})^{20}\text{F}$ $Q_{\text{m}} = -14.888$

Not reported.

28. $^{22}\text{Ne}(\text{p}, ^3\text{He})^{20}\text{F}$ $Q_{\text{m}} = -15.652$

At $E_{\text{p}} = 43.7$ to 45.0 MeV, analog states have been studied in ^{20}F and ^{20}Ne [the latter via $^{22}\text{Ne}(\text{p}, \text{t})^{20}\text{Ne}$]. The experimental cross-section ratio, R , for the transitions to $^{20}\text{Ne}^*(10.28)$ and $^{20}\text{F}(0)$ [2^+ ; $T = 1$] is 2.00 ± 0.20 ; R for the transitions to $^{20}\text{Ne}^*(12.25 \pm 0.03)$ and $^{20}\text{F}^*(1.82)$ [3^- ; $T = 1$] is 1.40 ± 0.15 (1969HA19). Angular distributions for the ^3He ions and the tritons corresponding to the first $T = 2$ states ($J^{\pi} = 0^+$) [$^{20}\text{Ne}^*(16.722 \pm 0.025)$ and $^{20}\text{F}^*(6.513 \pm 0.033)$] have also been compared. There is indication also for the excitation of the 2^+ ; $T = 2$ states [at $E_{\text{x}} = 8.05$ MeV in ^{20}F and at 18.5 MeV in ^{20}Ne (estimated errors ± 0.1 MeV)] (1964CE05, 1969HA38). See also (1970OL1B).

29. $^{22}\text{Ne}(\text{d}, \alpha)^{20}\text{F}$ $Q_{\text{m}} = 2.701$

At $E_{\text{d}} = 10$ MeV, α -particle groups are observed to a number of states below $E_{\text{x}} = 6.0$ MeV. $^{20}\text{F}^*(1.82)$ was more strongly excited than $^{20}\text{F}^*(1.84)$ (1970FO1H). The angular distributions (at $E_{\text{d}} = 10$ MeV) of the α -particles corresponding to $^{20}\text{F}^*(3.49, 3.53)$ have been measured. $^{20}\text{F}^*(3.49)$ is strongly populated and the angular distribution shows direct interaction features while $^{20}\text{F}^*(3.53)$ is weakly populated and the angular distribution is roughly symmetric about 90° . These results are consistent with the assignments $J^{\pi} = 1^+$ and 0^+ for $^{20}\text{F}^*(3.49, 3.53)$, respectively (1971FO14): see reactions 8 and 19. See also (1966LA14).

30. $^{23}\text{Na}(n, \alpha)^{20}\text{F}$

$$Q_m = -3.867$$

See (1960VA15, 1961WI22, 1962KA12, 1962KA26, 1963CS03, 1966WO03) and (1964FA04, 1964MA1N, 1964SA1B; theor.).

^{20}Ne
(Figs. 11 and 13)

GENERAL: (See also (1959AJ76):

Shell model: (1959BR1E, 1960SO1A, 1960TA1C, 1962TA14, 1963FL1A, 1963KE1B, 1963LE1C, 1963SA07, 1964BA1J, 1964BA1K, 1964IN03, 1964KE06, 1964MO1E, 1964PA1D, 1964TA1D, 1965BA1J, 1965DA1D, 1965DE1H, 1965YU1A, 1965ZA1B, 1966BA2E, 1966BA2H, 1966BA2C, 1966BO17, 1966IN01, 1966LO07, 1966RI1F, 1966ZA1E, 1967BA1W, 1967BA1K, 1967BA1V, 1967BO09, 1967EN01, 1967FE01, 1967FL13, 1967GU05, 1967HA1M, 1967LA1H, 1967MU02, 1967PA1P, 1967PA10, 1967PA12, 1967PI1B, 1967ST02, 1967ST1N, 1967ST25, 1967ST26, 1967ST1L, 1967SV1A, 1968AN1G, 1968AR02, 1968BA1L, 1969BO1Y, 1968CO11, 1968DR1B, 1968EL1C, 1968GU1E, 1968GU1C, 1968GU1G, 1968HA17, 1968HA26, 1968HA1P, 1968HE1H, 1968HI1H, 1968MO1G, 1968SA1J, 1968SA10, 1968ST06, 1968WO1A, 1969AB05, 1969AK02, 1969BA2G, 1969BA2J, 1969BA2K, 1969BE1V, 1969CO1M, 1969FA06, 1969FO1E, 1969FO04, 1969GU1E, 1969HA1Z, 1969HA45, 1969JA10, 1969KE1B, 1969KUZZ, 1969LA26, 1969LE1Q, 1969MA1T, 1969PA13, 1969SA19, 1969SA1F, 1969SA1A, 1969SV1A, 1969TE04, 1969WO1G, 1969WO06, 1970AN27, 1970AR21, 1970AS1F, 1970BO1J, 1970CA24, 1970EI06, 1970FA1F, 1970FR14, 1970GI1F, 1970GI11, 1970HA49, 1970KH01, 1970KR1D, 1970KU16, 1970MC1J, 1970MC23, 1970NG01, 1970PA15, 1970PA27, 1970RO1B, 1970RU1A, 1970ST1D, 1970SV1B, 1970TE1A, 1970TU01, 1970WA39, 1970WO12, 1971AR05, 1971AR25, 1971AR1R, 1971BO29, 1971FA08, 1971FO19, 1971GU08, 1971GU1N, 1971HA1U, 1971HO23, 1971JE02, 1971KE10, 1971LA13, 1971MA22, 1971RA16, 1971SC01, 1971ST01, 1971WI01, 1971WO13, 1971ZO03, 1972LE1L, 1972RE03).

Collective and deformed models: (1959BE1A, 1960BE1A, 1962TA14, 1963FL1A, 1963KE1B, 1963LE1C, 1964BA1J, 1964BA1K, 1964KE06, 1966AB04, 1966BO17, 1967BA1V, 1967BE1V, 1967DA1G, 1967FE01, 1967KE1H, 1967KR1E, 1967LA1G, 1967PA12, 1967PA1M, 1967RI1B, 1967ST25, 1967ST1L, 1968CO11, 1968HO1G, 1968RI1M, 1968SA10, 1968ST06, 1968UN1A, 1969AB05, 1969AK02, 1969BA2G, 1969BE1V, 1969DA14, 1969HA45, 1969LE1Q, 1969VO1E, 1969WO06, 1970FA1F, 1970FR14, 1970GO1Q, 1970KH01, 1970NG01, 1970PA15, 1970RI1C, 1970SV1B, 1970VO01, 1971AR1R, 1971BO29, 1971CH22, 1971DE1R, 1971HO19, 1971HO23, 1971KE10, 1971MI02, 1971PI02, 1971ST01, 1971ZO03, 1971ZO01, 1972AB1C, 1972LE1L).

Cluster and α -particle models: (1959BE1A, 1960BE1A, 1960SH1A, 1962BO23, 1962VA1D, 1964MA1G, 1965IN1A, 1965NE1C, 1966HA33, 1966KA1A, 1968AB1B, 1968PI1A, 1968TA1G, 1969AB1B, 1969BA2E, 1969BA2L, 1969CH1P, 1969HI1B, 1969SM1A, 1969WE1J, 1970BA2B, 1970BA2H, 1970BR35, 1970EI06, 1971CH22, 1971FR06, 1971HA44, 1971KH06, 1971MC1F, 1972AB1C).

Astrophysical questions: (1961GO1G, 1963DU1C, 1970BA1M).

Giant resonances (See also reactions 30 and 32.): (1963MI1B, 1965DA1D, 1965GR1H, 1966BA2F, 1967BE1V, 1969HA1Y, 1969HO1W).

Table 20.15: Energy levels of ^{20}Ne ^a

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$0^+; 0$	0^+		stable	1, 2, 3, 4, 5, 6, 8, 13, 14, 16, 19, 20, 26, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 45, 46, 47, 48, 49, 50, 51
1.6338 ± 0.4	$2^+; 0$	0^+	$\tau_m = 1.20 \pm 0.15$ psec	γ	3, 4, 8, 13, 14, 15, 16, 18, 19, 20, 26, 27, 28, 29, 36, 37, 39, 40, 42, 43, 44, 45, 46, 47, 50
4.2473 ± 1.8	$4^+; 0$	0^+	$\tau_m = 93 \pm 9$ fsec	γ	3, 4, 8, 13, 14, 16, 18, 19, 20, 26, 27, 28, 36, 37, 39, 40, 44, 45, 46, 47, 50
4.9682 ± 0.8	$2^-; 0$	2^-	$\tau_m = 4.8 \pm 0.5$ psec	γ	3, 4, 13, 19, 20, 26, 27, 28, 29, 36, 40, 44, 45, 47, 50
5.6217 ± 2.0	$3^-; 0$	2^-	$\tau_m = 200 \pm 50$ fsec	γ, α	3, 4, 8, 13, 14, 18, 19, 20, 26, 27, 28, 36, 37, 40, 45, 47, 50
5.785 ± 3	$1^-; 0$	0^-	$\Gamma > 1.3 \times 10^{-2}$	γ, α	3, 8, 13, 14, 18, 19, 20, 27, 28, 37, 40, 45, 50
6.722 ± 3	$0^+; 0$	0^+	15 ± 7	γ, α	3, 8, 9, 13, 19, 26, 27, 28, 45
7.0055 ± 2.8	$4^-; 0$	2^-	$\tau_m = 440 \pm 90$ fsec	γ	3, 4, 13, 27, 28
7.166 ± 4	$3^-; 0$	0^-	$\Gamma = 8$	α	3, 9, 13, 14, 27, 28, 40
7.196 ± 4	$0^+; 0$		4	γ, α	3, 8, 9, 13, 14, 27, 28, 37
7.424 ± 4	$2^+; 0$	0^+	8	α	3, 9, 26, 27, 36, 43
7.834 ± 4	$2^+; 0$		2	α	3, 9, 19, 26, 36, 43
8.447 ± 3	$5^-; 0$	2^-		γ, α	3, 8
≈ 8.6	$0^+; 0$		> 800	α	9
8.72 ± 20	$1^-; 0$		2.5	α	3, 9, 43
8.7750 ± 2.2	$6^+; 0$	0^+	0.110 ± 0.025	γ, α	3, 8, 9, 13, 19, 36

Table 20.15: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
≈ 8.8	$2^+; 0$		> 800	α	9
8.82	$(5^-); 0$		< 1	α	9
8.850 ± 5	$1^-; 0$		19	α	3, 9
9.040 ± 5	$4^+; 0$	0^+	≤ 3	γ, α	3, 8, 9, 19
9.117 ± 5	$3^-; 0$		3.2	α	3, 9, 26, 36
(9.34 ± 30)					3, 26
9.489 ± 9	$2^+; 0$		29 ± 15	γ, α	3, 8, 9, 43
9.950 ± 6	(1^+)		$\tau_m < 35$ fsec	γ	3, 19
9.99 ± 10	$4^+; 0$		$\Gamma = 150 \pm 50$	γ, α	8, 9, 19, 26, 36
10.257 ± 5	$5^-; 0$	0^-	141	α	9
10.26 ± 20	$2^+; 1$		$\lesssim 2$	γ, α	3, 8, 19, 26, 43
10.401 ± 5	$3^-; 0$		81	α	9
10.548 ± 5	$4^+; 0$		16	α	3, 9
10.579 ± 10	$2^+; 0$		24	α	3, 9
10.609 ± 7	$6^-; 0$	2^-	$\tau_m = 23 \pm 7$ fsec	γ	3
10.79 ± 100	$4^+; 0$		$\Gamma = 350$	α	9
10.836 ± 5	$2^+; 0$		13	α	9, 43
10.836	$3^-; 0$		45	α	9
10.853 ± 10	$(1, 2, 3)^+; 1$				19, 26
10.920 ± 7			$\tau_m < 30$ fsec	γ	3
10.97 ± 150	$0^+; 0$		$\Gamma = 580$	α	9
11.015 ± 10	$4^+; 0$		24	α	9
11.08 ± 20	$(4^+; 1)$		$\lesssim 3$	γ, α	8, 26
11.23 ± 30	$1^-; 0$		172	α	9
11.233 ± 10	$1^+; 1$			γ	20, 26, 34, 45
11.27 ± 30	2^+		$\lesssim 4$	γ, α	8, 43
11.324 ± 10	$2^+; 0$		53	α	9
11.528 ± 6	≤ 4		$\tau_m < 30$ fsec	γ, α	3, 8, 19
11.549 ± 10	$(T = 1)$				26, 34
11.871 ± 20	$2^+; 0$		$\Gamma = 46$	α	9

Table 20.15: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
11.925 \pm 5	4 ⁺ ; 0		0.44 \pm 0.15	α	9, 26
11.948 \pm 5	8 ⁺ ; 0	0 ⁺	(35 \pm 10) \times 10 ⁻³	α	3, 8, 9, 13
11.953 \pm 10	1 ⁻ ; 0		24	α	9
11.971 \pm 8	1 ⁻ ; 0		29 \pm 8	α	9
12.086 \pm 10	($T = 1$)			(α)	9, 26
12.150 \pm 10	6 ⁺ ; 0		< 40		3, 26
(12.200 \pm 10)	($T = 1$)				26
12.224 \pm 10	4 ⁺ ; 0		142	α	9
12.245 \pm 15	(2 ⁺); 1		40 \pm 16	γ, α	8, 20, 26
12.35 \pm 100	2 ⁺ ; 0			α	9
12.367 \pm 10	3 ⁻ ; 0		46 \pm 16	α	9, 26
12.410 \pm 5	0 ⁺ ; 0		\leq 8	α	9, 19, 26
12.559 \pm 10	6 ⁺ ; 0		101	α	3, 9
12.61 \pm 100				α	9
12.682 \pm 15	5 ⁻ ; 0		97	α	9
12.77 \pm 100	4 ⁺ ; 0		100	α	9
12.83 \pm 30			55	α	9, 19
12.98 \pm 75	(4 ⁺ ; 0)		60	α	9
13.060 \pm 4	2 ⁻		1.0	p, α	32
13.086 \pm 15	(4 ⁺ ; 0)		70	α	9, 26
13.168 \pm 1	1 ⁺ ; (1)		2.3 \pm 0.2	γ, p, α	20, 21, 22, 26
13.18 \pm 75	(4 ⁺ ; 0)		60	α	9
13.224	1 ⁻		95	p, α	22
13.224	0 ⁺		95	p, α	22
13.304 \pm 1	1 ⁺		0.9 \pm 0.1	γ, p, α	20, 21, 22
13.333 \pm 6	7 ⁻ ; 0	2 ⁻	(80 \pm 30) \times 10 ⁻³	α	3, 9
13.342	4 ⁺ ; 0		20	α	9
13.411	2 ⁻		35	γ, p, α	9, 20, 21, 22
13.42 \pm 140	(4 ⁺ ; 0)		110	α	9
13.462 \pm 20	1 ⁻		190	p, α	22

Table 20.15: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
13.479 \pm 1.5	1 ⁺ ; 1		7.1	γ, p, α	19, 20, 21, 22, 26
13.523	(1 ⁻)		30	p, α	21, 22
13.541	2 ⁺		63	p, α	22
13.584	2 ⁺		\approx 10	p, α	21, 22
13.63 \pm 30			7.9	p	19, 21
13.650 \pm 15	0 ⁺ ; 1		22	p, α	21, 22, 26
(13.66)	1 ⁻		115	p, α	22
13.673 \pm 1	2 ⁻		4.5 \pm 0.2	γ, p, α	20, 21, 22
13.699			4.6	p	21
13.7 \pm 400	(3, 7) ⁻		320	α	9
(13.73)	0 ⁺		170	p, α	22
13.733 \pm 1.5	1 ⁺		7.7 \pm 0.5	γ, p, α	20, 21, 22
13.775				γ, p	20
(13.87)	(1 ⁻)		190	p, α	22
13.88 \pm 30	(6 ⁺)		100	α	9, 13
13.882 \pm 15			\approx 1	γ, p	20, 21, 26
13.903	2 ⁺		48	p, α	32
13.924			3.5	p	21
13.946	0 ⁺		\approx 70	p, α	22
14.017	1 ⁻		\approx 70	p, α	21, 22
14.03	2 ⁺		\approx 140	p, α	22
14.065			18	γ, p	20, 21
14.098			3.8	γ, p	20
14.124 \pm 1.2	2 ⁻		4.7 \pm 0.7	γ, p, α	9, 20, 21, 22
14.134	2 ⁺		51	p, α	22
14.148 \pm 1.2	2 ⁻		11.8 \pm 1.0	γ, p, α	20, 21, 22
14.197	1 ⁺		13.9 \pm 1	γ, p	20, 21
14.3 \pm 300	6 ⁺		240	α	9, 13
14.373			\approx 5	p	21
14.421				p	21

Table 20.15: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
14.453 \pm 2			33 \pm 3	p, α	21, 22
14.467	0 ⁺		135	p, α	22
14.6 \pm 300	(4 ⁺)		240	α	9
14.604	1 ⁻		125	p, α	22
14.695 \pm 2.6	(1 ⁺)		38 \pm 10	p, α	21, 22
14.772 \pm 3.0			95 \pm 20	p, α	21, 22
14.85 \pm 150	(2 ⁺ , 4 ⁺)		\approx 100	p, α	9, 22
15.03 \pm 150	(2 ⁺)		\approx 90	p, α	9, 22
15.18 \pm 40	9 ⁻				3
15.23			28	p, α	22
15.26	(1 ⁻)		285	p, α	13, 22
15.30	(0 ⁺)		285	p, α	22
15.39			76	p, α	22
15.44			57	p, α	22
15.52			150	p, α	22
15.59				α	9
15.618	(8 ⁻)	2 ⁻			3
15.71			28	α	9, 22
15.9 \pm 40	5 ⁻			(α)	3, 9, 22
15.9 \pm 40	8 ⁺			(α)	3, 9, 22
16.02			100	p, α	22
16.24				α	9
16.34			(100)	p, α	9, 22
16.49				p, α	9, 22
16.59			125	p, α	9, 22
16.728 \pm 4	0 ⁺ ; 2		2.0 \pm 0.5	γ , p, α	19, 20, 21, 22, 45
16.77			100	p, α	22
16.98			100	n, p, α	22, 23
17.08			90	n, p, α	22, 23
b					

Table 20.15: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
17.30			60	n, p, α	22, 23
17.50			86	n, p, α	22, 23
17.58			38	n, p, α	22, 23
17.76			180	p, α	9, 22
18.08 \pm 180	(6 ⁺ , 7 ⁻)		140	n, p, α	3, 9, 23
18.31 \pm 300	(6 ⁺)		240	n, p, α	9, 23
18.426 \pm 5	2 ⁺ ; $T = 2$		10 \pm 3	γ , n, p, α	20, 21, 22, 23, 45
18.7	(6 ⁺)		600	α	9
19.16 \pm 250	(6 ⁺)		200	α	9
19.40 \pm 350	6 ⁺		280	α	9
19.84 \pm 350	6 ⁺		280	α	9
20.16 \pm 120	7 ⁻		250	α	9
20.4 \pm 180	6 ⁺		360	α	9
20.4 \pm 100	7 ⁻		200	α	9
20.68 \pm 60	9 ⁻		120	α	9
20.9				α	9
21.0	7 ⁻		200	α	9
21.08 \pm 40	9 ⁻		80	α	9
21.3	7 ⁻		300	α	9
21.8 \pm 200	7 ⁻		300	α	9
22.3 \pm 200	7 ⁻		500	α	9
22.7 \pm 250	9 ⁻		500	α	9
22.84 \pm 125	9 ⁻		250	α	9
23.4 \pm 250	8 ⁺		500	α	9
24.11 \pm 150	8 ⁺		350	α	9
25.0 \pm 300	8 ⁺		600	α	9
25.7 \pm 200			400	α	9
27.2 \pm 350			700	α	9
28.	8 ⁺		1600	^3He , α	9, 17
28.2 \pm 350			700	α	9

^a See also Tables 20.17 and 20.18.

^b Additional states with $E_x > 17.1$ MeV are reported in reaction 23: see Table 20.30.

Electromagnetic transitions: (1965GR1H, 1965NE1C, 1965ST22, 1967GU1D, 1967HS1A, 1967KA06, 1967KU1E, 1968AN1G, 1968EL1C, 1968GU1G, 1968HA17, 1969AB05, 1969AK02, 1969BE1V, 1969CU07, 1969FO1E, 1969HA1Y, 1969HA1Z, 1969HA1F, 1969HI1B, 1969JA10, 1969LA26, 1970AN27, 1970HA49, 1971AR05, 1971AR25, 1971FO19, 1971GU1N, 1971HA44).

Special levels: (1960SH1A, 1961BA1D, 1961TR1B, 1964BA1J, 1964BA1K, 1964EN1A, 1964MI16, 1964TA1D, 1966BA42, 1966BO17, 1966BR1H, 1966LO07, 1966ZA1E, 1967EN01, 1968AN1G, 1968AR02, 1968HA1P, 1968SA1J, 1969AK02, 1969BO1Z, 1969HA1Y, 1969HA1G, 1969HA1F, 1969JA10, 1969ST1J, 1969TE04, 1969WO06, 1970AN27, 1970AR21, BA70W, 1970GO1Q, 1970HO17, 1970RO1B, 1970RU1A, 1970TE1A, 1971AR1R, 1971BO1F, 1971FA08, 1971GU08, 1971LE30, 1971SE1C, 1971ST36).

Special reactions: (1960AN1A, 1960LE1A, 1970BE1D, 1970HU07, 1971AR02).

Pion and kaon capture and reactions: (1970GA1K).

Other topics: (1960EV1A, 1960SO1A, 1961BA1D, 1962CH12, 1963DA1C, 1963EV01, 1964AB1A, 1964BR1H, 1964EN1A, 1964IN03, 1964KA1A, 1964MO1E, 1965DE1H, 1965YO1A, 1966DA1E, 1966GI1A, 1966IN01, 1966SU1D, 1966YO1B, 1967AB03, 1967BA1W, 1967BR1G, 1967FL13, 1967GU05, 1967GU1D, 1967KA06, 1967KE1H, 1967PA1P, 1967ST02, 1967ST1N, 1967ST25, 1967ST26, 1968DA1P, 1968DR1B, 1968DW1A, 1968FL1C, 1968GU1E, 1968GU1C, 1968GU1G, 1968HE1H, 1968MO1G, 1968PA1Q, 1968SA10, 1968TA1G, 1968WO1A, 1969BA2G, 1969BA2J, 1969BA2K, 1969BO1Z, 1969DE15, 1969KE1B, 1969LA26, 1969MA1T, 1969RU04, 1969SH1E, 1969SO08, 1969WO06, 1970BO1J, 1970DA13, 1970FO1K, 1970GM1A, 1970GO1R, 1970PA15, 1970RU1A, 1970RU1B, 1970ST19, 1970TU01, 1970VO01, 1971FA13, 1971FO04, 1971FR1B, 1971GO36, 1971HA14, 1971HO19, 1971HO23, 1971JE02, 1971LA13, 1971MI02, 1971SC01, 1971SE1C, 1971ST01, 1971ZO03, 1972LE1L, 1972RE03).

Ground state: $J = 0$ (1960LU06);

$$\mu < 4 \times 10^{-4} \text{ nm (1960LU06); see also (1964LI14).}$$

See also (1967BA1V, 1967GU1D, 1967LA1G, 1967SH14, 1967ST26, 1968EL1C, 1968RI1M, 1969BA2J, 1969CO1M, 1969FO1E, 1969FO04, 1969FU11, 1969HI1B, 1969KE1B, 1969LA26, 1969SA19, 1969SA1F, 1970GO1R, 1970KH01, 1970RU1B, 1970ST19, 1970TU01, 1970VO01, 1971AR25, 1971BO29, 1971FO19, 1971GU1N, 1971LA13, 1971RA16, 1971SC01, 1971ST01, 1971TE1C, 1971WI01, 1971ZO03, 1972LE1L).

$$Q(^{20}\text{Ne}^*(1.63)) = -0.24 \pm 0.03 \text{ b (1970NA07);}$$

$$B(E2)(0^+ \rightarrow 2^+) = 0.048 \pm 0.007 \text{ e}^2 \cdot \text{b}^2 \text{ (1970NA07).}$$

1. ${}^9\text{Be}({}^{14}\text{N}, \text{t}){}^{20}\text{Ne}$ $Q_{\text{m}} = 6.304$

See (1958GO71).

2. ${}^{10}\text{B}({}^{14}\text{N}, \alpha){}^{20}\text{Ne}$ $Q_{\text{m}} = 19.534$

See (1970IS1A).

3. ${}^{12}\text{C}({}^{12}\text{C}, \alpha){}^{20}\text{Ne}$ $Q_{\text{m}} = 4.618$

Particle group and γ -ray energy measurements have been made for many states of ${}^{20}\text{Ne}$: see Table 20.16 (1961AL12, 1964PE02, 1966KU03, 1967KU04, 1970PA08, 1971HA26, 1971MI09, 1971MI1J, 1971PA1C, 1971SC12). Angular correlation (1961GO12, 1962KU03, 1964KU03, 1966KU03, 1967KU04, 1967SM04, 1971SC12) and γ -ray branching measurements [see Table 20.17] (1964BR18, 1967BR22, 1967SM04, 1971HA26) lead to the J^π assignments shown in Table 20.16, which also show level assignments to rotational bands. See also (1962BR35, 1964AL15, 1966BR1T).

At $E({}^{12}\text{C}) = 22.0$ to 35.0 MeV the α -spectra are dominated by the group corresponding to the 2^+ state at $E_{\text{x}} = 7.83$ MeV. Both ${}^{20}\text{Ne}^*(7.83)$ and ${}^{20}\text{Ne}^*(7.20)$ [$J^\pi = 0^+$] are much more strongly excited than the 0^+ and 2^+ states at $E_{\text{x}} = 6.72$ and 7.42 MeV. At $E({}^{12}\text{C}) = 25$ and 27 MeV, the distributions of the α -particles to ${}^{20}\text{Ne}^*(7.83)$ is barely oscillatory and is strongly forward peaked, indicating a direct interaction, while the distributions to ${}^{20}\text{Ne}^*(7.42)$ are compound nuclear in character. This is consistent with the hypothesis that ${}^{20}\text{Ne}^*(7.20, 7.83)$ are quartet states of the configuration (220) [two sd-shell α -particles outside a ${}^{12}\text{C}$ core], populated via a “semi-direct” mechanism that proceeds through the formation of doorway states in the incident channel (1971MI09). See also (1970AR21, 1971MI1J, 1971MI11). At $E({}^{12}\text{C}) = 50$ MeV, compound nucleus formation dominates (1971HA1D). Angular distributions are also reported at $E({}^{12}\text{C}) = 5.6$ to 6.2 MeV (1963AL07; $\alpha_0 \rightarrow \alpha_3, \alpha_{4+5}$), 10.6 to 12.4 MeV (1963KU05; α_0), 12.5 MeV (1961PO13; α_0, α_1), ≈ 20.0 to ≈ 28.4 MeV (1964BO35; α_0), 20.3 to 25.6 MeV (1964AL17, 1964VO03; α_0, α_1), 21.5 to 28.3 MeV (1964BO35; α_1) and 50 MeV (1970PA08, 1971PA1C; $\alpha_0, \alpha_5, \alpha_{10}$). See also (1972NO01; theor.).

(1969PA22) have studied the yield of this reaction to permit predictions of the cross section in the energy region of interest in stellar carbon burning. The importance of this reaction for astrophysical work is discussed by (1969AR1F). See also ${}^{24}\text{Mg}$ in (1973ENVA).

For lifetime measurements, see Table 20.18 (1962BR35, 1965EV03, 1971HA26). See also (1961CL06).

See also (1959AL1H, 1960AL05, 1961LI07, 1961LI17, 1961SE1C, 1968JA1F, 1970MA1D, 1971HA1D, 1971MA1E), (1966KU1D) and (1961LI06, 1965BO1H, 1966ER1A, 1967VO1B, 1968GA1J).

Table 20.16: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$

E_x (MeV \pm keV)							$\Gamma_{\text{c.m.}}$ (keV)	J^π ^e	K^π
(1961AL12) ^a	(1964PE02) ^a	(1967KU04) ^a	(1971HA26) ^b	(1971MI09) ^{a,c}	(1971PA1C) ^{a,d}	(1971SC12) ^a			
			1.6329 \pm 1.0					2 ⁺	0 ⁺
4.25 \pm 20	4.25 \pm 20		4.2456 \pm 2.5					4 ⁺	0 ⁺
4.97 \pm 20	4.97 \pm 20		4.9663 \pm 2.5					2 ⁻	2 ⁻
5.64 \pm 20	5.62 \pm 20		5.618 \pm 4					3 ⁻	2 ⁻
5.81 \pm 20	5.79 \pm 20							1 ⁻	0 ⁻
6.17 \pm 20									
6.74 \pm 20	6.71 \pm 20			6.722 \pm 4				0 ⁺	
6.87 \pm 20									
7.05 \pm 20	7.02 \pm 20		7.004 \pm 4	7.007 \pm 4				4 ⁻	2 ⁻
7.19 \pm 20	7.17 \pm 20			7.159 \pm 4		7.15 \pm 60		3 ⁻	0 ⁻
7.25 \pm 20	7.21 \pm 20			7.195 \pm 4				0 ⁺	
7.46 \pm 20	7.43 \pm 20			7.421 \pm 4				2 ⁺	
7.65 \pm 20									
7.86 \pm 20	7.85 \pm 20			7.833 \pm 4		7.83 \pm 60		2 ⁺	
7.93 \pm 20									
8.52 \pm 20	8.46 \pm 20		8.446 \pm 9					5 ⁻	2 ⁻
	8.71 \pm 20					8.73 \pm 60		1 ⁻	
	8.79 \pm 20								
8.92 \pm 20	8.87 \pm 20								
	9.04 \pm 20								
	9.11 \pm 20								
	(9.31 \pm 20)								
	9.48 \pm 20								
			9.950 \pm 6						

Table 20.16: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ (continued)

E_x (MeV \pm keV)							$\Gamma_{\text{c.m.}}$ (keV)	J^π ^e	K^π
(1961AL12) ^a	(1964PE02) ^a	(1967KU04) ^a	(1971HA26) ^b	(1971MI09) ^{a,c}	(1971PA1C) ^{a,d}	(1971SC12) ^a			
	10.24 \pm 20	10.57 \pm 40 (10.65)	10.609 \pm 7 10.920 \pm 7 11.528 \pm 6				< 40	(1 ⁺) 4 ⁺ 6 ⁻	2 ⁻
		11.99 ^{f,g} 12.19 \pm 40					< 40	8 ⁺ 6 ⁺	0 ⁺
		13.39 \pm 40				12.5 \pm 300 13.43 \pm 60	< 40	6 ⁺ 7 ⁻	2 ⁻
					15.18 \pm 40 15.62 \pm 30 15.9 \pm 40 15.9 \pm 40 17.4 18.15 \pm 40		\leq 23 \leq 40	9 ⁻ (8 ⁻) 5 ⁻ 8 ⁺	2 ⁻
							\leq 23 \leq 40	8 < J < 12 7 ⁻	

^a From measurements of particle groups.

^b From measurements of γ -rays.

^c See also (1971MI1J).

^d A.D. Panagiotou, private communication; see also (1970PA08).

^e From work done with this reaction.

^f (1966KU03).

^g See also (1971MA23).

Table 20.17: Radiative decays in ^{20}Ne ^a

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.
1.63	$2^+; 0$	0	100		17.8 ± 2.5	(1971HA26)
4.25	$4^+; 0$	1.63	≈ 100		21.9 ± 2.1	(1971HA26)
4.97 ^d	$2^-; 0$	0	0.6 ± 0.2		2×10^{-3}	(1967BR22)
		1.63	99		$(2.5 \pm 0.9) \times 10^{-3}$	(1971HA26)
					$(1.7 \pm 0.7) \times 10^{-2}$ ^b	(1971HA26)
					$(7.0 \pm 1.0) \times 10^{-6}$ ^c	(1971HA26)
5.62 ^e	$3^-; 0$	0	7.6 ± 1.0		10.6 ± 2.7	(1964BR18)
		1.63	87.6 ± 1.0			(1971HA26)
		4.97	4.8 ± 1.6		$(6.6 \pm 1.5) \times 10^{-6}$	(1964BR18)
5.79 ^f	$1^-; 0$	0	18 ± 5		35 ± 9	(1971HA26)
		1.63	82 ± 5	4.0	7.2×10^{-6}	(1965VA14)
6.72	$0^+; 0$	1.63	100	33	9.2×10^{-5}	(1965VA14)
7.01 ^g	$4^-; 0$	1.63	0.5 ± 0.2		3.8	(1965VA14)
		4.25	63.5			(1967BR22)
		4.97	11		$(1.7 \pm 0.4) \times 10^{-5}$	(1971HA26)
		5.62	25			(1967BR22)
					10.4 ± 1.8	(1971HA26)
					24 ± 5	(1967BR22)
7.20	$0^+; 0$	1.63				(1971HA26)
8.45	$5^-; 0$	5.62	100	13 ± 3	26 ± 6	(1969GR03)
8.78 ^h	$6^+; 0$	4.25	100	100 ± 15	20.4 ± 2.4	(1971RO33)
9.04	$4^+; 0$	1.63	(100)	0.38 ± 0.05	7.0	(1971DI08, 1971RO13)
9.49	$2^+; 0$	0		$\lesssim 0.06$	$\lesssim 0.3$	(1964PE05)
		1.63	(100)	0.26 ± 0.1	3.2 (E2), 0.03 (M1)	(1964PE05)
9.95	$(1^+); 0$	1.63	> 90			(1964PE05)
9.99	$4^+; 0$	0		$\lesssim 0.07$		(1964PE05)
		1.63	(100)	0.9 ± 0.4	6.9	(1964PE05)
10.26	$2^+; 1$	0		0.19 ± 0.05	0.6	(1964PE05)

Table 20.17: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.
10.61 ⁱ	$6^-; 0$	1.63		5.6 ± 0.6	0.41 (M1), 0.017 (E2)	(1964PE05)
		4.97		$\lesssim 0.44$	$\lesssim 0.006$	(1964PE05)
		7.01	95.5 ± 1.2		17_{-4}^{+7}	(1971HA26)
		8.45	4.5 ± 1.2		10 ± 4	(1971HA26)
10.92		1.63	25 ± 5			(1971HA26)
		4.25	75 ± 5			(1971HA26)
11.08	$(4^+; 1)$	1.63		$\lesssim 0.08$	$\lesssim 0.4$	(1964PE05)
		4.25	(100)	4.8 ± 0.5	0.72 (M1), 7.3 (E2)	(1964PE05)
		7.01		$\lesssim 1.4$	$\lesssim 0.04$	(1964PE05)
11.23	$1^+; 1$	0	(100)			(1964PE05)
11.53	≤ 4	4.25	34 ± 6			(1971HA26)
		4.97	66 ± 6			(1971HA26)
11.95	$8^+; 0$	8.78	100		7.5 ± 2.5	(1972AL05)
12.25	$(2^+); 1$	1.63	(100)			(1964PE05, 1968LA1H)
13.48	$1^+; 1$	1.63	95			(1961GO21)
		4.97	5			(1961GO21)
13.88		1.63	20			(1961GO21)
		4.97	80	^k		(1961GO21)
16.73	$0^+; 2$	11.23	(100)	≈ 5000		(1967KU06)
18.43	$2^+; 2$	12.25	(100)	≈ 300		(1968LA1H)

^a See also (1961CL06, 1962EI04, 1965EV03).

^b If M2.

^c If E2.

^d See also (1960KA18, 1961GO21).

^e See also (1964AL15, 1965VA14).

^f See also (1964AL15).

^g See also (1962BR35, 1966BR1T, 1967SM04).

^h See also (1967LI07).

ⁱ See also (1967SM04).

^j See also (1964PE05). Note: this footnote is not labeled in the tabular.

^k See Table 20.23.

Table 20.18: Lifetime measurements of some ^{20}Ne states ^a

$^{20}\text{Ne}^*$ (MeV)	τ_m	Refs.
1.63 ^b	1.23 ± 0.12 psec	(1965EV03)
	1.15 ± 0.20 psec	(1971HA26)
4.25 ^c	1.20 ± 0.15 psec	‘best’ value (1971HA26)
	134 ± 12 fsec	(1965EV03)
	150 ± 25 fsec	(1969AN08, 1969AN1J)
4.97 ^c	93 ± 9 fsec	(1971HA26)
	93 ± 9 fsec	‘best’ value
	$2.8_{-0.9}^{+2.4}$ psec	(1969AN08, 1969AN1J)
	$4.8_{-1.1}^{+1.7}$ psec	(1965EV03)
5.62	4.8 ± 0.5 psec	(1971HA26)
	4.8 ± 0.5 psec	‘best’ value
	350 ± 75 fsec	(1965EV03)
7.01	200 ± 50 fsec	(1971HA26)
	200 ± 50 fsec	‘best’ value
	405_{-90}^{+110} fsec	(1962BR35)
	470 ± 90 fsec	(1971HA26)
9.95	440 ± 90 fsec	mean value
	< 35 fsec	(1971HA26)
10.61	23 ± 7 fsec	(1971HA26)
10.92	< 30 fsec	(1971HA26)
11.53	< 30 fsec	(1971HA26)

^a See also (1964LI1B).

^b See also (1956DE22, 1959AL91, 1961CL06, 1969AN08, 1969AN1J, 1969GR03, 1969TH01).

^c See also (1961CL06).

4. $^{12}\text{C}(^{14}\text{N}, ^6\text{Li})^{20}\text{Ne}$ $Q_m = -4.182$

At $E(^{14}\text{N}) = 52$ and 60 MeV the reaction appears to proceed via the direct transfer of eight nucleons. Angular distributions have been reported for the ^6Li ions corresponding to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62, 7.01)$ (1971MA23). Assignment of observed states to rotational bands is discussed by (1971MA23, 1971NA17).

5. $^{12}\text{C}(^{16}\text{O}, ^8\text{Be})^{20}\text{Ne}$ $Q_m = -2.637$

See (1963GO27, 1971GO1M).

6. $^{12}\text{C}(^{19}\text{F}, ^{11}\text{B})^{20}\text{Ne}$ $Q_m = -3.112$

See (1967BO1P, 1971BO1V). See also (1969VO1D).

7. (a) $^{14}\text{N}(^6\text{Li}, \text{p})^{19}\text{F}$	$Q_m = 11.148$	$E_b = 23.994$
(b) $^{14}\text{N}(^6\text{Li}, \text{d})^{18}\text{F}$	$Q_m = 2.943$	
(c) $^{14}\text{N}(^6\text{Li}, \alpha)^{16}\text{O}$	$Q_m = 19.263$	
(d) $^{14}\text{N}(^6\text{Li}, \alpha\text{n})^{15}\text{O}$	$Q_m = 3.594$	

Yield curves have been measured for $E(^6\text{Li}) = 4.1$ to 6.3 MeV for the protons to $^{19}\text{F}^*(2.78)$, and the d_0 and α_0 groups: no structure is apparent (1968RI13). See also (1961NO05).

8. $^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}$ $Q_m = 4.730$

Observed resonances in the yield of capture γ -rays over the range $E_\alpha = 0.8$ to 10 MeV are displayed in Table 20.19 (1964PE05, 1965VA14, 1967LI07, 1969GR03, 1972AL05, 1971DI08, 1971RO13, 1971RO33, 1971TO06, 1971TO1C). Information on the character of the radiative decay is shown in Table 20.17.

No resonances have been observed below $E_\alpha = 1$ MeV: for $E_{\text{res}} < 0.85$ MeV, $(2J+1)\Gamma_\alpha\Gamma_\gamma/\Gamma \leq 0.044$ eV, and $(2J+1)\Gamma_\alpha\Gamma_{\gamma_1}/\Gamma \leq 0.031$ eV; for $E_{\text{res}} < 1.00$ MeV, $\omega\gamma \leq 0.036$ eV and $\omega\gamma_1 \leq 0.024$ eV (1971TO06, 1971TO1C). The astrophysical implications of this reaction are discussed in (1970TO1F, 1971TO06, 1971TO1C). See also (1957BU66) and (1965ME10, 1967WI1B).

The $J^\pi = 5^-$ states at $E_x = 8.45$ MeV [$E_\alpha = 4.65$ MeV] decays by an E2 transition [$|M|^2 =$

Table 20.19: Resonances in $^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\omega\gamma$ ^a (eV)	E_x (MeV)	$J^\pi; T$	Refs.
1.116 \pm 4	2.6×10^{-6} ^b	$(1.7 \pm 0.3) \times 10^{-3}$	5.623	$3^-; 0$	(1965VA14, 1971TO06, 1971TO1C)
1.319 \pm 3	$> 1.3 \times 10^{-2}$ ^b	$(14 \pm 3) \times 10^{-3}$	5.785	$1^-; 0$	(1965VA14, 1971TO06, 1971TO1C)
2.490 \pm 8	15 ± 7 ^b	$(38 \pm 10) \times 10^{-3}$	6.722	$0^+; 0$	(1965VA14, 1971TO06, 1971TO1C)
3.09			7.20	$0^+; 0$	(1969GR03)
4.647 \pm 3			8.447	$5^-; 0$	(1971RO33)
5.06	< 3	1.35 ± 0.15	8.775 ± 0.0032	$6^+; 0$	(1967LI07, 1971DI08, 1971RO13)
5.374 \pm 9	$\lesssim 3$	3.4 ± 0.4	9.028	$4^+; 0$	(1964PE05)
5.94 \pm 30	29 ± 15	1.3 ± 0.5	9.48	$2^+; 0$	(1964PE05)
6.61 \pm 30	155 ± 30	8 ± 3	10.02	$(4^+); 0$	(1964PE05)
6.930 \pm 10	≤ 2	28 ± 3	10.272	$2^+; 1$	(1964PE05)
7.94 \pm 20	≤ 3	46 ± 5	11.08	$(4^+; 1)$	(1964PE05)
8.17 \pm 30	≤ 4	0.9 ± 0.4	11.26	$(1^- \text{ or } 2^+)$	(1964PE05)
8.54 \pm 40		0.5 ± 0.3	11.56	≤ 4	(1964PE05)
^c		0.104 ± 0.035	11.948	$8^+; 0$	(1972AL05)
9.40 \pm 40	40 ± 16	7 ± 3	12.25	≤ 4	(1964PE05)

^a See also Table 20.17.

^b This is also Γ_α .

^c $E(^{16}\text{O}) \approx 36.05$ MeV.

26 ± 6 W.u.] to the 3^- state at $E_x = 5.62$ MeV (1971RO33). The $J^\pi = 6^+$ state at $E_x = 8.78$ MeV [$E_\alpha = 5.06$ MeV] decays by an E2 transition [$|M|^2 = 20.4 \pm 2.4$ W.u.] to the 4^+ state at $E_x = 4.25$ MeV (1971DI08, 1971RO13). The $J^\pi = 8^+$ state at $E_x = 11.95$ MeV decays by an E2 transition to the $E_x = 8.78$ MeV [$J^\pi = 6^+$] state which then decays via the 4^+ and 2^+ members of the ground state rotational band. The transition probability of the $8^+ \rightarrow 6^+$ transition is 7.5 ± 2.5 W.u. which establishes $^{20}\text{Ne}^*(11.95)$ as the 8^+ member of the ground state band. The experimental E2 transition probabilities in the ground state band deviate strongly from those predicted by the pure rotational model but agree reasonably well with simple shell model predictions (1972AL05).

The lifetime of $^{20}\text{Ne}^*(1.63)$, τ_m , is 0.84 ± 0.20 psec (1969GR03): see also Table 20.18.

- | | | |
|---|----------------|---------------|
| 9. (a) $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ | | $E_b = 4.730$ |
| (b) $^{16}\text{O}(\alpha, 2\alpha)^{12}\text{C}$ | $Q_m = -7.162$ | |
| (c) $^{16}\text{O}(\alpha, ^8\text{Be})^{12}\text{C}$ | $Q_m = -7.254$ | |

Excitation functions have been measured over a wide energy range for elastically scattered α -particles and γ -rays from the decay of $^{16}\text{O}^*(6.13, 6.92, 7.12)$: see Table 20.20. See also (1964BO1E, 1965CA02) and (1963DA1D). Angular distributions of various α -particle groups are reported for $E_\alpha = 5.0$ to 104 MeV: see Table 16.27 in (1971AJ02).

A number of anomalies are observed in the elastic scattering. Phase shift analyses lead to the results shown in Table 20.21 (1953CA44, 1960MC09, 1964PE05, 1965MC02, 1967HU06, 1967ME10, 1969JO18, 1971BE17, 1972HA07, 1971TA05). At higher energies, a prominent 2 MeV wide resonance is observed corresponding to $^{20}\text{Ne}^*(28.)$ with $J^\pi = 8^+$ (1969CO19, 1970CO13). In the range $E_\alpha = 37$ to 51 MeV, the cross section for elastic scattering appears to decrease monotonically (1965VA11). Identifying various of the observed states as members of different rotational bands is discussed by (1967HU06, 1972HA07): see also Table 20.15. The $J^\pi = 6^+$, 8^+ and 7^+ states at $E_x = 8.78$, 11.95 and 13.33 MeV are probable members of the lowest rotational bands with $K^\pi = 0^+$ and $K^\pi = 2^+$. The known reduced widths in both bands are small, indicating that the bands should be describable in terms of the spherical shell model. The reduced widths within each band are found to decrease sharply with increasing spin (1972HA07). See also (1963NO1C, 1968CE1B, 1969AG06) and (1968SH1G, 1969PI02, 1971MC03; theor.).

Reaction (b) can be described in terms of both sequential and one-step knockout processes. A state at $E_x \approx 24$ MeV in ^{20}Ne may be involved (1968PA12). Reaction (c) to $^{12}\text{C}^*(0, 4.4)$ has been studied for $E_\alpha = 35.5$ to 41.9 MeV. The results are qualitatively consistent with the statistical theory of compound nuclear reactions, with an average width of 1.1 MeV for the compound states (1965BR13). See also (1971GO1U). For spallation reactions see (1968JA1J, 1969JU03).

- | | | |
|--|-----------------|---------------|
| 10. $^{16}\text{O}(\alpha, n)^{19}\text{Ne}$ | $Q_m = -12.135$ | $E_b = 4.730$ |
|--|-----------------|---------------|

See (1962GO1J, 1963GO1J, 1965TS1A) and ^{19}Ne .

Table 20.20: Summary of $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ and $^{16}\text{O}(\alpha, ^8\text{Be})^{12}\text{C}$ excitation function measurements

E_α (MeV)	Detected	Refs.
0.9 – 4.0	α_0	(1953CA44)
3.7 – 6.5	α_0	(1960MC09)
5.0 – 11.0	α_0	(1972HA07)
5.8 – 10	α_0	(1967HU06) ^a
7.2 – 8.4	α_0	(1965MC02)
8.9 – 9.9	$\gamma_{6.13}$	(1964PE05)
10.0 – 19.0	$\alpha_0, \gamma_{6.13}, \gamma_{6.92+7.12}$	(1967ME10) ^a
13.0 – 17.4	^8Be	(1971GO1U)
13.5 – 30.5	α_0	(1963LU08)
14.5 – 22.7	α_0	(1962JO14)
18.9 – 30.0	α_0, α_{1+2}	(1971BE17)
20 – 24	α_0	(1968CE1B, 1969AG06, 1969FE10)
20 – 24.5	α_0	(1971TA05)
25 – 32	α_0	(1969CO19, 1970CO13)
35.5 – 41.9	^8Be	(1965BR13)
37 – 51	α_0	(1965VA11)

^a See also (JO68C, 1968ME1H).

$$11. \ ^{16}\text{O}(\alpha, p)^{19}\text{F} \qquad Q_m = -8.115 \qquad E_b = 4.730$$

See (1961YA02).

$$12. \ ^{16}\text{O}(\alpha, pn)^{18}\text{F} \qquad Q_m = -18.546 \qquad E_b = 4.730$$

See (1961FU01).

$$13. \ ^{16}\text{O}(^6\text{Li}, d)^{20}\text{Ne} \qquad Q_m = 3.257$$

Deuteron groups have been observed to all ^{20}Ne states below 7.2 MeV (1968HE08, 1968RO1K,

Table 20.21: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%)	E_x (MeV)	J^π	Refs.
2.490 ± 10	19	α_0	22	6.721	0^+	(1953CA44)
3.045 ± 10	8	α_0	36	7.165	3^-	(1953CA44)
3.090 ± 10	4	α_0	1.1	7.201	0^+	(1953CA44)
3.380 ± 10	8	α_0	4.7	7.433	2^+	(1953CA44)
3.885 ± 10	2	α_0	0.6	7.837	2^+	(1953CA44)
≈ 4.9	> 800	α_0	≈ 70	≈ 8.6	0^+	(1960MC09)
5.002	2.5	α_0	0.23	8.730	1^-	(1960MC09)
5.058 ± 3	$(110 \pm 25) \times 10^{-3}$	α_0	8.5 ± 1.5	8.775	6^+	(1972HA07)
≈ 5.1	> 800	α_0	≈ 95	≈ 8.8	2^+	(1960MC09)
5.11	< 1	α_0		8.82	(5^-)	(1960MC09)
5.152 ± 5	19	α_0	1.1	8.850	1^-	(1960MC09, 1969JO18)
5.395 ± 5	3	α_0	3.9	9.044	4^+	(1960MC09, 1969JO18)
5.486 ± 5	3.2	α_0	0.49	9.117	3^-	(1960MC09, 1969JO18)
5.955 ± 10	24	α_0	1.4	9.492	2^+	(1960MC09, 1967HU06, 1969JO18)
6.569 ± 10	97	α_0	17	9.983	4^+	(1967HU06, 1969JO18)
6.912 ± 5	141	α_0	66	10.257	5^-	(1967HU06, 1969JO18)
7.092 ± 5	81	α_0	4.8	10.401	3^-	(1967HU06, 1969JO18)
7.276 ± 5	16	α_0	1.8	10.548	4^+	(1969JO18)
7.314 ± 10	24	α_0	0.85	10.579	2^+	(1965MC02, 1967HU06, 1969JO18)
7.580 ± 100	349	α_0	33	10.79	4^+	(1967HU06, 1969JO18)
7.635 ± 5	13	α_0	0.42	10.836	2^+	(1965MC02, 1967HU06, 1969JO18)
7.636	45	α_0	2.1	10.836	3^-	(1969JO18)
(7.75)	80	α_0		(10.93)		(1967HU06)
7.80 ± 150	576	α_0	14	10.97	0^+	(1969JO18)

Table 20.21: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%)	E_x (MeV)	J^π	Refs.
7.860 ± 10	24	α_0	2.0	11.015	4^+	(1965MC02, 1967HU06, 1969JO18)
8.132 ± 30	172	α_0	4.2	11.233	1^-	(1969JO18)
8.246 ± 10	53	α_0	1.5	11.324	2^+	(1965MC02, 1967HU06, 1969JO18)
(≈ 8.6)	≈ 500	α_0		(≈ 11.6)	(2^+)	(1967HU06)
8.930 ± 20	46	α_0	1.1	11.871	2^+	(1969JO18)
8.997 ± 5	0.44 ± 0.15	$\alpha_0, \gamma_{6.13}$	0.04 ± 0.01	11.925	4^+	(1972HA07)
9.026 ± 5	$(35 \pm 10) \times 10^{-3}$	α_0	1.0 ± 0.3	11.948	8^+	(1972HA07)
9.033 ± 10	24	α_0	1.4	11.953	1^-	(1969JO18)
9.055 ± 8	29 ± 8	α_0		11.971	1^-	(1967HU06, 1972HA07)
(9.25 ± 40) ^b		α		(12.13)		(1964PE05)
9.365 ± 20	142	α_0	6.6	12.219	4^+	(1964PE05, 1967HU06, 1969JO18)
9.530 ± 100		α_0		12.35	2^+	(1969JO18)
9.550 ± 10 ^{b,c}	46 ± 16	α_0	1.2	12.367	3^-	(1964PE05, 1969JO18)
9.605 ± 5	≤ 8	α_0	< 0.15	12.411	0^+	(1969JO18)
9.790 ± 10	101	α_0	23	12.559	6^+	(1967HU06, 1969JO18)
9.860 ± 100		α_0		12.61		(1969JO18)
9.944 ± 15	97	α_0	7.3	12.682	5^-	(1969JO18)
10.050 ± 100 ^d	100	α_0		12.77	4^+	(1967ME10, 1969JO18)
10.14 ± 70	55	$\alpha_0, \gamma_{6.13}$		12.84		(1967ME10)
10.32 ± 75	60	$\alpha_0, \gamma_{6.13}$		12.98	(4^+)	(1967ME10)
10.43 ± 90	70	$\alpha_0, \gamma_{6.13}$		13.07	(4^+)	(1967ME10)
10.57 ± 75	60	$\alpha_0, \gamma_{6.13}$		13.18	(4^+)	(1967ME10)
10.759 ± 6	$(80 \pm 30) \times 10^{-3}$	α_0		13.333	7^-	(1972HA07)
10.770 ^e	20	$\alpha_0, \gamma_{6.13}$		13.342	4^+	(1967ME10, 1972HA07)

Table 20.21: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%)	E_x (MeV)	J^π	Refs.
10.83 \pm 50	40	$\gamma_{6.13}$		13.39		(1967ME10)
10.87 \pm 140	110	$\alpha_0, \gamma_{6.13}$		13.42	(4 ⁺)	(1967ME10)
11.20 \pm 400	320	$\alpha_0, \gamma_{6.13}$		13.7	(3, 7) ⁻	(1967ME10)
11.51 \pm 125	400	$\alpha_0, \gamma_{6.13}$		13.93	(6 ⁺)	(1967ME10)
11.77		$\alpha_0, \gamma_{6.9+7.1}$		14.14		(1967ME10)
11.97 \pm 300	240	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		14.3	6 ⁺	(1967ME10)
(12.06)		$\alpha_0, \gamma_{6.9+7.1}$		(14.37)		(1967ME10)
12.31 \pm 300	240	$\alpha_0, \gamma_{6.9+7.1}$		14.6	(4 ⁺)	(1967ME10)
12.66 \pm 150	120	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		14.85		(1967ME10)
12.86 \pm 150	120	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.01		(1967ME10)
13.165 \pm 150	120	$\alpha_0, \gamma_{6.13}$		15.26		(1967ME10)
13.22		α_0		15.30		(1967ME10)
13.37 \pm 470	380	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.4	7 ⁻	(1967ME10)
13.58		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.59		(1967ME10)
13.73		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.71		(1967ME10)
14.05		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.96		(1967ME10)
14.26		$\gamma_{6.13}, \gamma_{6.9+7.1}$		16.13		(1967ME10)
14.40		$\gamma_{6.13}$		16.24		(1967ME10)
14.53		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		(16.35)		(1967ME10)
14.69		$\alpha_0, \gamma_{6.9+7.1}$		16.48		(1967ME10)
14.782		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		16.58		(1967ME10)
(14.91)		α_0		(16.65)	(1 ⁻)	(1967ME10)
15.18 \pm 400	320	$\alpha_0, \gamma_{6.9+7.1}$		16.9	6 ⁺	(1967ME10)
15.673 \pm 340	270	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		17.3	2 ⁺	(1967ME10)

Table 20.21: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%)	E_x (MeV)	J^π	Refs.
16.01 \pm 230	180	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		17.3		(1967ME10)
16.30 \pm 230	180	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		17.76		(1967ME10)
(16.3)		α_0		(17.95)		(1967ME10)
16.70 \pm 180	140	$\alpha_0, \gamma_{6.13}$		18.08	6 ⁺	(1967ME10)
16.98 \pm 300	240	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		18.31	(6 ⁺)	(1967ME10)
17.45	600	$\alpha_0, \gamma_{6.13}$		18.7	(6 ⁺)	(1967ME10)
18.05 \pm 250	200	$\alpha_0, \gamma_{6.9+7.1}$		19.16	(6 ⁺)	(1967ME10)
18.35 \pm 350	280	α_0		19.40	6 ⁺	(1967ME10)
18.90 \pm 350	280	α_0		19.84	6 ⁺	(1967ME10)
19.30 \pm 120	250	α_0		20.16	7 ⁻	(1971BE17)
19.6 \pm 180	360	α_0		20.4	6 ⁺	(1971BE17)
19.6 \pm 100	200	α_0		20.4	7 ⁻	(1971BE17)
19.95 \pm 60	120	α_0		20.68	9 ⁻	(1971BE17)
20.18		α_0		20.9		(1971BE17)
20.4 \pm 100	200	α_0		21.0	7 ⁻	(1971BE17)
20.45 \pm 40	80	α_0		21.08	9 ⁻	(1971BE17)
20.70	300	α_0		21.3	7 ⁻	(1962JO14, 1971BE17, 1971TA05)
21.3 \pm 200	300	α_0		21.8	7 ⁻	(1971BE17, 1971TA05)
22.0 \pm 200	500	α_0		22.3	7 ⁻	(1971BE17, 1971TA05)
22.5 \pm 250	500	α_0		22.7	9 ⁻	(1971BE17)
22.65 \pm 125	250	α_0		22.84	9 ⁻	(1971BE17)
23.3 \pm 250	500	α_0		23.4	8 ⁺	(1971BE17, 1971TA05)
24.24 \pm 150	350	α_0		24.11	8 ⁺	(1971BE17, 1971TA05)
25.4 \pm 300	600	α_0		25.0	8 ⁺	(1971BE17)

Table 20.21: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%)	E_x (MeV)	J^π	Refs.
26.2 ± 200	400	α_0		25.7		(1971BE17)
28.1 ± 350	700	α_0		27.2		(1971BE17)
29.	1600	α_0		28.	8^+	(1969CO19, 1970CO13)
29.4 ± 350	700	α_0		28.2		(1971BE17)

^a See also (1959AJ76).

^b $^{20}\text{Ne}^*(12.11, 12.37)$ decays by α_2 to $^{16}\text{O}^*(6.13)$ with $\omega\Gamma_\alpha\Gamma'_\alpha/\Gamma = 100 \pm 50$ eV and $\omega\Gamma_\alpha\Gamma'_\alpha/\Gamma = 3 \pm 1$ keV, respectively (1964PE05).

^c $\omega\gamma \lesssim 1$ and $\lesssim 4$ eV to $^{20}\text{Ne}^*(0, 1.63)$, respectively; $\omega\Gamma_\alpha\Gamma'_\alpha/\Gamma = 3 \pm 1$ keV (1964PE05).

^d Values quoted are taken preferentially from the elastic scattering results (1967ME10).

^e $\theta_0^2 = 0.7 \pm 0.2$, $\theta_1^2 = 6 \pm 3$ % (1972HA07).

1969GO18). (1969BE1X) report the strong excitation of $^{20}\text{Ne}^*(8.78, 10.30, 12.64, 13.96, 14.35, 15.34)$ while $^{20}\text{Ne}^*(11.95)$, the 8^+ member of the ground state rotational band is comparatively weakly populated. Angular distributions have been measured at $E(^6\text{Li}) = 5.50, 5.70$ and 6.22 MeV (1968GR22; α_0, α_1 —the latter is isotropic), 20.0 MeV (1968HE08: all states below $E_x = 7$ MeV), 25.8 MeV (1969GO18, 1970OG1A: $E_x = 0, 1.6, 4.3, 5.8, 7.2, 8.8, 10.3, 12.7, 15.6$). The reaction proceeds to a large extent by compound nucleus formation. Even at $E(^6\text{Li}) = 34$ MeV, a large compound nuclear contribution is reported (1971HA1D). This is consistent also with $d_1 - \gamma$ angular correlation work at $E(^6\text{Li}) = 15$ MeV (1971BA24).


The lifetime of $^{20}\text{Ne}^*(1.63)$, τ_m , is 1.27 ± 0.24 psec, as determined in this reaction and in reaction 14 (1969TH01). See also (1968OG1A) and (1965ZE1B, 1969CH1K, 1969CH1Q, 1969JA08, 1969SM1A, 1970DO07; theor.).



At $E(^7\text{Li}) = 16$ MeV, this reaction excites only those rotational states which are allowed by the selection rules in the SU_3 scheme. The reaction appears to proceed mainly by a direct stripping mechanism. Relative maximum cross sections for formation of $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.62, 5.79)$ are 100, 272, 171, 39, 82 while the unresolved $^{20}\text{Ne}^*(7.17, 7.20)$ states are formed with a relative cross section of 486 (1968MI01). Angular distributions are also reported at $E(^7\text{Li}) = 15.0$ MeV (1970MI1E, 1970NE18; $t_0 \rightarrow t_2$), 20.0 MeV (1970NE18; t_2) and 30.3 MeV (1969GO18, 1970OG1A: same states as in reaction 13). Angular correlation work [$t_1 - \gamma$ and $t_2 - \gamma$ at $E(^7\text{Li}) = 13, 14$ MeV] shows some indication of competition between an α -particle transfer process and either multistep or compound nuclear processes (1971BA24). See also (1968OG1A, 1971HA1D) and (1969DA14, 1970DO07, 1971AR1R; theor.).



The lifetime of $^{20}\text{Ne}^*(1.63)$, τ_m , is ≈ 1 psec: see Table 20.18 (1969NI09).



See (1971BA33) and ^{24}Mg in (1973ENVA) for reaction (a). See (1971VA1H) for reaction (b).




Table 20.22: States of ^{20}Ne from $^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne}$ ^a

E_x (MeV \pm keV)	L	$J^\pi; T$	E_x (MeV \pm keV)	L	$J^\pi; T$
0	0	0^+	11.27 ± 50		
1.63 ± 160	2	2^+	11.59 ± 40		
4.22 ± 150	4	4^+	12.20 ± 30	2	2^+
4.96 ± 150			12.41 ± 30	0	0^+
5.73 ± 120			12.83 ± 30		
6.72 ± 100			13.10 ± 30	0	0^+
7.86 ± 100			13.34 ± 30		
8.79 ± 60			13.48 ± 30		
9.05 ± 60			13.63 ± 30		
9.98 ± 50			13.88 ± 30		
10.25 ± 50	2	$2^+; (1)$	14.22 ± 30		
10.88 ± 50			16.730 ± 6 ^b	0	$0^+; 2$

^a (1970GU08). See also (1970TA08).

^b $\Gamma < 20$ keV. This state is reported by (1969AD02).

The ground state excitation function has been measured for $E(^3\text{He}) = 7.0$ to 10.0 MeV: it shows a resonance corresponding to $^{20}\text{Ne}^*(28.)$. This resonance is also observed in the $^{16}\text{O}(\alpha, \alpha)$ elastic scattering. It is interpreted in terms of a quasi-molecular α -particle cluster model (1969CO19). See also (1965WA1D).

$$18. \ ^{17}\text{O}(\alpha, n)^{20}\text{Ne} \quad Q_m = 0.588$$

Angular distributions have been measured at $E_\alpha = 9.8$ to 12.3 MeV by (1967HA14; n_1, n_2, n_{4+5}). See (1970CL1C) for astrophysical implications.

$$19. \ ^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne} \quad Q_m = 13.119$$

Neutron groups have been observed to a number of ^{20}Ne states: see Table 20.22 (1969AD02, 1970GU08). In particular the first $T = 1$ state in ^{20}Ne is reported at $E_x = 10.25 \pm 0.05$ MeV ($J^\pi = 2^+$) (1970GU08) and the first $T = 2$ state is found at $E_x = 16.730 \pm 0.006$ MeV ($J^\pi = 0^+$) (1969AD02). See also (1969BA1Z).

Table 20.23: Resonances in $^{19}\text{F}(p, \gamma)^{20}\text{Ne}$ ^a

E_p (keV)	Γ_{lab} (keV)	Γ_{γ_0} (eV)	Γ_{γ_1} (eV)	$^{20}\text{Ne}^*$ (MeV)	Refs.
340		< 0.07	0.28 ± 0.06	13.167	(1962KE03)
484		≈ 0.05	0.42	13.304	(1963BE19)
597		< 0.6	12	13.411	(1963BE19)
668	7.5	1.0×10^{-2}	2.2^e	13.479	(1954SI07, 1955FA1A, 1959KU79, 1960KA18 ^f , 1961ET01, 1961GO21)
874				13.674	(1955FA1A)
935				13.732	(1955FA1A)
980				13.775	(1955FA1A)
1091	≈ 1			13.881	(1954SI07, 1955FA1A, 1961ET01, 1961GO21)
1280				14.060	(1955FA1A)
1320 ^b	4.0			14.098	(1954SI07, 1955FA1A)
1350				14.126	(1955FA1A)
1370				14.145	(1955FA1A)
1420	15.7			14.198	(1954SI07, 1955FA1A, 1961ET01)
4090 ± 5^c		$\Gamma_\gamma \approx 5 \text{ eV}$		16.728	(1967KU06)
5878 ± 5^d	11 ± 3	$\Gamma_\gamma \approx 0.3 \text{ eV}$		18.426	(1968LA1H)

^a See also Table 20.17.

^b See also (1961ET01).

^c $J^\pi = 0^+$; $T = 2$.

^d $J^\pi = 2^+$; $T = 2$.

^e Γ_γ to $^{20}\text{Ne}^*(4.97) = 0.12 \text{ eV}$ (1961GO21), 0.24 eV (1960KA18).

^f Assignment to this reaction probable but not certain.

Angular distributions have been measured at $E(^3\text{He}) = 2.80$ to 4.64 MeV (1971DI12), 3.1 MeV (1970GU08), 4 MeV (1970TA08) and 5.70 and 7.33 MeV (1969AD02). See also (1964DI1C, 1967AD01, 1967BE1W) and (1969TO1E; theor.).



Over the range $E_p = 2.9$ to 12.8 MeV, the γ_0 and γ_1 yields are dominated by the E1 giant resonance ($\Gamma \approx 6$ MeV) with the γ_1 giant resonance displaced upward in energy. Strong, well correlated structure is observed with a characteristic $\Gamma \approx 175$ keV. Angular distributions taken over the energy range do not vary greatly with energy. They are incompatible with γ_0 and γ_1 coming from the same levels in ^{20}Ne (1967SE02). See also (1959GE34, 1960BR35, 1964TA05, 1965TA1E, 1966ME1H, 1966PA1K).

The yield curve for 11.2 MeV γ -rays [from the decay of $^{20}\text{Ne}^*(11.23)$, $J^\pi = 1^+$; $T = 1$, to the ground state] displays a resonance at $E_p = 4.090 \pm 0.005$ MeV [$^{20}\text{Ne}^*(16.73)$]. The 11.2 MeV γ -rays are isotropic which is consistent with the presumed 0^+ character of this lowest $T = 2$ state in ^{20}Ne : $\Gamma_p\Gamma_\gamma/\Gamma \approx 0.5$ eV. Since Γ_p/Γ (from the elastic scattering) is ≈ 0.1 , $\Gamma_\gamma \approx 5$ eV (1967KU06). For $E_p = 5.65$ to 6.21 MeV, the $\gamma_0, \gamma_1, \gamma_2, \gamma_3, \gamma_4$ and γ_5 yields are not resonant but the yield of 10.6 MeV γ -rays is resonant at 5.878 ± 0.005 MeV [$\Gamma_{\text{lab}} = 11 \pm 3$ keV; $\Gamma_p\Gamma_\gamma/\Gamma \approx 0.055$ eV; $\Gamma_\gamma \approx 0.3$ eV]. The 10.6 MeV γ -ray is due to the cascade decay of $^{20}\text{Ne}^*(18.43)$, $J^\pi = 2^+$; $T = 2$, via $^{20}\text{Ne}^*(12.25)$, to the 2^+ state at 1.63 MeV (1968LA1H). See also (1968HA1T, 1968HA1U, 1969HA1Y).

These and other resonances observed at lower energies are displayed in Table 20.23 (1954SI07, 1955FA1A, 1960KA18, 1961ET01, 1961GO21, 1962KE03, 1963BE19). $^{20}\text{Ne}^*(13.48)$ ($E_p = 0.67$ MeV) decays predominantly to $^{20}\text{Ne}^*(1.63)$. [See Table 20.17 for branching ratios.] The radiation is M1 and the angular distribution of the 11.9 MeV γ -rays is approx. isotropic: $J^\pi = 1^+$ (1955FA1A, 1960KA18, 1961GO21). $^{20}\text{Ne}^*(13.88)$ [$E_p = 1.09$ MeV] decays predominantly to $^{20}\text{Ne}^*(4.97)$ (1961GO21). See also (1959AJ76) and (1969HO1W; theor.). For astrophysical considerations see (1969BA71).



The elastic scattering has been studied in the range $E_p = 500$ to 2000 keV by (1954DE1A, 1954PE1A, 1955BA1C, 1955WE44, 1956DE33, 1963BE19). Parameters for the observed resonances are exhibited in Tables 20.24 and 20.25 taken mainly from (1955BA1C). Some unresolved structure is observed at $E_p = 900, 1092$ and 1137 keV, in addition to a broad structure near $E_p = 1700$ keV (1955WE44). A sharp anomaly is observed in the elastic scattering at $E_p = 4.096 \pm 0.003$ MeV (1967BL19), 4.090 ± 0.005 MeV (1967KU06). It is an s-wave resonance corresponding to the 0^+ ; $T = 2$ state at $E_x = 16.73$ MeV (1967BL19, 1967KU06). There is

Table 20.24: Levels of ^{20}Ne from $^{19}\text{F}(p, p_0)^{19}\text{F}$ (1955BA1C)

E_p (keV)	Γ_{lab} (keV)	l	J^π	Γ_p/Γ	θ_p^2 (%)	$^{20}\text{Ne}^*$ (MeV)
340	2.9	0	1^+	0.016	3.8	13.167
483 ^b			1^+			13.303
598	37	1	2^-	0.0012	0.38	13.412
669	7.5	0	1^+	0.98	9.6	13.480
843	23	0	0^+	0.996	10.8	13.645
873	5.2	1	2^- ^a	0.21	1.5	13.674
935	8.0	0	1^+	0.17	0.44	13.732
1346	4.5	1	2^- ^a	0.067	0.07	14.123
1372	15	1	2^- ^a	0.17	0.52	14.147
1422	14.6	0	1^+	0.85	0.92	14.195
1694 ^c						14.453
1940 ^c		(0)	($0^+, 1^+$)			14.687
2030 ^c						14.772
4094 ^d	2.1 ± 0.5	0	0^+	0.062 ± 0.004		16.732
5880 ^e	11 ± 3	2	2^+	≈ 0.2		18.427

^a 1^- not excluded by elastic scattering alone.

^b (1963BE19).

^c (1956DE33).

^d (1967BL19, 1967KU06); $T = 2$.

^e (1968LA1H); $T = 2$.

Table 20.25: Resonance parameters in $^{19}\text{F} + \text{p}$ ^c (1955BA1C)

E_p (keV)	J^π	θ^2 ^a (%)					
		p_0	p_1	p_2	α_1	α_2	α_3
340	1^+	3.8	< 15		18.8	1.0	7.2
598	2^-	0.38	< 28	< 145	31	< 0.5	< 5.1
669	1^+	9.6	0.6	< 0.4	0.26	0.005	0.27
843	0^+	10.8	≈ 0.14	< 0.92			
873	2^-	1.5	< 0.07	2.7	1.05 ^b	1.45 ^b	3.4
935	1^+	0.44	5.0	< 0.8	3.3	0.34	2.3
1260	3^+						
1346	2^-	0.07 ^b	0.92	0.24	0.36	0.21 ^b	2.1
1372	2^-	0.52 ^b	1.93	0.56	1.7 ^b	0.34 ^b	0.86
1422	1^+	0.92	0.56	< 0.11	total < 0.034		

^a p_0, p_1, p_2 represent transitions to $^{19}\text{F}(0), (0.1), (0.2)$. $\alpha_1, \alpha_2, \alpha_3$ represent transitions to $^{16}\text{O}(6.1), (6.9), (7.1)$.

^b Assuming lowest possible values of l ; see (1957MA1A).

^c See also (1958IS11).

no indication of this resonance in the p_3, p_4 or p_5 yields (1967KU06). The amplitude of the $T = 1$ or $T = 0$ impurity in this state is $\approx 1.5\%$ (1967BL19). In the range $E_p = 5.65$ to 6.21 MeV, a single anomaly is seen in the elastic scattering at $E_p = 5.880 \pm 0.005$ MeV. The interference patterns show that the scattering is d-wave, corresponding to the excitation of the $J^\pi = 2^+; T = 2$ state at $E_x = 18.43$ MeV (1968LA1H). The parameters of these two $T = 2$ states are shown in Table 20.24. The elastic scattering has also been studied for $E_p = 4.2$ to 7.5 MeV by (1967TH06), and the polarization has been determined at $E_p = 10.2$ MeV by (1961RO05). The total reaction cross section is reported for $E_p = 24.6$ to 46.0 MeV by (1969MC1A).

Resonances for inelastic scattering involving $^{19}\text{F}^*(0.11) (J^\pi = \frac{1}{2}^-)$ and $^{19}\text{F}^*(0.197) (J^\pi = \frac{5}{2}^+)$ [p_1 and p_2] are listed in Table 20.26 (1955BA94, 1963BE19). See also (1958RA15). In general the resonances observed are identical with those reported from other $^{19}\text{F} + \text{p}$ reactions, although the relative intensities differ greatly. The p_2 scattering has been measured at $E_p = 5.6$ to 6.3 MeV (1967TH06). For a spallation study, see (1969EP1B). See also (1959TR1A, 1969LE08), (1969HA1Y, 1969MC1C, 1969TE1A), (1963MI1D, 1966AM1B, 1967AF01; theor.), (1959AJ76) and ^{19}F .

$$22. \ ^{19}\text{F}(p, \alpha)^{16}\text{O}$$

$$Q_m = 8.114$$

$$E_b = 12.845$$

Table 20.26: Resonances in $^{19}\text{F}(p, p')^{19}\text{F}^*$ (1955BA94)

E_p (keV)	J^π	Γ_{lab} (keV)	Γ_{p_1} (eV)	Γ_{p_2} (eV)	$\theta_{p_1}^2$ ^a (%)	$\theta_{p_2}^2$ ^a (%)	E_x in ^{20}Ne (MeV)
340	1 ⁺	2.9	< 0.5	< 0.1	< 15		13.167
483 ^b	1 ⁺	2.2	< 1.3	< 1.2			13.303
598 ^b	2 ⁻	37	< 100	< 60	< 28	< 145	13.412
669	1 ⁺	7.5	46	< 0.5	0.6	< 0.4	13.480
720		≈ 30	< 10000	< 10000			13.528
780		≈ 10	< 400	≈ 9000			13.585
831		8.3	< 6	≈ 2300			13.634
845	0 ⁺	23	≈ 50	< 10	≈ 0.14	< 0.92	13.647
873	2 ⁻	5.2	< 2	570	< 0.07	2.7	13.674
900		4.8	< 30	≈ 2200			13.699
935	1 ⁺	8.0	3000	< 20	5.0	< 0.8	13.732
1092		< 1.2					13.881
1137		3.7	< 40	≈ 2100			13.924
≈ 1250		≈ 80	≈ 70000	< 4000			14.03
1290		19	< 600	≈ 900			14.069
1346	2 ⁻	4.5	300	600	0.92	0.24	14.123
1372	2 ⁻	15	700	1400	1.93	0.56	14.147
1422	1 ⁺	14.6 ± 1	2200	≤ 35	0.56	≤ 0.11	14.195
1610		≈ 5					14.373
1660							14.421
1700							14.459

^a (1955BA1C).

^b (1963BE19).

Table 20.27: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(p, \alpha_0)^{16}\text{O}$

E_p (keV)	Γ_{lab} (keV)	θ_{α}^2 ^a (%)	$J\pi$	$^{20}\text{Ne}^*$ (MeV)
400 ^b	100		1^-	13.224
400 ^b	100		0^+	13.224
650 ± 20 ^b	200		1^-	13.462
710 ^{a,b}	35	0.6	(1^-)	13.519
733	66	1.0	2^+	13.541
778	≈ 10	0.02	2^+	13.583
843	23	0.16	2^+ ^g	13.645
≈ 860	120	2.1	1^-	13.66
≈ 930	≈ 180	2.9	0^+	13.73
≈ 1080	≈ 200	3.4	1^-	13.87
1115	50	0.55	2^+	13.903
1160	≈ 70	1.1	0^+	13.946
1235 ^{a,c}	≈ 70	1.2	1^-	14.017
≈ 1250 ^a	≈ 150	2.7	2^+	14.03
1358 ^{a,c,d}	54	0.49	2^+	14.134
1640 ^c	< 115			14.402
1709 ^{c,d}	140		0^+	14.467
1853 ^{c,d}	132		1^-	14.604
2110 ^{c,d,e}	75		$(2^+, 4^+)$	14.85
2310 ^{c,d,e}	90		(2^+)	15.04
2550 ^e	300		(1^-)	15.26
2590 ^{c,d,h}	300		(0^+)	15.30
2680 ^{c,h}	80			15.39
2730 ^e	60			15.44
2820 ^e	160			15.52
2940 ^h				(15.64)
3120 ^h	170			(15.81)
3340	105			16.02
3680	(100)			16.34
3860				16.51
3980	135			16.62

Table 20.27: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(\text{p}, \alpha_0)^{16}\text{O}$ (continued)

E_p (keV)	Γ_{lab} (keV)	θ_{α}^2 ^a (%)	J^{π}	$^{20}\text{Ne}^*$ (MeV)
4090 ⁱ			$0^+; T = 2$	16.73
4130	100			16.77
4360	100			16.98
4460	95			17.08
4690	65			17.30
4900	90			17.50
4990	40			17.58
5880 ± 5 ^f	11 ± 3		$2^+; T = 2$	18.427

^a (1958IS10, 1958IS11).

^b (1959BR67).

^c (1958RA15).

^d (1957CL42).

^e (1964BR12).

^f (1968LA1H). ^g $J = 0$ from $^{19}\text{F}(\text{p}, \text{p})^{19}\text{F}$; possibly $T = 1$ (1955BA1C, 1955BA94).

^h See, however, (1964BR12).

ⁱ (1967KU06). See [blue Note in Erratum](#).

For $E_p \approx 1$ to 3 MeV, five α -particle groups are reported. All show resonance effects with relative intensities varying greatly with bombarding energy. The long range group (α_0) leaves ^{16}O in the ground state ($J^{\pi} = 0^+$); the next longest (α_{π}) results in the formation of the $J^{\pi} = 0^+$ nuclear pair-emitting state at 6.05 MeV, while the three remaining groups ($\alpha_1, \alpha_2, \alpha_3$) lead to γ -ray emitting states at 6.13 ($J^{\pi} = 3^-$), 6.92 ($J^{\pi} = 2^+$) and 7.12 MeV ($J^{\pi} = 1^-$). At $E_p > 3$ MeV, excitation of higher ^{16}O levels occurs: see ^{16}O . Resonances for α_0 and α_{π} (Tables 20.27 and 20.28) are generally identical and different from those for $\alpha_1, \alpha_2, \alpha_3$ (Table 20.29). The resonances for α_0 and α_{π} are required to have even J , even π of odd J , odd π while the $\alpha_1, \alpha_2, \alpha_3$ resonances, insofar as their assignments are known, are all odd-even or even-odd.

Recent studies of the α_0 yield and of angular distributions have been carried out by (1959BR67: $E_p = 0.40$ to 0.72 MeV), (1964BR12: $E_p = 2.0$ to 3.3 MeV), (1966MO25: $E_p = 2.24$ to 3.35 MeV), (1963WA12: $E_p = 3.3$ to 12.2 MeV), (1965WA08: $E_p = 4.0$ to 12.0 MeV), (1960BR35: $E_p = 4.2$ to 8.8 MeV), (1968LA1H: $E_p \approx 5.9$ MeV), and (1959OG15: $E_p = 12.9$ to 14.1 MeV). See also (1969LE08). Resonances observed in the lower energy range are displayed in Table 20.27 (1957CL42, 1958IS10, 1958IS11, 1958RA15, 1959BR67, 1964BR12, 1968LA1H). Among these is a resonance at $E_p = 5.880 \pm 0.005$ MeV, $\Gamma_{\text{lab}} = 11 \pm 3$ keV which corresponds to the $J^{\pi} = 2^+; T = 2$ state at $E_x = 18.43$ MeV: the α_1 and α_2 channels also display it (1968LA1H). See also (1968HA1T, 1969HA1Y, 1969MC1C). (1965WA08) find that over the range $E_p = 4.0$ to 12.0

Table 20.28: Nuclear pair resonances (α_π) in $^{19}\text{F}(\text{p}, \alpha_\pi)^{16}\text{O}$

E_p (keV)	Γ_{lab} (keV)	σ^c (mb)	θ_α^2 (%)	J^π	$^{20}\text{Ne}^*$ (MeV)
710 ^a	35	≈ 0.2	2	1^-	13.519
780	≈ 10	≈ 0.2	0.15	2^+	13.585
842	23	3.4	0.27	2^+ ^d	13.644
1115	50	1.5	3.6	2^+	13.903
1236 ^{a,b}	≈ 70	3	1.0	1^-	14.018
1367 ^{a,b}	30	6.0	0.29	2^+	14.143
1630 ^b	60				14.39
1720	95	≈ 18			14.48
1880	170				14.63
2170	95				14.91
2330	≈ 100				15.06
2600	100				15.31
2680	100				15.39
2820	125				15.52
3120	145				15.81
3340	100				16.02
(3500)	(80)				(16.17)
(3590)	(115)				(16.25)
3960	200				16.61
4360	95				16.98
4690	< 150				17.30
4900	115				17.50
4990	40				17.58
5170	220				17.75

^a See (1950CH53, 1951PH1A, 1954DE36, 1958IS11).

^b (1958RA15).

^c (1958IS11): resonant cross sections derived from analysis: see also (1950CH53).

^d See footnote ^g in Table 20.27.

MeV, direct interaction mechanisms are predominant although compound nuclear contributions cannot be ignored at certain energies. The continuing structure in the α_0 yield at the higher energies has been interpreted by (1964TE1F) in terms of fluctuations with a coherence energy $\Gamma = 160$ keV. See also (1966MO25).

Resonances in the $^{19}\text{F}(p, \alpha_\pi)^{16}\text{O}$ yield have been investigated by (1950CH53, 1951PH1A, 1954DE36, 1958IS11, 1958RA15): see Table 20.28. Resonance locations and absolute reduced widths appear to correspond closely to those for (p, α_0) , although some exceptions occur. In the work of (1958RA15) only 6 of the 23 (p, α_0) resonances have no clear counterpart in $\sigma(p, \alpha_\pi)$. For resonances at $E_p = 1.35, 1.72, 1.88$ and 2.33 MeV, $\theta_{\alpha_0}^2 = \theta_{\alpha_\pi}^2$ within about 10%; at $E_p = 2.17$ MeV, a large difference occurs, possibly to be ascribed to superposition of several resonances (1958RA15). Below $E_p = 1.3$ MeV, several fairly large differences occur (1958IS11).

Recent yield measurements of α -groups to excited states of ^{16}O and of the de-excitation γ -rays are reported by (1965AS07: $E_p = 0.9$ to 2.6 MeV; $\alpha_1, \alpha_2, \alpha_3$), (1969OS1B: $E_p = 1.9$ to 4.2 MeV; σ_t for $6.13, 6.92, 7.12, 8.87$ γ), (1963WA12: $E_p = 10$ to 12 MeV; $\alpha_{1+2}, \alpha_{3+4}$) and (1971GO1U: $E_p = 2$ to 6.2 MeV; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4$). See also (1961AS02). Resonances are displayed in Table 20.29 (1950AR1A, 1950BA1A, 1950CH1A, 1952WI27, 1955BA94, 1955HU1A, 1955KI28, 1959BO14, 1959KU79, 1959LI51, 1960HU11, 1962KE03, 1962RY01, 1963BE19, 1964SE1A, 1965AS07, 1966MA60, 1967KU06). Anomalies are observed in the $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ and α_4 yields at $E_p = 4.09$ and 5.88 MeV, corresponding to the $T = 2$ state at $E_x = 16.73$ and 18.43 MeV. [P. Gorodetzky, private communication.] See also (1967KU06).

A detailed discussion of the evidence leading to many of the J^π assignments shown in Table 20.29 is given in (1959AJ76).

The total cross section has been measured for $E_p = 5.1$ to 6.5 MeV by (1960TE03). For spallation measurements see (1963VA1C). See also (1959TR1A, 1960GO12, 1961ET01, 1968KO05, 1970WI13, 1971GU23) and (1963MI1D; theor.). For astrophysical considerations, see (1969BA71). See also ^{16}O in (1971AJ02) and (1959AJ76).

23. $^{19}\text{F}(p, n)^{19}\text{Ne}$ $Q_m = -4.021$ $E_b = 12.845$

Yield measurements are reported for $E_p = 4.23$ to 4.93 MeV (1959GI47; σ_t), 4.23 to 6.01 MeV (1968RI08; σ_t), 4.4 to 6.1 MeV (1969BL02; γ_1, γ_2), 4.9 to 11.0 MeV (1963JE04; σ_t) and 6.2 to 6.9 MeV (1962FR09; n_0, n_{1+2}, n_3, n_4). See also (1965VA23, 1968LA1H). Observed resonances are displayed in Table 20.30 (1952WI27, 1963JE04, 1968RI08). A narrow anomaly is reported in the n_0 and n_1 yield at $E_p = 5.879 \pm 0.004$ MeV corresponding to a state at $E_x = 18.427$ MeV in ^{20}Ne , presumed to be the second $T = 2$ state in ^{20}Ne (1968AD1C). See also (1959AJ76) and ^{19}Ne .

24. $^{19}\text{F}(p, pn)^{18}\text{F}$ $Q_m = -10.431$ $E_b = 12.845$

See (1961GU1A, 1962GU10, 1963VA1C, 1965VA23).

Table 20.29: Resonances for 6 – 7 MeV γ -rays ($\alpha_1, \alpha_2, \alpha_3$) in $^{19}\text{F}(p, \alpha)^{16}\text{O}$

E_p (keV)	Γ_{lab} (keV)	Γ_{α_1} (eV)	Γ_{α_2} (eV)	Γ_{α_3} (eV)	J^π	$^{20}\text{Ne}^*$ (MeV)
226.9 ± 3.4 ^{a,b}	1.0	1000	< 2.5	< 2.5	2^-	13.060
340.46 ± 0.04 ^{c,d}	2.4 ± 0.2	2800	16	75	1^+	13.168
483.8 ± 0.3 ^e	0.9 ± 0.1	700	19	190	1^+	13.304
872.11 ± 0.20 ^{f,g}	4.7 ± 0.2	2200	620	180	2^-	13.673
935.4 ± 1.3 ^g	8.1 ± 0.5	2900	110	720	1^+	13.733
1347.7 ± 1.0 ^h	4.9 ± 0.7	2250	650	1200	2^-	14.124
1373.0 ± 1.0 ⁱ	12.4 ± 1.0	6650	700	300	2^-	14.148
1694 ± 1.7 ^j	35 ± 3					14.453
1949 ± 2.5 ^j	40 ± 10					14.695
2030 ± 3.0 ^j	120 ± 20					14.772
2320 ^k	85					15.05
2510	30					15.23
2630	90					15.34
2800	60					15.50
3020	30					15.71
3190	80					15.87
3490	40					16.16
3920	30					16.57
4000	110					16.64
4090 ^l					$0^+; T = 2$	16.73
4290	50					16.92
4490	30					17.11
4570	30					17.18
4710	30					17.32
4780	35					17.38
4990	20					17.58
5070	35					17.66
5200	70					17.78

- ^a (1959KU79; assignment to this reaction probable but not certain). See also (1959AJ76).
- ^b (1962KE03). $\Gamma_{\alpha_0} < 100$ eV.
- ^c (1955BA94, 1959BO14, 1964SE1A). See also (1959KU79, 1960HU11, 1962KE03) and (1959AJ76).
- ^d (1950AR1A, 1950BA1A, 1950CH1A). Values listed for E_p and Γ are those recommended by (1966MA60).
- ^e (1959BO14, 1959KU79, 1960HU11, 1963BE19). $\Gamma_{\alpha_0} < 25$ eV (1963BE19).
- ^f Values listed for E_p and Γ are those recommended by (1966MA60). Other values are $E_{\text{res}} = 872.4 \pm 0.4$ keV (1959BO14), 873.5 ± 0.7 keV (1960HU11), 872.3 ± 0.5 keV (1961BE13), 871.80 ± 0.25 keV (1962RY01), 872.5 ± 1.1 keV (1965AS07). See also Table 20.13 in (1959AJ76) for earlier references.
- ^g (1965AS07).
- ^h Recently reported resonance energies are $E_p = 1344.5 \pm 1.0$ keV (1959LI51), 1347.7 ± 1.0 keV (1960HU11), 1347.7 ± 1.8 keV (1965AS07). See Table 20.13 in (1959AJ76) for earlier references.
- ⁱ Recently reported resonance energies are $E_p = 1373.0 \pm 1.0$ keV (1959LI51), 1373.7 ± 1.2 keV (1960HU11), 1374.5 ± 1.8 keV (1965AS07). See Table 20.13 in (1959AJ76) for earlier references.
- ^j (1952WI27, 1955HU1A).
- ^k (1952WI27); these values should be reduced by about 0.2%: see (1955KI28).
- ^l Resonance in α_2 yield: see text (1967KU06).

25. $^{19}\text{F}(p, ^8\text{Be})^{12}\text{C}$

$$Q_m = 0.861$$

$$E_b = 12.845$$

The excitation curves show strong resonant behavior (cross sections up to 1.5 mb/sr) for $E_x = 15.3$ to 18.7 MeV, over which region 28 angular distributions have been measured. Twelve states with $J^\pi \leq 4^+$ have been observed.

The strongly populated states are in better agreement with those reported in the (p, α_1) yield to $^{16}\text{O}^*(6.05)$ [$J^\pi = 0^+$] than those reported in the (p, α_0) yield (1969GO1B, 1971GO1U). It is suggested that most of the observed states are of 8p-4h and 12p-8h configurations (1971GO1U).

26. $^{19}\text{F}(d, n)^{20}\text{Ne}$

$$Q_m = 10.621$$

Levels of ^{20}Ne derived from reported neutron groups are listed in Table 20.31 (1958MO02, 1963FE1B, 1964SA09, 1966RI05, 1968LA03, 1969RI01); those derived from threshold measurements are displayed in Table 20.32 (1960BU07). Gamma-ray measurements have been reported by (1960KR02, 1960RA23, 1965PE01) and angular correlation measurements by (1964SA09, 1966RI05). For a survey of the earlier work see (1959AJ76).

Angular distributions have been measured at $E_d = 0.5$ to 0.7 MeV (1964SA09; n to $^{20}\text{Ne}^*(10.31)$), 1.0 MeV (1966RI05; n to $^{20}\text{Ne}^*(10.31, 11.03)$), 1.0 to 1.3 MeV (1962FI02; n_0, n_1), 1.0 to 2.5 MeV (1966WA17; n_0, n_1), 2.17 MeV (1958MO02; $n_0 \rightarrow n_4$), 2.5 and 3.0 MeV (1963FE1B; n_0 ,

n_1 and n to $^{20}\text{Ne}^*(5.80, 6.75, 8.76)$), 2.51 to 3.31 MeV (1964SI09; $n_0 \rightarrow n_3, n_{4+5}, n_6$), 2.98 MeV (1968LA03: see Table 20.31), 3.06 to 6.07 MeV (1967BA40; n_0, n_1), 3.57 MeV (1961BE10; $n_0 \rightarrow n_3, n_{4+5}, n_6$) and 5.10 MeV (1969RI01: see Table 20.31). See (1959AJ76) for earlier references. See also (1963FE01, 1964AL14, 1964JO04, 1965BA1K), (1966WE1B) and (1966AR1D, 1966IN01, 1968WA1M, 1970BO1K; theor.).

Table 20.30: Resonances in $^{19}\text{F}(p, n)^{19}\text{Ne}$

E_p (MeV)		Γ_{lab} (keV)	$^{20}\text{Ne}^*$ (MeV)
(1968RI08) ^a	(1963JE04) ^b	(1952WI27)	
4.30		45	16.93
4.46		80	17.08
4.52		20	17.14
4.61		60	17.22
4.72		25	17.33
4.75		45	17.36
4.87			17.47
4.95	4.96	20	17.55
5.03	5.03		17.62
5.11	5.11		17.70
5.23			17.81
5.26	5.26 ^c		17.84
5.37	5.37		17.94
(5.44)			(18.01)
5.50			18.07
5.57			18.13
(5.62)			(18.18)
(5.69)			(18.25)
5.72	5.73		18.28
5.77			18.32
5.84			18.39
	5.879 ± 0.004 ^d	≈ 9 ^d	18.427
5.90			18.45
6.00	6.03		18.54
	6.15		18.68
	6.35 ^e		18.87

Table 20.30: Resonances in $^{19}\text{F}(p, n)^{19}\text{Ne}$ (continued)

E_p (MeV)		Γ_{lab} (keV)	$^{20}\text{Ne}^*$ (MeV)
(1968RI08) ^a	(1963JE04) ^b	(1952WI27)	
	6.53		19.04
	6.81 ^e		19.31
	7.14		19.62
	7.27		19.75
	7.41		19.88
	7.52		19.98
	7.74		20.19
	8.02		20.46
	8.15		20.58
	8.28		20.71
	8.37		20.79
	8.70		21.10
	8.82		21.22
	9.08		21.47
	9.2		21.6
	9.5		21.9
	9.8		22.1
	10.2		22.5

^a ± 5 keV.

^b ± 20 keV, except for the last four values.

^c See also (1969BL02).

^d (1968AD1C): $T = 2$.

^e See also (1962FR09).

27. $^{19}\text{F}(^3\text{He}, d)^{20}\text{Ne}$

$$Q_m = 7.351$$

At $E(^3\text{He}) = 9.5$ to 10.0 MeV, angular distributions have been measured for the deuterons corresponding to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62, 5.79, 6.72, 7.01, 7.17+7.20, 7.42)$. The distributions to $^{20}\text{Ne}^*(0, 1.63, 5.79, 6.72, 7.42)$ have been fitted with $l = 0, 2, 1, 0$ and 2 , respectively. Spectroscopic factors were also obtained. The results are consistent with the description of the states

in terms of rotational bands (1965SI18). Angular distributions are also reported at $E(^3\text{He}) = 13.0$ MeV (1963JA01; d_0, d_1).

Table 20.31: Neutron groups from $^{19}\text{F}(d, n)^{20}\text{Ne}$ ^a

E_x (MeV \pm keV)				l_p ^a	$J^\pi; T$
(1958MO02) ^b	(1963FE1B)	(1968LA03)	(1969RI01)		
0		c	c	0	0 ⁺
1.74 \pm 30				2	2 ⁺
4.20 \pm 40					
4.96 \pm 50					
5.62 \pm 40					
6.80 \pm 10		c	c	0	0 ⁺
7.16 \pm 90					
7.41 \pm 50					
7.90 \pm 40					
(8.71 \pm 10)					
9.15 \pm 40	9.17 \pm 30				
	9.37 \pm 20				
(9.50 \pm 40)	9.50 \pm 20				
	9.60 \pm 20				
	9.92 \pm 30				
10.01 \pm 30	10.00 \pm 30		c		
	10.30 \pm 30	c	d,i	h	h
			10.59 ^c		
	11.00 \pm 20	10.853 \pm 10 ^j	10.879 \pm 40	2	$T = 1$ ^f
			11.03 \pm 80 ⁱ		
	11.32 \pm 20	11.233 \pm 10	11.26 \pm 40	0	1 ⁺ ; (1) ^{e,f}
	11.66 \pm 30	11.549 \pm 10	11.568 \pm 35	2	($T = 1$) ^f
			11.915 \pm 30		
		12.086 \pm 10		g	($T = 1$) ^f
		12.150 \pm 10	12.179 \pm 25	g	($T = 0$) ^f
		12.200 \pm 10		g	($T = 1$) ^f
		12.245 \pm 10		2	$T = 1$ ^f
		12.379 \pm 10	12.397 \pm 20	0	$T = 0$ ^f

Table 20.31: Neutron groups from $^{19}\text{F}(d, n)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV)				l_p ^a	$J^\pi; T$
(1958MO02) ^b	(1963FE1B)	(1968LA03)	(1969RI01)		
			13.086 \pm 15		
			13.170 \pm 15	0	1 ⁺ ; (1)
			13.481 \pm 15	0	1 ⁺ ; 1 ^e
			13.650 \pm 15	0	(0 ⁺); 1 ^e
			13.882 \pm 15		

^a See also Table 20.16 in (1959AJ76).

^b Evidence for some other states is also reported.

^c Observed but no parameters reported.

^d $E_x = 10.31 \pm 0.07$ MeV (1964SA09), $E_x = 10.33 \pm 0.05$ MeV (1966RI05).

^e (1969RI01).

^f (1968LA03).

^g Weak group.

^h See (1964SA09, 1966RI05).

ⁱ This state decays to $^{20}\text{Ne}^*(1.63)$ (1966RI05).

^j Data of (1968LA03) are adjusted downward by 26 keV: see (1969RI01).

Gamma-ray measurements lead to $E_x = 1635.3 \pm 1.8$, 4249 ± 2.5 , 4968 ± 3 , 5623 ± 3 and 7174 ± 4 keV and to $\tau_m = 1500_{-400}^{+900}$, 150 ± 25 and 2800_{-900}^{+2400} fsec, respectively, for $^{20}\text{Ne}^*(1.63, 4.25, 4.97)$ (1969AN08, 1969AN1J; see also Table 20.18). See also (1964HE1D, 1966AR1D, 1966IN01, 1970BO1K; theor.).

$$28. \ ^{19}\text{F}(\alpha, t)^{20}\text{Ne} \quad Q_m = -6.969$$

At $E_\alpha = 28.5$ MeV, angular distributions are reported for the tritons to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62 + 5.79, 7.02 + 7.17 + 7.20)$. All are typical of stripping reactions except that for $^{20}\text{Ne}^*(4.97)$. The distributions have been analyzed by DWBA. $^{20}\text{Ne}^*(6.72)$ is very weakly populated (1967HA23). Angular distributions have also been reported at $E_\alpha = 18.5$ MeV (1963JA01; t_0, t_1) and 28.4 MeV (1965KA14; $t_0, t_1, t_2, t_3, t_{4+5}$). See also (1964EL1B; theor.).

$$29. \ ^{20}\text{F}(\beta^-)^{20}\text{Ne} \quad Q_m = 7.026$$

The decay is principally to $^{20}\text{Ne}^*(1.63)$ with a half-life of 11.03 ± 0.06 sec: see Table 20.5 for a listing of half-life measurements and Table 20.33 for the branching to various ^{20}Ne states.

Table 20.32: Levels of ^{20}Ne from $^{19}\text{F}(\text{d}, \text{n})^{20}\text{Ne}$ thresholds (1960BU07)

$E_{\text{thresh.}}$ (keV)	$^{20}\text{Ne}^*$ ^a (MeV)
510 ± 20	11.082
(600 ± 20)	(11.163)
760 ± 20	11.308
(850 ± 20)	(11.389)
1150 ± 20	11.660
1350 ± 20	11.841
1700 ± 20	12.157
1790 ± 20	12.239
2060 ± 20	12.483

^a All these states are reported to decay principally to the ground state except for $^{20}\text{Ne}^*(11.11, 11.19, 12.27)$ which decay to $^{20}\text{Ne}^*(1.63)$. See, however, Table 20.15.

The 0.02% branching to $^{20}\text{Ne}^*(4.97)$ [$J^\pi = 2^-$] is consistent with the assignment $J^\pi = 2^+$ to the ground state of ^{20}F (1969GA05), as are measurements of the $\beta - \gamma$ circularly polarized correlation (1961FR02, 1965MA28).

$E_{\beta^-}(\text{max}) = 5.419 \pm 0.013$ MeV (1954WO23), 5.416 ± 0.015 MeV (1959AL06). The energy of the subsequent γ -ray is 1634.8 ± 0.6 keV (1967VA08), 1632.6 ± 0.8 keV (1966AL12). The γ -ray from the $(4.97 \rightarrow 1.63)$ transition has $E_\gamma = 3334.3 \pm 0.7$ keV. When recoil-corrected, $\Delta E = 3334.6 \pm 0.7$ keV, and using the E_x for $^{20}\text{Ne}^*(1.63)$ shown in Table 20.15, E_x for $^{20}\text{Ne}^*(4.97) = 4968.4 \pm 0.8$ keV (1969GA05). See also (1960SC01, 1961OK1A, 1968SP01, 1971TO1K), (1959AJ76) and (1967YA04, 1968KR10, 1969BL1D, 1969SU12, 1970AN27, 1970MC23, 1970YA01, 1971LI1H, 1971WI18, 1971YA04; theor.).

$$30. \ ^{20}\text{Ne}(\gamma, \text{n})^{19}\text{Ne} \quad Q_{\text{m}} = -16.866$$

A giant resonance is observed at $E_\gamma = 21.5$ MeV, $\Gamma = 6.6$ MeV, $\sigma(\text{max}) = 7.3$ mb (1954FE16). See also (1962GO1E).

$$31. \ ^{20}\text{Ne}(\gamma, \text{pn})^{18}\text{F} \quad Q_{\text{m}} = -23.275$$

See (1959RE1A).

Table 20.33: Branching in $^{20}\text{F}(\beta^-)^{20}\text{Ne}$

Decay to $^{20}\text{Ne}^*$ (MeV)	J^π	Branch (%)	$\log ft^a$	Refs.
0	0^+	< 0.032		(1954WO23)
1.63	2^+	99.8	4.98	
4.25	4^+	< 0.015		(1969GA05)
4.97	2^-	0.017 ± 0.003	6.88	(1969GA05)
5.62	3^-	< 0.048		(1963GL01)
5.79	1^-	< 0.1		(1963GL01)
6.72	0^+	< 0.67		(1963GL01)
7.01	4^-	< 0.2		(1963GL01)

^a I am grateful to B. Zimmerman for calculating these values.

$$32. \ ^{20}\text{Ne}(\gamma, p)^{19}\text{F} \quad Q_m = -12.845$$

For cross section measurements see (1969HO16, 1963FI1B). See also (1962DO1A, 1963HA1E), (1959AJ76, 1959HA1C, 1961DO08, 1962GO1E) and (1965SH1D; theor.).

$$33. \ ^{20}\text{Ne}(\gamma, \alpha)^{16}\text{O} \quad Q_m = -4.730$$

See (1959HA1C).

$$34. \ (a) \ ^{20}\text{Ne}(e, e)^{20}\text{Ne} \\ (b) \ ^{20}\text{Ne}(e, ep)^{19}\text{F} \quad Q_m = -12.845$$

The ^{20}Ne charge radius, $r_{\text{rms}} = 3.116 \pm 0.025$ fm (using a Born approx.) (1971MO15).

At $E_e = 39$ and 56 MeV, the 180° inelastic scattering is dominated by the transition to a $J^\pi = 1^+$; $T = 1$ state at $E_x = 11.22 \pm 0.05$ MeV with $\Gamma_{\gamma_0} = 11.2_{-1.8}^{+2.1}$ eV. A subsidiary peak is observed corresponding to $E_x = 11.58 \pm 0.03$ MeV [if $J^\pi = 1^+$ or 2^+ , $\Gamma_{\gamma_0} = 0.65 \pm 0.18$ or 0.40 ± 0.13 eV]. A number of small peaks are also reported corresponding to $E_x \approx 12.0, 12.9, 13.9, 15.8, 16.9, 18.0$ and 19.0 MeV (1971BE18). See also (1963BA19, 1963GO04) and (1971FO19, 1971HO20; theor.). A study of reaction (b) at $E_e = 30$ MeV shows strong resonances (assuming ground state transitions) at $E_x = 17.70, 18.87, 19.87$ and 21.02 MeV [$\Gamma = 0.50, 0.58, 0.54$ and

0.49 MeV, respectively], as well as some weaker structures (1962DO1A). See also (1961DO08) and (1970PA1H; theor.).

35. $^{20}\text{Ne}(n, n)^{20}\text{Ne}$

See ^{21}Ne in (1973ENVA) and (1959CO82).

36. (a) $^{20}\text{Ne}(p, p)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(p, p\alpha)^{16}\text{O}$ $Q_m = -4.730$

Angular distributions have been measured at $E_p = 2.15$ and 2.72 MeV (1961SO02; p_1, γ_1), 3.65 to 4.35 MeV (1963HU1D; p_0, p_1), 4.95 to 5.50 MeV (1959OD08; p_0, p_1), 5.20 to 6.23 MeV (1962HU12; p_0, p_1), 6.86 to 7.15 MeV (1969MC14; p_0), 7.75 to 14.2 MeV (1960OD01; p_0, p_1), 17.4 MeV (1962SC12; p_1, p_2, p_3 , and p to states with $E_x = 5.63 \pm 0.07, 7.45 \pm 0.08, 7.85 \pm 0.08, 9.20 \pm 0.09, 10.0 \pm 0.1$ MeV), 18.03 MeV (1969LE1P; p_0), 24.5 MeV (1969DE15; p_0, p_1, p_2, p to $^{20}\text{Ne}^*(8.78)$) and 41.8 MeV (1970FA17; p_0). A large hexadecapole deformation, β_4 , is needed to reproduce both the shapes and the intensities of the angular distributions to the $2^+, 4^+$ and 6^+ members of the ground state rotational band [$^{20}\text{Ne}^*(1.63, 4.25, 8.78)$] (1969DE15). Polarization measurements have been carried out at $E_p = 3.8$ to 5.1 MeV (1967BE25; p_0), 8.0 MeV (1961RO13; p_0), 14.5 MeV (1966RO1R; p_0) and 24.5 MeV (1970BA2F, 1971BA1T; p_0, p_1, p_2, p_3).

37. $^{20}\text{Ne}(d, d)^{20}\text{Ne}$

Angular distributions are reported at $E_d = 10.95$ MeV (1960TA08; d_0), 11.6 MeV (1965JA13; d_0, d_1, d_2, d_{4+5}), 11.8 MeV (1966JA1J; d_0, d_1) and 52 MeV (1968HI09, 1968HI14; d_0, d_1, d_2, d_{4+5} , and d to $^{20}\text{Ne}^*(7.2, 10.5)$). The second excited state has $E_x = 4.250 \pm 0.008$ MeV (1960FR04). See also (1960EL09, 1966WE04, 1968ME1E, 1971DU12; theor.).

38. $^{20}\text{Ne}(t, t)^{20}\text{Ne}$

The elastic scattering has been studied at $E_t = 1.80$ and 2.00 MeV (1969HE08).

39. $^{20}\text{Ne}(^3\text{He}, ^3\text{He})^{20}\text{Ne}$

Angular distributions of elastically scattered ^3He particles have been measured at $E(^3\text{He}) = 10$ and 15 MeV (1969BA62), 15 MeV (1969ZU02), 17.83 MeV (1971KE11), 28 MeV (1960CA1C) and 35 MeV (1969AR08, 1969AR10). See also (1960CA10, 1962AG01). At $E(^3\text{He}) = 17.83$ MeV, angular distributions of the ^3He 's corresponding to $^{20}\text{Ne}^*(1.63, 4.25)$ [$J^\pi = 2^+$ and 4^+ , respectively] have been analyzed by a coupled channels calculation. For $^{20}\text{Ne}^*(1.63)$, β_2 is in the range 0.42 to 0.48; for $^{20}\text{Ne}^*(4.25)$, $\beta_4 < 0.05$ (1971KE11). See also (1964VE1A, 1968HO1C; theor.).

40. (a) $^{20}\text{Ne}(\alpha, \alpha)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(\alpha, ^{12}\text{C})^{12}\text{C}$ $Q_m = -4.618$

Angular distributions have been measured at $E_\alpha = 12.7$ to 15.2 MeV (1967IV02, 1970BU25, 1970IV04; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_{4+5}$), 16.8 MeV (1970FR1F; $\alpha_0, \alpha_1, \alpha_2$), 18.0 MeV (1958SE51, 1966LU02; α_0, α_1), 20.2 to 23.3 MeV (1970AG08, 1970FE09, 1970PI1G; α_0, α_1), 20.5 to 23.2 MeV (1971TA20; α_0), 20.9 to 24.0 MeV (1966SE1G, 1967GH1A; $\alpha_0, \alpha_1, \alpha_2$), 21.6 MeV (1968TA1Q; α_0), 22 MeV (1962EI04; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_{4+5}$), 22.5 MeV (1970LA20; α_0, α_1), 27.3 MeV (1964KO02, 1965KO07, 1965KO1A, 1967KO1J; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_{4+5}$), 31.8 MeV (1959MC1A; $\alpha_0, \alpha_1, \alpha_2$), 33.0 MeV (1967RE1B, 1968RE1F; $\alpha_0 \rightarrow \alpha_5$ and α to $^{20}\text{Ne}^*(7.17)$), 44 MeV (1968FA1A; α_0), 50.9 MeV (1965SP02, 1965SP1E, 1967RE1B, 1968RE1F; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha$ to $^{20}\text{Ne}^*(7.17)$), 80.8 MeV (1967RE1B, 1968RE1F; $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_{4+5}$) and 104 MeV (1969HA14, 1970SP01, 1971RE09; $\alpha_0, \alpha_1, \alpha_2$). At $E_\alpha = 104$ MeV the quadrupole and hexadecapole deformation lengths have been determined (1969HA14, 1970SP01, 1971RE09). The γ -decay of several ^{20}Ne states is reported by (1962EI04): see Table 20.17. See also (1963CR01, 1963MI1C, 1968SP04) and (1959BL31, 1961EI1A, 1962HU1A, 1964GR1L, 1964VE1A, 1965FA1D, 1965SA1G, 1967RA1E, 1971GO28, 1971OB1A; theor.).

Reaction (b) has been studied for $E_\alpha = 12$ to 20 MeV (1963LA08). See also ^{24}Mg in (1973ENVA).

41. $^{20}\text{Ne}(^{16}\text{O}, ^{16}\text{O})^{20}\text{Ne}$

See (1961NE04).

42. $^{20}\text{Ne}(^{32}\text{S}, ^{32}\text{S})^{20}\text{Ne}$

The static quadrupole moment of $^{20}\text{Ne}^*(1.63)$ has been determined to be $Q_0 = +0.94 \pm 0.38$ b (1969SC08).

Table 20.34: Decay of ^{20}Na ^a

Decay to $^{20}\text{Ne}^*$ (MeV \pm keV)	J^π	Branching ratio (%)	$\log ft$
1.6332 \pm 1	2 ⁺	90.0	4.93
4.97	2 ⁻	^b	> 6.1
5.62	3 ⁻	^b	> 6.5
5.79	1 ⁻	^b	> 6.5
7.43 \pm 10	2 ⁺	8.1	4.48
7.84 \pm 30	2 ⁺	0.38	5.65
8.74 \pm 30	1 ⁻	0.024	6.46
9.48 \pm 20	2 ⁺	0.11	5.41
10.28 \pm 10	2 ⁺ ; $T = 1$	1.38	3.79 \pm 0.05 ^d
10.86 \pm 20	2 ⁺	0.097	4.46
11.28 \pm 40	2 ⁺	0.032	4.50

^a (1967PO11, 1967SU05). See also (1964MA44).

^b Not observed. See discussion in (1967SU05).

^c B. Zimmerman, private communication. Note: this footnote is not labeled in the tabular.

^d (1971GO18).

43. $^{20}\text{Na}(\beta^+)^{20}\text{Ne}$ $Q_m = 13.892$

^{20}Na decays to a number of states of ^{20}Ne : see Table 20.34 (1967PO11, 1967SU05, 1964MA44). The half-life of ^{20}Na is $\tau_{1/2} = 446 \pm 3$ msec [see ^{20}Na]. The ratio of the mirror decays $^{20}\text{Na} \rightarrow ^{20}\text{Ne}^*(1.63)$ and $^{20}\text{F} \rightarrow ^{20}\text{Ne}^*(1.63)$, $(ft)^+/(ft)^- = 1.054 \pm 0.023$. If this number is directly interpreted in terms of second-class currents, the induced tensor coupling constant $g_{IT} = (1.5 \pm 0.6) \times 10^{-3}$ (1970WI05, 1971WI07).

The line shapes of the α -spectra from the decay of $^{20}\text{Ne}^*(7.42, 10.26)$ are fitted by Gamow-Teller and by Fermi decay theory. The former indicates some evidence for longitudinal nuclear alignment of β -recoils (1971MA09). See also (1970OA01, 1971TO1K, 1971TO1L) and (1971HO1D, 1971LI1H; theor.).

44. $^{21}\text{Ne}(p, d)^{20}\text{Ne}$ $Q_m = -4.536$

At $E_p = 20$ MeV, the angular distribution of the deuterons to $^{20}\text{Ne}^*(1.63)$ [very strongly populated] is characterized by $l_n = 2$ while that of the d_2 group (to $^{20}\text{Ne}^*(4.25)$) is suggestive of a weak $l_n = 2$ component. All of the observed $l_n = 1$ pick up strength is associated with $^{20}\text{Ne}^*(4.97)$ (1970HO1R). See also (1969HE1T).

$$45. \ ^{22}\text{Ne}(p, t)^{20}\text{Ne} \quad Q_m = -8.644$$

Angular distributions have been reported at $E_p = 26.9, 35.1$ and 42.4 MeV (1971FA07; $t_0, t_1, t_2, t_3, t_{4+5}, t_6$) and at 43.7 MeV (1964CE05). At the higher energy the distributions of the tritons to the ground state of ^{20}Ne and to the first $0^+; T = 2$ state [$E_x = 16.722 \pm 0.025$ MeV (1969HA38)] have been fitted by $L = 0$ and the tritons to $^{20}\text{Ne}^*(18.5)$ by $L = 2$. The latter is the first $2^+; T = 2$ state (1964CE05). The $0^+; T = 2$ state [$^{20}\text{Ne}^*(16.73)$] decays by α_0 [$-6 \pm 5\%$], $\alpha_1 + \alpha_2$ [$35 \pm 12\%$], $\alpha_3 + \alpha_4$ [$29 \pm 12\%$], $p_0 + p_1 + p_2$ [$14 \pm 9\%$], and $p_3 + p_4 + p_5$ [$13 \pm 8\%$] [measured branching ratios in percent are given in the brackets] to final states in ^{16}O and ^{19}F (1970MC04). The ratios of the cross section for formation of the analog states $^{20}\text{Ne}^*(10.26)/^{20}\text{F}^*(0)$ and $^{20}\text{Ne}^*(12.25 \pm 0.03)/^{20}\text{F}^*(1.85)$ are 2.00 ± 0.20 and 1.40 ± 0.15 , respectively, at $E_p = 45$ MeV (1969HA19). See also (1970OL1B) and (1969TO1E; theor.).

$$46. \ ^{22}\text{Ne}(\alpha, ^6\text{He})^{20}\text{Ne} \quad Q_m = -16.156$$

Angular distributions have been obtained at $E_\alpha = 42$ MeV for the transitions to $^{20}\text{Ne}^*(0, 1.63, 4.25)$ (1970AR1H).

$$47. \ ^{23}\text{Na}(p, \alpha)^{20}\text{Ne} \quad Q_m = 2.377$$

$$Q_0 = 2.373 \pm 0.008 \text{ (1967SP09).}$$

Alpha-particle groups have been observed to the ground state and to ^{20}Ne states with $E_x = 1.634 \pm 0.004$ MeV (1953DO04), 1.635 ± 0.006 , 4.248 ± 0.006 , 4.969 ± 0.006 and 5.631 ± 0.006 MeV (1957BU36). An attempt has been made to see if $^{20}\text{Ne}^*(4.97)$ could consist of unresolved states: the result was negative (1965ME10). Angular distributions have been measured at $E_p = 10.0$ MeV (1963WA10; $\alpha_0 \rightarrow \alpha_3$) and at 45.5 MeV (1969KO09; α_0).

The cross over transitions from $^{20}\text{Ne}^*(4.25, 4.97)$ are $\leq 9\%$ and $\leq 4\%$, respectively, of the cascade transition via $^{20}\text{Ne}^*(1.63)$ (1959KR66, 1960KR02): see Table 20.17. See also (1960RA23). The lifetime, τ_m , of $^{20}\text{Ne}^*(1.63) = 0.76 \pm 0.33$ psec (1956DE22): see also Table 20.18. See also (1959AJ76) and (1961AD01).

For papers dealing with resonances in the compound nucleus, see (1963FI09, 1963KU24, 1964NO09, 1966VE09, 1967KUI1, 1968VA20, 1968VA21, 1969BR1M, 1970KR1K, 1970LU17).

48. $^{23}\text{Na}(^3\text{He}, ^6\text{Li})^{20}\text{Ne}$ $Q_m = -1.642$

At $E(^3\text{He}) = 40.7$ MeV, the ground state angular distribution has been measured by (1969OH1B).

49. $^{24}\text{Mg}(p, \alpha)^{21}\text{Na}^*(p)^{20}\text{Ne}$ $Q_m = -9.315$

See (1970LI08).

50. $^{24}\text{Mg}(d, ^6\text{Li})^{20}\text{Ne}$ $Q_m = -7.840$

At $E_d = 55$ MeV, $^{20}\text{Ne}^*(0, 1.63, 4.25)$ [$J^\pi = 0^+, 2^+,$ and 4^+ , respectively] and $^{20}\text{Ne}^*(4.97, 5.62)$ [$J^\pi = 2^-$ and 3^- , respectively] are strongly populated, while $^{20}\text{Ne}^*(5.79)$ is weakly populated, as predicted by the SU_3 model. The results are similar to those from $^{16}\text{O}(^7\text{Li}, t)^{20}\text{Ne}$ (reaction 14) (1971MC04).

51. $^{25}\text{Mg}(d, ^7\text{Li})^{20}\text{Ne}$ $Q_m = -7.921$

See (1971MC04).

52. $^{26}\text{Mg}(p, t)^{24}\text{Mg}(\alpha)^{20}\text{Ne}$ $Q_m = -19.258$

See (1967MC06).

^{20}Na
(Figs. 11, 12 and 13)

Mass of ^{20}Na : From the threshold energy of the $^{20}\text{Ne}(p, n)^{20}\text{Na}$ reaction, $E_{\text{thresh.}} = 15.419 \pm 0.006$ MeV, the atomic mass excess of ^{20}Na is 6.850 ± 0.006 MeV (1971GO18, 1971WI07). See also (1964GA1C, 1966GA25, 1966KE16, 1969HA38).

1. $^{20}\text{Na}(\beta^+)^{20}\text{Ne}$ $Q_m = 13.892$

Table 20.35: Energy levels of ^{20}Na

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 446 \pm 3$ msec	β^-	1, 2, 3, 4
0.591 ± 12				2, 3
0.768 ± 8				2, 3
(0.85 ± 50)				3
0.958 ± 8				2, 3
(1.010 ± 14)				2
1.310 ± 10				2, 3
1.92 ± 40				3
2.89 ± 50		a		3
4.33 ± 100		a		3

^a Broad or unresolved.

Table 20.36: States of ^{20}Na from $^{20}\text{Ne}(p, n)^{20}\text{Na}$ and $^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$

E_x (MeV \pm keV)	
$^{20}\text{Ne}(p, n)^{20}\text{Na}$ (1971MO34)	$^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$ (1965DO04, 1965PE04)
0	0
0.591 ± 12	0.65 ± 50 ^a
0.768 ± 8	0.75 ± 50 ^a
	0.85 ± 50 ^a
0.958 ± 8	0.95 ± 50 ^a
(1.010 ± 14)	
1.310 ± 10	1.27 ± 50
	1.92 ± 40
	2.89 ± 50
	4.33 ± 100

^a These states are not fully resolved.

^{20}Na decays by positron emission to $^{20}\text{Ne}^*(1.63)$ and to a number of excited states which decay by α -emission to the ground state of ^{16}O : see Table 20.34 (1967PO11, 1967SU05). The half-life of ^{20}Na is $\tau_{1/2} = 408 \pm 6$ msec (1967SU05), 448 ± 4 msec (1970OA01), 442 ± 5 msec (1971GO18, 1971WI07), 446 ± 8 msec (1972MO08). See also (1959AJ76) and reaction 43 in ^{20}Ne .

The character of the β^+ decay, which takes place by allowed transitions to 2^+ states of ^{20}Ne , sets $J^\pi = 2^+$ for the ground state of ^{20}Na .

$$2. \ ^{20}\text{Ne}(p, n)^{20}\text{Na} \quad Q_m = -14.674$$

$$Q_0 = -14.674 \pm 0.006 \text{ (1971GO18)}.$$

Observed neutron groups at $E_p = 22.9$ MeV are displayed in Table 20.36 (1971MO34). See also (1970OA01, 1971BE46, 1971RU1F, 1971WI07).

$$3. \ ^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na} \quad Q_m = -13.910$$

$$Q_0 = -14.04 \pm 0.08 \text{ (1965PE04)};$$

$$Q_0 = -13.89 \pm 0.06 \text{ (1965DO04)}.$$

At $E(^3\text{He}) = 32$ MeV, triton groups are observed to nine states of ^{20}Na : see Table 20.36 (1965DO04, 1965PE04).

$$4. \ ^{24}\text{Mg}(p, \alpha n)^{20}\text{Na} \quad Q_m = -23.989$$

See (1967PO11, 1967SU05).

^{20}Mg
(Not illustrated)

^{20}Mg has not been observed [see, however, (1964MA44)]. The mass excess of ^{20}Mg is calculated to be 17.509 ± 0.002 MeV (using the seniority scheme) and 17.510 ± 0.002 MeV (using the supermultiplet scheme). ^{20}Mg would then be stable with respect to breakup into $^{19}\text{Na} + p$ by 2.75 MeV (1969HA38). See also (1962GO31, 1964GA1C, 1964GO1G, 1966GO1B, 1966GO1K, 1966KE16).

^{20}Al
(Not illustrated)

^{20}Al has not been observed: see (1966KE16).

References

(Closed 31 December 1971)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1950AR1A Arnold, Phys. Rev. 79 (1950) 170; Erratum Phys. Rev. 80 (1950) 34
- 1950BA1A Barnes, French and Devons, Nature 166 (1950) 145
- 1950CH1A Chao, Phys. Rev. 80 (1950) 1035
- 1950CH53 C.Y. Chao, A.V. Tollestrup, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 79 (1950) 108
- 1950HE92 R.L. Henkel and H.H. Barschall, Phys. Rev. 80 (1950) 145
- 1951PH1A Phillips and Heydenburg, Phys. Rev. 83 (1951) 184
- 1952AJ38 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 24 (1952) 321
- 1952WI27 H.B. Willard, J.K. Bair, J.D. Kington, T.M. Hahn, C.W. Snyder and F.P. Green, Phys. Rev. 85 (1952) 849
- 1953CA44 J.R. Cameron, Phys. Rev. 90 (1953) 839
- 1953DO04 D.J. Donahue, K.W. Jones, M.T. McEllistrem and H.T. Richards, Phys. Rev. 89 (1953) 824
- 1954DE1A Dearnaley, Phil. Mag. 45 (1954) 1213
- 1954DE36 S. Devons, G. Goldring and G.R. Lindsey, Proc. Phys. Soc. A67 (1954) 134
- 1954FE16 G.A. Ferguson, J. Halpern, R. Nathans and P.F. Yergin, Phys. Rev. 95 (1954) 776
- 1954PE1A Peterson, Barnes, Fowler and Lauritsen, Phys. Rev. 94 (1954) 1075
- 1954SI07 R.M. Sinclair, Phys. Rev. 93 (1954) 1082
- 1954WO23 C. Wong, Phys. Rev. 95 (1954) 761
- 1955BA1C Baranger, Phys. Rev. 99 (1955) 145
- 1955BA94 C.A. Barnes, Phys. Rev. 97 (1955) 1226
- 1955FA1A Farney, Givin, Kern and Hahn, Phys. Rev. 97 (1955) 720
- 1955FR1B Freeman, Phys. Rev. 99 (1955) 1446
- 1955HU1A Hunt and Firth, Phys. Rev. 99 (1955) 786
- 1955KI28 J.D. Kington, J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 99 (1955) 1393
- 1955MB01 J.B. Marion and R.M. Brugger, Phys. Rev. 100 (1955) 69

1955WE44 T.S. Webb, F.B. Hagecorn, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 99 (1955) 138
1956AH1A Ahnlund, Ark. Fys. 10 (1956) 425
1956DE22 S. Devons, G. Manning and J.H. Towle, Proc. Phys. Soc. A69 (1956) 173
1956DE33 G. Dearnaley, Phil. Mag. 1 (1956) 821
1956EL1A El-Bedewi, Proc. Phys. Soc. A69 (1956) 221
1957BO04 N.A. Bostrom, E.L. Hudspeth and I.L. Morgan, Phys. Rev. 105 (1957) 1545
1957BU36 W.W. Buechner and A. Sperduto, Phys. Rev. 106 (1957) 1008
1957BU66 E.M. Burbidge, G.R. Burbidge, W.A. Fowler and F. Hoyle, Rev. Mod. Phys. 29 (1957) 547
1957CL42 R.L. Clarke and E.B. Paul, Can. J. Phys. 35 (1957) 155
1957MA1A Martin, Fowler, Lauritsen and Lauritsen, Phys. Rev. 106 (1957) 1260
1957PE1B Peterson, Bratenahl and Stoering, TID-7547 (1957)
1957TH14 P. Thieberger, J. Inorg. Nucl. Chem. 4 (1957) 367
1958GO71 C.D. Goodman and J.L. Need, Phys. Rev. 110 (1958) 676
1958GR01 L.V. Groshev, A.M. Demidov, V.N. Lutsenko and V.I. Pelekhov, Atlas of Thermal Neutron Capture Gamma Rays, Atomizdat, Moscow (1958)
1958IS10 A. Isoya, H. Ohmura and T. Momota, Nucl. Phys. 7 (1958) 116
1958IS11 A. Isoya, Nucl. Phys. 7 (1958) 126
1958MO02 S. Morita and K. Takeshita, J. Phys. Soc. Jpn. 13 (1958) 1241
1958RA15 W.A. Ranken, T.W. Bonner and J.H. McCrary, Phys. Rev. 109 (1958) 1646
1958SA1A Sawicki, Bull. Acad. Pol. Sci. 6 (1958) 37
1958SE51 L. Seidlitz, E. Bleuler and D.J. Tendam, Phys. Rev. 110 (1958) 682
1958WI36 J.E. Wills, Jr., J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 109 (1958) 891
1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
1959AL06 D.E. Alburger, A. Gallmann and D.H. Wilkinson, Phys. Rev. 116 (1959) 939
1959AL1H Alkhazov et al., Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1959) 1466
1959AL91 D.G. Alkhazov, A.P. Grinberg, I.K. Lemberg and V.V. Rozhdestvenskii, Zh. Eksp. Teor. Fiz. 36 (1959) 322; Sov. Phys. JETP 9 (1959) 222
1959AM13 S. Amiel and R.E. Segel, J. Inorg. Nucl. Chem. 10 (1959) 4
1959BE1A Bedreag, Bull. Inst. Polytech. IASI (Romania) 5 (1959) 143
1959BL31 J.S. Blair, Phys. Rev. 115 (1959) 928
1959BO14 R.O. Bondelid and C.A. Kennedy, Phys. Rev. 115 (1959) 1601
1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205

- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959BR67 G. Breuer, Z. Phys. 154 (1959) 339
- 1959CO82 H.O. Cohn and J.L. Fowler, Phys. Rev. 114 (1959) 194
- 1959GA08 F. Gabbard, R.H. Davis and T.W. Bonner, Phys. Rev. 114 (1959) 201
- 1959GE34 D.S. Gemmell, A.H. Morton and W.I.B. Smith, Nucl. Phys. 10 (1959) 45
- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959GO1C Gorodetzky, Muller, Bergdolt and Post, Congress Int. Phys. Nucl., Paris, 1958 (1959) 885
- 1959HA1C Hay and Warren, Can. J. Phys. 37 (1959) 1153
- 1959HO1D Honda and Nagasaki, Proc. Phys. Soc. A74 (1959) 517
- 1959JA01 N. Jarmie and M.G. Silbert, Phys. Rev. Lett. 3 (1959) 50
- 1959KR66 T.H. Kruse, Thesis, Columbia Univ. (1959); CU-181 (1959)
- 1959KU79 J. Kuperus, P.J.M. Smulders and P.M. Endt, Physica 25 (1959) 600
- 1959LI51 A. Lippold, J.W. Muller, H.H. Staub and H. Winkler, Helv. Phys. Acta 32 (1959) 268
- 1959MA1C MacGregor, Congress Int. Phys. Nucl., Paris, 1958 (1959) 612
- 1959MC1A McKee and Rowe, Congress Int. Phys. Nucl., Paris, 1958 (1969) 501
- 1959OD08 Y. Oda, M. Takeda, C. Hu and S. Kato, J. Phys. Soc. Jpn. 14 (1959) 396
- 1959OG15 H. Ogata, J. Phys. Soc. Jpn. 14 (1959) 707
- 1959ON26 K. Ono, J. Shimada, K. Kuroda, O. Tanaka, H. Kamitsubo, A. Ito, S. Tanaka and M. Imaizumi, J. Phys. Soc. Jpn. 14 (1959) 117
- 1959RE1A Reid et al., Congres Int. Phys. Nucl., Paris, 1958 (1959) 700
- 1959SI1A Sitenko, Usp. Fiz. Nauk 67 (1959) 377
- 1959TR1A Trail and Raboy, Bull. Amer. Phys. Soc. 4 (1959) 220
- 1959VA10 S.S. Vasilev and L.Y. Shavtvalov, Zh. Eksp. Teor. Fiz. 36 (1959) 317; Sov. Phys. JETP 9 (1959) 218
- 1960AL05 E. Almqvist, D.A. Bromley and J.A. Kuehner, Phys. Rev. Lett. 4 (1960) 515
- 1960AM03 G. Amsel and O. Smulkowski, Compt. Rend. 251 (1960) 950
- 1960AN1A Andreev, Erokhina and Lemberg, Bull. Acad. Sci. USSR 24 (1960) 1470
- 1960BE1A Bedreag, Analele Stiint. Univ. Al. I. Cuza Iasi (Ser. Nine) I (Romania) 6 (1960) 129
- 1960BR35 C. Broude and H.E. Gove, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds., D.A. Bromley and E.W. Vogt (1960) 754
- 1960BU07 J.W. Butler, Phys. Rev. 118 (1960) 222
- 1960BU1C Bullock and Moore, Phys. Rev. 119 (1960) 721

- 1960CA10 J. Catala, A.G. Rodriguez and J. Aguilar, *An. Real. Soc. Espan. Fis. y Quim.* A56 (1960) 35
- 1960CA1C Catala, Rodriguez and Aguilar, *Nucl. Forces and the Few Nucleon Prob.*, Pergamon (1960) 693
- 1960EL09 M. El-Nadi and M. Wafik, *Proc. Phys. Soc.* 76 (1960) 185
- 1960EV1A Everling, *Z. Naturforsch.* A15 (1960) 84
- 1960FR04 J.M. Freeman, *Phys. Rev.* 120 (1960) 1436
- 1960GO12 S. Gorodetzky, G. Sutter, P. Chevallier, F. Scheibling and R. Armbruster, *Comp. Rend.* 250 (1960) 1028
- 1960HU11 S.E. Hunt, R.A. Pope, D.V. Freck and W.W. Evans, *Phys. Rev.* C12 (1960) 1740
- 1960JA13 N. Jarmie and M.G. Silbert, *Phys. Rev.* 120 (1960) 914
- 1960KA18 J.V. Kane, R.E. Pixley and D.H. Wilkinson, *Phys. Rev.* 120 (1960) 952
- 1960KR02 T.H. Kruse, R.D. Bent and L.J. Lidofsky, *Phys. Rev.* 119 (1960) 289
- 1960LE1A Lemberg, *Bull. Amer. Phys. Soc.* 5 (1960) 473
- 1960LU06 A. Lurio, G. Weinreich, C.W. Drake, V.W. Hughes and J.A. White, *Phys. Rev.* 120 (1960) 153
- 1960MC05 J.H. McCrary and I.L. Morgan, *Bull. Amer. Phys. Soc.* 5 (1960) 246, HA5
- 1960MC09 L.C. McDermott, K.W. Jones, H. Smotrich and R.E. Benenson, *Phys. Rev.* 118 (1960) 175
- 1960NA1A Nagarajan and Banerjee, *Nucl. Phys.* 17 (1960) 341
- 1960OD01 Y. Oda, M. Takeda, N. Takano, T. Yamazaki, C. Hu, K. Kikuchi, S. Kobayashi, K. Matsuda and Y. Nagahara, *J. Phys. Soc. Jpn.* 15 (1960) 760
- 1960PE25 J.M. Peterson, A. Bratenahl and J.P. Stoering, *Phys. Rev.* 120 (1960) 521
- 1960RA23 T.A. Rabson, W.D. Barfield, D.L. Bernard, T.W. Bonner and W.W. Givens, *Nucl. Phys.* 21 (1960) 43
- 1960RI05 J. Rickards, *Rev. Mex. Fis.* 9 (1960) 35
- 1960SC01 G. Scharff-Goldhaber, A. Goodman and M.G. Silbert, *Phys. Rev. Lett.* 4 (1960) 25
- 1960SH1A Sheline and Wildermuth, *Nucl. Phys.* 21 (1960) 196
- 1960SM03 D.M. Smith, N.A. Bostrom and E.L. Hudspeth, *Phys. Rev.* 117 (1960) 514
- 1960SO1A Solovev, *Sov. Phys. Dokl.* 5 (1960) 298
- 1960TA08 M. Takeda, *J. Phys. Soc. Jpn.* 15 (1960) 557
- 1960TA1C Talmi and Unna, *Ann. Rev. Nucl. Sci.* 10 (1960) 353
- 1960TE03 I.B. Teplov, O.P. Shevchenko and E.K. Ruuge, *Zh. Eksp. Teor. Fiz.* 39 (1960) 923; *Sov. Phys. JETP* 12 (1961) 640

- 1960TS02 K. Tsukada and T. Fuse, J. Phys. Soc. Jpn. 15 (1960) 1994
- 1960VA15 L. Varga, Nucl. Phys. 20 (1960) 487; Rept. Central Res. Inst. Phys. Hung. Acad. Sci. 8 (1960) 7
- 1960VE06 G.E. Velyukhov, A.N. Prokofev and S.V. Starodubtsev, Zh. Eksp. Teor. Fiz. 39 (1960) 563; Sov. Phys. JETP 12 (1961) 395
- 1960ZE03 Ya.B. Zeldovich, Zh. Eksp. Teor. Fiz. 38 (1960) 1123; Sov. Phys. JETP 11 (1960) 812
- 1961AD01 H.S. Adams, J.D. Fox, N.P. Heydenburg and G.M. Temmer, Bull. Amer. Phys. Soc. 6 (1961) 250, GA14
- 1961AL12 E. Almqvist and J.A. Kuehner, Can. J. Phys. 39 (1961) 1246
- 1961AS02 L. Ask, Ark. Fys. 19 (1961) 219
- 1961BA1C Baz, Goldanskii and Zeldovich, Sov. Phys. Uspekhi 3 (1961) 729
- 1961BA1D Badalyan and Baz, Zh. Eksp. Teor. Fiz. 40 (1961) 549; Sov. Phys. JETP 13 (1961) 383
- 1961BE10 R.E. Benenson, H.Y. Chen and L.J. Lidofsky, Phys. Rev. 122 (1961) 874
- 1961BE13 E.H. Beckner, R.L. Bramblett, G.C. Phillips and T.A. Eastwood, Phys. Rev. 123 (1961) 2100
- 1961BR1A Brill', Vlasov, Kalinin and Sokolov, Dokl. Akad. Nauk SSSR 136 (1961) 55; Sov. Phys., Dokl. 6 (1961) 24
- 1961CL06 M.A. Clark, H.E. Gove and A.E. Litherland, Can. J. Phys. 39 (1961) 1241
- 1961DA16 E.A. Davis, F. Gabbard, T.W. Bonner and R. Bass, Nucl. Phys. 27 (1961) 448
- 1961DO08 W.R. Dodge, Thesis, Stanford Univ. (1961); HEPL-246 (1961); Nucl. Sci. Abstr. 16, 750, Abstr. 5921 (1962)
- 1961EI1A Eisberg and Porter, Rev. Mod. Phys. 33 (1961) 190
- 1961ET01 K. Etoh, K. Hisatake, T. Kobayashi, K. Horie, H. Ohmura and Y. Nogami, J. Phys. Soc. Jpn. 16 (1961) 2091
- 1961FR02 E. Freiberg and V. Soergel, Z. Phys. 162 (1961) 114
- 1961FU01 M. Furukawa and S. Tanaka, J. Phys. Soc. Jpn. 16 (1961) 129
- 1961GO12 H.E. Gove, A.E. Litherland and M.A. Clark, Can. J. Phys. 39 (1961) 1243
- 1961GO1G Gove, Litherland and Clark, Nature 191 (1961) 1381
- 1961GO21 H.E. Gove, A.E. Litherland and A.J. Ferguson, Phys. Rev. 124 (1961) 1944
- 1961GU1A Gusakow, Albouy, Poffe and Riehl, J. Phys. Rad. 22 (1961) 636
- 1961JA1E Jarmie, Nucl. Phys. 28 (1961) 426
- 1961LI06 A.E. Litherland, Can. J. Phys. 39 (1961) 1245
- 1961LI07 A.E. Litherland, M.A. Clark and H.E. Gove, Can. J. Phys. 39 (1961) 1249

- 1961LI17 A.E. Litherland, J.A. Kuehner, H.E. Gove, M.A. Clark and E. Almqvist, Phys. Rev. Lett. 7 (1961) 98
- 1961NE04 E. Newman, P.G. Roll and F.E. Steigert, Phys. Rev. 122 (1961) 1842
- 1961NO05 E. Norbeck, Phys. Rev. 121 (1961) 824
- 1961OK1A Okada, Nucleonics 19 (1961) 79
- 1961PO13 A.J.P.L. Policarpo, W.R. Phillips and J.C. Lisle, Proc. Rutherford Jub. Int. Conf. (1961) 607
- 1961RA06 L.A. Rayburn, Phys. Rev. 122 (1961) 168
- 1961RO05 L. Rosen, J.E. Brolley, Jr. and L. Stewart, Phys. Rev. 121 (1961) 1423
- 1961RO13 L. Rosen, J.E. Brolley, Jr., M.L. Gursky and L. Stewart, Phys. Rev. 124 (1961) 199
- 1961SE1C Seaman and Quinton, Bull. Amer. Phys. Soc. 6 (1961) 470
- 1961SO02 P.V. Sorokin, A.I. Popov, V.E. Storizhko and A.Ya. Taranov, Zh. Eksp. Reor. Fiz. 40 (1961) 1253; Sov. Phys. JETP 13 (1961) 883
- 1961TR1B True and Warburton, Nucl. Phys. 22 (1961) 426
- 1961WA03 A.H. Wapstra and D.W. Connor, Nucl. Phys. 22 (1961) 336
- 1961WI1A Wilkinson, Wollan and Koehler, Ann. Rev. Nucl. Sci. 11 (1961) 303
- 1961WI1C Williamson and Picard, J. Phys. Rad. 22 (1961) 651
- 1961WI22 C.F. Williamson, Phys. Rev. 122 (1961) 1877
- 1961YA02 H. Yamaguchi, J. Phys. Soc. Jpn. 16 (1961) 583
- 1962AG01 J. Aguilar, J.B.A. England and A. Garcia, An. Real. Soc. Espan. Fis. y Quim. A58 (1962) 319
- 1962BI1C Bird, Gibbons and Good, Phys. Lett. 1 (1962) 262
- 1962BO23 M. Bouten, Nuovo Cim. 26 (1962) 63
- 1962BR35 C. Broude, M.A. Clark and A.E. Litherland, Phys. Lett. 3 (1962) 118
- 1962CE1B Cevolani and Petralia, Nuovo Cim. 26 (1962) 1328
- 1962CH12 E. Chacon and M. Moshinsky, Phys. Lett. 1 (1962) 330
- 1962DA06 T. Dazai, Prog. Theor. Phys. 27 (1962) 433
- 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
- 1962EI04 W.W. Eidson and R.D. Bent, Phys. Rev. 128 (1962) 1312
- 1962FI02 R.W. Finlay and G.E. Owen, Phys. Rev. 128 (1962) 2309
- 1962FR09 J.M. Freeman and D. West, Nucl. Phys. 38 (1962) 89
- 1962GO1E Gorbunov et al., Zh. Eksp. Teor. Fiz. 42 (1962) 747; Sov. Phys. JETP 15 (1962) 520
- 1962GO1J Gorshkov, Ziyabkin and Tsvetkov, Atomn. Energ. (USSR) 13 (1962) 65

- 1962GO31 V.I. Goldanskii, Dokl. Akad. Nauk SSSR 146 (1962) 1309; Sov. Phys. Dokl. 7 (1963) 922
- 1962GU10 M. Gusakow, Ann. Phys. 7 (1962) 67
- 1962HI06 S. Hinds, H. Marchant and R. Middleton, Nucl. Phys. 38 (1962) 81
- 1962HU12 H. Hulubei, A. Berinde, I. Neamu, J. Franz, N. Martalogu and M. Ivascu, Nucl. Phys. 39 (1962) 686
- 1962HU1A Huizenga and Igo, Nucl. Phys. 29 (1962) 462
- 1962JO14 J.C. Jodogne, P.C. Macq and J. Steyaert, Phys. Lett. 2 (1962) 325
- 1962KA12 O.N. Kaul, Nucl. Phys. 33 (1962) 177
- 1962KA26 O.N. Kaul, Nucl. Phys. 39 (1962) 325
- 1962KA37 J. Kantele and D.G. Gardner, Nucl. Phys. 35 (1962) 353
- 1962KE03 L. Keszthelyi, I. Berkes, I. Demeter and I. Fodor, Nucl. Phys. 29 (1962) 241
- 1962KU03 J.A. Kuehner, Phys. Rev. 125 (1962) 1650
- 1962MA38 S. Malmskog and J. Konijn, Nucl. Phys. 38 (1962) 196
- 1962MO08 R. Moreh and G. Davis, Nucl. Phys. 36 (1962) 315
- 1962OT01 P.S. Otstavnov and V.I. Popov, Zh. Eksp. Teor. Fiz. 43 (1962) 385; Sov. Phys. JETP 16 (1963) 276
- 1962RY01 A. Rytz, H.H. Staub, H. Winkler and W. Zych, Helv. Phys. Acta 35 (1962) 341
- 1962SC12 G. Schrank, E.K. Warburton and W.W. Daehnick, Phys. Rev. 127 (1962) 2159
- 1962TA14 J.D. Talman, Can. J. Phys. 40 (1962) 1730
- 1962TA1B Talmi and Unna, Nucl. Phys. 30 (1962) 280
- 1962VA1D Van Leuven and Bouten, AERE-R-4131 (1962)
- 1963AL07 E. Almqvist, D.A. Bromley, J.A. Kuehner and B. Whalen, Phys. Rev. 130 (1963) 1140
- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963BA1F Bartolini, Donaldson and Passell, Bull. Amer. Phys. Soc. 8 (1963) 477
- 1963BE19 I. Berkes, I. Dezsi, I. Fodor and L. Keszthelyi, Nucl. Phys. 43 (1963) 103
- 1963BO1C Borecka and Neuert, in Padua (1963) 556A
- 1963CH20 A. Chatterjee, Nucl. Phys. 47 (1963) 511
- 1963CR01 J.G. Cramer, Jr. and W.W. Eidson, Bull. Amer. Phys. Soc. 8 (1963) 26, F10
- 1963CS03 J. Csikai, B. Gyarmati and I. Hunyadi, Nucl. Phys. 46 (1963) 141
- 1963DA1C Davis, Bull. Amer. Phys. Soc. 8 (1963) 26
- 1963DA1D Davis, 3rd Conf. on Reactions between Complex Nuclei (1963) 61

1963DU1C Duorah and Kushwaha, *Nuovo Cim.* 27 (1963) 530
 1963EV01 F. Everling, *Nucl. Phys.* 40 (1963) 670
 1963FE01 A.T.G. Ferguson, G.C. Morrison, N. Gale and R.E. White, *Bull. Amer. Phys. Soc.* 8 (1963) 47, NA2
 1963FE1B Ferguson, Gale, Morrison and White, in Padua (1963) 510
 1963FI09 T.R. Fisher and W. Whaling, *Phys. Rev.* 131 (1963) 1723
 1963FI1B Finck et al., *Z. Phys.* 174 (1963) 337
 1963FL1A Flamm, Levinson and Meshkov, *Phys. Rev.* 129 (1963) 297
 1963GI1D Gibbons, *Prog. in Fast Neutron Phys.* (1963) 193
 1963GL01 S.S. Glickstein and R.G. Winter, *Phys. Rev.* 129 (1963) 1281
 1963GO04 J. Goldemberg and Y. Torizuka, *Phys. Rev.* 129 (1963) 312
 1963GO1J Gorshkov and Tsvetkov, *Atomn. Energ. (USSR)* 14 (1963) 550
 1963GO27 G. Goldring, H. Winkler and C. Zaidins, *Bull. Amer. Phys. Soc.* 8 (1963) 598, F10
 1963HA1E Hayward, *Rev. Mod. Phys.* 35 (1963) 324
 1963HU1C Husain and Taylor, in Padua (1963) 559A
 1963HU1D Hulubei et al., *Zh. Eksp. Teor. Fiz.* 45 (1963) 1822; *Sov. Phys. JETP* 18 (1964) 1248
 1963JA01 R. Jahr, *Phys. Rev.* 129 (1963) 320
 1963JE04 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, *Nucl. Phys.* 44 (1963) 453
 1963KE1B Kelson, *Phys. Rev.* 132 (1963) 2189
 1963KU05 J.A. Kuehner, J.D. Prentice and E. Almqvist, *Phys. Lett.* 4 (1963) 332
 1963KU19 D. Kurath, *Phys. Rev.* 132 (1963) 1146
 1963KU24 J. Kuperus, P.W.M. Glaudemans and P.M. Endt, *Physica* 29 (1963) 1281
 1963LA08 N.O. Lassen and J.S. Olsen, *Kgl. Danske Videnskab. Selskab., Mat.-Fys. Medd.* 33, No.13 (1963)
 1963LE1C Levinson, *Phys. Rev.* 132 (1963) 2184
 1963LE1D Levkovskii, *Zh. Eksp. Teor. Fiz.* 45 (1963) 305; *Sov. Phys. JETP* 18 (1964) 213
 1963LU08 H.F. Lutz, J.B. Mason and C.T. Paulson, *Nucl. Phys.* 43 (1963) 405
 1963MA58 R.L. Macklin, J.H. Gibbons and T. Inada, *Nucl. Phys.* 43 (1963) 353
 1963MA60 R.L. Macklin, J.H. Gibbons and T. Inada, *Phys. Rev.* 129 (1963) 2695
 1963MI1B Mihailovic and Rosina, *Nucl. Phys.* 40 (1963) 252
 1963MI1C Mikumo, Nonaka, Yamaguchi and Maki, in Padua (1963) 1088A
 1963MI1D Micu, Padua (1963) 339A; *Acta Phys. Pol.* 23 (1963) 305

1963NA08 E.G. Nadjakov, Nucl. Phys. 48 (1963) 492
 1963NO1C Nonaka et al., INSJ-56 (1963)
 1963PA03 S.P. Pandya, Nucl. Phys. 43 (1963) 636
 1963PI1B J. Picard and C. Williamson, J. Phys. (France) 24 (1963) 813
 1963RA1A Rayburn, in Padua (1963) 322
 1963RO13 D. Robson, Nucl. Phys. 42 (1963) 592
 1963RO21 V.M. Rout, W.M. Jones and D.G. Waters, Nucl. Phys. 45 (1963) 369
 1963SA07 K. Sasaki, Prog. Theor. Phys. 29 (1963) 383
 1963TS01 T. Tsang and D. Connor, Phys. Rev. 132 (1963) 1141
 1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163
 1963WA10 K.L. Warsh, G.M. Temnar and H.R. Blieden, Nucl. Phys. 46 (1963) 45
 1963WA12 K.L. Warsh, G.M. Temnar and H.R. Blieden, Phys. Rev. 131 (1963) 1690
 1964AB1A Abecasis, Heras and Bosh, in Paris (1964) 453
 1964AL14 T.K. Alexander, J.D. Pearson, A.E. Litherland and C. Broude, Phys. Rev. Lett. 13 (1964) 86
 1964AL15 E. Almqvist and J.A. Kuehner, Nucl. Phys. 55 (1964) 145
 1964AL17 E. Almqvist, J.A. Kuehner, D. McPherson and E.W. Vogt, Phys. Rev. 136 (1964) B84
 1964BA1J Bassichis, Levinson and Kelson, Phys. Rev. 136 (1964) B380
 1964BA1K Bassichis and Kelson, Phys. Rev. 136 (1964) B385
 1964BO1E Boschitz, Vincent, Berchaw and Priest, Phys. Rev. Lett. 13 (1964) 442
 1964BO35 J. Borggreen, B. Elbek and R.B. Leachman, Kgl. Danske Videnskab. Selskab., Mat.-Fys. Medd. 34, No. 9 (1964)
 1964BR12 G. Breuer and L. Jahnke, Z. Naturforsch. A19 (1964) 471
 1964BR18 C. Broude, A.E. Litherland and J.D. Pearson, Phys. Lett. 11 (1964) 321
 1964BR1H Brown, in Paris (1964) 129
 1964CE05 J. Cerny, R.H. Pehl and G.T. Garvey, Phys. Lett. 12 (1964) 234
 1964CH21 P.R. Chagnon, Nucl. Phys. 59 (1964) 257
 1964CO24 S. Cohen, R.D. Lawson, M.H. Macfarlane and M. Soga, Phys. Lett. 9 (1964) 180
 1964DI1C Din, Unpublished Thesis, Rice Univ. (1964)
 1964EL01 A.Z. El-Behay, M.A. Farouk, V.J. Gontchar, V.A. Lutsik, M.H. Nassef and I.I. Zasloubovsky, Nucl. Phys. 56 (1964) 224
 1964EL02 A.J. Elwyn, J.E. Monahan, R.O. Lane and A. Langsdorf, Jr., Nucl. Phys. 59 (1964) 113

1964EL1B El-Nadi and Riad, Nucl. Phys. 50 (1964) 33
 1964EN1A Engeland and Kallio, Nucl. Phys. 59 (1964) 211
 1964FA04 U. Facchini, E. Saetta-Menichella, F. Tonolini and L. Tonolini-Severgnini, Nucl. Phys. 51 (1964) 460
 1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49
 1964GA1C Garvey, Cerny and Pehl, Phys. Rev. Lett. 13 (1964) 548
 1964GO1G Goldsanskii, Dokl. Akad. Nauk SSSR 157 (1964) 321; Sov. Phys. Dokl. 9 (1965) 566
 1964GR1L Greider, Private Communication (1964)
 1964HE18 I. Heertje, I. Delvenne, W. Nagel and A.H.W. Aten, Jr., Physica 30 (1964) 1762
 1964HE1D Henley and Yu, Phys. Rev. 135 (1964) B1152
 1964HI04 C.T. Hibdon, Phys. Rev. 133 (1964) B353
 1964IN03 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, Nucl. Phys. 59 (1964) 1
 1964JO04 C.H. Johnson and C.C. Trail, Phys. Rev. 133 (1964) B1183
 1964KA1A Kawai, Terasawa and Izumo, Nucl. Phys. 59 (1964) 289
 1964KE06 I. Kelson and C.A. Levinson, Phys. Rev. 134 (1964) B269
 1964KO02 J. Kokame, K. Fukunaga, N. Inoue and H. Nakamura, Phys. Lett. 8 (1964) 342
 1964KU03 J.A. Kuehner and J.D. Pearson, Can. J. Phys. 42 (1964) 477
 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 379
 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 411
 1964LO1A Lopez, Rickards and Mazari, Nucl. Phys. 51 (1964) 321
 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
 1964MA1N Massot, El'Baz and Lafoucriere, Nucl. Phys. 58 (1964) 273
 1964MA44 R.D. Macfarlane and A. Siivola, Nucl. Phys. 59 (1964) 168
 1964ME1B Meyerhof, Phys. Lett. 8 (1964) 267
 1964MI05 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 63
 1964MI16 F.C. Michel, Phys. Rev. 133 (1964) B329
 1964MO1E Moshinsky, Congress Int. Phys. Nucl., Paris, 1964 (1964) 638
 1964NO09 R. Nordhagen and H.B. Steen, Physica Norvegica 1 (1964) 239; Phys. Abstr. 68, 72, Abstr. 804 (1965)
 1964PA1D Pandya and Green, Nucl. Phys. 57 (1964) 658
 1964PE02 J.D. Pearson, E. Almqvist and J.A. Kuehner, Can. J. Phys. 42 (1964) 489

1964PE05 J.D. Pearson and R.H. Spear, Nucl. Phys. 54 (1964) 434
 1964SA09 D.G. Sargood and G.D. Putt, Aust. J. Phys. 17 (1964) 56
 1964SA1B Saetta-Menichella, Tonolini and Tonolini-Severgnini, Nucl. Phys. 51 (1964) 449
 1964SE1A Seagrave, Brolley and Beery, Rev. Sci. Instrum. 35 (1964) 1290
 1964SI09 R.H. Siemssen, R. Felst, M. Cosack and J.L. Weil, Nucl. Phys. 52 (1964) 273
 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
 1964ST1F Strohal, Kulisic, Kolar and Cindro, Phys. Lett. 10 (1964) 104
 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Edition, Suppl. Vol. 1 (1964)
 1964TA05 N.W. Tanner, G.C. Thomas and E.D. Earle, Nucl. Phys. 52 (1964) 29
 1964TA1D Talman, Phys. Rev. 135 (1964) B1302
 1964TE1F Temmer, Phys. Rev. Lett. 12 (1964) 330
 1964TR1A Trlifaj, Nucl. Phys. 52 (1964) 114
 1964VE1A Venter and Frahn, Ann. Phys. 27 (1964) 385, 401
 1964VO03 E.W. Vogt, D. McPherson, J.A. Kuehner and E. Almqvist, Phys. Rev. 136 (1964) B99
 1965AS07 L. Ask, Ark. Fys. 29 (1965) 195
 1965BA1J Bassichis, Giraud and Ripka, Phys. Rev. Lett. 15 (1965) 980
 1965BA1K Barrows, Unpublished Thesis, Univ. of Kentucky (1965)
 1965BI03 J.R. Bird, J.A. Biggerstaff, J.H. Gibbons and W.M. Good, Phys. Rev. 138 (1965) B20
 1965BO1H Bondorf and Leachman, Kgl. Danske Videnskab. Selskab., Mat.-Fys. Medd. 34, No. 10 (1965)
 1965BO42 M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Buttner, A. Lindner and H. Meldner, Nucl. Phys. 63 (1965) 438
 1965BR13 R.E. Brown, J.S. Blair, D. Bodansky, N. Cue and C.D. Kavaloski, Phys. Rev. 138 (1965) B1394
 1965CA02 E.B. Carter, P.H. Stelson, M.K. Mehta and D.L. Bernard, Nucl. Phys. 63 (1965) 575
 1965DA1D Danos and Fuller, Ann. Rev. Nucl. Sci. 15 (1965) 29
 1965DE1H De Llano et al., Nucl. Phys. 72 (1965) 379
 1965DE1M De Llano, NRL 6256 (1965)
 1965DO04 P.F. Donovan and P.D. Parker, Phys. Rev. Lett. 14 (1965) 147
 1965EV03 H.C. Evans, M.A. Eswaran, H.E. Gove, A.E. Litherland and C. Broude, Can. J. Phys. 43 (1965) 82
 1965FA1D Fayard, Lamot, El'baz and Lafoucriere, Compt. Rend. 261 (1965) 1663

1965FE02 P. Federman and I. Talmi, Phys. Lett. 19 (1965) 490
1965GO1C Goloskie, Niiler and Wooten, Bull. Amer. Phys. Soc. 10 (1965) 1135
1965GR1H Grechukhin, Nucl. Phys. 62 (1965) 273
1965IN1A Inopin, Kresnin and Tishchenko, Yad. Fiz. 2 (1965) 802
1965JA13 R. Jahr and G. Mairle, Nucl. Phys. 70 (1965) 383
1965KA14 S. Kakigi, J. Phys. Soc. Jpn. 20 (1965) 1967
1965KO07 J. Kokame, K. Fukunaga, H. Nakamura and N. Inoue, J. Phys. Soc. Jpn. 20 (1965) 475
1965KO1A Kokame, Fukunaga, Nakamura and Inoue, Suppl. Bull. Inst. Chem. Res. Kyoto Univ. (1965) 26
1965MA28 L.G. Mann and S.D. Bloom, Phys. Rev. 139 (1965) B540
1965MC02 G.J. McCallum and A.J. Ferguson, Can. J. Phys. 43 (1965) 111
1965ME10 L. Meyer-Schutzmeister and R.E. Segel, Nucl. Phys. 71 (1965) 343
1965MO19 R. Moreh, Nucl. Phys. 70 (1965) 293
1965NA1C Nagel and Aten, Physica 31 (1965) 1091
1965NE07 R.W. Newsome, Jr., Nucl. Phys. 71 (1965) 353
1965NE1C Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 25
1965PE01 J.D. Pearson, T.K. Alexander, C. Broude, A.E. Litherland, J.A. Kuehner and E. Almqvist, Bull. Amer. Phys. Soc. 10 (1965) 37, CC6
1965PE04 R.H. Pehl and J. Cerny, Phys. Lett. 14 (1965) 137
1965PI03 J. Picard and C.F. Williamson, Nucl. Phys. 63 (1965) 673
1965SA1G Satchler, Nucl. Phys. 70 (1965) 177; Erratum Nucl. Phys. 89 (1966) 707
1965SH1D Shoda, Nucl. Phys. 72 (1965) 305
1965SI18 R.H. Siemssen, L.L. Lee, Jr. and D. Cline, Phys. Rev. 140 (1965) B1258
1965SO1A Sorriaux, Compt. Rend. 260 (1965) 3045
1965SP02 A. Springer and B.G. Harvey, Phys. Rev. Lett. 14 (1965) 316
1965SP1E Springer, UCRL 11681 (1965)
1965ST22 P.H. Stelson and L. Grodzins, Nucl. Data A1 (1965) 21
1965TA1E Tanner, Nucl. Phys. 63 (1965) 383
1965TS1A Tsenter and Silin, Atomn. Energ. (USSR) 19 (1965) 48
1965VA11 W.T.H. Van Oers, G.J.C. Van Niftrik, H.L. Jonkers and K.W. Brockman, Jr., Nucl. Phys. 74 (1965) 469
1965VA14 C. van der Leun, D.M. Sheppard and P.J.M. Smulders, Phys. Lett. 18 (1965) 1344

1965VA23 L. Valentin, Nucl. Phys. 62 (1965) 81
1965WA08 K.L. Warsh and S. Edwards, Nucl. Phys. 65 (1965) 382
1965WA1D Waggoner and Jaffe, Nucl. Phys. 69 (1965) 305
1965WY1B Wynchank, Garg, Rainwater and Havens, Bull. Amer. Phys. Soc. 10 (1965) 514
1965YO1A You-Wen, Tsung-Hua and Min, Acta Phys. Sin. 21 (1965) 542
1965YU1A Yu, Chang and Yu, Acta Phys. Sin. 21 (1965) 545
1965ZA1B Zamick, Phys. Lett. 19 (1965) 580
1965ZE1B Zelenskaya, Mailing, Neudachin and Smirnov, Yad. Fiz. 2 (1965) 427
1966AB04 C. Abulaffio, Nucl. Phys. 81 (1966) 71
1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743
1966AM1B Amos, Nucl. Phys. 77 (1966) 225
1966AR10 A. Armigliato, F. Brandolini, F. Pellegrini and E. Crescenti, Nuovo Cim. B45 (1966) 92
1966AR1D Arima, 2nd Symp. Struct. Low-Medium Mass Nuclei (1966) 77
1966BA2C Bassichis, Giraud and Ripka, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintmeister; ZFK-122 (1966) 354
1966BA2E Bassichis, Giraud and Ripka, J. Phys. (Paris) 27 (1966) C1-46
1966BA2F Bassichis and Scheck, Phys. Rev. 145 (1966) 771
1966BA2H Bartouv, Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch (1966) 453
1966BA42 F.J. Bartis, Nuovo Cim. B45 (1966) 113
1966BH05 S.M. Bharathi, U.T. Raheja and E. Kondaiah, Nucl. Phys. 83 (1966) 407
1966BO17 G.J. Borse and J.M. Eisenberg, Phys. Lett. 22 (1966) 630
1966BR04 G.E. Brown and A.M. Green, Nucl. Phys. 75 (1966) 401
1966BR1H Brink, Struct. of Complex Nuclei, 1966; Ed., N.N. Bogolyubov (1969) 69
1966BR1T Broude, Nucl. Spin Parity Assignments; Ed., Gove (1966) 162
1966CA14 J. Cabe, M. Laurat and P. Yvon, J. Phys. (France) 27 (1966) 755
1966CH1G Chacon, Bull. Amer. Phys. Soc. 11 (1966) 762
1966DA1E Dalidchik and Sayasov, Yad. Fiz. 3 (1966) 820
1966EL1C Elwyn et al., Antwerp 1965 Neutron Conf. (1966) 537
1966ER1A Ericson and Mayer-Kuckuk, Ann. Rev. Nucl. Sci. 16 (1966) 183
1966GA1K Garg et al., Antwerp 1965 Neutron Conf. (1966) 219
1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197

1966GI1A Gillet, Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch (1966) 43
1966GO1B Goldanskii, Ann. Rev. Nucl. Sci. 16 (1966) 1
1966GO1K Goldanskii, Sov. Phys. Dokl. 11 (1966) 235
1966HA33 P.S. Hauge and G.H. Duffey, Phys. Rev. 152 (1966) 1023
1966IN01 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, Nucl. Phys. 85 (1966) 184
1966JA1J Jahr and Santo, Z. Naturforsch. A21 (1966) 964
1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
1966KA1A Kadenskii, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 138; Bull. Acad. Sci. USSR Phys. Ser. 30 (1966) 145
1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
1966KN02 T. Knellwolf and J. Rossel, Helv. Phys. Acta 39 (1966) 376
1966KU03 J.A. Kuehner and R.W. Ollerhead, Phys. Lett. 20 (1966) 301
1966KU1D Kuehner, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 146
1966LA14 M. Lambert, G. Dumazet and H. Beaumevieille, Compt. Rend. B263 (1966) 513
1966LO07 P. Loncke and J. Pradal, Phys. Lett. 22 (1966) 320
1966LU02 B.T. Lucas, S.W. Cospers and O.E. Johnson, Phys. Rev. 144 (1966) 972
1966MA60 J.B. Marion, Rev. Mod. Phys. 38 (1966) 660
1966ME1H Meyer-Schutzmeister, 2nd Symp. Struct. Low-Medium Mass Nuclei (1966) 195
1966MI14 B. Mitra and A.M. Ghose, Nucl. Phys. 83 (1966) 157
1966MO25 S. Morita, T. Nakagawa, C.-C. Hsu and S.-M. Lee, J. Phys. Soc. Jpn. 21 (1966) 2435
1966NE1C Newsome, Bull. Amer. Phys. Soc. 11 (1966) 44
1966PA1K Parker, Whitehead and Shute, Aust. J. Phys. 19 (1966) 147
1966PI1B Picard and De Pinho, J. Phys. (Paris) 27 (1966) C1-42
1966PR14 R. Prasad, D.C. Sarkar and C.S. Khurana, Nucl. Phys. 85 (1966) 476
1966RI05 R.C. Ritter, Nucl. Phys. 80 (1966) 377
1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (1966) 237
1966RO1R Rosen, Antwerp 1965 Neutron Conf. (1966) 379
1966SC09 J.P. Schiffer, L.L. Lee, Jr., A. Marinov and C. Mayer-Boricke, Phys. Rev. 147 (1966) 829
1966SE1G Sen Gupta and Khan, AECD/EP/10 (1966)
1966SU1D Suzuki and Nagata, Prog. Theor. Phys. 36 (1966) 37
1966TR02 L. Trlifaj, M. Gmitro and M. Sotona, Nucl. Phys. 86 (1966) 681

1966VE09 W.A.M. Veltman and F.C. Erne, Phys. Lett. 23 (1966) 139
 1966VE1B Velkley, Brandenberger and McEllistrem, Bull. Amer. Phys. Soc. 11 (1966) 832
 1966WA17 S.I. Warshaw, D.A. Goldberg and G.E. Owen, Phys. Rev. 151 (1966) 834
 1966WE04 J.J. Wesolowski, L.F. Hansen, J.G. Vidal and M.L. Stelts, Phys. Rev. 148 (1966) 1063
 1966WE1B Weinberg, Antwerp 1965 Neutron Conf. (1966) 37
 1966WO03 G. Wolfer and M. Bormann, Z. Phys. 194 (1966) 75
 1966YO1B Yoccoz, J. Phys. (Paris) 27 (1966) C1-3
 1966ZA1E Zamick, Bull. Amer. Phys. Soc. 11 (1966) 56
 1967AB03 C. Abulaffio, Phys. Rev. 161 (1967) 925
 1967AD01 E. Adelberger and A.B. McDonald, Phys. Lett. B24 (1967) 270; Erratum Phys. Lett. B24 (1967) 618
 1967AF01 I.R. Afnan, Phys. Rev. 163 (1967) 1016
 1967BA1K W.H. Bassichis and J.P. Svenne, Phys. Rev. Lett. 18 (1967) 80
 1967BA1V J. Bar-Touv, Phys. Rev. 164 (1967) 1241
 1967BA1W J. Bar-Touv and C.A. Levinson, Phys. Rev. 153 (1967) 1099
 1967BA40 A.W. Barrows, Jr., F. Gabbard and J.L. Weil, Phys. Rev. 161 (1967) 928
 1967BE1V Bertsch, Bull. Amer. Phys. Soc. 12 (1967) 576
 1967BE1W Benenson, Taylor and Bernard, Bull. Amer. Phys. Soc. 12 (1967) 1191
 1967BE25 P. Bem, J. Habanec, O. Karban and J. Nemeec, Nucl. Phys. A96 (1967) 529
 1967BE36 I. Bergqvist, J.A. Biggerstaff, J.H. Gibbons and W.M. Good, Phys. Rev. 158 (1967) 1049
 1967BI01 G.A. Bissinger, R.M. Mueller, P.A. Quin and P.R. Chagnon, Nucl. Phys. A90 (1967) 1
 1967BL04 D.E. Blatchley and G.E. Thomas, Bull. Amer. Phys. Soc. 12 (1967) 53, CF15
 1967BL19 R. Bloch, R.E. Pixley and P. Truol, Phys. Lett. B25 (1967) 215
 1967BL1J Block, Bartolome, Moyer and Tatarczuk, Bull. Amer. Phys. Soc. 12 (1967) 512
 1967BO09 M.C. Bouten, J.P. Elliott and J.A. Pullen, Nucl. Phys. A97 (1967) 113
 1967BO1P Bock, Grosse-Schulte and Von Oertzen, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 174
 1967BO24 M. Bormann and I. Riehle, Z. Phys. 207 (1967) 64
 1967BR1G B. Bremond, Nucl. Phys. A90 (1967) 241
 1967BR22 C. Broude, A.E. Litherland, R.W. Ollerhead and T.K. Alexander, Can. J. Phys. 45 (1967) 3837

- 1967CH1L Chase, Nucl. Research with Low Energy Accelerators; Eds., Marion and Van Patter (1967) 445
- 1967CS02 J. Csikai and G. Peto, Acta Phys. Acad. Sci. Hung. 23 (1967) 87
- 1967CS03 J. Csikai and S. Nagy, Nucl. Phys. A91 (1967) 222
- 1967DA1G S. Das Gupta and A. Van Ginneken, Phys. Rev. 164 (1967) 1320
- 1967DO1B F. Donau, K. Hehl, C. Riedel, R.A. Broglia and P. Federman, Nucl. Phys. A101 (1967) 495
- 1967EL1F Elwyn, Monahan and Mooring, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 282
- 1967EN01 T. Engeland and P.J. Ellis, Phys. Lett. B25 (1967) 57
- 1967FE01 P. Federman, Nucl. Phys. A95 (1967) 443
- 1967FE1F Feshbach, Kerman and Lemmer, Private Communication (1967)
- 1967FL13 J. Flores and R. Perez, Phys. Lett. B26 (1967) 55
- 1967FL16 F. Flesch and P. Hille, Oesterr. Akad. Wiss. Math-Naturw. Kl. Sitzber. Abt. II, 176 (1967) 45
- 1967GH1A Ghosh, Gupta and Khan, Sci. Res. (Pakistan) 4 (1967) 207
- 1967GU05 M.R. Gunye and C.S. Warke, Phys. Rev. 156 (1967) 1087
- 1967GU14 A.D. Gulko, S.S. Trostin and A. Hudoklin, Yad. Fiz. 6 (1967) 657; Sov. J. Nucl. Phys. 6 (1967) 477
- 1967GU1D M.R. Gunye and C.S. Warke, Phys. Rev. 159 (1967) 885
- 1967HA14 L.F. Hansen, J.D. Anderson, J.W. McClure, B.A. Pohl, M.L. Stelts, J.J. Wesolowski and C. Wong, Nucl. Phys. A98 (1967) 25
- 1967HA1M Halbert, Kim, McGrory and Kuo, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 531
- 1967HA23 L.F. Hansen, H.F. Lutz, M.L. Stelts, J.G. Vidal and J.J. Wesolowski, Phys. Rev. 158 (1967) 917
- 1967HE1D Hershberger, Cummings and Donahue, Bull. Amer. Phys. Soc. 12 (1967) 535
- 1967HS1A L.S. Hsu, Phys. Lett. B25 (1967) 588
- 1967HU06 W.E. Hunt, M.K. Mehta and R.H. Davis, Phys. Rev. 160 (1967) 782
- 1967IV02 M. Ivascu, G. Semencescu, M. Titirici and E. Marincu, Rev. Roum. Phys. 12 (1967) 525
- 1967KA06 C.S. Kalman, J.-P. Bernier and M. Harvey, Can. J. Phys. 45 (1967) 1297
- 1967KE1H I. Kelson, Phys. Rev. 160 (1967) 775
- 1967KO1J Kokame, Fukunaga and Nakamura, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 153

- 1967KR1E H.J. Krappe and H.G. Wahsweiler, Nucl. Phys. A104 (1967) 633
- 1967KU04 J.A. Kuehner and E. Almqvist, Can. J. Phys. 45 (1967) 1605
- 1967KU06 H.M. Kuan, D.W. Heikkinen, K.A. Snover, F. Riess and S.S. Hanna, Phys. Lett. B25 (1967) 217
- 1967KU1E D. Kurath and R.D. Lawson, Phys. Rev. 161 (1967) 915
- 1967KU1L Kuperus, NP 16461 (1967)
- 1967LA1G A. Lande, Phys. Rev. Lett. 18 (1967) 496
- 1967LA1H Lawson and Soper, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 511
- 1967LI07 A.E. Litherland, R.W. Ollerhead, P.J.M. Smulders, T.K. Alexander, C. Broude, A.J. Ferguson and J.A. Kuehner, Can. J. Phys. 45 (1967) 1901
- 1967MC06 R.L. McGrath, S.W. Cospers and J. Cerny, Phys. Rev. Lett. 18 (1967) 243
- 1967ME10 M.K. Mehta, W.E. Hunt and R.H. Davis, Phys. Rev. 160 (1967) 791
- 1967ME22 H.O. Menlove, K.L. Coop, H.A. Grench and R. Sher, Phys. Rev. 163 (1967) 1308
- 1967MO1M J.E. Monahan and A.J. Elwyn, Phys. Rev. 153 (1967) 1148
- 1967MU02 R. Muthukrishnan, Nucl. Phys. A93 (1967) 417
- 1967PA10 M.K. Pal and A.P. Stamp, Phys. Rev. 158 (1967) 924
- 1967PA12 J.C. Parikh, Phys. Lett. B25 (1967) 181
- 1967PA1M Parikh, Lect. on Nucl. Many Body Problems, Herceg Novi (1967)
- 1967PA1P M.K. Pal and A.P. Stamp, Nucl. Phys. A99 (1967) 228
- 1967PA27 A. Pasquarelli, Nucl. Phys. A93 (1967) 218
- 1967PI1B Pinkston, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 419
- 1967PO11 R.M. Polichar, J.E. Steigerwalt, J.W. Sunier and J.R. Richardson, Phys. Rev. 163 (1967) 1084
- 1967QU01 P.A. Quin, A.A. Rollefson, G.A. Bissinger, C.P. Browne and P.R. Chagnon, Phys. Rev. 157 (1967) 991
- 1967RA1E Rahman, Khan and Sen Gupta, Nuovo Cim. B50 (1967) 40
- 1967RA24 N.C. Rasmussen, V.J. Orphan and Y. Hukai, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada 1967 (1967) 278
- 1967RE1B Rees, Harvey and Hendrie, Bull. Amer. Phys. Soc. 12 (1967) 912
- 1967RI1B Ripka, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 833
- 1967SE02 R.E. Segel, Z. Vager, L. Meyer-Schutzmeister, P.P. Singh and R.G. Allas, Nucl. Phys. A93 (1967) 31
- 1967SH14 V.S. Shirley, UCRL 17990 (1967)

- 1967SM04 P.J.M. Smulders, C. Broude and A.E. Litherland, *Can. J. Phys.* 45 (1967) 2133
- 1967SP09 A. Sperduto, *Proc. 3rd Int. Conf. on Atomic Masses, Winnipeg, Canada, 1967* (1967) 657
- 1967ST02 J.E. Stover, *Nucl. Phys.* A92 (1967) 209
- 1967ST1L Stephenson, *Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter* (1967) 433
- 1967ST1N K.W.C. Stewart, *Nucl. Phys.* A100 (1967) 74
- 1967ST25 A.P. Stamp, *Nucl. Phys.* A105 (1967) 627
- 1967ST26 A.P. Stamp, *Phys. Lett.* B26 (1967) 58
- 1967SU05 J.W. Sunier, A.J. Armini, R.M. Polichar and J.R. Richardson, *Phys. Rev.* 163 (1967) 1091
- 1967SV1A Svenne, *Lect. on Nucl. Many Body Problems, Herceg Novi* (1967)
- 1967TH06 W.J. Thompson, S. Edwards and T. Tamura, *Nucl. Phys.* A105 (1967) 678
- 1967VA08 G. van Middelkoop, *Nucl. Phys.* A97 (1967) 209
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, *Nucl. Phys.* A98 (1967) 305
- 1967VO1B Vogt, *Int. Nucl. Phys. Conf., Gatlinburg 1966* (1967) 748
- 1967WI1B Wiedling, *Ark. Fys.* 34 (1967) 467
- 1967YA04 C.T. Yap, *Nucl. Phys.* A100 (1967) 619
- 1967YU01 H.P. Yule, *Nucl. Phys.* A94 (1967) 442
- 1968AB1B Abe, Saito, Endo and Hiura, in *Tokyo* (1968)
- 1968AD1C Adelberger, Nero and Dietrich, *Bull. Amer. Phys. Soc.* 13 (1968) 883
- 1968AN1G Ananthanarayanan, *Nucl. Phys. Solid State Phys. Symp., Powai, India* (1968)
- 1968AR02 A. Arima, S. Cohen, R.D. Lawson and M.H. Macfarlane, *Nucl. Phys.* A108 (1968) 94
- 1968BA1L W.H. Bassichis, B.A. Pohl and A.K. Kerman, *Nucl. Phys.* A112 (1968) 360
- 1968BE1U Becker, *3rd Symp. Struct. Low-Medium Mass Nuclei* (1968) 1
- 1968BI09 M. Bister, A. Fontell and P. Vikberg, *Ann. Acad. Sci. Fennicae, Ser. A VI, No. 268* (1968) 2
- 1968BL1C Blatchley and Thomas, *ANL-7282* (1968) 71
- 1968CA1E Calvi, Cavallaro, Figuera and Sandoli, *Nuovo Cim.* B53 (1968) 485
- 1968CE01 J. Cerny, *Ann. Rev. Nucl. Sci.* 18 (1968) 27
- 1968CE1B Cervera, Garcia and Senent, *An. Real. Soc. Espan. Fis. y Quim.* 64 (1968) 33
- 1968CO11 S. Cohen, E.C. Halbert and S.P. Pandya, *Nucl. Phys.* A114 (1968) 353
- 1968CO1N S. Cohen, R.D. Lawson and S.P. Pandya, *Nucl. Phys.* A114 (1968) 541

- 1968DA1P S. Das Gupta and A. Van Ginneken, Phys. Rev. 174 (1968) 1316
- 1968DR1B R. Dreizler, P. Federman, B. Giraud and E. Osnes, Nucl. Phys. A113 (1968) 145
- 1968DW1A M. Dworzecka and C.S. Warke, Phys. Rev. 175 (1968) 1346
- 1968EL1C Elliott and Wilsdon, Proc. Roy. Soc. 302 (1968) 509
- 1968FA1A J.C. Faivre, H. Krivine and A.M. Papiou, Nucl. Phys. A108 (1968) 508
- 1968FL1C Flores, 3rd Symp. Struct. Low-Medium Mass Nuclei (1968) 188
- 1968GA1J E. Gadioli, I. Iori and L. Zetta, Phys. Rev. 174 (1968) 1140
- 1968GR22 M.W. Greene, Phys. Rev. 176 (1968) 1204
- 1968GU1C M.R. Gunye, Nucl. Phys. A118 (1968) 174
- 1968GU1E M.R. Gunye, Phys. Lett. B27 (1968) 136
- 1968GU1G M.R. Gunye, Nucl. Phys. A120 (1968) 691
- 1968HA17 E.C. Halbert, J.B. McGrory and B.H. Wildenthal, Phys. Rev. Lett. 20 (1968) 1112
- 1968HA1P Halbert, 3rd Symp. Struct. Low-Medium Mass Nuclei (1968) 128
- 1968HA1T Hanna et al., 3rd Symp. Struct. Low-Medium Mass Nuclei (1968) 88
- 1968HA1U Hanna et al., Nucl. Struct., Dubna Symp., 1968 (1968) 266
- 1968HA26 M. Harvey and T. Sebe, Nucl. Phys. A117 (1968) 289
- 1968HE08 D.W. Heikkinen and W. Feldman, Nucl. Phys. A113 (1968) 57
- 1968HE1H K.T. Hecht, P.J. Ellis and T. Engeland, Phys. Lett. B27 (1968) 479
- 1968HI09 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, G.J. Wagner and P. Turek, Nucl. Phys. A115 (1968) 570
- 1968HI14 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, G.J. Wagner and P. Turek, Nucl. Phys. A111 (1968) 265
- 1968HI1H Hinterberger et al., in Tokyo (1968) 274
- 1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, 1967 (1968) 41
- 1968HO1G Horiuchi and Ikeda, Prog. Theor. Phys. 40 (1968) 277; Prog. Theor. Phys. 41(1969) 1386
- 1968HU17 I. Hunyadi and I. Angeli, Nucl. Phys. A119 (1968) 525
- 1968JA1F Jacmart et al., in Tokyo (1968) 209
- 1968JA1J Jacquot et al., Compt. Rend. B266 (1968) 963
- 1968KO05 M. Kobayakawa, H. Niwa, S. Tanemura, K. Suzuki and K. Shinzi, J. Phys. Soc. Jpn. 24 (1968) 645
- 1968KR10 D.E. Kraft and Y.V. Naumov, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 744; Bull. Acad. Sci. USSR Phys. Ser. 32 (1968) 684

1968LA03 B.T. Lawergren, A.T.G. Ferguson and G.C. Morrison, Nucl. Phys. A108 (1968) 325
 1968LA1H Latshaw, O'Connell, Heikkinen and Kuan, Bull. Amer. Phys. Soc. 13 (1968) 883
 1968LE1P F.S. Levin, Nucl. Phys. A115 (1968) 449
 1968ME1E Meyer, Audebert, Elbaz and Lafoucriere, Compt. Rend. B266 (1968) 969
 1968ME1H Meyers, Aldridge and Davis, Bull. Amer. Phys. Soc. 13 (1968) 1713
 1968MI01 R. Middleton, B. Rosner, D.J. Pullen and L. Polsky, Phys. Rev. Lett. 20 (1968) 118
 1968MO1G Moshinsky, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 105
 1968OG1A Ogloblin, Nucl. Struct., Dubna Symp., 1968 (1968) 204
 1968PA12 A.W. Parker, J.S. Allen, R.L. Yerke and V.G. Stotland, Phys. Rev. 174 (1968) 1093
 1968PA1Q Pascual and Flores, Rev. Mex. Fis. 17 (1968) 86
 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
 1968PI1A Pipiraite and Shugurov, Lietuvos Fizikos Rinkinys 8 (1968) 775
 1968RE07 D. Rendic, B. Antolkovic, G. Paic, M. Turk and P. Tomas, Nucl. Phys. A117 (1968) 113
 1968RE1F Reed, UCRL 18414 (1968)
 1968RI08 F.L. Riffle, J.D. Goss, D.R. Parsignault and J.C. Harris, Nucl. Phys. A115 (1968) 120
 1968RI13 L.R. Rice and R.T. Carpenter, Nucl. Phys. A120 (1968) 220
 1968RI1M Ripka, 3rd Symp. Struct. Low-Medium Mass Nuclei (1968) 167
 1968RO1K Rosner, Middleton, Pullen and Polsky, Bull. Amer. Phys. Soc. 13 (1968) 652
 1968SA10 L. Satpathy, Phys. Rev. 174 (1968) 1324
 1968SA1J L. Satpathy and S.C.K. Nair, Phys. Lett. B26 (1968) 257
 1968SH1G R. Shanta and C.S. Shastri, Phys. Rev. 176 (1968) 1254
 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, Nucl. Phys. A113 (1968) 395
 1968SP04 R.H. Spear and I.F. Wright, Aust. J. Phys. 21 (1968) 307
 1968ST06 A.P. Stamp and M.B. Spencer, Nucl. Phys. A111 (1968) 353
 1968SU1C K. Sugawara, Nucl. Phys. A110 (1968) 305
 1968TA1G Takada, Suppl. Prog. Theor. Phys. (1968) 222
 1968TA1Q Takeda, Kato and Yamazaki, in Tokyo (1968) 280
 1968UN1A Une and Yoshida, in Tokyo (1968) 63

- 1968VA20 S.S. Vasilev, T.N. Mikhaleva and D.L. Chuprunov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 32 (1968) 587; *Bull. Acad. Sci. USSR Phys. Ser.* 32 (1969) 539
- 1968VA21 S.S. Vasilev, T.N. Mikhaleva and D.L. Chuprunov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 32 (1968) 594; *Bull. Acad. Sci. USSR Phys. Ser.* 32 (1969) 545
- 1968VO1B Vonach, Vonach, Munzer and Schramel, *Neutron Cross Sections Tech.*, NBS Special Pub. 299 (1968) 885
- 1968WA1M Warke, in *Tokyo* (1968) 62
- 1968WO1A S.S.M. Wong, *Nucl. Phys.* A116 (1968) 609
- 1969AB05 Y. Abgrall, G. Baron, E. Caurier and G. Monsonego, *Nucl. Phys.* A131 (1969) 609
- 1969AB1B A.Y. Abul-Magd, *Nucl. Phys.* A129 (1969) 610
- 1969AD02 E.G. Adelberger, A.B. McDonald and C.A. Barnes, *Nucl. Phys.* A124 (1969) 49
- 1969AG06 J. Aguilar, C.F. Pineda, J. Catala, J.L. Ferrero and A. Garcia, *An. Fis.* 65 (1969) 201
- 1969AK02 Y. Akiyama, A. Arima and T. Sebe, *Nucl. Phys.* A138 (1969) 273
- 1969AN08 J.H. Anderson and R.C. Ritter, *Nucl. Phys.* A128 (1969) 305
- 1969AN1E I. Angeli, J. Csikai and I. Hunyadi, *Phys. Lett.* B29 (1969) 36
- 1969AN1J Anderson, Thesis, Univ. of Virginia (1969)
- 1969AR08 K.P. Artemov, V.Z. Goldberg and V.P. Rudakov, *Yad. Fiz.* 9 (1969) 1173; *Sov. J. Nucl. Phys.* 9 (1969) 686
- 1969AR10 K.P. Artemov, V.Z. Goldberg and V.P. Rudakov, *Yad. Fiz.* 9 (1969) 266; *Sov. J. Nucl. Phys.* 9 (1969) 157
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, *Nucl. Phys.* A137 (1969) 348
- 1969AR1F Arnett and Truran, *Astrophys. J.* 157 (1969) 339
- 1969BA1P Bacon et al., *Acta Cryst.* A25 (1969) 391
- 1969BA1Z Barnes, *Nucl. Isospin, Proc. 1969 Asilomar Conf.* (1969) 179
- 1969BA2E A.I. Baz and V.I. Manko, *Phys. Lett.* B28 (1969) 541
- 1969BA2G M.K. Banerjee, D. d'Oliveira and G.J. Stephenson, *Phys. Rev.* 181 (1969) 1404
- 1969BA2J M.K. Banerjee, C.A. Levinson and G.J. Stephenson, *Phys. Rev.* 178 (1969) 1709
- 1969BA2K J. Bar-Touv, A. Goswami, A.L. Goodman and G.L. Struble, *Phys. Rev.* 178 (1969) 1670
- 1969BA2L Baz, Zhukov and Khrylin, *Yad. Fiz.* 10 (1969) 271
- 1969BA2M Baldeweg, Beckert and Schneidereit, *Zentralin. Kernf. Ross. Dresden, Rept. No. ZFK* 183 (1969)

- 1969BA62 D.J. Baugh, J. Nurzynski, D.M. Rosalky and C.H. Osman, *Aust. J. Phys.* 22 (1969) 555
- 1969BA71 N.A. Bahcall and W.A. Fowler, *Astrophys. J.* 157 (1969) 645
- 1969BE1V H.G. Benson and B.H. Flowers, *Nucl. Phys.* A126 (1969) 305
- 1969BE1X Bethge, Middleton and Scholz, *Bull. Amer. Phys. Soc.* 14 (1969) 530
- 1969BL02 J. Bleck, D.W. Haag, W. Leitz, R. Michaelsen, W. Ribbe and F. Sichelschmidt, *Nucl. Phys.* A123 (1969) 65
- 1969BL1D Blin-Stoyle, *Isospin in Nucl. Phys.*; Ed., D.H. Wilkinson (1969) 115
- 1969BO1Y Boffi and Miglietta, *Nuovo Cim.* B58 (1968) 313
- 1969BO1Z Borse, *Lett. Nuovo Cim.* 2 (1969) 659
- 1969BR1F Bruninx and Crombeen, *Int. J. Appl. Rad. Isotopes* 20 (1969) 255
- 1969BR1M Broude, *Proc. Int. Conf., Montreal* (1969) 221
- 1969CH1K Il-T. Cheon and H.M. Sen Gupta, *Phys. Lett.* B29 (1969) 268
- 1969CH1N S. Chatterjee and A. Chatterjee, *Nucl. Phys.* A125 (1969) 593
- 1969CH1P F.C. Chang, *Phys. Rev.* 178 (1969) 1725
- 1969CH1Q Il-T. Cheon and H.M. Sen Gupta, *Nucl. Phys.* A136 (1969) 567
- 1969CO19 A.A. Cowley and G. Heymann, *Phys. Lett.* B30 (1969) 618
- 1969CO1M Cohenca and Hara, *Contrib., Montreal* (1969) 214
- 1969CU07 R.Y. Cusson, *Can. J. Phys.* 47 (1969) 2835
- 1969DA14 V.V. Davydov and I.M. Pavlichenkov, *Phys. Lett.* B29 (1969) 551
- 1969DE15 R. de Swiniarski, C. Glashausser, D.L. Hendrie, J. Sherman, A.D. Bacher and E.A. McClatchie, *Phys. Rev. Lett.* 23 (1969) 317
- 1969EP1B M. Epherre, E. Gradsztajn, R. Klapisch and H. Reeves, *Nucl. Phys.* A139 (1969) 545
- 1969FA06 A. Faessler and H.H. Wolter, *Z. Phys.* 223 (1969) 192
- 1969FE10 J.L. Ferrero, A. Garcia, J. Milio and F. Senent, *An. Fis.* 65 (1969) 307
- 1969FE1A Federman, *Cargese Lect. in Phys.*; Ed., M. Jean, Vol. 3 (1969) 21
- 1969FO04 W.F. Ford and R.C. Braley, *Nucl. Phys.* A134 (1969) 265
- 1969FO1E W.F. Ford and R.C. Braley, *Nucl. Phys.* A126 (1969) 671
- 1969FU11 G.H. Fuller and V.W. Cohen, *Nucl. Data Tables* A5 (1969) 433
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, *Phys. Rev.* 179 (1969) 921
- 1969GO18 V.Z. Goldberg, V.B. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, *Izv. Akad. Nauk SSSR Ser. Fiz.* 33 (1969) 579; *Bull. Acad. Sci. USSR Phys. Ser.* 33 (1970) 536
- 1969GO1B Gorodetzky, Rudolf, Scheibling and Chevallier, *Contrib., Montreal* (1969) 302

- 1969GR03 H. Grawe and K.P. Lieb, Nucl. Phys. A127 (1969) 13
- 1969GU1E M.R. Gunye, Nucl. Phys. A128 (1969) 457
- 1969HA04 R. Hardell and A. Hasselgren, Nucl. Phys. A123 (1969) 215
- 1969HA14 G. Hauser, R. Lohken, H. Rebel, G. Schatz, G.W. Schweimer and J. Specht, Nucl. Phys. A128 (1969) 81
- 1969HA19 J.C. Hardy, H. Brunnader and J. Cerny, Phys. Rev. Lett. 22 (1969) 1439
- 1969HA1F Hanna, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 591
- 1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
- 1969HA1Y Hanna, Nucl. Isospin Proc. 1969 Asilomar Conf. (1969) 3
- 1969HA1Z Harvey, Proc. Int. Conf., Montreal (1969) 313
- 1969HA38 J.C. Hardy, H. Brunnader, J. Cerny and J. Janecke, Phys. Rev. 183 (1969) 854
- 1969HA45 M. Harvey and T. Sebe, Nucl. Phys. A136 (1969) 459
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
- 1969HE1P Hevezi, Kloplic, Darden and Lietz, Bull. Amer. Phys. Soc. 14 (1969) 186
- 1969HE1T Heikkinen, Lutz, Curtis and Bartolini, Bull. Amer. Phys. Soc. 14 (1969) 852
- 1969HE20 R.L. Hershberger, M.J. Wozniak, Jr. and D.J. Donahue, Phys. Rev. 186 (1969) 1167
- 1969HI1B Hiura, Abe, Saito and Endo, Prog. Theor. Phys. 42 (1969) 555
- 1969HO16 W. Hofmann, R. Kosiek, G. Kraft and R. Mundhenke, Z. Phys. 225 (1969) 303
- 1969HO1W Ho-Kim, Contrib., Montreal (1969) 207
- 1969HO20 T. Holtebekk, S. Tryti and G. Vamraak, Nucl. Phys. A134 (1969) 353
- 1969HO32 R.J.W. Hodgson, Can. J. Phys. 47 (1969) 2269
- 1969JA08 M. Jameel, Phys. Lett. B30 (1969) 155
- 1969JA10 H.U. Jager, Nucl. Phys. A136 (1969) 641
- 1969JO18 J. John, J.P. Aldridge and R.H. Davis, Phys. Rev. 181 (1969) 1455
- 1969JU03 M. Jung, C. Jacquot, C. Baixeras-Aiguabella, R. Schmitt, H. Braun and L. Girardin, Phys. Rev. 188 (1969) 1517
- 1969KE1B Kerman, Cargese Lect. in Phys.; Ed., M. Jean, Vol. 3 (1969) 395
- 1969KO09 C.J. Kost and B. Hird, Nucl. Phys. A132 (1969) 611
- 1969KU1G Kulkarni and Pandya, Nuovo Cim. B60 (1969) 199
- 1969KUZZ D.R. Kulkarni and S.P. Pandya, Nucl. Phys. Solid State Phys. Symp., Abstracts, Roorkee, India (1969) 89
- 1969LA26 H.A. Lamme and E. Boeker, Nucl. Phys. A136 (1969) 609; Erratum Nucl. Phys. A149 (1970) 684

1969LE08 G.M. Lerner and J.B. Marion, Nucl. Instrum. Meth. 69 (1969) 115
 1969LE1P Leonard, Baron, Stewart and Jett, Bull. Amer. Phys. Soc. 14 (1969) 1218
 1969LE1Q Le Tourneux, Proc. Int. Conf., Montreal (1969) 591
 1969LI1F Lindsay, Evans, Toews and Veit, Bull. Amer. Phys. Soc. 14 (1969) 851
 1969MA1T MacFarlane, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 457
 1969MC14 A.B. McDonald, J.R. Patterson and H. Winkler, Nucl. Phys. A137 (1969) 545
 1969MC1A McGill, UCLA P-83 (1969)
 1969MC1C McGrath, Nucl. Isospin Proc. 1969 Ailomar Conf. (1969) 29
 1969NI09 R.J. Nickles, Nucl. Phys. A134 (1969) 308
 1969OH1B Oh, Zaidins and Hayakawa, Bull. Amer. Phys. Soc. 14 (1969) 1227
 1969OS1B Osman, Rosalky and Baugh, Rept. ANU-P/450 (1969)
 1969PA13 R. Padjen and G. Ripka, Phys. Lett. B29 (1969) 548
 1969PA22 J.R. Patterson, H. Winkler and C.S. Zaidins, Astrophys. J. 157 (1969) 367
 1969PI02 J. Pigeon, J. Barguil and E. El Baz, Nucl. Phys. A130 (1969) 49
 1969RI01 R.C. Ritter, J.T. Parson and D.L. Bernard, Phys. Lett. B28 (1969) 588
 1969RU04 F.R. Ruehl, Jr., Nucl. Phys. A136 (1969) 241
 1969SA19 L. Satpathy, D. Goss and M.K. Banerjee, Phys. Rev. 183 (1969) 887
 1969SA1A Sauer, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 717
 1969SA1F P.U. Sauer, A. Faessler, H.H. Wolter and M.M. Stingl, Nucl. Phys. A125 (1969) 257
 1969SC08 D. Schwalm and B. Povh, Phys. Lett. B29 (1969) 103
 1969SC14 J.P. Schiffer, J.A. Nolen, Jr. and N. Williams, Phys. Lett. B29 (1969) 399
 1969SH1E C.S. Shastri, Phys. Rev. 187 (1969) 1737
 1969SM1A Smirnov, Bochum Conf. STI/PUB/232 IAEA (1969) 153
 1969SO08 B. Sorensen, Nucl. Phys. A134 (1969) 1
 1969ST1J Struble and Tewari, Contrib., Montreal (1969) 199
 1969SU12 M. Subotowicz, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 1631; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1496
 1969SV1A Svenne, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 760
 1969TE04 S.N. Tewari, Phys. Lett. B29 (1969) 5
 1969TE1A Temmer, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 693
 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011

- 1969TO1E Towner and Hardy, *Adv. Phys.* 18 (1969) 401
- 1969VO1D Voos et al., *The Phys. and Product. of Heavy Ions, La Plagne, France* (1969)
- 1969VO1E Vogt, *Proc. Int. Conf., Montreal* (1969) 5
- 1969WE1J Weiguny, *Bochum Conf. STI/PUB/232 IAEA* (1969) 284
- 1969WO06 S.K.M. Wong, J. LeTourneux, N. Quang-Hoc and G. Saunier, *Nucl. Phys.* A137 (1969) 318
- 1969WO1G Wong and Davies, *Bull. Amer. Phys. Soc.* 14 (1969) 604
- 1969ZU02 R.W. Zurmuhle and C.M. Fou, *Nucl. Phys.* A129 (1969) 502
- 1970AG08 J. Aguilar, C.F. Pineda, C.R. Bauza, J.L. Ferrero, A. Garcia and F. Senent, *An. Fis.* 66 (1970) 279
- 1970AN1F Angeli, Csikai and Hunyadi, *Acta Phys. Acad. Sci. Hung.* 28 (1970) 629
- 1970AN27 K. Ananthanarayanan, *Nucl. Phys.* A159 (1970) 45
- 1970AR1D Artukh et al., *Nucl. Reactions Included by Heavy Ions, Heidelberg, 1969* (1970) 140
- 1970AR1H Arnold and Bercaw, *Bull. Amer. Phys.* 15 (1970) 630
- 1970AR21 A. Arima, V. Gillet and J. Ginocchio, *Phys. Rev. Lett.* 25 (1970) 1043
- 1970AS1F Ascutto, Braley and Ford, *Bull. Amer. Phys. Soc.* 15 (1970) 1599
- 1970BA1M Bahcall and Fowler, *Astrophys. J.* 161 (1970) 119
- 1970BA2B Baz and Manko, *Sov. J. Nucl. Phys.* 10 (1970) 46
- 1970BA2F Bacher et al., *Bull. Amer. Phys. Soc.* 15 (1970) 543
- 1970BA2H Baz, Zhukov and Khrylin, *Sov. J. Nucl. Phys.* 10 (1970) 155
- 1970BA66 H. Baba, *Nucl. Phys.* A159 (1970) 625
- 1970BE1D Bethge, *Ann. Rev. Nucl. Sci.* 20 (1970) 255
- 1970BO08 T.I. Bonner, *Phys. Rev. C* 1 (1970) 1699
- 1970BO1J Bouten, *Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249* (1970) 361
- 1970BO1K Borbei and Dolinski, *Sov. J. Nucl. Phys.* 10 (1970) 173
- 1970BR35 D.M. Brink, H. Friedrich, A. Weiguny and C.W. Wong, *Phys. Lett.* B33 (1970) 143
- 1970BU25 D. Bucurescu, M. Ivascu, G. Semenescu and M. Titirici, *Rev. Roum. Phys.* 15 (1970) 1093
- 1970CA24 B. Castel, I.P. Johnstone, B.P. Singh and J.C. Parikh, *Nucl. Phys.* A157 (1970) 137
- 1970CL03 R.L. Clarke and P. Perrin, *Nucl. Phys.* A147 (1970) 174
- 1970CL1C Clayton, *2nd Int. Conf. on Nucl. Data for Reactors, Helsinki* (1970) 35
- 1970CO13 A.A. Cowley and G. Heymann, *Nucl. Phys.* A146 (1970) 465
- 1970DA13 M. Danos and B.M. Spicer, *Z. Phys.* 237 (1970) 320

- 1970DO07 E.I. Dolinsky, V.V. Turovtsev and R. Yarmukhamedov, Phys. Lett. B33 (1970) 147
- 1970EI06 J. Eichler and A. Faessler, Nucl. Phys. A157 (1970) 166
- 1970FA01 W.R. Falk, A. Huber, U. Matter, R.W. Benjamin and P. Marmier, Nucl. Phys. A140 (1970) 548
- 1970FA17 W.R. Falk, R.J. Kidney, P. Kulisic and G.K. Tandon, Nucl. Phys. A157 (1970) 241
- 1970FA1F A. Faessler and B. Slavov, Phys. Lett. B31 (1970) 350
- 1970FE09 C. Fernandez Pineda, J. de la Rubia and C. Ruiz Bauza, An. Fis. 66 (1970) 367
- 1970FO1H Fortune, Garrett, Neogy and Middleton, Bull. Amer. Phys. Soc. 15 (1970) 36
- 1970FO1J Fortune et al., Bull. Amer. Phys. Soc. 15 (1970) 483
- 1970FO1K Ford and Braley, Bull. Amer. Phys. Soc. 15 (1970) 1599
- 1970FR14 M.I. Friedman, Nucl. Phys. A159 (1970) 422
- 1970FR1F Frickey and Davis, Bull. Amer. Phys. Soc. 15 (1970) 653
- 1970GA1A Garber et al., BNL 400, 3rd Edition, Vol. 1 (1970)
- 1970GA1K Garrett et al., Nucl. Phys. B24 (1970) 653
- 1970GI11 B. Giraud and P.U. Sauer, Nucl. Phys. A154 (1970) 587
- 1970GI1F B. Giraud, J. Le Tourneux and S. K. M. Wong, Phys. Lett. B32 (1970) 23
- 1970GM1A Gmitro, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 697
- 1970GO1Q P. Goode and S.S.M. Wong, Phys. Lett. B32 (1970) 89
- 1970GO1R D. Goss, Phys. Rev. C2 (1970) 1168
- 1970GU08 K. Gul, B.H. Armitage and B.W. Hooton, Nucl. Phys. A153 (1970) 390
- 1970HA49 M. Harvey and F.C. Khanna, Nucl. Phys. A155 (1970) 337
- 1970HO05 T. Holtebekk, R. Stromme and S. Tryti, Nucl. Phys. A142 (1970) 251
- 1970HO17 A.J. Howard, R.G. Hirko, D.A. Bromley, K. Bethge and J.W. Olness, Phys. Rev. C1 (1970) 1446
- 1970HO1R Howard, Pronko and Hirko, Bull. Amer. Phys. Soc. 15 (1970) 483
- 1970HU07 J. Hudis, T. Kirsten, R.W. Stoenner and O.A. Schaeffer, Phys. Rev. C1 (1970) 2019
- 1970IS1A Ischenko, Reichardt, Voit and Held, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 105
- 1970IV04 M. Ivascu, Stud. Cerc. Fiz. (Rumania) 22 (1970) 477
- 1970KH01 S.B. Khadkikar and M.R. Gunye, Nucl. Phys. A144 (1970) 289
- 1970KR1D S.J. Krieger, Phys. Rev. C1 (1970) 76
- 1970KR1K Krone, Cockburn and Stark, COO-1120-92 (1970)
- 1970KU16 D.R. Kulkarni and S.P. Pandya, Lett. Nuovo Cim. 4 (1970) 133

- 1970LA20 E. Labie, P. Leleux and P. Macq, Ann. Soc. Sci. Brux. Ser. II, 84 (1970) 240
- 1970LI08 R.A. Lindgren, J.G. Pronko, A.J. Howard and D.A. Bromley, Phys. Lett. B32 (1970) 103
- 1970LU17 A. Luukko, A. Anttila, M. Bister and M. Piiparinen, Phys. Scr. 2 (1970) 159
- 1970MA1D Mazarakis and Stephens, Bull. Amer. Phys. Soc. 15 (1970) 629
- 1970MA42 H.B. Mak, H. Spinka and H. Winkler, Phys. Rev. C2 (1970) 1729
- 1970MC04 R.L. McGrath, J. Cerny, J.C. Hardy, G. Goth and A. Arima, Phys. Rev. C1 (1970) 184
- 1970MC1J J.B. McGrory, Phys. Lett. B31 (1970) 339
- 1970MC23 J.B. McGrory, Phys. Lett. B33 (1970) 327
- 1970MI1E Middleton, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 263
- 1970NA07 K. Nakai, F.S. Stephens and R.M. Diamond, Nucl. Phys. A150 (1970) 114
- 1970NE18 P. Neogy, W. Scholz, J. Garrett and R. Middleton, Phys. Rev. C2 (1970) 2149
- 1970NG01 W.Y. Ng, L.E.H. Trainor and S.K.M. Wong, Phys. Lett. B33 (1970) 545
- 1970NI03 R.W. Nightingale, J.A. Becker, R.E. McDonald and D. Kohler, Phys. Rev. C1 (1970) 893
- 1970OA01 N.S. Oakey and R.D. Macfarlane, Phys. Rev. Lett. 25 (1970) 170
- 1970OG1A Ogloblin, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 231
- 1970OL1B Olsen, Thesis, Univ. of Minnesota (1970)
- 1970PA08 A.D. Panagiotou, Phys. Lett. B31 (1970) 361
- 1970PA15 R. Padjen and G. Ripka, Nucl. Phys. A149 (1970) 273
- 1970PA1H Pasichny, Acad. Sci. Ukrainian, SSR, Inst., Theor. Phys. Preprint ITF 70 (1970) 42
- 1970PA27 J.C. Parikh, B. Castel, K.W.C. Stewart and A. Lande, Nucl. Phys. A156 (1970) 501
- 1970PI1G Pineda, De La Rubia and Bauza, An. Fiz. (Spain) 66 (1970) 367
- 1970QU04 P.A. Quin, G.A. Bissinger and P.R. Chagnon, Nucl. Phys. A155 (1970) 495
- 1970QU1A Quin and Vigdor, Bull. Amer. Phys. Soc. 15 (1970) 1686
- 1970RA1A Raisbeck et al., PUC-937-369 (1970)
- 1970RI1C Ripka and Padjen, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 427
- 1970RO06 A.A. Rollefson, P.F. Jones and R.J. Shea, Phys. Rev. C1 (1970) 1761
- 1970RO1B Rowe, Wong and Parikh, Bull. Amer. Phys. Soc. 15 (1970) 527
- 1970RU1A Ruehl, Thesis, Univ. of California, Los Angeles (1970)
- 1970RU1B Ruehl, Chin. J. Phys. 8 (1970) 53
- 1970SC14 B. Schlenk and J. Zimanyi, Acta Phys. Acad. Sci. Hung 28 (1970) 5

- 1970SP01 J. Specht, H. Rebel, G. Schatz, G.W. Schweimer, G. Hauser and R. Lohken, Nucl. Phys. A143 (1970) 373
- 1970ST19 A.P. Stamp, Phys. Lett. B33 (1970) 257
- 1970ST1D Stingl, Sauer, Faessler and Wolter, Nucl. Phys. A145 (1970) 177
- 1970SV1B Svenne, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 413
- 1970TA08 R. Tamisier, C. Lebrun, D. Ardouin, Y. Deschamps, L.-H. Rosier and P. Avignon, Comp. Rend. B270 (1970) 1309
- 1970TE1A S.N. Tewari and G.L. Struble, Phys. Rev. C1 (1970) 1156
- 1970TO1F Toevs, Thesis, CalTech (1970)
- 1970TU01 D.R. Tuerpe, W.H. Bassichis and A.K. Kerman, Nucl. Phys. A142 (1970) 49
- 1970VO01 A.B. Volkov, Nucl. Phys. A141 (1970) 337
- 1970WA39 A. Watt, Nucl. Phys. A159 (1970) 202
- 1970WI05 D.H. Wilkinson and D.E. Alburger, Phys. Rev. Lett. 24 (1970) 1134
- 1970WI13 K.-J. Willkomm and H.V. Klapdor, Z. Phys. 236 (1970) 424
- 1970WO12 C.Y. Wong, Phys. Lett. B32 (1970) 668
- 1970YA01 C.T. Yap, Nucl. Phys. A142 (1970) 161
- 1970ZA02 V.M. Zabegai, A.E. Melenevskii, O.F. Nemets and V.N. Nemykin, Yad. Fiz. 11 (1970) 277; Sov. J. Nucl. Phys. 11 (1970) 155
- 1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AR05 A. Arima and D. Strottman, Nucl. Phys. A162 (1971) 423
- 1971AR1P Artukh et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 193
- 1971AR1R A. Arima and I. Hamamoto, Ann. Rev. Nucl. Sci. 21 (1971) 55
- 1971AR25 A. Arima, M. Sakakura and T. Sebe, Nucl. Phys. A170 (1971) 273
- 1971BA1T Bacher et al., Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haerberli (1971) 708
- 1971BA24 D.P. Balamuth, Phys. Rev. C3 (1971) 1565
- 1971BA33 D.P. Balamuth, J.E. Holden, J.W. Noe and R.W. Zurmuhle, Phys. Rev. Lett. 26 (1971) 1271
- 1971BE17 C. Bergman and R.K. Hobbie, Phys. Rev. C3 (1971) 1729
- 1971BE18 W.L. Bendel, L.W. Fagg, S.K. Numrich, E.C. Jones, Jr. and H.F. Kaiser, Phys. Rev. C3 (1971) 1821

- 1971BE19 R.R. Betts, H.T. Fortune, J.D. Garrett, R. Middleton, D.J. Pullen and O. Hansen, Phys. Rev. Lett. 26 (1971) 1121
- 1971BE1G Beghian, Rogers and Clikeman, Bull. Amer. Phys. Soc. 16 (1971) 579
- 1971BE46 R.F. Bentley, J.D. Carlson, D.A. Lind, R.B. Perkins and C.D. Zafiratos, Phys. Rev. Lett. 27 (1971) 1081
- 1971BO1F Bolsterli, LA-DC 12417 (1971)
- 1971BO1V Bock et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 494
- 1971BO29 M. Bouten, M.C. Bouten, H. Depuydt and L. Schotsmans, Phys. Lett. B35 (1971) 464
- 1971BU02 V.E. Bunakov, K.A. Gridnev and L.V. Krasnov, Phys. Lett. B34 (1971) 27
- 1971BU1E Butler, Poskanzer and Landis, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 210
- 1971CH22 F.C. Chang, Phys. Rev. C3 (1971) 2113
- 1971CU1B Cuzzocrea, Perillo and Notarrigo, Nuovo Cim. A4 (1971) 251
- 1971DE1R de Swiniarski and Raynal, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971DI08 W.T. Diamond, T.K. Alexander and O. Hausser, Can. J. Phys. 49 (1971) 1589
- 1971DI12 G.U. Din and J.L. Weil, Aust. J. Phys. 24 (1971) 139
- 1971DU12 A. Dudek and D.J. Edens, Phys. Lett. B36 (1971) 309
- 1971FA07 W.R. Falk, P. Kulisic and A. McDonald, Nucl. Phys. A167 (1971) 157
- 1971FA08 A. Faessler, K. Goeke and L. Satpathy, Phys. Lett. B35 (1971) 193
- 1971FA13 M.A. Faessler, B. Povh and D. Schwalm, Ann. Phys. 63 (1971) 577
- 1971FO04 W.F. Ford and R.C. Braley, Nucl. Phys. A162 (1971) 513
- 1971FO14 H.T. Fortune, J.D. Garrett, J.R. Powers and R. Middleton, Phys. Rev. C4 (1971) 850; Erratum Phys. Rev. C5 (1972) 1438
- 1971FO19 W.F. Ford, R.C. Braley and J. Bar-Touv, Phys. Rev. C4 (1971) 2099
- 1971FO1P D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 604
- 1971FO24 D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 576
- 1971FR06 H. Friedrich and A. Weiguny, Phys. Lett. B35 (1971) 105
- 1971FR1B J.B. French and J.C. Parikh, Phys. Lett. B35 (1971) 1
- 1971GA13 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. C4 (1971) 98
- 1971GI05 V. Gillet and J.M. Normand, Nucl. Phys. A176 (1971) 225; Erratum Nucl. Phys. A239 (1975) 530
- 1971GO18 D.R. Goosman, K.W. Jones, E.K. Warburton and D.E. Alburger, Phys. Rev. C4 (1971) 1800

- 1971GO1M Gobbi et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 424
- 1971GO1U Gorodetzky, Rudolf, Scheibling and Chevallier, Symp., Saclay (1971)
- 1971GO28 V.Y. Gonchar and K.S. Zheltonog, Yad. Fiz. 13 (1971) 84; Sov. J. Nucl. Phys. 13 (1971) 48
- 1971GO36 V.Y. Gonchar and A.V. Yushkov, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 830; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 764
- 1971GU08 M.R. Gunye, C.S. Warke and S.B. Khadkikar, Phys. Rev. C3 (1971) 1936
- 1971GU1N M.R. Gunye, Phys. Lett. B37 (1971) 125
- 1971GU23 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Semenescu, Phys. Rev. C4 (1971) 1092
- 1971HA14 M. Harvey and A.S. Jensen, Nucl. Phys. A164 (1971) 641
- 1971HA1D Harar, Panagiotou and Gove, Bull. Amer. Phys. 16 (1971) 490
- 1971HA1U Hausser, Private Communication (1971)
- 1971HA26 O. Hausser, T.K. Alexander, A.B. McDonald, G.T. Ewan and A.E. Litherland, Nucl. Phys. A168 (1971) 17
- 1971HA44 P.S. Hauge, S.A. Williams and G.H. Duffey, Phys. Rev. C4 (1971) 1044
- 1971HO19 B.R. Holstein, Phys. Rev. C4 (1971) 740; Erratum Phys. Rev. C9 (1974) 437
- 1971HO1D B.R. Holstein, Phys. Rev. C4 (1971) 764
- 1971HO20 Y. Horikawa, Y. Torizuka, A. Nakada, S. Mitsunobu, Y. Kojima and M. Kimura, Phys. Lett. B36 (1971) 9
- 1971HO23 Q. Ho-Kim, Phys. Rev. C4 (1971) 1077
- 1971JE02 A.S. Jensen and M. Harvey, Can. J. Phys. 49 (1971) 1837
- 1971JO01 I.P. Johnstone, B. Castel and P. Sostegno, Phys. Lett. B34 (1971) 34
- 1971KA18 S. Kardonsky, H.L. Finston and E.T. Williams, Phys. Rev. C4 (1971) 846
- 1971KE10 I. Kelson, Phys. Rev. Lett. 26 (1971) 1386
- 1971KE11 K.W. Kemper, D.S. Haynes and N.R. Fletcher, Phys. Rev. C4 (1971) 108
- 1971KH06 S.B. Khadkikar, Phys. Lett. B36 (1971) 451
- 1971LA13 K.R. Lassey and A.B. Volkov, Phys. Lett. B36 (1971) 4
- 1971LA1D Lang, Z. Phys. 242 (1971) 179
- 1971LA1F Laribi et al., Bull. Amer. Phys. Soc. 16 (1971) 1176
- 1971LE1H T.Y. Lee, S.T. Hsieh, W.W. Yeh and C.T. Chen-Tsai, Phys. Rev. C4 (1971) 64
- 1971LE30 T.Y. Lee, C.T. Chen-Tsai, W.W. Yeh and S.T. Hsieh, Phys. Rev. C4 (1971) 1991
- 1971LI1H H.J. Lipkin, Phys. Rev. Lett. 27 (1971) 432

- 1971MA09 R.D. Macfarlane, N.S. Oakey and R.J. Nickles, Phys. Lett. B34 (1971) 133
- 1971MA1E Mazarakis, DeBolt and Stephens, Bull. Amer. Phys. Soc. 16 (1971) 600
- 1971MA22 M.H. Macfarlane and A.P. Shukla, Phys. Lett. B35 (1971) 11
- 1971MA23 N. Marquardt, W. Von Oertzen and R.L. Walter, Phys. Lett. B35 (1971) 37
- 1971MC03 K.W. McVoy, Phys. Rev. C3 (1971) 1104
- 1971MC04 R.L. McGrath, D.L. Hendrie, E.A. McClatchie, B.G. Harvey and J. Cerny, Phys. Lett. B34 (1971) 289
- 1971MC1F McVoy, Bull. Amer. Phys. Soc. 16 (1971) 1421
- 1971MI02 M.V. Mihailovic, E. Kujawski and J. Lesjak, Nucl. Phys. A161 (1971) 252
- 1971MI09 R. Middleton, J.D. Garrett and H.T. Fortune, Phys. Rev. Lett. 27 (1971) 950
- 1971MI11 R. Middleton, J.D. Garrett, H.T. Fortune and R.R. Betts, Phys. Rev. Lett. 27 (1971) 1296
- 1971MI1J R. Middleton, J.D. Garrett, H.T. Fortune and R.R. Betts, J. Phys. (Paris) 32 (1971) C6-39
- 1971MO15 J.R. Moreira, R.P. Singhal and H.S. Caplan, Can. J. Phys. 49 (1971) 1434
- 1971MO34 C.E. Moss and A.B. Comiter, Nucl. Phys. A178 (1971) 241
- 1971NA17 K. Nagatani, M.J. Le Vine, T.A. Belote and A. Arima, Phys. Rev. Lett. 27 (1971) 1071
- 1971OB1A Ober, Bull. Amer. Phys. Soc. 16 (1971) 192
- 1971PA1C Panagiotou, Gove and Harar, Bull. Amer. Phys. Soc. 16 (1971) 490
- 1971PI02 S. Pittel, Phys. Lett. B34 (1971) 555
- 1971PR09 R. Prasad and D.C. Sarkar, Nuovo Cim. A3 (1971) 467
- 1971PR10 J.G. Pronko and R.W. Nightingale, Phys. Rev. C4 (1971) 1023
- 1971RA16 J.A. Rabbat and G. Do Dang, Phys. Lett. B35 (1971) 480
- 1971RE09 H. Rebel, G.W. Schweimer, J. Specht, G. Schatz, R. Lohken, D. Habs, G. Hauser and H. Klewe-Nebenius, Phys. Rev. Lett. 26 (1971) 1190
- 1971RO06 A.A. Rollefson and J.A. Aymar, Phys. Rev. C3 (1971) 1704
- 1971RO13 D.W.O. Rogers, J.H. Aitken, A.E. Litherland, W.R. Dixon and R.S. Storey, Can. J. Phys. 49 (1971) 1397
- 1971RO33 D.W.O. Rogers, K.W. Allen, H.C. Evans, N.A. Jelley, A.E. Litherland and B.Y. Underwood, Phys. Lett. B37 (1971) 65
- 1971RU1F Rudolph, Bentley and Moss, Bull. Amer. Phys. Soc. 16 (1971) 1154
- 1971SC01 D. Schutte, Nucl. Phys. A161 (1971) 73
- 1971SC12 D. Schwalm, K. Traxel, H. Weiss, M. Morando and B. Povh, Phys. Rev. C4 (1971) 10

1971SE1C D. Sen and C.S. Shastry, Phys. Rev. C4 (1971) 349
 1971ST01 A.P. Stamp, Nucl. Phys. A161 (1971) 81
 1971ST36 D. Strottman, Phys. Lett. B36 (1971) 321
 1971TA05 M. Takeda, S. Kato and T. Yamazaki, J. Phys. Soc. Jpn. 30 (1971) 56
 1971TA20 M. Takeda, S. Kato and T. Yamazaki, J. Phys. Soc. Jpn. 31 (1971) 625
 1971TE06 C.S. Tee and C.T. Yap, Nucl. Phys. A170 (1971) 445
 1971TE1C S.N. Tewari, Phys. Lett. B34 (1971) 366
 1971TO06 J.W. Toevs, Nucl. Phys. A172 (1971) 589
 1971TO1C Toevs, Fowler, Barnes and Lyons, Astrophys. J. 169 (1971) 421
 1971TO1K Torgerson, Wien and MacFarlane, Bull. Amer. Phys. Soc. 16 (1971) 1178
 1971TO1L Torgerson and MacFarlane, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 288
 1971VA1H Vandenbosch, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 337
 1971WI01 WildeB.H. Wildenthal, J.B. McGrory and P.W.M. Glaudemans, Phys. Rev. Lett. 26 (1971) 96
 1971WI07 D.H. Wilkinson, D.E. Alburger, D.R. Goosman, K.W. Jones, E.K. Warburton, G.T. Garvey and R.L. Williams, Nucl. Phys. A166 (1971) 661.
 1971WI18 D.H. Wilkinson, Phys. Rev. Lett. 27 (1971) 1018
 1971WO13 S.K.M. Wong and A.P. Zuker, Phys. Lett. B36 (1971) 437
 1971YA04 C.T. Yap and C.S. Tee, Nucl. Phys. A165 (1971) 497
 1971ZO01 J. Zofka and G. Ripka, Phys. Lett. B34 (1971) 10
 1971ZO03 J. Zofka and G. Ripka, Nucl. Phys. A168 (1971) 65
 1972AB1C Abgrall and Caurier, Private Communication (1972)
 1972AL05 T.K. Alexander, O. Hausser, A.B. McDonald, A.J. Ferguson, W.T. Diamond and A.E. Litherland, Nucl. Phys. A179 (1972) 477
 1972HA07 O. Hausser, A.J. Ferguson, A.B. McDonald, I.M. Szoghy, T.K. Alexander and D.L. Disdier, Nucl. Phys. A179 (1972) 465
 1972HA2H Hardy, Lumpkin and Lee, Bull. Amer. Phys. Soc. 17 (1972) 90
 1972LA1N Laribi et al., Private Communication (1972)
 1972LE1L Lee and Cusson, Private Communication (1972)
 1972MO08 C.E. Moss, C. Detraz, C.S. Zaidins and D.J. Frantsvog, Phys. Rev. C5 (1972) 1122
 1972NO01 J.V. Noble, Phys. Rev. Lett. 28 (1972) 111; Erratum Phys. Rev. Lett. 46 (1981) 1045
 1972RE03 N.E. Reid, G.J. Stephenson and Jr., M.K. Banerjee, Phys. Rev. C5 (1972) 287

1973ENVA P.M. Endt and C. Van der Leun, Nucl. Phys. A214 (1973) 1; Erratum Nucl. Phys.
A248 (1975) 153

BA70W Unknown Source

JO68C Unknown Source

