

Table 11.9 from (2012KE01): ^{10}Be and ^{11}Be γ -ray transition intensities (per 100 decays) observed following ^{11}Li β -decay: measurements prior to DBLA analysis

E_γ (keV)	$^{10}\text{Be}(E_i \rightarrow E_f)$	I^a	I^b	I^c	I^d	I^e
219.2	6180.3 \rightarrow 5961.1	0.78 ± 0.06	0.55 ± 0.10	0.5		0.95 ± 0.35
320.0	$^{11}\text{Be}^*(0.320 \rightarrow 0)$	7.8 ± 0.8	6.3 ± 0.6	7.6 ± 0.8	9.2 ± 0.7	5.2 ± 1.4
2590.3	5958.4 \rightarrow 3368.0	6 ± 1	8.0 ± 1.2	8.5		3.5 ± 1.0
2592.7	5961.1 \rightarrow 3368.0					
2811.8	6180.3 \rightarrow 3368.0	2.8 ± 0.3	0.8 ± 0.2	1.0		1.6 ± 0.7
2896.0	6264.5 \rightarrow 3368.0					
3367.1	3368.0 \rightarrow 0	33 ± 3	29 ± 3	33.3	35 ± 3	21 ± 6
5956.5	5958.4 \rightarrow 0					
5959.2	5961.1 \rightarrow 0					

^a (1997MO35).

^d (1981BJ01).

^b (1997BO01).

^e (1980DE39).

^c (1997AO01, 1997AO04).