

Table 11.17 from (1980AJ01):  
States of  $^{11}\text{B}$  from  $^{11}\text{B}(p, p')^{11}\text{B}^*$   
and  $^{13}\text{C}(d, \alpha)^{11}\text{B}$  <sup>a</sup>

(1971BR41, 1974KA15) <sup>b</sup>	(1970BR23) <sup>c</sup>
$E_x$ (keV)	
0	0
$2124.7 \pm 0.5$	$2125.4 \pm 1.4$
$4445.2 \pm 0.5$	$4444.5 \pm 1.6$
$5021.1 \pm 0.6$	$5020.2 \pm 1.9$
$6743.0 \pm 0.7$ <sup>d</sup>	$6745.8 \pm 3.4$
$6792.6 \pm 1.6$	$6795 \pm 3.0$
$7285.6 \pm 1.5$	
$7978.0 \pm 1.7$	
$8559.4 \pm 1.9$	$8520 \pm 70$
$8920.2 \pm 2.0$	$8910 \pm 60$
$9185.0 \pm 2.0$	
$9274.4 \pm 2.0$	
$10450 \pm 150$ <sup>e</sup>	
$11650 \pm 150$	
$12850 \pm 100$	
$15200 \pm 150$	
$16400 \pm 150$	

<sup>a</sup> See also Table 11.18 in (1975AJ02).

<sup>b</sup>  $^{11}\text{B}(p, p')^{11}\text{B}$ .

<sup>c</sup>  $^{13}\text{C}(d, \alpha)^{11}\text{B}$ .

<sup>d</sup> Values below are normalized to  $E_x = 4445.3, 5020.0$  and  $6743.4$  keV.

<sup>e</sup> This value and the values below are from (1969SU03).