

Table 16.6 from (1982AJ01):
Excited states in ^{16}N from $^{14}\text{C}(^3\text{He}, \text{p})^{16}\text{N}$

E_x (MeV \pm keV)		Γ (keV)	$J^\pi; T$
(1966GA08)	(1968HE03, 1972FR05)		
0.121 \pm 6			0 ⁻
0.298 \pm 6			3 ⁻
0.396 \pm 7			
3.348 \pm 7			1 ⁺
3.517 \pm 7			2 ⁺ , (3) ⁺
3.958 \pm 7			(2) ⁺ , 3 ⁺
4.313 \pm 9			1 ⁺
4.386 \pm 9			
4.768 \pm 11			
5.052 \pm 9			
5.137 \pm 9			
5.234 \pm 9			(1, 2, 3) ⁺
	5.512 \pm 5		(1, 2, 3) ⁺
	5.724 \pm 5		5 ⁺
	6.168 \pm 5		
	6.843 \pm 5		
	7.113 \pm 5		
	7.570 \pm 5		
	7.636 \pm 5		
	7.673 \pm 5		
	8.205 \pm 5		
	9.760 \pm 10	15 \pm 8	$T = 1$
	9.813 \pm 10		$T = 1$
	9.928 \pm 7	< 12	0 ⁺ ; 2
	11.701 \pm 7	< 12	1 ⁻ , 2 ⁺ ; 2