

Table 16.8 from (1986AJ04): Resonances in  $^{15}\text{N}(n, n)^{15}\text{N}$  <sup>a,b</sup>

| $E_n$<br>(MeV $\pm$ keV)   | $\Gamma_{\text{lab}}$ (keV) | $E_x$ (MeV) | $J^\pi$                     |
|----------------------------|-----------------------------|-------------|-----------------------------|
| 0.921                      | 14                          | 3.354       | 1 <sup>+</sup> <sup>c</sup> |
| 1.095                      | 3                           | 3.517       | 1                           |
| 1.563                      | $\leq 2$                    | 3.955       | 1                           |
| 1.944                      | 29                          | 4.312       | 1 <sup>+</sup> <sup>d</sup> |
| 2.038                      | 56                          | 4.400       | 1 <sup>-</sup> <sup>d</sup> |
| 2.30 $\pm$ 70 <sup>e</sup> | 410 $\pm$ 100 <sup>e</sup>  | 4.65        | 1 <sup>-</sup> <sup>d</sup> |
| 2.399                      | 107                         | 4.738       | 2 <sup>+</sup> <sup>d</sup> |
| 2.732                      | 35                          | 5.050       | 1 <sup>-</sup>              |
| 2.830                      | 12                          | 5.142       | 3 <sup>(-)</sup>            |
| 2.84 $\pm$ 70 <sup>f</sup> | 710 $\pm$ 100 <sup>f</sup>  | 5.15        | 2 <sup>-</sup> <sup>d</sup> |
| 2.915                      | 4                           | 5.222       | $\geq 2$                    |
| 2.93                       | 260                         | 5.24        | 1 <sup>+</sup>              |
| 3.225                      |                             | 5.512       |                             |
| 3.454                      | 24                          | 5.727       | 1 <sup>+</sup>              |
| 3.69                       | 297                         | 5.95        | 1 <sup>-</sup>              |
| 3.987                      | 88                          | 6.226       | (1 <sup>+</sup> )           |
| 4.126                      | 78                          | 6.356       | (3 <sup>-</sup> )           |
| 4.252                      | 113                         | 6.474       | (2 <sup>+</sup> )           |
| 4.64                       | $> 150$                     | 6.84        | $\geq 2$                    |
| 4.80                       | 37                          | 6.99        | $\geq 1$                    |
| 5.055                      | 25                          | 7.227       | $\geq 2$                    |
| 5.43                       | 30                          | 7.58        | $\geq 3$                    |
| 5.56                       |                             | 7.70        |                             |
| 5.73                       | 165                         | 7.86        | $\geq 4$                    |
| 5.90                       |                             | 8.02        |                             |
| 6.28                       |                             | 8.37        | $\geq 1$                    |
| 6.42                       |                             | 8.51        | $\geq 1$                    |
| 6.65                       | 45                          | 8.72        | $\geq 1$                    |
| 6.76                       |                             | 8.82        |                             |
| 7.10                       | 110                         | 9.14        | $\geq 2$                    |

Table 16.8 from (1986AJ04): Resonances in  $^{15}\text{N}(n, n)^{15}\text{N}$  <sup>a,b</sup>(continued)

| $E_n$<br>(MeV $\pm$ keV) | $\Gamma_{\text{lab}}$ (keV) | $E_x$ (MeV) | $J^\pi$  |
|--------------------------|-----------------------------|-------------|----------|
| 7.31                     |                             | 9.34        |          |
| 7.44                     | 105                         | 9.46        | $\geq 2$ |
| 7.71                     | 150                         | 9.71        | $\geq 2$ |
| 8.07                     | 30                          | 10.05       | $\geq 3$ |
| 8.30                     | 175                         | 10.27       | $\geq 2$ |
| 8.77                     | 130                         | 10.71       | $\geq 2$ |
| 9.61                     |                             | 11.49       | $\geq 3$ |
| 9.77                     |                             | 11.64       | $\geq 3$ |
| 10.25                    |                             | 12.09       |          |
| 10.64                    |                             | 12.46       |          |
| 11.09                    |                             | 12.88       |          |
| 11.41                    |                             | 13.12       |          |
| 12.10                    |                             | 13.83       |          |

<sup>a</sup> For references see [Table 16.7 in \(1977AJ02\)](#).

<sup>b</sup> Below  $E_n = 4.5$  MeV, the multilevel  $R$ -matrix formalism was used to determine  $E_\lambda$ ,  $\Gamma_\lambda$  and whenever possible  $J^\pi$  by a  $\chi^2$  fitting and minimization technique. Above this energy the  $2J + 1$  dependence was used; the parity cannot be determined because no marked interference effects are observed between resonance and potential scattering. Above 5.65 MeV all  $J$ -values are lower limits because the inelastic channel is open. [A channel radius  $a = 4.69$  fm was used.]

<sup>c</sup> Parity determined from angular distribution.

<sup>d</sup>  $J^\pi$  also obtained by phase-shift analysis.

<sup>e</sup> The phase-shift analysis indicates that the resonance is at  $E_n = 2.42 \pm 0.08$  MeV with  $\Gamma = 250 \pm 50$  keV. This is one of two ( $d_{3/2}p_{1/2}^{-1}$ ) single-particle resonances.

<sup>f</sup> The phase-shift analysis finds  $E_\lambda = 2.94 \pm 0.1$  MeV,  $\Gamma = 320 \pm 80$  keV. This is the other ( $d_{3/2}p_{1/2}^{-1}$ ) single-particle resonance.