

Table 17.14 from (1986AJ04):  
Some inelastic groups observed in  $^{17}\text{O}(e, e)^a$

$E_x$ (MeV)	$\Gamma$ (keV)	$E_x$ (MeV)	$\Gamma$ (keV)
$11.71 \pm 0.05^b$	narrow	$14.76 \pm 0.10^b$	$> 300$
$11.95 \pm 0.05^b$	$\approx 250$	$15.24 \pm 0.10^b$	$\approx 200$
$12.22 \pm 0.02^c$	$\leq 20$	$16.52 \pm 0.05^b$	$\approx 300$
$12.66 \pm 0.05^b$	$\approx 90$	$17.92 \pm 0.02^c$	$98 \pm 16$
$12.96 \pm 0.05^b$	$\approx 200$	$18.72 \pm 0.02^c$	$87 \pm 33$
$13.56 \pm 0.05^b$	$\approx 150$	22.0 <sup>b,d</sup>	
$14.14 \pm 0.10^b$	$\approx 100$	23.0 <sup>b,d</sup>	
$14.72 \pm 0.02^c$	$35 \pm 11$		

<sup>a</sup> See also [Table 17.13](#) for other inelastic groups.

<sup>b</sup> (1977NO06).

<sup>c</sup> (1986MA48) and D.M. Manley, private communication. I am very indebted to Dr. Manley for his many useful comments.

<sup>d</sup> C1.