

Table 18.6 from (1978AJ03): States in  $^{18}\text{O}$  from  $^{16}\text{O}(t, p)^{18}\text{O}$

$E_x$ (MeV $\pm$ keV)		$L^b$	$J^\pi$
(1960JA17)	(1962HI06)		
0	0	0	$0^+$
$1.979 \pm 5$	$1.980 \pm 5$	2	$2^+$
$3.552 \pm 5$	$3.549 \pm 5$	3 or 4	$3^-$ or $4^+$
$3.634 \pm 5$	$3.627 \pm 5$	0	$0^+$
$3.915 \pm 5$	$3.915 \pm 5$	2	$2^+$
$4.448 \pm 5$	$4.449 \pm 5$	(3)	<sup>c</sup>
	$5.090 \pm 5$	3	$3^-$
	$5.247 \pm 7$	2	$2^+$
	$5.329 \pm 7$	0	$0^+$
	$5.368 \pm 10$	(2)	$(2^+)^d$
	$5.521 \pm 10$		
	$6.189 \pm 10$		
	$6.341 \pm 10$		
	$6.391 \pm 10$		

<sup>a</sup> Observed but not measured.

<sup>b</sup> From PWBA analysis of angular distributions: see (1964MI05). See also (1960JA17).

<sup>c</sup>  $J^\pi = 1^-$ . See, e.g. discussion in (1964MI05).

<sup>d</sup> See, however, [reaction 19](#).