

Table 18.15 from (1995TI07): States in ^{18}O from $^{16}\text{O}(t, p)$ ^a

E_x (keV)	L	J^π	E_x (keV)	E_x (keV)
0	0	0^+	7623 ± 18	9713 ± 7
1986 ± 4	2	2^+	7782 ± 6	9890 ± 11
3556 ± 2	4	4^+	7871 ± 2 ^d	10120 ± 40
3634 ^b	0	0^+	7983 ± 3 ^d	10300 ± 20
3915 ± 2	2	2^+	8046 ± 7	10400 ± 10
4458 ± 3	1	1^-	8140 ± 10	10610 ± 20
5105 ± 2	3	3^-	8233 ± 9	
5258 ± 6	2	2^+	8294 ± 5 ^d	
5340 ± 4	0	0^+	8430 ± 12	
5382 ± 4			8521 ± 3 ^d	
5530 ± 4			8660 ± 6	
6197 ± 3	1	1^-	9030 ± 15 ^e	
6356 ± 7	1, 2	$(1^-, 2^+)$ ^c	9362 ± 5 ^d	
6399 ± 3	3	3^-	9420 ± 20	
6885 ± 9			9480 ± 30	
7123 ± 7	4	4^+	9671 ± 8	

^a (1981CO13): $E_t = 15$ MeV; DWBA analysis. See also Table 18.6 in (1978AJ03).

^b Nominal energy.

^c See, however, Table 18.18.

^d Comparisons of E_x shown here with those displayed in Table 18.3 for $^{18}\text{O}^*(3.92, 5.10, 6.40, 7.77)$ suggest that the uncertainty shown may be low: ± 6 keV was arbitrarily used in calculating the best value for E_x for this state in Table 18.3 of (1987AJ02).

^e This is the “average” of several unresolved levels. (1985FO11) states that the main components are at 8.96 and 9.03 MeV. [Comment: It is not clear whether these states are actually resolved (1987AJ02).]