

Table 19.4 from (1983AJ01):
Levels of ^{19}O from $^{17}\text{O}(t, p)$ and $^{18}\text{O}(d, p)$

E_x (MeV \pm keV) ^a	$\Gamma_{c.m.}$ (keV) ^a	l_n ^b	L ^c	S ^d	J^π
0		2	0	0.57	$\frac{5}{2}^+$
0.0960 ± 0.5		2	2		$\frac{3}{2}^+$
1.4719 ± 0.5		0	2	1.00	$\frac{1}{2}^+$
2.3715 ± 1.0		2	(2 + 4)		$\frac{5}{2}^+$
2.7790 ± 0.9		(2)	2		$\frac{7}{2}^+$
3.0671 ± 2.6		2	(0 + 2)	(0.06)	$\frac{5}{2}^+$
3.1535 ± 2.4		a			$\frac{3}{2}^+$
3.237 ± 5		1		0.11	$\frac{3}{2}^-$
3.944 ± 3	< 15	2	(2)	0.03	$\frac{3}{2}^+$
4.118 ± 5	< 15				
4.333 ± 12					
4.402 ± 12	75 ± 5	1		0.15	$\frac{3}{2}^-$
4.584 ± 12	< 15	2	a	0.02	$\frac{5}{2}^+$
4.707 ± 12	< 15				
4.998 ± 12	< 15	2	a	0.08	$\frac{5}{2}^+$
5.150 ± 10	320 ± 25	2	(2 + 4)	0.85	$\frac{3}{2}^+$
5.455 ± 10	45				
5.502 ± 12	< 15	2		0.17	$(\frac{3}{2})^+$
5.714 ± 12	< 15	3		0.13	$(\frac{7}{2})^-$
6.280 ± 12					
6.480 ± 15					
6.560 ± 15					
6.899 ± 15					
6.997 ± 15					
7.117 ± 15					
7.248 ± 15					

^a For references and other values see [Table 19.3 in \(1978AJ03\)](#).

^b $^{18}\text{O}(d, p)^{19}\text{O}$.

^c $^{17}\text{O}(t, p)^{19}\text{O}$.

^d $E_{\bar{d}} = 14.8$ MeV: polarization and differential cross section measurements. The spectroscopic factors for the states with $E_x > 4.1$ MeV have been calculated in the weakly bound approximation.