

Table 19.15 from (1978AJ03): Resonances in $^{18}\text{O}(p, \gamma)^{19}\text{F}$ ^a

E_p (keV)	Γ_{lab} (keV)	J^π	E_x (MeV)	Refs.
629.6 ± 0.3	2.0 ± 0.3	$\frac{3}{2}^+$	8.5896	A
848 ± 2	40 ± 5	$\frac{3}{2}$	8.796	A
1166.5 ± 0.4	$(25 \pm 24) \times 10^{-3}$	$\frac{7}{2}^+$ ^b	9.0980	A
1398 ± 3	4		9.317	A
1685 ± 5 ^d	< 15		(9.589)	(1959BU05)
1769 ± 2	4.0 ± 1.0	$\frac{3}{2}^+$	9.669	(1959BU05, 1962NE03)
1778			(9.677)	(1962NE03)
1790			(9.688)	(1962NE03)
1928.4 ± 0.6 ^c	0.3 ± 0.05	$\frac{5}{2}^f$	9.819	A
2263.0 ± 0.7	5.0 ± 1.0	$\frac{3}{2}^-$	10.136	(1962NE03, 1969DU1A, 1971WO12, 1972WO15)
2.36 ^e			(10.23)	(1962NE03)
2.39	47 ± 10		(10.26)	(1962NE03)
2.41	10 ± 5		(10.28)	(1962NE03)
2.44			(10.30)	(1962NE03)
(2.60)			(10.46)	(1962NE03)
(2.66)			(10.51)	(1962NE03)
(2.68)			(10.53)	(1962NE03)
(2.73)			(10.58)	(1962NE03)
(2.77)			(10.62)	(1962NE03)
(2.80)			(10.64)	(1962NE03)
(2.84)			(10.68)	(1962NE03)

A: See references for this state in [Table 19.12 in \(1972AJ02\)](#).

^a See also [Table 19.7](#).

^b Most probable value, although $J^\pi = \frac{9}{2}^+$ is also possible: see text. $T = \frac{3}{2}$ ([1965AL20](#)).

^c Γ_γ and Γ_p are \lesssim few eV ([1969DU1A](#)).

^d See, however, ([1962NE03](#)).

^e See ([1962NE03](#)) for additional resonant structure between $E_p = 2.33$ and 2.78 MeV.

^f From γ -ray angular distributions (I.E. Wright, private communication).