

Table 20.1 from (1972AJ02): Energy levels of  $^{20}\text{O}$

| $E_x$ (MeV $\pm$ keV) | $J^\pi; T$ | $\tau_{1/2}$ (sec) | Decay      | Reactions |
|-----------------------|------------|--------------------|------------|-----------|
| 0                     | $0^+; 2$   | $13.57 \pm 0.1$    | $\beta^-$  | 1, 2, 3   |
| $1.672 \pm 5$         | $2^+$      |                    | $\gamma$   | 2         |
| $3.568 \pm 5$         | $4^+$      |                    | $(\gamma)$ | 2         |
| $4.065 \pm 5$         | $2^+$      |                    | $(\gamma)$ | 2         |
| $4.446 \pm 7$         | $0^+$      |                    | $(\gamma)$ | 2         |
| $4.838 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $4.997 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $5.220 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $5.298 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $5.382 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $5.603 \pm 7$         |            |                    | $(\gamma)$ | 2         |
| $(5.83 \pm 20)$       |            |                    | $(\gamma)$ | 2         |