

Table 20.10 from (1972AJ02):
States of ^{20}F involved in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

| E_x (keV) | |
|--------------|--------------|
| (1968SP01) | (1969HA04) |
| 0 | 0 |
| 656.3 ± 0.3 | 656 ± 1 |
| 822.9 ± 0.3 | |
| 983.8 ± 0.3 | 984 ± 1 |
| 1057.2 ± 0.3 | 1057 ± 1 |
| 1309.1 ± 0.3 | 1309 ± 1 |
| 1843.4 ± 0.3 | 1843 ± 1 |
| 1970.6 ± 0.3 | |
| 2044.2 ± 0.4 | 2044 ± 1 |
| 2194.5 ± 0.6 | 2194 ± 2 |
| 2965.8 ± 0.5 | 2966 ± 2 |
| 3488.3 ± 0.3 | 3488 ± 2 |
| 3526.0 ± 0.5 | |
| 3587.3 ± 0.3 | 3588 ± 2 |
| 3681.0 ± 0.4 | 3681 ± 2 |
| | 3967 ± 2 |
| 4082.2 ± 0.5 | 4085 ± 2 |
| 4276.7 ± 0.5 | 4275 ± 2 |
| 5413.1 ± 0.6 | |
| 5554.7 ± 6 | |
| | 5713 ± 2 |
| 5936.0 ± 0.3 | 5937 ± 1 |
| 6017.3 ± 0.3 | 6018 ± 1 |
| 6044.6 ± 0.4 | |
| 6601.1 ± 0.3 | 6602.0 ± 0.6 |

^a For measurements of I_γ , see (1968SP01, 1969HA04).