

Table 20.10 from (1983AJ01):  
States of  $^{20}\text{F}$  involved in  $^{19}\text{F}(n, \gamma)^{20}\text{F}$  <sup>a</sup>

$E_x$ (keV)	
0	$3526.0 \pm 0.5$
$656.3 \pm 0.3$ <sup>b</sup>	$3587.3 \pm 0.3$
$822.9 \pm 0.3$	$3681.0 \pm 0.4$
$983.8 \pm 0.3$ <sup>b</sup>	$3967 \pm 2$ <sup>c</sup>
$1057.2 \pm 0.3$	$4082.2 \pm 0.5$
$1309.1 \pm 0.3$	$4276.7 \pm 0.5$
$1843.4 \pm 0.3$	$5413.1 \pm 0.6$
$1970.6 \pm 0.3$	$5554.7 \pm 0.6$
$2044.2 \pm 0.4$	$5713 \pm 2$ <sup>c</sup>
$2194.5 \pm 0.6$	$5936.0 \pm 0.3$
$2965.8 \pm 0.5$	$6017.3 \pm 0.3$
$3488.3 \pm 0.3$	$6044.6 \pm 0.4$
	$6601.1 \pm 0.3$ <sup>d</sup>

<sup>a</sup> (1968SP01).

<sup>b</sup>  $656.1 \pm 0.3$  and  $983.4 \pm 0.4$  keV (1972OP01).

<sup>c</sup> (1969HA04).

<sup>d</sup>  $6602.0 \pm 0.6$  keV (1969HA04).