

Table 20.11 from (1998TI06): States of ^{20}F involved in $^{19}\text{F}(\text{n}, \gamma)^{20}\text{F}$ ^a

E_x (keV)	E_x (keV)	E_x (keV)	E_x (keV)
0	2864.86 ± 0.10	4082.17 ± 0.04	5555.34 ± 0.04
656.02 ± 0.03	2966.11 ± 0.03	4277.09 ± 0.04	5623.13 ± 0.06
822.73 ± 0.03	3171.69 ± 0.14	4371.47 ± 0.11	5810.1 ± 0.4
983.59 ± 0.03	3488.41 ± 0.03	4591.72 ± 0.07	5936.13 ± 0.03
1056.82 ± 0.03	3526.31 ± 0.04	4892.76 ± 0.17	5939.10 ± 0.10
1309.19 ± 0.03	3586.54 ± 0.03	5226.1 ± 0.4	6017.78 ± 0.03
1843.80 ± 0.03	3589.80 ± 0.04	5282.79 ± 0.10	6044.92 ± 0.03
1970.83 ± 0.04	3680.17 ± 0.04	5319.17 ± 0.04	6299.1 ± 0.3
2043.98 ± 0.03	3965.07 ± 0.04	5465.89 ± 0.17	6601.35 ± 0.03
2194.30 ± 0.03			

^a (1996RA04). For the earlier work see Tables 20.11 in (1978AJ03) and 20.8 in (1987AJ02).