

Table 2 from (1990AJ01): Some electromagnetic transitions in  $A = 11 - 12$  <sup>a</sup>

Nucleus	$E_{\text{xi}} \rightarrow E_{\text{xf}}$ (MeV)	$J_1^\pi \rightarrow J_f^\pi$ <sup>b</sup>	$\Gamma_\gamma$ (eV)	Mult.	$S$ (W.u.)
<sup>11</sup> Be	0.32 $\rightarrow$ 0	$\frac{1}{2}^- \rightarrow \frac{1}{2}^+$	$(3.97 \pm 0.36) \times 10^{-3}$	E1	$0.360 \pm 0.033$
<sup>11</sup> B <sup>c</sup>	2.125 $\rightarrow$ 0	$\frac{1}{2}^- \rightarrow \frac{3}{2}^-$	$0.120 \pm 0.009$	M1	$0.60 \pm 0.04$
	4.445 $\rightarrow$ 0	$\frac{5}{2}^- \rightarrow \frac{3}{2}^-$	$0.54 \pm 0.02$	M1	$0.29 \pm 0.01$
	5.020 $\rightarrow$ 0 $\rightarrow$ 2.125	$\frac{3}{2}^- \rightarrow \frac{3}{2}^-$	$1.68 \pm 0.06$	M1	$0.63 \pm 0.02$
		$\rightarrow \frac{1}{2}^-$	$0.28 \pm 0.01$	M1	$0.55 \pm 0.02$
	6.743 $\rightarrow$ 0 $\rightarrow$ 4.445	$\frac{7}{2}^- \rightarrow \frac{3}{2}^-$	$(2.1 \pm 0.5) \times 10^{-2}$	E2	$1.3 \pm 0.3$
		$\rightarrow \frac{5}{2}^-$	$(9.0 \pm 2.3) \times 10^{-3}$	M1	$(3.5 \pm 0.9) \times 10^{-3}$
	6.792 $\rightarrow$ 0 $\rightarrow$ 2.125	$\frac{1}{2}^+ \rightarrow \frac{3}{2}^-$	$0.26 \pm 0.03$	E1	$(2.5 \pm 0.3) \times 10^{-3}$
		$\rightarrow \frac{1}{2}^-$	$0.110 \pm 0.014$	E1	$(3.21 \pm 0.41) \times 10^{-3}$
	$\rightarrow$ 5.020	$\rightarrow \frac{3}{2}^-$	$(1.54 \pm 0.21) \times 10^{-2}$	E1	$(8.2 \pm 1.1) \times 10^{-3}$
		$\rightarrow \frac{5}{2}^-$	$1.00 \pm 0.08$	E1	$(7.7 \pm 0.6) \times 10^{-3}$
	7.286 $\rightarrow$ 0 $\rightarrow$ 4.445	$\frac{5}{2}^+ \rightarrow \frac{3}{2}^-$	$(6.3 \pm 1.2) \times 10^{-2}$	E1	$(8.2 \pm 1.6) \times 10^{-3}$
		$\rightarrow \frac{5}{2}^-$	$(8.0 \pm 1.2) \times 10^{-2}$	E1	$(2.1 \pm 0.3) \times 10^{-2}$
	7.978 $\rightarrow$ 0 $\rightarrow$ 2.125	$\frac{3}{2}^+ \rightarrow \frac{3}{2}^-$	$0.53 \pm 0.07$	E1	$(3.1 \pm 0.4) \times 10^{-3}$
		$\rightarrow \frac{1}{2}^-$	$0.61 \pm 0.08$	E1	$(9.1 \pm 1.2) \times 10^{-3}$
	$\rightarrow$ 7.286	$\rightarrow \frac{1}{2}^+$	$(9.8 \pm 1.4) \times 10^{-3}$	M1	$1.4 \pm 0.2$
		$\rightarrow \frac{3}{2}^-$	$0.53 \pm 0.05$	M1	$(4.0 \pm 0.4) \times 10^{-2}$
	8.560 $\rightarrow$ 0 $\rightarrow$ 2.125	$\frac{3}{2}^- \rightarrow \frac{1}{2}^-$	$0.28 \pm 0.03$	M1	$(5.1 \pm 0.5) \times 10^{-2}$
		$\rightarrow \frac{5}{2}^-$	$(4.7 \pm 1.1) \times 10^{-2}$	M1	$(3.2 \pm 0.7) \times 10^{-2}$
	$\rightarrow$ 4.445 $\rightarrow$ 5.020	$\rightarrow \frac{3}{2}^-$	$(8.5 \pm 1.2) \times 10^{-2}$	M1	$(9.1 \pm 1.3) \times 10^{-2}$
		$\rightarrow \frac{5}{2}^-$	$4.10 \pm 0.20$	M1	$0.28 \pm 0.01$
8.920 $\rightarrow$ 0 $\rightarrow$ 4.445	$\frac{5}{2}^- \rightarrow \frac{5}{2}^-$	$(5.0 \pm 3.6) \times 10^{-2}$	E2	$0.7 \pm 0.5$	
	$\rightarrow \frac{5}{2}^-$	$0.22 \pm 0.02$	M1	$0.12 \pm 0.01$	
9.185 $\rightarrow$ 0 $\rightarrow$ 4.445	$\frac{7}{2}^+ \rightarrow \frac{3}{2}^-$	$(2.7 \pm 1.2) \times 10^{-3}$	M2	$0.56 \pm 0.25$	
	$\rightarrow \frac{5}{2}^-$	$0.25 \pm 0.09$	E1	$(7.0 \pm 2.5) \times 10^{-3}$	
$\rightarrow$ 6.743	$\rightarrow \frac{7}{2}^-$	$(3.8 \pm 1.3) \times 10^{-2}$	E1	$(7.8 \pm 2.7) \times 10^{-3}$	
	$\rightarrow \frac{5}{2}^-$	$(6.40 \pm 0.45) \times 10^{-2}$	M1	$0.38 \pm 0.03$	
<sup>11</sup> C	2.000 $\rightarrow$ 0	$\frac{1}{2}^- \rightarrow \frac{3}{2}^-$	$0.26 \pm 0.06$	M1	$(2.3 \pm 0.5) \times 10^{-2}$
	8.105 $\rightarrow$ 0	$\frac{3}{2}^- \rightarrow \frac{1}{2}^-$	$(9.1 \pm 2.3) \times 10^{-2}$	M1	$(1.9 \pm 0.5) \times 10^{-2}$
	8.420 $\rightarrow$ 0	$\frac{5}{2}^- \rightarrow \frac{3}{2}^-$	$3.1 \pm 1.3$	M1	$0.25 \pm 0.10$
<sup>12</sup> B <sup>d</sup>	0.953 $\rightarrow$ 0	$2^+ \rightarrow 1^+$	$(2.53 \pm 0.40) \times 10^{-3}$	M1	$0.14 \pm 0.02$
<sup>12</sup> C <sup>e</sup>	4.439 $\rightarrow$ 0	$2^+ \rightarrow 0^+$	$(1.08 \pm 0.06) \times 10^{-2}$	E2	$4.65 \pm 0.26$
	7.654 $\rightarrow$ 4.439	$0^+ \rightarrow 2^+$	$(3.7 \pm 0.5) \times 10^{-3}$	E2	$8.0 \pm 1.1$
	9.641 $\rightarrow$ 0	$3^- \rightarrow 0^+$	$(3.1 \pm 0.4) \times 10^{-4}$	E3	$12 \pm 2$
	12.71 $\rightarrow$ 0	$1^+ \rightarrow 0^+$	$0.35 \pm 0.05$	M1	$(8.1 \pm 1.2) \times 10^{-3}$

Table 2 from (1990AJ01): Some electromagnetic transitions in  $A = 11 - 12$  <sup>a</sup> (continued)

Nucleus	$E_{xi} \rightarrow E_{xf}$ (MeV)	$J_i^\pi \rightarrow J_f^\pi$ <sup>b</sup>	$\Gamma_\gamma$ (eV)	Mult.	$S$ (W.u.)
	$\rightarrow 4.439$	$\rightarrow 2^+$	$(5.3 \pm 1.0) \times 10^{-2}$	M1	$(4.5 \pm 0.8) \times 10^{-3}$
	15.11 $\rightarrow 0$	$1^+; 1 \rightarrow 0^+; 0$	$38.5 \pm 0.8$	M1	$0.531 \pm 0.011$
	$\rightarrow 4.439$	$\rightarrow 2^+; 0$	$0.96 \pm 0.13$	M1	$(3.8 \pm 0.5) \times 10^{-2}$
	$\rightarrow 7.654$	$\rightarrow 0^+; 0$	$1.09 \pm 0.14$	M1	$0.13 \pm 0.02$
	$\rightarrow 12.71$	$\rightarrow 1^+; 0$	$0.59 \pm 0.17$	M1	$2.0 \pm 0.6$
	16.11 $\rightarrow 0$	$2^+; 1 \rightarrow 0^+; 0$	$0.59 \pm 0.11$	E2	$0.40 \pm 0.08$
	$\rightarrow 4.439$	$\rightarrow 2^+; 0$	$12.8 \pm 1.5$	M1	$0.38 \pm 0.05$
	$\rightarrow 9.641$	$\rightarrow 3^-; 0$	$0.31 \pm 0.06$	E1	$(3.2 \pm 0.6) \times 10^{-3}$
	$\rightarrow 12.71$	$\rightarrow 1^+; 0$	$0.19 \pm 0.04$	M1	$0.23 \pm 0.05$
	16.57 $\rightarrow 0$	$2^-; 1 \rightarrow 0^+; 0$	$(4.80 \pm 0.08) \times 10^{-2}$	M2	$0.489 \pm 0.008$

<sup>a</sup> The last column gives the  $\gamma$ -ray strengths expressed in Weisskopf units (see D.H. Wilkinson, in *Nuclear Spectroscopy Part B*, ed. F. Ajzenberg-Selove (Academic Press, NY, 1960)). The Weisskopf estimates ( $\Gamma_w$  in eV,  $E_\gamma$  in MeV) are:

$$\begin{aligned} \Gamma_w(E1) &= 6.8 \times 10^{-2} A^{2/3} E_\gamma^3, & \Gamma_w(E2) &= 4.9 \times 10^{-8} A^{4/3} E_\gamma^5, \\ \Gamma_w(E3) &= 2.3 \times 10^{-14} A^2 E_\gamma^7, & \Gamma_w(M1) &= 2.1 \times 10^{-2} E_\gamma^3, \\ \Gamma_w(M2) &= 1.5 \times 10^{-8} A^{2/3} E_\gamma^5. \end{aligned}$$

The values for these  $\gamma$ -ray strengths are occasionally different from those listed in other tables of this paper because different values of  $r_0$  were used. In this table  $r_0 = 1.2$  fm is used consistently. The multiplicities in the next to the last column were used to calculate the  $\Gamma_w$ . See also (1979EN05).

<sup>b</sup>  $T$  shown in usual convention [ $J^\pi; T$ ] only if transitions from the initial state involve a change in  $T$ .

<sup>c</sup> See Table 11.14 and (1982MI08).

<sup>d</sup> See Table 2 in (1980AJ01) for additional information.

<sup>e</sup> See Table 12.7.

<sup>f</sup> Assumed to be  $\frac{3}{2}^-$  since it is probably the analog to  $^{11}\text{C}^*(8.10)$ .