

Adopted Levels

$S(p)=-2730$ *syst* (2021Wa16)

$\Delta S(p)=500$ (syst,2021Wa16).

$S_{2p}=-3090$ 500 (sys.) (2021Wa16).

The ${}^{13}\text{F}$ nucleus is unbound to proton emission. A resonance is observed in (2021Ch19); however, since this resonance is reported at a higher energy than expected for the ground state, it is believed to correspond to an excited state. See also (2023Th03).

In (2013Ti01), the Kelson-Garvey mass systematics (1966Ga25) are improved to provide reliable predictions of the masses of proton-rich nuclei based on analysis of their neutron-rich mirror nuclei. For ${}^{13}\text{F}$, the predicted binding energy $E_{\text{bind}}=53.666$ MeV *16* and mass excess $\Delta M=43.386$ MeV *16* are based on comparison with ${}^{13}\text{Be}$.

In (2012Fo22) a potential model is developed that is informed by ${}^{12,13}\text{Be}$ and ${}^{12}\text{O}$ properties. The expected $J^\pi=1/2^+$ ground state is predicted near the resonance energy $E_p=2.4$ MeV with $\Gamma\approx 0.6$ MeV, relative to ${}^{12}\text{O}_{\text{g.s.}}$. The $J^\pi=5/2^+$ excited state is predicted near $E_p\approx 5$ MeV with $\Gamma\approx 0.35$ MeV. The analysis is updated in (2013Fo22) with similar results.

See (1993Po11) for earlier shell-model analysis of ground-state binding energies, including ${}^{13}\text{F}$.

 ${}^{13}\text{F}$ LevelsCross Reference (XREF) Flags

A ${}^9\text{Be}({}^{13}\text{O}, {}^{13}\text{F})$

<u>$E(\text{level})^a$</u>	<u>J^π</u>	<u>Γ</u>	<u>E_T (MeV)^b</u>	<u>XREF</u>	<u>Comments</u>
x	(5/2 ⁺)	1.01 MeV 27	7.06 9	A	%p=100 Analysis of the invariant-mass spectrum is consistent with sequential proton decay via ${}^{12}\text{O}(0:0_1^+, 2.2 \text{ MeV}; 2_1^+, 4.8 \text{ MeV}; 2_2^+)$ with intensities of 40% <i>16</i> , 28% <i>18</i> and 32% <i>16</i> , respectively.

^a The authors suggest the $J^\pi=5/2^+$ excited state has been observed. The $1/2^+$ g.s. is expected ≈ 3 MeV lower in energy.

^b ${}^{10}\text{C}+3\text{p}$ Invariant Mass.

${}^9\text{Be}({}^{13}\text{O}, {}^{13}\text{F})$ 2021Ch19

2021Ch19: XUNDL dataset compiled by TUNL (2021).

The authors analyzed the excitation spectra of particle-unbound nuclides produced in ${}^{13}\text{O}$ reactions on a ${}^9\text{Be}$ target. The first observation of a ${}^{13}\text{F}$ resonance, produced via charge-exchange reactions, was reported.

A beam of 69.5 MeV/nucleon ${}^{13}\text{O}$ ions, from the NSCL/A1900 fragment separator, was purified in the Radio Frequency Fragment Separator before impinging on a 1 mm thick ${}^9\text{Be}$ target. ${}^{13}\text{F}$ nuclides were produced and decayed in-flight via ${}^{13}\text{F} \rightarrow \text{p} + {}^{12}\text{O} \rightarrow 3\text{p} + {}^{10}\text{C}$ reactions. All decay products were detected using the HiRA High-Resolution position sensitive $\Delta\text{E-E}$ telescope array, which covered the polar angles $\theta_{\text{lab}} = 2.1^\circ$ to 12.4° . The invariant-mass spectrum, E_{T} , was deduced by analyzing the momenta of the $3\text{p} + {}^{10}\text{C}$ decay products. As in the past study of ${}^{11}\text{O}$ (2020We08), structures in the excitation spectrum are enhanced by analyzing events where ${}^{10}\text{C}$ is ejected perpendicular to the beam direction in the ${}^{13}\text{F}$ center-of-mass system.

The invariant-mass spectrum is dominated by a $\Gamma \approx 1$ MeV peak around $E_{\text{T}} \approx 7$ MeV that sits on top of a broad smooth background. For perspective, (2012Fo22) predicts a $J^\pi = 1/2^+$ ground state at around $E_{\text{T}} = 4$ MeV and a $J^\pi = 5/2^+$ first excited state near $E_{\text{T}} = 7$ MeV. No evidence is seen for a lower energy resonance; the authors suggest the lower state may have a broad width that prevents its observation in the present measurement. Comparison with the mirror ${}^{13}\text{Be}$ nucleus ground state provides ambiguous results for the measured $1/2^+$ width, leaving uncertainty on the expectation for ${}^{13}\text{F}$.

A Monte Carlo simulation was developed to interpret the observed features, since the level structures of ${}^{12}\text{O}$ and ${}^{10}\text{C}$ are convoluted in the spectrum. Involvement of the ${}^{10}\text{C}^*(3353 \text{ keV})$ first excited state is generally inconsistent with the observed invariant-mass spectrum so the analysis is limited to final population of ${}^{10}\text{C}_{\text{g.s.}}$. Hence the analysis focused on branching ratios for sequential decay of a $(5/2^+)$ ${}^{13}\text{F}$ resonance through ${}^{12}\text{O}$ states that subsequently 2p decay via processes described in (2019We11), and the total decay energy was assumed to be carried in the $3\text{p} + {}^{10}\text{C}$ kinetic energies.

Sequential decays via low-lying ${}^{12}\text{O}$ states are considered in the Monte Carlo simulation. See the text for discussion and comparison of expectations based on the mirror ${}^{13}\text{Be}$ nucleus. The branching ratios to the $J^\pi = 0_1^+$ and 2_1^+ and 2_2^+ ${}^{12}\text{O}$ states are found as 40% 16, 28% 18 and 32% 16, respectively. An alternate and reasonable fit to the data assumed zero contribution from the 2_2^+ state; in this case the relative ratios for the 0_1^+ and 2_1^+ states remained consistent with the initial three-level fit.

 ${}^{13}\text{F}$ Levels

<u>$E(\text{level})^a$</u>	<u>J^π</u>	<u>Γ</u>	<u>$E_{\text{T}} (\text{MeV})^b$</u>	<u>Comments</u>
x	$(5/2^+)$	1.01 MeV 27	7.06 9	Analysis of the invariant-mass spectrum is consistent with sequential proton decay via ${}^{12}\text{O}(0:0_1^+, 2.2 \text{ MeV}:2_1^+, 4.8 \text{ MeV}:2_2^+)$ with intensities of 40% 16, 28% 18 and 32% 16, respectively.

^a The authors suggest the $J^\pi = 5/2^+$ excited state has been observed. The $1/2^+$ g.s. is expected 3 MeV lower in energy.

^b Invariant Mass.

REFERENCES FOR A=13

- 1966Ga25 G.T.Garvey, I.Kelson - Phys.Rev.Letters 16, 197 (1966).
New Nuclidic Mass Relationship.
- 1993Po11 N.A.F.M.Poppelier, A.A.Wolters, P.W.M.Glaudemans - Z.Phys. A346, 11 (1993).
Properties of Exotic Light Nuclei.
- 2012Fo22 H.T.Fortune, R.Sherr - Phys.Rev. C 86, 034301 (2012).
Predictions for the first two positive-parity states of ^{13}F .
- 2013Fo22 H.T.Fortune - Phys.Rev. C 88, 024309 (2013).
Mirror energy differences of $2s_{1/2}$ single-particle states: Masses of ^{10}N and ^{13}F .
- 2013Ti01 J.Tian, N.Wang, C.Li, J.Li - Phys.Rev. C 87, 014313 (2013).
Improved Kelson-Garvey mass relations for proton-rich nuclei.
- 2019We11 T.B.Webb, R.J.Charity, J.M.Elson, D.E.M.Hoff et al. - Phys.Rev. C 100, 024306 (2019); Erratum Phys.Rev. C 102, 019904 (2020).
Particle decays of levels in $^{11,12}\text{N}$ and ^{12}O investigated with the invariant-mass method.
- 2020We08 T.B.Webb, R.J.Charity, J.M.Elson, D.E.M.Hoff et al. - Phys.Rev. C 101, 044317 (2020).
Invariant-mass spectrum of ^{11}O .
- 2021Ch19 R.J.Charity, T.B.Webb, J.M.Elson, D.E.M.Hoff et al. - Phys.Rev.Lett. 126, 132501 (2021).
Observation of the Exotic Isotope ^{13}F Located Four Neutrons beyond the Proton Drip Line.
- 2021Wa16 M.Wang, W.J.Huang, F.G.Kondev, G.Audi, S.Naimi - Chin.Phys.C 45, 030003 (2021).
The AME 2020 atomic mass evaluation (II). Tables, graphs and references.
- 2023Th03 M.Thoennessen - Int.J.Mod.Phys. E32, 2330001 (2023).
2022 Update of the discoveries of nuclides.