

# Energy Levels of Light Nuclei $A = 7$

F. Ajzenberg-Selove<sup>a</sup> and T. Lauritsen<sup>b</sup>

<sup>a</sup> *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

<sup>b</sup> *California Institute of Technology, Pasadena, California*

**Abstract:** An evaluation of  $A = 5-24$  was published in *Nuclear Physics* 11 (1959), p. 1. This version of  $A = 7$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 01, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 7$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides:  [\${}^7\text{He}\$](#) ,  [\${}^7\text{Li}\$](#) ,  [\${}^7\text{Be}\$](#)

B. Tables of Recommended Level Energies:

[Table 7.1](#): Energy levels of  ${}^7\text{Li}$

[Table 7.4](#): Energy levels of  ${}^7\text{Be}$

C. [References](#)

D. Figures:  [\${}^7\text{Li}\$](#) ,  [\${}^7\text{Be}\$](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

**<sup>7</sup>He**  
(Not illustrated)

Not observed: see (1955AJ61).

**<sup>7</sup>Li**  
(Fig. 5)

GENERAL:

*Theory:* See (1955AU1A, 1955DA1A, 1955LA1D, 1956AB1A, 1956FE1A, 1956KU1A, 1956ME1A, 1957FE1A, 1957FR1B, 1957LE1E, 1957MA1E, 1957MA57, 1957SO1A, 1958HA1D, 1958SK1A).

1.  ${}^3\text{H}(\alpha, \gamma){}^7\text{Li}$   $Q_m = 2.465$

For  $E_\alpha = 0.5$  to  $1.9$  MeV, capture radiation is observed to  ${}^7\text{Li}(0)$  and  ${}^7\text{Li}^*(0.48)$ , with intensity ratio  $5 : 2$ . The smooth rise of the cross section suggests a direct capture process. The angular distribution is not isotropic, indicating  $l > 0$  (1958RI34); see also (1958LI1A). In the range  $E_\alpha = 0.48$  to  $1.13$  MeV, the cross section increases from  $0.6$  to  $1.2 \mu\text{b}$  (1959HO03).

2.  ${}^4\text{He}(t, t){}^4\text{He}$   $E_b = 2.465$

Differential cross sections have been measured for  $E_t = 1.2$  to  $2.2$  MeV (1956HE16). At  $E_t = 1.677$  MeV,  $\theta = 30^\circ$ ,  $d\sigma/d\Omega$  (lab) =  $875$  mb/sr (1958AL05).

3.  ${}^4\text{He}(\alpha, p){}^7\text{Li}$   $Q_m = -17.347$

At  $E_\alpha = 38.5$  MeV, two groups of protons are observed, corresponding to the ground and  $0.48$ -MeV states of  ${}^7\text{Li}$ . Absolute c.m. differential cross sections are given as  $(1.53 \pm 0.09) + (1.93 \pm 0.15) \cos^2 \theta$  and  $(1.32 \pm 0.07) + (0.66 \pm 0.23) \cos^2 \theta$  mb/sr for the ground state and excited state groups, respectively. The cross sections are in fair agreement with those calculated from  ${}^7\text{Li}(p, \alpha){}^4\text{He}$  by the principle of detailed balancing. The partial inelastic cross sections are estimated to be  $11 \pm 1$  mb for s-waves,  $33 \pm 3$  mb for d-waves (1958BU38).

4.  ${}^6\text{Li}(n, \gamma){}^7\text{Li}$   $Q_m = 7.252$

Table 7.1: Energy levels of  ${}^7\text{Li}$ 

| $E_x$ in ${}^7\text{Li}$ (MeV) | $J^\pi$                          | $\tau_m$ (sec) or $\Gamma$ (keV)         | Decay                     | Reactions  |
|--------------------------------|----------------------------------|--|---------------------------|--|
| 0                              | $\frac{3}{2}^-$                  |  | stable                    | 1, 3, 4, 9, 10, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31 |
| $0.4780 \pm 0.0003$            | $\frac{1}{2}^-$                  | $\tau_m = (0.9 \pm 0.1) \times 10^{-13}$ | $\gamma$                  | 1, 3, 4, 9, 10, 16, 17, 19, 23, 24, 25, 28, 30, 31                 |
| $4.630 \pm 0.01$               | $(\leq \frac{7}{2}^-)$           | $93 \pm 8$                               | t, $\alpha$               | 9, 18, 19, 23, 30, 31  |
| $6.54 \pm 0.02$                | $(\frac{1}{2}^+, \frac{3}{2}^+)$ | broad                                    | t, $\alpha$               | (5), 9, 18, 19   |
| $7.47 \pm 0.01$                | $\frac{5}{2}^-$                  | $141 \pm 20$                             | n, $\alpha$ , t           | 5, 6, 9, 18, 19, 30  |
| (9.6)                          |                                  |  | $\gamma$ , n, t, $\alpha$ | 12, 15, 18, 19   |
| (10.8)                         |                                  |  | $\gamma$ , n, t, $\alpha$ | 12, 15   |
| 12.4                           |                                  |  | $\gamma$ , n              | 12   |
| 14.0                           |                                  |  | $\gamma$ , n              | 12   |
| 16.2                           |                                  |  | $\gamma$ , t, $\alpha$    | 15   |
| 17.5                           |                                  |  | $\gamma$ , n              | 12, 19   |
| (21.5)                         |                                  |  | $\gamma$ , t, $\alpha$    | 15, 19   |
| (23.5)                         |                                  |  | $\gamma$ , t, $\alpha$    | 15   |

Two  $\gamma$ -rays with  $E_\gamma = 7.26 \pm 0.03$  and  $6.78 \pm 0.05$  MeV, and relative intensities 10 and  $7.5 \pm 2.0$ , corresponding to transitions to the first two states of  ${}^7\text{Li}$  are observed. The total radiative capture cross section is  $28 \pm 8$  mb (1957BA18).

### 5. ${}^6\text{Li}(n, n){}^6\text{Li}$

$$E_b = 7.252$$

The total cross section has been measured from  $E_n = 4$  eV to 10 MeV (1954JO17, 1954NE1A, 1955HU1B, 1956GO62, 1957HU1D, 1957KA1B, 1958HU18), from  $E_n = 7$  to 14 MeV by (1958BR16) and at 14 MeV by (1952CO41, 1957KA1B). A pronounced resonance occurs at  $E_n = 255$  keV (see Table 7.2) with a peak cross section of 10.3 b (1954JO17). Angular distributions for  $E_n = 0.2$  to 0.4 MeV have been studied by (1956WI04): all observations are consistent with p-wave formation of a  $J = \frac{5}{2}^-$  level,  ${}^7\text{Li}^*(7.47)$ . The s-wave potential scattering has a statistical channel spin mixture with negative phase shift (1956WI04). Resonance parameters are compared with those of the mirror level ( ${}^7\text{Be}^*$ ) in Table 7.2. The 7.47-MeV level has been assumed to be the upper member of the  ${}^2F_{7/2, 5/2}$  doublet of which  ${}^7\text{Li}^*(4.6)$  is the lower. It is pointed out by (1956ME1A, 1957MA57, 1957SO1A) that it may in fact be a component of the  ${}^4P$  multiplet and that the  $F_{5/2}$  is to be found at a lower energy. This assumption would explain the large values of  $\theta_p^2$

and  $\theta_n^2$  and would make the large difference of  $\theta_\alpha^2$  for the mirror levels less disturbing (1957MA57). See also (1954LA1A, 1956DA1A, 1958HA1D; theor.).

No other clearly defined resonance is observed, although the cross section exhibits a broad maximum near  $E_n = 5$  MeV (1954JO17, 1957HU1D, 1958HU18).

The large coherent thermal scattering length suggests the existence of at least one bound s-level (1954LA1A).

$$6. \text{}^6\text{Li}(n, \alpha)^3\text{H} \qquad Q_m = 4.787 \qquad E_b = 7.252$$

The isotropic thermal capture cross section is 945 b (1958HU18); up to  $E_n \approx 70$  eV, the cross section varies as  $1/v$ . Cross sections and angular distributions are summarized in (1955HU1B, 1956HU1A, 1957HU1D, 1958HO1C, 1958HU18). A strong resonance is observed at  $E_n = 0.25$  MeV with a peak cross section of 2.9 b, attributed to p-wave formation of a  $J = \frac{5}{2}^-$  level (see  ${}^6\text{Li}(n, n){}^6\text{Li}$  and Table 7.2). In the range  $E_n = 9$  to 90 keV the cross section is higher than the  $1/v$  law would indicate (1956GO1E, 1959BA1H). Above the resonance, the cross section appears to exhibit a broad hump,  $\sigma \approx 250$  mb, at  $E_n = 1.5$  to 2 MeV, and then decreases smoothly to 26 mb at 14 MeV. The decrease at 2 MeV is attributed to competing reactions (1956RI34, 1958MU07; see, however, (1958HO1C)). (1958KE30) find an almost linear decrease from 34.3 mb at  $E_n = 12.5$  MeV to 17.5 mb at  $E_n = 18.3$  MeV. For  $E_n \gtrsim 2$  MeV the reaction appears to proceed by a pickup process. See  ${}^6\text{Li}(n, t){}^4\text{He}$  in  ${}^6\text{Li}$ . See also (1955PE1B, 1956DA1A, 1957BE71, 1957EL1C, 1957KO1A) and (1959BA1H).

$$7. \text{}^6\text{Li}(n, p){}^6\text{He} \qquad Q_m = -2.753 \qquad E_b = 7.252$$

The cross section at  $E_n = 14$  MeV is 6.7 mb (1953BA04, 1954FR03).

$$8. \text{}^6\text{Li}(n, d){}^5\text{He} \qquad Q_m = -2.429 \qquad E_b = 7.252$$

The total cross section is about 200 mb at  $E_n = 5.5, 6.5$  and 14 MeV (1956RI34). For the  ${}^5\text{He}$  ground-state peak,  $\sigma = 89$  mb for the continuum,  $\sigma = 77$  mb at  $E_n = 14$  MeV (1954FR03). The reaction appears to proceed by pickup: see  ${}^6\text{Li}(n, d){}^5\text{He}$  in  ${}^6\text{Li}$ .

$$9. \text{}^6\text{Li}(d, p){}^7\text{Li} \qquad Q_m = 5.027$$

The weighted mean value of the excitation energy of the first excited state is  $477 \pm 2$  keV (see (1955AJ61)). Levels are also observed at  $E_x = 4.630$  MeV,  $\Gamma = 93 \pm 8$  keV (1957BR97),  $4.61 \pm$

Table 7.2: Resonance parameters for 7.2 – 7.5 MeV level in  ${}^7\text{Li}$ - ${}^7\text{Be}$ 

|  | ${}^6\text{Li}(n, n){}^6\text{Li}$ , ${}^6\text{Li}(n, \alpha){}^3\text{H}$ |                  |                    |                    | ${}^6\text{Li}(p, \alpha){}^3\text{He}$ |                    |
|--|---|------------------|--------------------|--------------------|---|--------------------|
|  | (1951BA79)  | (1954JO17)       | (1956WI04)         | (1956MA91)         | (1951BA79)                              | (1956MA91)         |
| $E_r$ (keV, lab) <sup>a</sup>          | 250   | 255              | 255                |                    | 1820                                    | 1850               |
| $\Gamma$ (keV, lab)                    | 100   | 100              | 100                |                    | 500                                     |                    |
| $E_\lambda$ (keV above g.s.)           | 7540  | 7700             | 6640 <sup>b</sup>  | 7680               | 7310                                    | 7800               |
| $\Gamma_{n,p}(E_r)$ (keV, c.m.)        | 50  | 114              | 82                 | 114                | 400                                     | 670                |
| radius (n, p) in $10^{-13}$ cm         | 4.0   | 2.54             | 3.94               | 4.08               | 4.0                                     | 4.08               |
| $\gamma_{n,p}^2$ ( $10^{-13}$ MeV-cm)  | 2.4   | 11.8             | 3.5 <sup>b</sup>   | 4.6                | 2.6                                     | 4.2                |
| $\theta_{n,p}^2$                       | [0.13]  | 0.42             | 0.19 <sup>b</sup>  | 0.26               | [0.14]                                  | 0.24               |
| $\Gamma_\alpha(E_r)$ (keV, c.m.)       | 36  | 60               | 43                 | 64                 | 30                                      | 50                 |
| radius ( $\alpha$ ) in $10^{-13}$ cm   |   | 3.5              |                    | 4.39               |   | 4.39               |
| $\gamma_\alpha^2$ ( $10^{-13}$ MeV-cm) |   | 0.4 <sup>c</sup> | 0.12 <sup>b</sup>  | 0.14               |   | 0.79               |
| $\theta_\alpha^2$                      |   | [0.04]           | 0.014 <sup>b</sup> | 0.017 <sup>d</sup> |   | 0.095 <sup>d</sup> |

<sup>a</sup>  $E_r = 265$  keV,  $\sigma = 9.5 \pm 0.2$  b (1956GO62); see also (1958HU18, 1959BA1H).

<sup>b</sup> With same convention as other references,  $E_\lambda = 7580$  (H.B. Willard, private communication).

<sup>c</sup>  $\gamma_n^2/\gamma_\alpha^2 = 30$ ; see (1956MA91: footnote, p. 1406).

<sup>d</sup> Corrected value: see (1957MA57).

0.02 MeV (1952GE07),  $4.46 \pm 0.02$  MeV (1955KH35) and  $E_x = 6.54 \pm 0.02$  MeV (1955KH31, 1955KH35), 6.56 MeV (1955LE24: see, however, (1958HA22)) and  $E_x = 7.457$  MeV,  $\Gamma = 91 \pm 8$  keV (1957BR97). A search for a level at 5.5 MeV was unsuccessful at  $\theta = 30^\circ$ ,  $70^\circ$  and  $90^\circ$ ,  $E_d = 7.0$  and 7.5 MeV (1957BR97). Angular distributions of the protons to the ground state and the 0.48-MeV level, at  $E_d = 8$  MeV (1953HO48) and  $E_d = 14.5$  MeV (1955LE24), analyzed by stripping theory, show odd parity,  $J \leq \frac{5}{2}$  for both states: see also (1954NI10). The ratio of reduced widths  $\theta^2(\text{g.s.})/\theta^2(0.48) = 0.69$  and 0.66 from these two experiments (1956RE04);  $\theta^2(\text{g.s.}) \approx 0.060$  (1957FR1B). The angular correlation between the protons and the 0.48-MeV  $\gamma$ -rays is isotropic (see (1955AJ61)) indicating  $J = \frac{1}{2}$  for  ${}^7\text{Li}^*$ . The mean life of this state as determined in the present reaction and in  ${}^9\text{Be}(d, \alpha){}^7\text{Li}$  is  $(7.7 \pm 0.8) \times 10^{-14}$  sec (1956BU83). At  $E_d = 15$  MeV, the angular distribution of the protons to the 4.6-MeV state has been measured for  $\theta = 30^\circ$  to  $100^\circ$  and can be reasonably accounted for by stripping theory with  $l_n = 2$ ,  $J \leq \frac{7}{2}^-$  (1956HA90); on the other hand, (1958HA22) report the distribution nearly isotropic at  $E_d = 15$  MeV. At  $E_d = 24$  MeV the distribution is in poor agreement with stripping theory (1957CO68). For the 7.5-MeV level, the proton distribution at  $E_d = 15$  MeV indicates  $l_n = 1$  and a reduced width equal to that of the ground state (1958HA22). A search for 4.1 MeV  $\gamma$ -rays from a possible (4.6  $\rightarrow$  0.48) transition was unsuccessful: at  $E_d = 1.8$  MeV,  $I_{4.1} < 0.1$  ( $I_{0.48}$ ) (1957WI24). See also  ${}^8\text{Be}$ , (1955AU1A, 1955LA1D, 1957FR1B; theor.) and (1952AJ38).

10.  ${}^6\text{Li}(t, d){}^7\text{Li}$

$$Q_m = 0.994$$

$$Q_0 = 0.986 \pm 0.007 \text{ (1954AL35)}.$$

At  $E_t = 240$  keV, the reaction has been observed to the ground state and to the 0.48-MeV level (1954AL35). See also (1955CU17).

11.  ${}^6\text{Li}(\alpha, {}^3\text{He}){}^7\text{Li}$

$$Q_m = -13.325$$

Not reported.

12.  ${}^7\text{Li}(\gamma, n){}^6\text{Li}$

$$Q_m = -7.252$$

The total cross section for neutron production (including the  $(\gamma, np)$  and  $(\gamma, 2n)$  processes) has a broad maximum ( $\sigma = 2.3$  mb) at  $E_\gamma = 16.8$  MeV,  $\Gamma = 9.3$  MeV (1958RY77:  ${}^{\text{nat}}\text{Li}$ ). For the  $(\gamma, n)$  process alone, the peak cross section is estimated as  $\approx 2.7$  mb at  $E_\gamma \approx 14$  MeV (1954GO1A). Discontinuities reported in the integral yield curve are listed in Table 7.3. Integrated cross sections are tabulated by (1954GO39). See also (1953MO1B, 1955HE51, 1958BE1C, 1958TI1A) and (1953PE1A, 1957BA1H; theor.). (It is to be noted that the total cross section in  ${}^6\text{Li}(n, n){}^6\text{Li}$  exhibits no evidence of sharp structure in the region  $E_x = 9$  to 12 MeV: see (1954JO17, 1957HU1D, 1958HU18).)

13.  ${}^7\text{Li}(\gamma, p){}^6\text{He}$

$$Q_m = -10.005$$

According to (1953TI02) and (1954TI16), the cross section in the range  $E_\gamma = 12$  to 20 MeV exhibits a single broad maximum at  $\approx 15.6$  MeV with a width of  $\approx 4$  MeV. At  $E_\gamma = 14.8$  MeV the cross section is  $2.2 \pm 0.25$  mb; according to (HA55A), the integrated cross section is 11 MeV-mb. There appears to be some disagreement on the shape of this excitation function. It is suggested that a  $\gamma$ -emitting state of  ${}^6\text{He}$  at  $\approx 1.6$  MeV could account for the discrepancy between the Canberra and Saskatoon results (1954TI16, 1955TI1A). Some evidence for fine structure is reported by (HA55A). See also (1953TU1A, 1954RU27, 1954TU1A, 1958SM1A, 1958ST1A, 1958TI1A, 1958WH35).

14.  ${}^7\text{Li}(\gamma, d){}^5\text{He}$

$$Q_m = -9.681$$

See (1958BA1C, 1958PA1B, 1958TI1A, 1958WH35).

Table 7.3: Levels of  ${}^7\text{Li}$  (MeV) from  ${}^7\text{Li}(\gamma, t){}^4\text{He}$  and  ${}^7\text{Li}(\gamma, n){}^6\text{Li}$

| $(\gamma, t)$ |            |            |            | $(\gamma, n)$    |
|---------------|------------|------------|------------|------------------|
| (1953TI02)    | (1953ST27) | (1954ST89) | (1955MI55) | (1954GO39)       |
|               |            | 4.7        |            |                  |
|               | 5.25       | 5.5        |            |                  |
|               |            | 6.8        |            |                  |
|               | 7.25       |            | 7.6        |                  |
|               |            |            | 8.6        |                  |
| 9.3           | (9.25)     |            | 9.6        | 9.6 <sup>a</sup> |
|               |            |            | 11.7       | 10.8             |
|               |            |            |            | 12.4             |
|               |            |            | (13.5)     | 14.0             |
| (16.7)        |            |            | 16.2       |                  |
|               |            |            |            | 17.5             |
| (21.5)        |            |            |            |                  |
| (23.5)        |            |            |            |                  |

<sup>a</sup>  $9.96 \pm 0.04$  MeV (1958RY77).

15.  ${}^7\text{Li}(\gamma, t){}^4\text{He}$   $Q_m = -2.465$

Reported peaks in the excitation function are listed in Table 7.3. Observations are made in nuclear emulsions, mainly under continuous-spectrum irradiation. It is pointed out by (1953PE1A) that  $T = \frac{3}{2}$  levels of  ${}^7\text{Li}$  will not be observed in the present reaction in the region 0 to 15 MeV above threshold: compare  ${}^7\text{Li}(\gamma, n){}^6\text{Li}$ . See also (1952NA1A, 1953TI1A, 1953TU1A, 1954ER1A, 1954TU1A, 1955TI1A, 1958TI1A) and (1955CZ1A, 1956CZ1A; theor.).


16.  ${}^7\text{Li}(\gamma, \gamma'){}^7\text{Li}^*$

Measurement of resonance scattering and absorption in Li yield  $\tau(\text{mean}) = (1.1 \pm 0.3) \times 10^{-13}$  sec (1958BE10),  $(1.15 \pm 0.14) \times 10^{-13}$  sec (1958SW65, 1959SW63) for the 0.48-MeV state.


17.  ${}^7\text{Li}(e, e){}^7\text{Li}$



Elastic and inelastic (0.47-MeV state) scattering have been studied at  $E_e = 187$  MeV by (1955ST85). Elastic scattering results indicate an r.m.s. radius of about  $2.1 \times 10^{-13}$  cm (1957HO1E: see also (1956HO93, 1957ME1B)).



At  $E_n = 14$  MeV, there is evidence for  ${}^7\text{Li}$  states at  $4.6 \pm 0.25$ ,  $\approx 6.5 \pm 0.25$ ,  $7.5 \pm 0.25$  and  $9.25$  (?) MeV (1954AL24: Li-loaded photoplates). See also (1954FR03).



Elastic scattering at  $E_p > 10$  MeV is characterized by direct interaction processes: see (1956KI54, 1957HI56). Inelastic proton groups corresponding to the states at 0.48, 4.63, 6.54 and 7.46 MeV have been observed at bombarding energies up to 18.3 MeV: see (1952AJ38). A careful check on the isotropy of the 477-keV radiation from the  $J = \frac{1}{2}^-$  level yields  $W(\theta) = 1 + \alpha \cos \theta$ , with  $\alpha = (2.9 \pm 6.2) \times 10^{-4}$ ; the result implies an upper limit for the intensity of a parity non-conserving part of the wave function  $F^2 < 1 \times 10^{-4}$  (1958WI38).

At  $E_p = 12$  MeV, the angular distribution of the protons leaving  ${}^7\text{Li}$  in the 4.63-MeV state can be accounted for by a mixture of direct interaction with  $l = 0$ , and compound nucleus formation, indicating  $J = \frac{1}{2}^-$ ,  $\frac{3}{2}^-$ , or  $\frac{5}{2}^-$ . It is also possible to fit the data under the assumption of compound nucleus formation with s, p, and d-waves, permitting  $J = \frac{7}{2}^-$  (1957CO53). Angular distributions at  $E_p = 17.5$  MeV are reported by (1957MA04). The relative intensities observed in this work for protons leading to the 4.6, 6.6, and 7.5-MeV states are consistent with assignments  ${}^{22}\text{F}_{7/2}$ ,  ${}^{22}\text{F}_{5/2}$ ,  ${}^{24}\text{P}_{5/2}$ ; it is presumed that the 6.6-MeV state in question here is not the broad even-parity state at 6.54 MeV (1957LE1E). At 31.8 MeV, angular distributions of protons leading to the 4.6-MeV and 6.6-MeV levels fit the direct interaction theory with  $l = 1$  ( $J = \frac{1}{2}^+$ ,  $\frac{3}{2}^+$ ,  $\frac{5}{2}^+$ ,  $\frac{7}{2}^+$ ). A weak group to a 9.6-MeV level is also reported (1957SI36). Levels at  $18 \pm 1.4$  (?) and  $22 \pm 1.4$  (?) MeV are observed with 96-MeV protons (1956ST30). See also (1957TY35, 1958CH26, 1958MA1B, 1958TY46, 1958TY49).



The ground-state reduced width  $\theta_n^2 = 0.05$  and  $0.035$  for  ${}^6\text{Li}(0)$  and  ${}^6\text{Li}^*(2.2)$ , respectively: see  ${}^6\text{Li}$ .



$\theta_n^2 = 0.11$  and  $0.061$  for  ${}^6\text{Li}(0)$  and  ${}^6\text{Li}^*(2.2)$ , respectively: see  ${}^6\text{Li}$ .

22.  ${}^7\text{Li}(d, {}^3\text{He}){}^6\text{He}$   $Q_m = -4.512$

$\theta_p^2 = 0.055$  and  $0.017$  for  ${}^6\text{He}(0)$  and  ${}^6\text{He}^*(1.7)$ , respectively: see  ${}^6\text{He}$ .

23.  ${}^7\text{Li}(d, d'){}^7\text{Li}^*$

Inelastic deuteron groups have been observed to the states at 0.48 and 4.6 MeV: see (1952AJ38) and (1955KH31, 1955KH35). (1955LE24) report  $Q = -4.62 \pm 0.04$ ,  $\Gamma = 0.3 \pm 0.1$  MeV, for the latter state;  $\theta_\alpha^2 = 0.5 \pm 0.2$  (1957MA57). At  $E_d = 14.4$  MeV, the angular distribution of the deuterons corresponding to the 0.48-MeV state is moderately well accounted for by the theory of (1952HU1B), with  $l = 2$  (1955LE24). Some discrepancies appear to exist between reported angular distributions of deuterons corresponding to the 4.6-MeV level at  $E_d = 14$  to 15 MeV: compare (1955LE24) and (1956HA90). See also (1956SO21, 1956SO33).

24.  ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}^*$

${}^7\text{Li}$  states at 0.48 and 4.6 MeV have been observed at various energies up to 48 MeV: see (1955AJ61). At  $E_\alpha = 31.8$  and 48 MeV, the angular distribution of the  $\alpha$ -particles to the 4.6-MeV state exhibits a prominent peak in the forward direction, suggesting a direct interaction process (1956CO61, 1957SI36). (1954ZH1A) finds  $E_\gamma = 478 \pm 2$  keV and a lifetime for the first excited state  $< 1.3 \times 10^{-13}$  sec (see also  ${}^6\text{Li}(d, p){}^7\text{Li}$  and  ${}^9\text{Be}(d, \alpha){}^7\text{Li}$ ). At  $E_\alpha = 1.9$  MeV, the angular distribution of the 0.48-MeV radiation is isotropic within 10% (1954LI48). See also (1955BR1A, 1957NE1B).

25.  ${}^7\text{Be}(\epsilon){}^7\text{Li}^\dagger$   $Q_m = 0.863$

$^\dagger$  The symbol ( $\epsilon$ ) denotes orbital electron capture.

The decay proceeds to the ground and 0.48-MeV states. The fraction to the excited state is  $0.115 \pm 0.01$  (see (1955AJ61)). The  $\gamma$ -ray energy is  $477.8 \pm 0.3$  keV (weighted mean of values quoted in (1955AJ61), including the value  $477.3 \pm 0.4$  reported by (1957DU37)). The weighted mean value of the half-life is  $53.37 \pm 0.11$  days (1949SE20, 1953KR16, 1956BO36, 1957WR37),  $\log ft = 3.25$  for the ground state transition and 3.44 for the excited state. Both transitions are super-allowed (1954MA1D, 1956CH1B). A measurement of the internal conversion coefficient for the 0.48-MeV  $\gamma$ -radiation indicates that the radiation is 55% to 100% M1 (1958LE20). See also (1955LA1D; theor.).

26.  ${}^7\text{Be}(n, p){}^7\text{Li}$   $Q_m = 1.646$

See  ${}^8\text{Be}$ .

27. (a)  ${}^9\text{Be}(\gamma, d){}^7\text{Li}$   $Q_m = -16.693$

(b)  ${}^9\text{Be}(\gamma, np){}^7\text{Li}$   $Q_m = -18.919$

See (1955AJ61) and  ${}^9\text{Be}$ .

28.  ${}^9\text{Be}(n, t){}^7\text{Li}$   $Q_m = -10.435$

See (1957SC12, 1957VA12) and  ${}^{10}\text{Be}$ .

29.  ${}^9\text{Be}(p, {}^3\text{He}){}^7\text{Li}$   $Q_m = -11.200$

Not observed.

30.  ${}^9\text{Be}(d, \alpha){}^7\text{Li}$   $Q_m = 7.152$

A number of  $\alpha$ -groups have been observed with deuteron energies of up to 14 MeV. These correspond to levels at  $480 \pm 2$  keV (1948BU31, 1953CO02),  $4.62 \pm 0.02$  MeV (1953GE01: see also (1951GO47, 1952AS1A, 1956JU1D)), and  $7.5 \pm 0.17$  (?) MeV (1951GO47). There is no evidence in this reaction of a state at 5.5 MeV suggested in  ${}^7\text{Li}(\gamma, t){}^4\text{He}$  (see (1955AJ61)): at  $E_d = 0.5$  and  $0.7$  MeV ( $\theta = 70^\circ$ ), the intensity of a group leading to such a state is less than 5% of the intensity of the ground-state alpha group (1956GE1A: see also (1955CU16, 1956JU1D)). Observation of the continuous distribution of alpha-particles and tritons indicates production and breakup of  ${}^7\text{Li}$  levels at 4.6 and 6.6 MeV (1955CU16, 1956JU1D).

The  $(\alpha, \gamma)$  angular correlation has been observed for  $E_d = 0.40$  and  $0.84$  MeV (1953UE01, 1954CO17). There is no significant departure from isotropy, in agreement with  $J = \frac{1}{2}$  for the 0.48-MeV level. The mean life of this state is  $(7.7 \pm 0.8) \times 10^{-14}$  sec (1956BU83: Doppler shift; mean of observations in this reaction and in  ${}^6\text{Li}(d, p){}^7\text{Li}$ ); compare  ${}^7\text{Li}(\gamma, \gamma){}^7\text{Li}$ . This value is considerably shorter than that given by a shell-model calculation (1955LA1D). See also  ${}^{10}\text{B}(p, \alpha){}^7\text{Be}$  in  ${}^7\text{Be}$ .

See also (1956GR37, 1956TU1A) and  ${}^{11}\text{B}$ .

31.  ${}^{10}\text{B}(n, \alpha){}^7\text{Li}$   $Q_m = 2.793$

With thermal neutrons, two groups of  $\alpha$ -particles are observed, corresponding to  ${}^7\text{Li}^*(0, 0.48)$ ; the fraction of transitions leading to the ground state is about 6%; see (1952AJ38, 1954DE38, 1958BU02) and  ${}^{11}\text{B}$ . The  $\gamma$ -ray energy is  $478.5 \pm 1.5$  keV (1948EL1A),  $478 \pm 4$  keV (1956DA23); the mean life is  $(7.5 \pm 2.5) \times 10^{-14}$  sec (1949EL07). The  $(\alpha\text{-}\gamma)$  correlation is isotropic (1950RO1A: pile neutrons). See also (1955JA18).

For neutrons in the range  $E_n = 5.6$  to 19.5 MeV, the reaction appears to proceed mainly through the 4.6-MeV level or through direct three-body disintegration (1956FR18). See also (1955AJ61).

32.  ${}^{11}\text{B}(\gamma, \alpha){}^7\text{Li}$

$$Q_m = -8.670$$

Not reported.

**<sup>7</sup>Be**  
(Fig. 6)

GENERAL:

*Theory:* See (1957FR1B, 1957MA57, 1958SK1A).

1.  ${}^7\text{Be}(\epsilon){}^7\text{Li}$   $Q_m = 0.863$

The decay is complex; see  ${}^7\text{Li}$ .

2.  ${}^4\text{He}({}^3\text{He}, \gamma){}^7\text{Be}$   $Q_m = 1.584$

In the range  $E({}^3\text{He}) = 0.48$  to  $1.32$  MeV, the capture cross section increases from  $0.04$  to  $1.2$   $\mu\text{b}$ . At  $E({}^3\text{He}) = 1.32$  MeV, about 50% of the transitions involve the  $0.43$ -MeV state (1959HO03). See also (1958BA59, 1958HE1F). The significance of this reaction for energy generation in stars is discussed by (1958CA1C, 1958FO1A).

3.  ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$   $E_b = 1.584$

Elastic scattering studies for  $E({}^3\text{He}) = 3$  to  $5.5$  MeV ( $\theta = 54^\circ$  to  $126^\circ$ ) indicate a prominent resonance at  $E({}^3\text{He}) = 5.17$  MeV,  $\Gamma = 0.180$  MeV ( ${}^7\text{Be}^* = 4.53$  MeV,  $\Gamma_{\text{c.m.}} = 0.102$  MeV). Phase-shift analysis indicates that the state is  $F_{7/2}$ , with  $\gamma_\alpha^2 = 3.0 \times 10^{-13}$  MeV-cm,  $\theta_\alpha^2 = 0.36$ ,  $R = 4.4 \times 10^{-13}$  cm. The behavior of the p-wave phase shifts suggests that the ground state, and possibly  ${}^7\text{Li}^*(0.48)$ , have widths which approach  $\theta_\alpha^2 = 1$ . The s-wave phase shift is consistent with hard-sphere scattering, with  $R = 2.8 \times 10^{-13}$  cm (1958MI92). See also (1958CH35).

4.  ${}^4\text{He}(\alpha, n){}^7\text{Be}$   $Q_m = -18.993$

A search for this reaction at  $E_\alpha = 39$  MeV,  $0.5$  MeV (c.m.) above threshold, led to a negative result; the cross section is  $< 0.7$  mb. This result is consistent with the assumption that  ${}^7\text{Be}$  has negative parity (1952WA31).

5.  ${}^6\text{Li}(p, \gamma){}^7\text{Be}$   $Q_m = 5.607$   
 $Q_0 = 5.66 \pm 0.05$  (1956WA03).

Table 7.4: Energy levels of  ${}^7\text{Be}$

| $E_x$ in ${}^7\text{Be}$ (MeV) | $J^\pi$              | $\tau$ or $\Gamma$ (keV)                     | Decay                              | Reactions           |
|--------------------------------|----------------------|--|------------------------------------|---------------------|
| 0                              | $\leq \frac{3}{2}^-$ | $\tau_{1/2} = 53.37 \pm 0.11$ d              | $\epsilon$                         | 1, 5, 9, 13, 16, 18 |
| $0.431 \pm 0.002$              | $\frac{1}{2}^-$      | $\tau_m = (2.7 \pm 1.0) \times 10^{-13}$ sec | $\gamma$                           | 5, 9, 13, 18        |
| $4.53 \pm 0.02$                | $\frac{7}{2}^-$      | 102  | ${}^3\text{He}, \alpha$            | 3, 13, 18           |
| $6.35 \pm 0.1$                 | $(\frac{3}{2}^+)$    | $\approx 1000$                               | p, ${}^3\text{He}, \alpha(\gamma)$ | 5, 7, 9, 18         |
| $7.18 \pm 0.06$                | $(\frac{5}{2}^-)$    | 575  | p, ${}^3\text{He}, \alpha$         | 7, 9, 13, 18        |
| $(14.6 \pm 0.3)$               |                      |  |                                    | 18                  |

Gamma transitions to the ground state and to the 0.43-MeV state are observed. The yield shows no evidence of resonance for  $E_p = 0.4$  to 1.0 MeV, and the branching ratio remains approximately constant at  $62 \pm 5\%$ , (1955BA59: 65%) to the ground state, 38% to the 0.43-MeV state, (1955BA59:  $< 4\%$  to the 4.6-MeV state). The  $90^\circ$ -differential cross section at 750 keV is  $0.02 \mu\text{b}/\text{sr}$  (1956WA03). (1955BA59) estimate  $0.2 \mu\text{b}/\text{sr}$  from observation of  $(430 \pm 20)$ -keV radiation. (Because of the extremely low cross section of the present reaction, small amounts of contaminants may have a strong influence; in particular, earlier reports of a resonance at  $E_p \approx 1.6 - 1.8$  MeV may be due to  ${}^{10}\text{B}(p, \alpha){}^7\text{Be}$ : see (1956WA03).)

The angular distribution of both  $\gamma$ -rays (to the ground state and the 0.43-MeV level) is the same at  $E_p = 750$  keV,  $W(\theta) = 1 + (1.05 \pm 0.15) \cos^2 \theta$ . Neither s- or d-wave capture yields such a distribution, and p-wave,  $J = \frac{3}{2}^-$ , is indicated; a direct non-resonant capture process seems also possible (1956WA03).

6.  ${}^6\text{Li}(p, n){}^6\text{Be}$

$$Q_m = -5.2$$

$$E_b = 5.607$$

At  $E_p = 9$  MeV, the cross section for production of ground state neutrons is  $d\sigma/d\Omega = 0.19 + 0.23 \cos \theta + 0.70 \cos^2 \theta$  mb/sr;  $\sigma = 5 \pm 1$  mb (1957BO1F).

7. (a)  ${}^6\text{Li}(p, \alpha){}^3\text{He}$

$$Q_m = 4.022$$

$$E_b = 5.607$$

(b)  ${}^6\text{Li}(p, p){}^6\text{Li}$

The excitation function follows the Gamow function from  $E_p = 19$  to 250 keV (1953SA1A, 1957JA37). The total cross section exhibits two resonances, one at  $\approx 1$  MeV (broad) and one at 1.83 MeV (1951BA79, 1956MA91). (There is some discrepancy in the reported absolute cross sections.) In the range  $E_p = 1.3$  to 3.1 MeV, the elastically scattered protons,  $\theta = 164^\circ$ , exhibit a single maximum at  $E_p = 1.75$  MeV, with  $\Gamma = 0.5$  MeV (1951BA79): see Table 7.2. Preliminary

results of a study at  $\theta = 156^\circ$  for  $E_p = 0.45$  to 1.8 MeV clearly show the 1.75-MeV anomaly, but no marked effect near 1 MeV (1958MC65). Analysis of  $\alpha$ -particle angular distributions in the range  $0.6 \leq E_p \leq 2.5$  MeV is consistent with  $J = \frac{3}{2}^+, \frac{5}{2}^-$  for the two states at 6.35 and 7.18 MeV. A large, non-resonant background may indicate other states or some direct interaction (1956MA91): see also  ${}^6\text{Li}(p, \gamma){}^7\text{Be}$ . At  $E_p = 15$  and 18 MeV, the angular distributions of  ${}^3\text{He}$ -particles are sharply peaked forward (1956LI37: see  ${}^6\text{Li}$ ). See also (1955SA1C; theor.).

8.  ${}^6\text{Li}(p, d){}^5\text{Li}$   $Q_m = -3.269$

See  ${}^5\text{Li}$ .

9. (a)  ${}^6\text{Li}(d, n){}^7\text{Be}$   $Q_m = 3.380$

(b)  ${}^6\text{Li}(d, n){}^3\text{He} + {}^4\text{He}$   $Q_m = 1.796$

Two neutron groups are reported, corresponding to the ground- and 0.43-MeV states. The  $\gamma$ -ray energy is given as  $428.9 \pm 2$  keV (corrected for Doppler shift): the  ${}^7\text{Li}^* - {}^7\text{Be}^*$  difference is  $48.5 \pm 1.0$  keV (1952TH24). At  $E_d = 3.5$  MeV, the angular distributions of the neutron groups, analyzed by stripping theory indicate  $l_p = 1$ ,  $J \leq \frac{5}{2}^-$  for both states. Reaction (b) is also observed (1952AJ1B). Pronounced stripping patterns for neutrons leading to the 0.43-MeV state, observed at  $E_d = 0.6$  to 1.5 MeV, also indicate  $l_p = 1$ . Neutron-gamma correlations, with the neutron counter at a fixed angle, are isotropic; taken together with the stripping results, this observation indicates  $J = \frac{1}{2}^-$  for the excited state (1956NE13, 1958NE38, 1958SA17); see also (1951TH1A, 1953TH1B). A search for neutron groups leading to levels of  ${}^7\text{Be}$  in the range 0.3 to 2.0 MeV revealed no others with intensities  $> 10\%$  of the main group (1956NE13).

Broad maxima observed in the ratio of low energy to high-energy neutrons at  $E_d = 4.2$  and 5.1 MeV are attributed to  ${}^7\text{Be}$  state at 6.5 MeV ( $\Gamma_{c.m.} = 1.2$  MeV) and 7.2 MeV ( $\Gamma_{c.m.} = 0.5$  MeV). The 4.6-MeV state produces no such effect, presumably because of the high angular momentum of the neutrons (1957SL01): see  ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$ . At  $E_d = 1.8$  MeV, a search for 4.2-MeV  $\gamma$ -rays (from the  $4.58 \rightarrow 0.43$  transition) was unsuccessful:  $I(4.2\text{-MeV } \gamma) < 0.1 I(0.4\text{-MeV } \gamma)$  (1957WI24). See also (1955CA1A).

10.  ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$   $Q_m = 0.113$

Not observed.

11.  ${}^6\text{Li}(\alpha, t){}^7\text{Be}$   $Q_m = -14.206$

Not observed.

$$12. {}^6\text{Li}({}^6\text{Li}, n\alpha){}^7\text{Be} \quad Q_m = 1.909$$

See (1957NO17).

$$13. {}^7\text{Li}(p, n){}^7\text{Be} \quad Q_m = -1.646$$

A weighted mean of measurements of the ground state threshold gives  $1.8811 \pm 0.0005$  MeV (1954JO10). Using a two-meter absolute electrostatic analyzer, (1958BO76) obtain  $E_{\text{thresh.}} = 1.8812 \pm 0.0009$  keV.

The weighted mean value of five independent observations for the energy of the excited state is  $431 \pm 5$  keV; see Table II (7) in (1952AJ38). A neutron threshold determination gives  $434 \pm 4$  keV (1955MA84). At  $E_p = 9$  MeV, neutrons corresponding to  ${}^7\text{Be}^*(4.6)$  are reported by (1957BO1F). At  $E_p = 18.3$  MeV,  ${}^7\text{Be}$  levels are observed at  $4.6 \pm 0.2$  and  $7.1 \pm 0.2$  MeV (1952TH1C).

$$14. {}^7\text{Li}({}^3\text{He}, t){}^7\text{Be} \quad Q_m = -0.881$$

Not observed.

$$15. {}^9\text{Be}(\gamma, 2n){}^7\text{Be} \quad Q_m = -20.565$$

See  ${}^9\text{Be}$ .

$$16. {}^9\text{Be}(p, t){}^7\text{Be} \quad Q_m = -12.081$$

See (1954CO02) and (1956BE14).

$$17. {}^9\text{Be}(d, tn){}^7\text{Be} \quad Q_m = -14.307$$

See (1955HE83) and  ${}^{11}\text{B}$ .



18.  $^{10}\text{B}(\text{p}, \alpha)^7\text{Be}$

$$Q_m = 1.147$$

The weighted mean of five determinations of the energy of the first excited state gives  $430.3 \pm 2$  keV: see Table III (7) in (1952AJ38), and (1952CR30);  $E_\gamma = 432 \pm 3$  keV (1954DA20). The mean lifetime of this state is  $(2.7 \pm 1.0) \times 10^{-13}$  sec (1956BU83: Doppler shift). This value agrees with a shell model calculation by (1955LA1D).

At  $E_p = 18$  MeV,  $\alpha$ -groups are reported corresponding to  $^7\text{Be}^* = 0, 0.49 \pm 0.10, 4.72 \pm 0.08, 6.27 \pm 0.10, 7.21 \pm 0.10$  and  $14.6 \pm 0.3$  MeV. The last group is ten times as intense as any of the others. It is not completely excluded that it may be due to  $^{10}\text{B}(\text{p}, ^3\text{He})^8\text{Be}$  (1955RE16).

## References

(Closed December 01, 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1948BU31 W.W. Buechner, E.N. Strait, C.G. Stergiopoulos and A. Sperduto, Phys. Rev. 74 (1948) 1569
- 1948EL1A Elliott and Bell, Phys. Rev. 74 (1948) 1869
- 1949EL07 L.G. Elliott and R.E. Bell, Phys. Rev. 76 (1949) 168
- 1949SE20 E. Segre and C.E. Wiegand, Phys. Rev. 75 (1949) 39; Erratum Phys. Rev. 81 (1951) 284
- 1950RO1A Rose and Wilson, Phys. Rev. 78 (1950) 68
- 1951BA79 S. Bashkin and H.T. Richards, Phys. Rev. 84 (1951) 1124
- 1951GO47 H.E. Gove and J.A. Harvey, Phys. Rev. 82 (1951) 658
- 1951TH1A Thirion, Compt. Rend. 233 (1951) 37
- 1952AJ1B Ajzenberg, Phys. Rev. 87 (1952) 205A
- 1952AJ38 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 24 (1952) 321
- 1952AS1A Ashmore and Raffle, Proc. Phys. Soc. (London) A65 (1952) 296
- 1952CO41 J.H. Coon, E.R. Graves and H.H. Barschall, Phys. Rev. 88 (1952) 562
- 1952CR30 D.S. Craig, D.J. Donahue and K.W. Jones, Phys. Rev. 88 (1952) 808
- 1952GE07 R.W. Gelinas and S.S. Hanna, Phys. Rev. 86 (1952) 253
- 1952HU1B Huby and Newns, Phil. Mag. 42 (1952) 1442
- 1952NA1A Nabholz, Stoll and Waffler, Helv. Phys. Acta 25 (1952) 701
- 1952TH1C Thomson, Phys. Rev. 88 (1952) 954
- 1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
- 1952WA31 D. Walker, W.T. Link and W.I.B. Smith, Proc. Phys. Soc. (London) A65 (1952) 861
- 1953BA04 M.E. Battat and F.L. Ribe, Phys. Rev. 89 (1953) 80
- 1953CO02 E.R. Collins, C.D. McKenzie and C.A. Ramm, Proc. Roy. Soc. A216 (1953) 242
- 1953GE01 R.W. Gelinas and S.S. Hanna, Phys. Rev. 89 (1953) 483
- 1953HO48 J.R. Holt and T.N. Marsham, Prog. Phys. Soc. A66 (1953) 1032

1953KR16 J.J. Kraushaar, E.D. Wilson and K.T. Bainbridge, Phys. Rev. 90 (1953) 610  
1953MO1B Montalbetti, Katz and Goldemberg, Phys. Rev. 91 (1953) 659  
1953PE1A Peaslee and Telegdi, Phys. Rev. 92 (1953) 126  
1953SA1A Sawyer and Phillips, Los Alamos Rept.1578 (1953)  
1953ST27 P. Stoll and M. Wachter, Nuovo Cim. 10 (1953) 347  
1953TH1B Thirion, Ann. Phys. 8 (1953) 489  
1953TI02 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A66 (1953) 194  
1953TI1A Titterton and Brinkley, Proc. Phys. Soc. (London) A66 (1953) 579  
1953TU1A Tucker and Gregg, Phys. Rev. 91 (1953) 1579  
1953UE01 R.G. Uebergang and N.W. Tanner, Aust. J. Phys. 6 (1953) 53  
1954AL24 D.L. Allan, Nature 174 (1954) 267  
1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684  
1954CO02 B.L. Cohen and T.H. Handley, Phys. Rev. 93 (1954) 514  
1954CO17 L. Cohen, S.S. Hanna and C.M. Class, Phys. Rev. 94 (1954) 419  
1954DA20 R.B. Day and T. Huus, Phys. Rev. 95 (1954) 1003  
1954DE38 J.A. De Juren and H. Rosenwasser, Phys. Rev. 93 (1954) 831  
1954ER1A Erdos, Stoll, Wachter and Wataghin, Nuovo Cim. 12 (1954) 639  
1954FR03 G.M. Frye Jr., Phys. Rev. 93 (1954) 1086  
1954GO1A Goldemberg and Katz, Can. J. Phys. 32 (1954) 49  
1954GO39 J. Goldemberg and L. Katz, Phys. Rev. 95 (1954) 471  
1954JO10 K.W. Jones, R.A. Douglas, M.T. McEllistrem and H.T. Richards, Phys. Rev. 94 (1954) 947  
1954JO17 C.H. Johnson, H.B. Willard and J.K. Bair, Phys. Rev. 96 (1954) 985  
1954LA1A Lane, AERE-T/R-1289 (1954)  
1954LI48 C.W. Li and R. Sherr, Phys. Rev. 96 (1954) 389  
1954MA1D Major and Biedenharn, Rev. Mod. Phys. 26 (1954) 321  
1954NE1A Nereson et al., LA 1655 (1954)  
1954NI10 W.E. Nickell, Phys. Rev. 95 (1954) 426  
1954RU27 R. Rubin and M. Walter, Helv. Phys. Acta 27 (1954) 163  
1954ST89 P. Stoll, Helv. Phys. Acta 27 (1954) 395  
1954TI16 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A67 (1954) 469  
1954TU1A Tucker and Gregg, Phys. Rev. 93 (1954) 362A

1954ZH1A Zhpinel, *Izv. Akad. Nauk* 18 (1954) 65  
1955AJ61 F. Ajzenberg and T. Lauritsen, *Rev. Mod. Phys.* 27 (1955) 77  
1955AU1A T. Auerbach and J.B. French, *Phys. Rev.* 98 (1955) 1276  
1955BA59 S. Bashkin and R.R. Carlson, *Phys. Rev.* 97 (1955) 1245  
1955BR1A R.J. Breen and M.R. Hertz, *Phys. Rev.* 98 (1955) 599  
1955CA1A Catala, Aguilar and Senent, *An. Real. Soc. Espan. Fis. y Quim.* A51 (1955) 173  
1955CU16 P. Cuer and J. Jung, *J. Phys. Rad.* 16 (1955) 385  
1955CU17 P. Cuer, D. Magnac-Valette and G. Baumann, *Compt. Rend.* 240 (1955) 1880  
1955CZ1A Czyz, *Bull. Acad. Polon. Sci.* 3 (1955) 365  
1955DA1A Dabrowski and Sawicki, *Phys. Rev.* 97 (1955) 1002  
1955HE51 F. Heinrich and R. Rubin, *Helv. Phys. Acta* 28 (1955) 185  
1955HE83 R.E. Heft and W.F. Libby, *Phys. Rev.* 100 (1955) 799  
1955HU1B Hughes and Harvey, BNL-325 (1955)  
1955JA18 D.B. James, W. Kubelka, S.A. Heiberg and J.B. Warren, *Can. J. Phys.* 33 (1955) 219  
1955KH31 L.M. Khromchenko and V.A. Blinov, *Zh. Eksp. Teor. Fiz.* 28 (1955) 741; *JETP (Sov. Phys.)* 1 (1955) 596  
1955KH35 L.M. Khromchenko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 19 (1955) 277; *Columbia Tech. Transl.* 19 (1956) 252  
1955LA1D Lane, *Proc. Phys. Soc. (London)* A68 (1955) 189  
1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, *Phys. Rev.* 97 (1955) 1249  
1955MA84 J.B. Marion, T.W. Bonner and C.F. Cook, *Phys. Rev.* 100 (1955) 91  
1955MI55 M. Miwa, *J. Phys. Soc. Jpn.* 10 (1955) 173  
1955PE1B Peshkin, Roberts and Telegdi, *Phys. Rev.* 98 (1955) 1204A  
1955RE16 J.B. Reynolds, *Phys. Rev.* 98 (1955) 1289  
1955SA1C Sawicki, *Nuovo Cim.* 1 (1955) 957  
1955ST85 J.F. Streib, *Phys. Rev.* 100 (1955) 1797A  
1955TI1A Titterton, *Prog. Nucl. Phys.* 4 (1955) 1  
1956AB1A Abraham, *Nucl. Phys.* 1 (1956) 415  
1956BE14 J. Benveniste, R.G. Finke and E.A. Martinelli, *Phys. Rev.* 101 (1956) 655  
1956BO36 P. Bouchez, J. Tobaillem, J. Robert, R. Muxart, R. Mellet and P. Daud, *J. Phys. Rad.* 17 (1956) 363  
1956BU83 D.S.T. Bunbury, S. Devons, G. Manning and J.H. Towle, *Proc. Phys. Soc. (London)* A69 (1956) 165

1956CH1B Cheng, Rev. Mod. Phys. 28 (1956) 45  
 1956CO61 H.E. Conzett and R. Silver, Bull. Amer. Phys. Soc. 1 (1956) 388, L7  
 1956CZ1A Czyz, Acta Phys. Pol. 15 (1956) 129  
 1956DA1A Davaney, Los Alamos Rept. 1960 (1956)  
 1956DA23 R.B. Day, Phys. Rev. 102 (1956) 767  
 1956FE1A Feingold, Phys. Rev. 101 (1956) 258  
 1956FR18 G.M. Frye Jr. and J.H. Gammel, Phys. Rev. 103 (1956) 328  
 1956GE1A Gelinas and Hanna, Phys. Rev. 104 (1956) 1681  
 1956GO1E Gorlov, Gokhberg, Morozov and Otroshchenko, Dokl. Akad. Nauk SSSR 111 (1956) 791  
 1956GO62 G.V. Gorlov, B.M. Gokhberg, V.M. Morozov and G.A. Otroshchenko, Dokl. Akad. Nauk SSSR 110 (1956) 963; Sov. Phys. Dokl. 1 (1956) 598  
 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28  
 1956HA90 J.W. Haffner, Phys. Rev. 103 (1956) 1398  
 1956HE16 A. Hemmendinger, Bull. Amer. Phys. Soc. 1 (1956) 96, N7  
 1956HO93 R. Hofstadter, Rev. Mod. Phys. 28 (1956) 214  
 1956HU1A Hughes and Carter, BNL-400 (1956)  
 1956JU1D Jung and Cuer, J. Phys. Rad. 17 (1956) 555  
 1956KI54 B.B. Kinsey and T. Stone, Phys. Rev. 103 (1956) 975  
 1956KU1A Kurath, Phys. Rev. 101 (1956) 216  
 1956LI37 J.G. Likely and F.P. Brady, Phys. Rev. 104 (1956) 118  
 1956MA91 J.B. Marion, G. Weber and F.S. Mozer, Phys. Rev. 104 (1956) 1402  
 1956ME1A Meshkov and Ufford, Phys. Rev. 101 (1956) 734  
 1956NE13 G.C. Neilson and J.B. Warren, Phys. Rev. 103 (1956) 1758  
 1956RE04 J.B. Reynolds and K.G. Standing, Phys. Rev. 101 (1956) 158  
 1956RI34 F.L. Ribe, Phys. Rev. 103 (1956) 741  
 1956SO21 I.L. Sokolov, M.M. Sulkovskaia, E.I. Karpushkina and E.A. Albitskaia, Zh. Eksp. Teor. Fiz. 30 (1956) 1007; JETP (Sov. Phys.) 3 (1956) 740  
 1956SO33 I.L. Sokolov, M.M. Sulkovskaia, E.A. Albitskaia and E.I. Karpushkina, Dokl. Akad. Nauk SSSR 111 (1956) 1219; Sov. Phys. Dokl. 1 (1957) 773  
 1956ST30 K. Strauch and F. Titus, Phys. Rev. 104 (1956) 191  
 1956TU1A Tulinov, Zh. Eksp. Teor. Fiz. 31 (1956) 698; JETP (Sov. Phys.) 4 (1957) 596  
 1956WA03 J.B. Warren, T.K. Alexander and G.B. Chadwick, Phys. Rev. 101 (1956) 242

1956WI04 H.B. Willard, J.K. Bair, J.D. Kington and H.O. Cohn, Phys. Rev. 101 (1956) 765  
 1957BA18 G.A. Bartholomew and P.J. Campion, Can. J. Phys. 35 (1957) 1347  
 1957BA1H Barker, Phil. Mag. 2 (1957) 780  
 1957BE71 A.A. Bergman, A.I. Isakov, I.P. Popov and F.L. Shapiro, Zh. Eksp. Teor. Fiz. 33 (1957) 9; JETP (Sov. Phys.) 6 (1958) 6  
 1957BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Sov. J. At. Energy 3 (1957) 987  
 1957BR97 C.P. Browne, Bull. Amer. Phys. Soc. 2 (1957) 350, N3  
 1957CO53 H.E. Conzett, Phys. Rev. 105 (1957) 1324  
 1957CO68 H.E. Conzett and D.G. Hoffman, Bull. Amer. Phys. Soc. 2 (1957) 382, J10  
 1957DU37 J.W.M. DuMond, Ann. Phys. 2 (1957) 283  
 1957EL1C Elpidinskii, Shapiro and Stranikh, Sov. J. At. Energy, Suppl. 5 (1957) 56  
 1957FE1A Feingold, Bull. Amer. Phys. Soc. 2 (1957) 392  
 1957FR1B French and Fujii, Phys. Rev. 105 (1957) 652  
 1957HI56 N.M. Hintz, Phys. Rev. 106 (1957) 1201  
 1957HO1E Hofstadter, Ann. Rev. Nucl. Sci., Vol. 7 (1957)  
 1957HU1D Hughes and Schwartz, BNL-325, Suppl. 1 (1957)  
 1957JA37 N. Jarmie, J.D. Seagrave et al., LA-2014 (1957)  
 1957KA1B Katsaurov, Musaelian and Popov, Sov. J. At. Energy, Suppl. 5 (1957) 67  
 1957KO1A Kopaleishvili, Zh. Eksp. Teor. Fiz. 33 (1957) 788; JETP (Sov. Phys.) 6 (1958) 606  
 1957LE1E Levinson and Banerjee, Ann. Phys. 2 (1957) 471  
 1957MA04 D.R. Maxson and E.F. Bennett, Bull. Amer. Phys. Soc. 2 (1957) 180, E5  
 1957MA1E Mamasakhlisov and Chilashvili, Zh. Eksp. Teor. Fiz. 32 (1957) 806; JETP (Sov. Phys.) 5 (1957) 661  
 1957MA57 J.B. Marion, Nucl. Phys. 4 (1957) 282; Erratum Nucl. Phys. 4 (1957) 492  
 1957ME1B Meyer-Berkhout and Hofstadter, Bull. Amer. Phys. Soc. 2 (1957) 390  
 1957NE1B Nemilov and Pisarevskii, Zh. Eksp. Teor. Fiz. 32 (1957) 139; JETP (Sov. Phys.) 5 (1957) 115  
 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754  
 1957SC12 C.D. Schrader, J. Benveniste and J.H. Zenger, Bull. Amer. Phys. Soc. 2 (1957) 309, H9  
 1957SI36 R. Silver, Thesis, University of California (1957); UCRL 3887 (1957)  
 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809  
 1957SO1A Soper, Phil. Mag. 2 (1957) 1219

- 1957TY35 H. Tyren, T.A.J. Maris and P. Hillman, *Nuovo Cim.* 6 (1957) 1507
- 1957VA12 S.S. Vasilev, V.V. Komarov and A.M. Popova, *Zh. Eksp. Teor. Fiz.* 33 (1957) 527; *JETP (Sov. Phys.)* 6 (1958) 411
- 1957WI24 D.H. Wilkinson, *Phil. Mag.* 2 (1957) 83
- 1957WR37 H.W. Wright, E.I. Wyatt, S.A. Reynolds, W.S. Lyon and T.H. Handle, *Nucl. Sci. Eng.* 2 (1957) 427
- 1958AL05 R.C. Allen and N. Jarmie, *Phys. Rev.* 111 (1958) 1129
- 1958BA1C Barton and Smith, *Phys. Rev.* 110 (1958) 1143
- 1958BA59 J.B. Ball, A.W. Fairhall and I. Halpern, *Bull. Amer. Phys. Soc.* 3 (1958) 322, H6
- 1958BE10 O. Beckman and R. Sandstrom, *Nucl. Phys.* 5 (1958) 595
- 1958BE1C Benade, *Conf. on Photonucle. Reactions, National Bureau of Standards* (1958)
- 1958BO76 R.O. Bondelid and C.A. Kennedy, *NRL Rept.* 5083 (1958)
- 1958BR16 A. Bratenahl, J.M. Peterson and J.P. Steering, *Phys. Rev.* 110 (1958) 927
- 1958BU02 E. Bujdoso, *Nucl. Phys.* 6 (1958) 107
- 1958BU38 W.E. Burcham, G.P. McCauley, D. Bredin, W.M. Gibson, D.J. Prowse and J. Rotblat, *Nucl. Phys.* 5 (1958) 141
- 1958CA1C Cameron, *Bull. Amer. Phys. Soc.* 3 (1958) 227
- 1958CH26 S.W. Chen, *Bull. Amer. Phys. Soc.* 3 (1958) 189, K14
- 1958CH35 R. Chiba, H.E. Conzett, H. Morinaga, N. Mutsuro, K. Shoda and M. Kimura, *Bull. Amer. Phys. Soc.* 3 (1958) 419, N10
- 1958FO1A Fowler, *Astrophys. J.* 127 (1958) 551
- 1958HA1D Haig, *Nucl. Phys.* 7 (1958) 429
- 1958HA22 E.W. Hamburger and J.R. Cameron, *Bull. Amer. Phys. Soc.* 3 (1958) 381, T7
- 1958HE1F Henley, *Conf. on Photonucle. Reactions, National Bureau of Standards* (1958)
- 1958HO1C Howerton, *UCRL* 5226 (1958)
- 1958HU18 D.J. Hughes and R.B. Schwartz, *BNL-325, 2nd Ed.* (1958); *BNL-325, 2nd Ed., Suppl. I* (1960)
- 1958KE30 B.D. Kern and W.E. Kreger, *Phys. Rev.* 112 (1958) 926
- 1958LE20 R.D. Leamer, *Bull. Amer. Phys. Soc.* 3 (1958) 61, VA1
- 1958LI1A Livesey, *Conf. on Photonucle. Reactions, National Bureau of Standards* (1958)
- 1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, *Nucl. Phys.* 7 (1958) 1
- 1958MC65 J.A. McCray, *Bull. Amer. Phys. Soc.* 3 (1958) 418, N5
- 1958MI92 P.D. Miller and G.C. Phillips, *Phys. Rev.* 112 (1958) 2048

1958MU07 R.B. Murray and H.W. Schmitt, Bull. Amer. Phys. Soc. 3 (1958) 381, T5  
1958NE38 G.C. Neilson, W.K. Dawson and J.T. Sample, Bull. Amer. Phys. Soc. 3 (1958) 323, H11  
1958PA1B Palfrey and Tautfest, Bull. Amer. Phys. Soc. 3 (1958) 173  
1958RI34 P.J. Riley, J.B. Warren and G.M. Griffiths, Bull. Amer. Phys. Soc. 3 (1958) 330, N3  
1958RY77 T.W. Rybka and L. Katz, Phys. Rev. 110 (1958) 1123  
1958SA17 J.T. Sample, W.K. Dawson and G.C. Neilson, Bull. Amer. Phys. Soc. 3 (1958) 323, H13  
1958SK1A Skyrme, Proc. Rehovoth Conf. (North-Holland Publishing Co., Amsterdam, 1958)  
1958SM1A Smith and Baker, Bull. Amer. Phys. Soc. 3 (1958) 172  
1958ST1A Stewart, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
1958SW65 C.P. Swann, F.R. Metzger and V.K. Rasmussen, Bull. Amer. Phys. Soc. 3 (1958) 187, K3  
1958TI1A Titterton, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
1958TY46 H. Tyren and T.A.J. Maris, Nucl. Phys. 6 (1958) 82  
1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10  
1958WH35 C. Whitehead, W.R. McMurray, M.J. Aitken, N. Middlemas and C.H. Collie, Phys. Rev. 110 (1958) 941  
1958WI38 D.H. Wilkinson, Phys. Rev. 109 (1958) 1614  
1959BA1H Bame and Cubitt, Phys. Rev. (1959)  
1959HO03 H.D. Holmgren and R.L. Johnston, Phys. Rev. 113 (1959) 1556  
1959SW63 C.P. Swann, V.K. Rasmussen and F.R. Metzger, Phys. Rev. 114 (1959) 862  
HA55A Unknown Source



