

Energy Levels of Light Nuclei $A = 7$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A227* (1974), p. 1. This version of $A = 7$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 31, 1973)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 7$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^7\text{H}\$](#) , [\${}^7\text{He}\$](#) , [\${}^7\text{Li}\$](#) , [\${}^7\text{Be}\$](#) , [\${}^7\text{B}\$](#)

B. Tables of Recommended Level Energies:

[Table 7.1](#): Energy levels of ${}^7\text{Li}$

[Table 7.5](#): Energy levels of ${}^7\text{Be}$

C. [References](#)

D. Figures: [\${}^7\text{Li}\$](#) , [\${}^7\text{Be}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

${}^7\text{H}$

(Not illustrated)

A search for ${}^7\text{H}$ in ${}^7\text{Li}(\pi^-, \pi^+){}^7\text{H}$ was unsuccessful (1965GI10). See also (1968CE01).

${}^7\text{He}$

(Fig. 10)

Mass of ${}^7\text{He}$: From the Q of the ${}^7\text{Li}(t, {}^3\text{He}){}^7\text{He}$ reaction, the atomic mass excess of ${}^7\text{He}$ is 26.11 ± 0.03 MeV. ${}^7\text{He}$ is unbound with respect to ${}^6\text{He} + n$ by 0.44 ± 0.03 MeV (1968ST1J): $\Gamma < 0.2$ MeV (1973LI02).

GENERAL:

See (1960GO1B, 1965BO1C, 1965LA1B, 1967CO36, 1970LO1E, 1972CA37, 1972GA1L, 1972PN1A, 1973JU2A) and (1966LA04).

1. ${}^7\text{Li}(t, {}^3\text{He}){}^7\text{He}$ $Q_m = -11.18$
 $Q_0 = -11.18 \pm 0.03$ (1968ST1J).

The ${}^3\text{He}$ particles to the ground state of ${}^7\text{He}$ have been observed at $E_t = 22$ MeV. The width of the ground state is 160 ± 30 keV; for a radius of 2.2 fm and $l_n = 1$, this width is 0.22 of the Wigner limit. The angular distribution is peaked in the forward direction. No other states of ${}^7\text{He}$ were observed for $E_x < 2.4$ MeV (1967ST04, 1968ST1J). See also (1968CE01).

2. ${}^7\text{Li}(n, p){}^7\text{He}$ $Q_m = -10.42$

At $E_n = 14.8$ MeV, a proton group is reported corresponding to ${}^7\text{He}_{\text{g.s.}}$: $\Gamma < 0.2$ MeV (1973LI02). See also (1967ME11, 1971KO24) and ${}^8\text{Li}$.

3. ${}^9\text{Be}(n, {}^3\text{He}){}^7\text{He}$ $Q_m = -17.20$

Not reported.

⁷Li
(Figs. 8 and 10)

GENERAL: (See also (1966LA04).)

Shell model: (1961KO1A, 1965CO25, 1965KU09, 1965VO1A, 1966BA26, 1966HA18, 1966WI1E, 1967BO1C, 1967BO22, 1967CO32, 1967FA1A, 1969GU03, 1969TA1H, 1969VA1C, 1970ZO1A, 1971CO28, 1972LE1L, 1973HA49, 1973KU03).

Cluster model: (1965NE1B, 1968HA1G, 1968KU1B, 1969ME1C, 1969SM1A, 1969VE1B, 1969WI21, 1970BA1Q, 1972HA06, 1972HI16, 1972JA23, 1972KU12, 1972LE1L, 1973KU03, 1973KU12).

Rotational and deformed models: (1965VO1A, 1966EL08).

Special levels: (1966BA26, 1966EL08, 1967BO22, 1967CO32, 1967FA1A, 1969GU03, 1969HA1G, 1969HA1F, 1970FR1C, 1971CO28, 1972BB26, 1973AS02, 1973FE1J, 1973MA1K).

Electromagnetic transitions: (1965CO25, 1965KU09, 1966BA26, 1966EL08, 1967BO22, 1968EL06, 1968KU1D, 1969HA1G, 1969HA1F, 1969VA1C, 1973AS02, 1973HA49, 1973HA1V, 1973SU1C).

Astrophysical questions: (1967DA1C, 1967MI1A, 1968HA1C, 1969BA2A, 1970BA1M, 1972CL1A, 1972KO1E, 1972RA30, 1973AU1H, 1973LA19, 1973RE1G, 1973SA1J, 1973SC1T, 1974AU1A).

Special reactions: (1965GR1C, 1965ZH1A, 1967AU1B, 1968YI01, 1969GA18, 1969YI1A, 1972HA06, 1972RA30, 1972VO06, 1973KO1D, 1973KU03, 1973LA19, 1973OS1C, 1973PF02).

Muon capture: (1965LO1B, 1969WU1A, 1970FA15, 1971DE2D, 1973MU11).

Pion capture and reactions: (1966DA1A, 1966DE1G, 1968BO32, 1968BO1T, 1968KU1B, 1968LO1A, 1968NO1A, 1968PE1B, 1968WI1B, 1969BU1C, 1969KO30, 1969MI10, 1969MI1G, 1969MO1E, 1970BA1E, 1970BO1V, 1970JA23, 1971CA01, 1971CA1J, 1971FA09, 1971GO22, 1971KO02, 1972HU1A, 1972SW1A, 1973BA2R, 1973BA2V, 1973BA2G, 1973DO1F, 1973NY04, 1973PE1E, 1973SQ01).

Kaon reactions: (1973BA1Y).

Other topics: (1965CO25, 1965VO1A, 1966DE1E, 1966HA18, 1966WI1E, 1966YO1B, 1967BO1C, 1967CA17, 1967FA1A, 1968EL06, 1969GU03, 1969HE1N, 1969HO1M, 1970HO1J, 1970ZO1A, 1971BA2Y, 1971CH1H, 1971GR16, 1971ZA1D, 1972AB14, 1972AN05, 1972BB26, 1972CA37, 1972DA21, 1972FR09, 1972GA1L, 1972LE1L, 1972PN1A, 1973BA1Y, 1973CL09, 1973JU2A, 1973KU03, 1973MA48, 1973RO1R).

Ground state properties: (1965CO25, 1965HU13, 1965KU09, 1965MO17, 1965PR04, 1965VO1A, 1966BA26, 1966CA1F, 1966EL08, 1966WI1E, 1967BO22, 1967PA1G, 1967SH05, 1967SH14, 1968PE16, 1968PE1B, 1969GU03, 1969VA1C, 1972FR09, 1972LE1L, 1973DO1F, 1973MA1K).

$$\mu = +3.2564 \text{ nm (1969FU11, 1971SH26);}$$

Table 7.1: Energy levels of ${}^7\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	—	stable	1, 4, 6, 12, 13, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 27, 28, 29, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54
0.477611 ± 0.012	$\frac{1}{2}^-; \frac{1}{2}$	$\tau_m = 105 \pm 5$ fsec	γ	6, 12, 13, 16, 18, 19, 20, 22, 23, 24, 27, 30, 33, 35, 38, 42, 43, 45, 47, 49, 50, 51, 53, 54
4.633 ± 8	$\frac{7}{2}^-; \frac{1}{2}$	$\Gamma = 93 \pm 8$ keV	t, α	3, 12, 18, 19, 20, 21, 22, 23, 24, 33, 35, 38, 39, 42
6.675 ± 54	$\frac{5}{2}^-; \frac{1}{2}$	875^{+200}_{-100} keV	t, α	3, 18, 20, 21, 24, 35
7.467 ± 4	$\frac{5}{2}^-; \frac{1}{2}$	89 ± 7 keV	n, t, α	2, 3, 7, 11, 12, 18, 20, 24, 33, 35, 42
9.61 ± 81	$\frac{7}{2}^-; \frac{1}{2}$	broad	n, t, α	2, 3, 12, 35
10.25 ± 100	$\frac{3}{2}^-; \frac{1}{2}$	1.40 ± 0.10 MeV	n, α	7, 18
11.245 ± 31	$\frac{3}{2}^-; \frac{3}{2}$	258 ± 33 keV	n, p	7, 8, 18, 33
16.8		9.3 MeV	γ , n	17

$$\mu = 3.255985 (2) \text{ nm [quoted in (1973CO1P)];}$$

$$Q = -0.058 \pm 0.010 \text{ b (1966IS01);}$$

$$Q = -0.04 \text{ b (1969FU11); see also (1971SH26, 1972BA77).}$$

$$B(\text{E2: } \frac{3}{2}^- \rightarrow \frac{1}{2}^-) = 8.3 \pm 0.6 \text{ e}^2 \cdot \text{fm}^4 \text{ (1972HA06, 1973HA47);}$$

$$= 7.4 \pm 0.1 \text{ e}^2 \cdot \text{fm}^4 \text{ (1972BA77) [see also (1973HA47)].}$$

1. ${}^3\text{H}(\alpha, \gamma){}^7\text{Li}$

$$Q_m = 2.4668$$

Excitation functions and angular distributions have been studied for $E_\alpha = 0.5$ to 1.9 MeV (1959HO03, 1961GR27). The cross section rises smoothly as expected for a direct capture process: at $E_\alpha = 1.32$ MeV, $\sigma = 3.58 \pm 0.06 \mu\text{b}$ and the corresponding reduced cross section factor $S = 0.064 \pm 0.016 \text{ keV} \cdot \text{b}$ (1961GR27). Cross sections of (1961GR27) are 2 to 2.5 times higher

Table 7.2: ${}^7\text{Li}$ levels from ${}^3\text{H} + {}^4\text{He}$

E_x (MeV \pm keV)	J^π	l_α	LS term	R (fm)	θ_α^2 ^a	$\theta_{n_0}^2$	$\theta_{n_1}^2$ ^b	Refs.
4.65 \pm 50	$\frac{7}{2}^-$	3	${}^2\text{F}_{7/2}$	4.0	0.57 \pm 0.04			(1967SP10)
4.65 \pm 20	$\frac{7}{2}^-$	3	${}^2\text{F}_{7/2}$	4.4	0.37			(1968IV01)
6.64 \pm 100	$\frac{5}{2}^-$	3	${}^2\text{F}_{5/2}$	4.0	1.36 \pm 0.13	0.000 \pm 0.002		(1967SP10)
6.79 \pm 90	$\frac{5}{2}^-$	3	${}^2\text{F}_{5/2}$	4.4	0.52			(1968IV01)
7.47 \pm 30	$\frac{5}{2}^-$	3	${}^4\text{P}_{5/2}$	4.0	0.011 \pm 0.001	0.26 \pm 0.02		(1967SP10)
9.67 \pm 100	$\frac{7}{2}^-$	3	${}^4\text{D}_{7/2}$	4.0	0.53 \pm 0.22		2.3 \pm 0.7	(1967SP10)

^a $\gamma^2 / (\frac{3}{2}\hbar^2 / \mu a^2)$.

^b To ${}^6\text{Li}^*(2.19)$.

than those of (1959HO03). See (1966LA04) for further comments and (1972BA77, 1972HA06, 1972SM02).

2. ${}^3\text{H}(\alpha, n){}^6\text{Li}$

$$Q_m = -4.784$$

$$E_b = 2.4668$$

The cross section for this reaction has been measured for $E_\alpha = 11$ to 18 MeV: the data show the effect of ${}^7\text{Li}^*(7.47)$ and indicate a broad resonance near $E_\alpha = 16.8$ MeV [${}^7\text{Li}^*(9.6)$]. The level parameters derived from this reaction and from reaction 3 (1967SP10) are displayed in Table 7.2. See also (1972BB26; theor.).

3. ${}^3\text{H}(\alpha, \alpha){}^3\text{H}$

$$E_b = 2.4668$$

The elastic scattering has been studied for $E_\alpha = 3.6$ to 18.2 MeV. The excitation curves show the effects of ${}^7\text{Li}^*(4.63, 6.68, 7.47, 9.61)$. The derived level parameters are displayed in Table 7.2. Polarization parameters are calculated (1967SP10, 1968IV01). Polarization measurements are reported for $E_t = 6.0$ to 12.3 MeV (1968KE03, 1971AR1K) and angular distributions have also been measured for $E_\alpha = 2.13$ to 2.98 MeV (1971CH42). See also (1970LI06) and (1968BR1H, 1971KU22, 1972BB26, 1972CL1C, 1972NE17, 1973KO1Q; theor.).

4. ${}^4\text{He}(\alpha, p){}^7\text{Li}$

$$Q_m = -17.348$$

See (1958BU38).

5. ${}^6\text{He}(p, t){}^4\text{He}$

$$Q_m = 7.511$$

$$E_b = 9.978$$

See (1971PO1A).

6. ${}^6\text{Li}(n, \gamma){}^7\text{Li}$

$$Q_m = 7.2506$$

$$Q_0 = 7250.0 \pm 0.5 \text{ keV (1968SP01);}$$

$$Q_0 = 7250.3 \pm 0.9 \text{ keV (1972OP01);}$$

$$Q_0 = 7251.0 \pm 1.0 \text{ keV (1967RA24);}$$

$$Q_0 = 7250.6 \pm 1.5 \text{ keV (E.T. Journey, private communication).}$$

The total radiative capture cross section for thermal neutrons is 38.5 ± 3.0 mb (E.T. Journey, private communication). Two γ -rays with $E_\gamma = 7247 \pm 2$ and 6769 ± 2 keV are observed corresponding to transitions to ${}^7\text{Li}^*(0, 0.48)$ with branching ratios of 61 ± 3 and $39 \pm 2\%$, respectively[†]. Gamma rays with $E_\gamma = 4.63$, 4.15 and 2.62 MeV corresponding to the decays of, and the transition to, ${}^7\text{Li}^*(4.63)$ are not observed: upper limits are 2, 2 and 6%, respectively (1967TH05), 1% (E.T. Journey, private communication). See also (1968SP01). For astrophysical implications of this reaction see (1968FOZY).

7. ${}^6\text{Li}(n, n){}^6\text{Li}$

$$E_b = 7.2506$$

The total cross section has been measured for $E_n = 4$ eV to 29 MeV: see (1960HU08, 1964ST25, 1966LA04). Recent measurements have been carried out at $E_n = 10$ to 1236 keV (1968HI1E), 50 to 650 keV (1968FA1D), 100 to 1500 keV (1972ME17), 0.7 to 30 MeV (1973GO2B, and C.A. Goulding, private communication) and 2.5 to 15 MeV (1971FO1P, 1971FO24). A pronounced resonance occurs at $E_n = 255$ keV with a peak cross section of about 11.0 b (1960HU08, 1968HI1E, 1972ME17). See also (1968FA1D). The elastic contribution is 7.2 b (1961LA1A). No other clearly defined resonance is observed although the total cross section exhibits its a broad maximum at $E_n \approx 5$ MeV (1954JO17, 1960HU08). The coherent scattering length (thermal, bound) is $1.8 + 0.25i$ fm (1969BA1P, 1973MU14).

Angular distributions are tabulated by (1970GA1A) and in reaction 11 of ${}^6\text{Li}$. All observations near the 255 keV resonance are consistent with p-wave formation of a $J^\pi = \frac{5}{2}^-$ level [${}^7\text{Li}^*(7.47)$]. Table 7.3 gives the resonance parameters compared with those for ${}^7\text{Be}^*(7.18)$. These states are believed to have a ${}^4P_{5/2}$ character, in agreement with their large θ_n^2 and θ_p^2 (1959GA08, 1963MC09).

The excitation function for 3.56 MeV γ -rays exhibits an anomaly, also seen in the (n, p) reaction [reaction 6]. The data are well fitted assuming $E_{\text{res}} = 3.50$ and 4.60 MeV [$E_x = 10.25 \pm 0.10$ and

[†] $E_\gamma = 477.6 \pm 0.5$, 6770.4 ± 1.5 and 7246.6 ± 1.5 keV, and the branching ratios are 61 ± 1 and $39 \pm 1\%$ to the ground state and to $E_x = 477.6 \pm 0.5$ keV (E.T. Journey, private communication).

Table 7.3: Resonance parameters for 7.5 – 7.2 MeV levels in ${}^7\text{Li}$ and ${}^7\text{Be}$

Reaction	${}^6\text{Li} + \text{n}$		${}^6\text{Li} + \text{p}$
	a	b	c
Refs.			
E_r (keV, lab)	262	255	1840
$\Gamma(E_r)$ (keV, c.m.)	154	65	836
E_λ (keV above g.s.)	7700		7580
$\Gamma_{\text{n,p}}(E_r)$ (keV, c.m.)	118	46.8	798
radius (n, p) in fm	3.94		4.08
$\gamma_{\text{n,p}}^2$ (MeV · fm)	4.85		5.02
$\theta_{\text{n,p}}^2$	0.26		0.28
$\Gamma_\alpha(E_r)$ (keV, c.m.)	36	18.5	38
radius (α) in fm	4.39		4.39
γ_α^2 (MeV · fm)	0.101		0.101
θ_α^2	0.012		0.012

^a (1959GA08: see (1963MC09)). See also (1965SC07).

^b (1972ME17).

^c (1963MC09).

11.19 ± 0.05 MeV]. $T = \frac{1}{2}$ and $\frac{3}{2}$, $\Gamma_{\text{cm}} = 1.40 \pm 0.10$ and 0.27 ± 0.05 MeV, respectively. Both states have $J^\pi = \frac{3}{2}^-$. The reduced widths for the $T = \frac{3}{2}$ state [${}^7\text{Li}^*(11.19)$] are $\theta_{\text{n}}^2 = 2 \times 10^{-4}$, $\theta_{\text{n}''}^2 = 0.16$ [to ${}^6\text{Li}^*(3.56)$] and $\theta_{\text{p}}^2 = 0.09$ (1969PR04). Cross-section measurements are also reported at $E_{\text{n}} = 1$ to 100 keV (1970AS1E), 4.8, 5.7 and 7.5 MeV (1968HO03), 10 MeV (1967CO01) and 14 MeV (1966ME1C). See also (1966AG1A, 1971RE07, 1973LA26; theor.).

Polarization measurements are reported at $E_{\text{n}} = 3.4$ MeV (1968WO1F) and at 4.4 MeV (1966ST09). See also (1967BE1F; theor.) and (1966LA04) for earlier measurements.

8. ${}^6\text{Li}(\text{n}, \text{p}){}^6\text{He}$

$$Q_{\text{m}} = -2.727$$

$$E_{\text{b}} = 7.2506$$

The excitation function, measured from threshold to $E_{\text{n}} = 8.9$ MeV, exhibits an anomaly at $E_{\text{n}} = 4.6$ MeV: see reaction 6 (1969PR04).

See also (1966JE1B, 1971CU1B, 1971PR09, 1972ED01, 1973BO1Y).

9. ${}^6\text{Li}(\text{n}, \text{d}){}^5\text{He}$

$$Q_{\text{m}} = -2.36$$

$$E_{\text{b}} = 7.2506$$

See (1969LI1F, 1973BO1Y) and (1966LA04).

$$10. \text{}^6\text{Li}(n, 2n)\text{}^5\text{Li} \quad Q_m = -5.66 \quad E_b = 7.2506$$

See (1963AS01).

$$11. \text{}^6\text{Li}(n, \alpha)\text{}^3\text{H} \quad Q_m = 4.7839 \quad E_b = 7.2506$$

$$Q_0 = 4.794 \pm 0.012 \text{ (1967DE15)}.$$

Excitation functions and angular distributions are summarized in (1960HU08, 1964ST25, 1966LA04, 1970GA1A). More recent cross-section measurements are reported by (1967CO1N: $E_n = 10.7$ to 102 keV), (1966BA1V: 25, 67 and 100 keV), (1970FO1E: 81.8 to 517 keV) [see below for thermal measurements]. Recent angular distributions are given in (1971OV1A: 0.10 to 1.80 MeV) and in (1966RO1L: 0.25, 0.39 and 0.60 MeV).

The isotropic thermal cross section is 938 ± 6 b (1970ME1F), 940 ± 4 b (1970SO1A) [the value listed in (1966LA04) is in error: it should have been 945 b]. Below 5 keV, the total cross section is given by $\sigma = (149.5/\sqrt{E(\text{eV})}) + 0.696$ b (1970SO1A). (1970SO1A) have also measured the ratio of (n, α) cross sections for ${}^6\text{Li}$ and ${}^{10}\text{B}$ in the range $E_n = 10$ eV to 80 keV.

A resonance occurs at $E_n = 258$ keV, with $\sigma_{\text{max}} = 2.75$ b (1959BA46), 2.80 ± 0.22 b (1959GA08). The resonance is formed by p-waves, $J^\pi = \frac{5}{2}^-$, and has a large neutron width and a small α -width: see Table 7.3 (1959GA08). Above the resonance the cross section decreases monotonically to $E_n = 18$ MeV, except for a slight bump near $E_n = 1.6$ to 2.1 MeV (1959GA08, 1959MU25). See also (1970MC1A, 1971MA1Y, 1972ZV1A, 1973BO1Y, 1974BA1K), (1966JE1B, 1968GI1D, 1970DE1H) and (1966BL1C, 1966MA1L, 1968FA1D, 1968SE1A, 1972LA1F). For astrophysical implications, see (1964FO1A, 1965BU1C).

$$12. \text{(a) } \text{}^6\text{Li}(d, p)\text{}^7\text{Li} \quad Q_m = 5.0260$$

$$\text{(b) } \text{}^6\text{Li}(d, np)\text{}^6\text{Li} \quad Q_m = -2.22464$$

$$Q_0 = 5.024 \pm 0.007 \text{ (1967SP09)}.$$

Angular distribution measurements have been recently carried out at $E_d = 1$ to 2 MeV (1966BR25; p_0, p_1), 1.48 to 2.94 MeV (1968TU1A; p_0, p_1), 1.5 MeV (1965RI09; p_0, p_1, p_2), 2.9 MeV (1966RO1J; p_0, p_1), 4.5 to 5.5 MeV (1970PO03; p_0, p_1) and 12 MeV (1967SC29; p_0, p_1). See also (1969HO39, 1969VI06) and (1966LA04) for earlier references. The p_0 and the p_1 [$E_x = 477 \pm 2$ keV (1959AJ76)] groups show stripping patterns with $l_n = 1$. The p_2 [$E_x = 4.630 \pm 0.009$ MeV, $\Gamma_{\text{cm}} = 93 \pm 8$ keV (1966LA04)] angular distribution is isotropic (1960HA14). In addition the excitation of ${}^7\text{Li}^*(7.47)$ with $E_x = 7.464 \pm 0.010$ MeV, $\Gamma_{\text{cm}} = 91 \pm 8$ keV, is also reported

(1957BR97). See also (1966LA04). Ratios of observed θ_n^2 [see Table 7.3 in (1966LA04)] are consistent with assignments ^{22}P to $^7\text{Li}^*(0, 0.48)$ and ^{24}P to $^7\text{Li}^*(7.47)$ (1960HA14, 1960MA32). At $E_d = 12$ MeV spectroscopic factors for $^7\text{Li}^*(0, 0.48)$ [$S = 0.90, 1.15$] derived by DWBA are in good accord with the shell model calculations of (1967CO32, 1967SC29). The angular correlation between p_1 and the 0.48 MeV γ -rays is isotropic (1959AJ76): $^7\text{Li}^*(0.48)$ has $J^\pi = \frac{1}{2}$.

Using vector polarized deuterons with $E_d = 10$ MeV, (1970FI07) determined the probabilities p_j for transfer of a neutron with total angular momentum j . The results are in quite good agreement with the shell-model calculations of (1965CO25, 1967CO32). The circular polarization of the 0.48 MeV γ -rays has been determined by (1966SC1E). See also ^8Be (1966AU1A, 1966BE1E, 1968BE1P, 1969LE22) and (1967OG1A).

A kinematically complete study of reaction (b) at $E_d = 10$ MeV shows pronounced final state interactions via $^7\text{Li}^*(7.47)$ and possibly $^7\text{Li}^*(9.6)$ [$\Gamma = 0.5 \pm 0.1$ MeV] (1971VO07). See also ^8Be .

13. $^6\text{Li}(t, d)^7\text{Li}$ $Q_m = 0.9930$

See (1954AL35, 1961HO21, 1967BI1E).

14. $^6\text{Li}(\alpha, ^3\text{He})^7\text{Li}$ $Q_m = -13.3279$

Not reported.

15. (a) $^6\text{Li}(^6\text{Li}, p\alpha)^7\text{Li}$ $Q_m = 3.552$
 (b) $^6\text{Li}(^9\text{Be}, ^8\text{Be})^7\text{Li}$ $Q_m = 5.586$

See (1966LA04), ^{12}C in (1968AJ02) and ^{15}N in (1970AJ04).

16. $^7\text{Li}(\gamma, \gamma)^7\text{Li}$

For a listing of lifetime measurements of $^7\text{Li}^*(0.48)$, see Table 7.4 in (1966LA04): $\tau_m = 0.107 \pm 0.005$ psec.

17. (a) $^7\text{Li}(\gamma, n)^6\text{Li}$ $Q_m = -7.2506$
 (b) $^7\text{Li}(\gamma, 2n)^5\text{Li}$ $Q_m = -12.91$

(c) ${}^7\text{Li}(\gamma, p){}^6\text{He}$	$Q_m = -9.978$
(d) ${}^7\text{Li}(\gamma, pn){}^5\text{He}$	$Q_m = -11.84$
(e) ${}^7\text{Li}(\gamma, d){}^5\text{He}$	$Q_m = -9.61$
(f) ${}^7\text{Li}(\gamma, t){}^4\text{He}$	$Q_m = -2.4668$

Reports on the structure of the (γ, n) cross section [reaction (a)] differ widely. The total photoneutron cross section rises sharply from 10 MeV to reach a broad plateau of about 1.5 mb from 14 to 20 MeV, decreases more slowly to about 0.5 mb at 25 MeV and then remains approximately constant to 30 MeV. There are indications of weak structure through the entire region particularly at $E_\gamma = 7.5$ and 10.5 MeV (1973BR1M; monoenergetic photons). See also (1954GO1A, 1958RY77, 1966BR1M). There are many reports of fine structure [see Table 7.5 in (1966LA04) and (1965HA19, 1965WA19, 1966BA1W)]. See also (1970HA1F) and (1968KA38, 1968RA1E, 1969MU1C). The integrated cross sections from threshold to 32 MeV are 20 MeV · mb and 10 MeV · mb for the total neutron yield and for reaction (b), respectively (1966BR1M). See also (1971KA70, 1973AH1A).

The cross section for the (γ, p) reaction (reaction (c)) shows a maximum at ≈ 15.6 MeV with a width of ≈ 4 MeV (1954TI16, 1962GR08) [however a number of authors claim the existence of many additional peaks: see (1966LA04)]. The energy distribution of the photoprotons has been measured with bremsstrahlung radiation $E_{\text{bs}} = 50$ to 34 MeV (1970SA14) and 100 MeV (1968MA19) and at $E_\gamma = 60$ MeV (1973GA16). The polarization of the protons produced in the interaction between high-energy γ -rays and ${}^7\text{Li}$ has been studied by (1969AN20, 1970TO09). See also ${}^6\text{He}$ and (1966MA17, 1967DE11, 1969AN1H, 1969MU1C, 1970AN05, 1970WO10, 1971AN04, 1973DO13) and (1970HA1F).

For reaction (d) see (1967SM1A). For reaction (e) see (1967DE11, 1969AN1H, 1971AN04, 1971AN15). A number of peaks have been reported in the (γ, t) cross section: see Table 7.5 in (1966LA04) and (1970SE1D). See also (1965DA06, 1966DZ07, 1966MA17, 1967DE11, 1969HU1E, 1969DE28, 1970SE1A). See also (1967SH1E, 1973AR1L, 1973CO1N) and (1968EL06, 1970RA1H, 1973AS02; theor.).

18. (a) ${}^7\text{Li}(e, e'){}^7\text{Li}^*$	
(b) ${}^7\text{Li}(e, ep){}^6\text{He}$	$Q_m = -9.978$

The electric form factor measurements for $E_e = 100$ to 600 MeV are well accounted for by a simple harmonic oscillator shell model with a quadrupole contribution described by an undeformed p-shell: $R_{\text{rms}} = 2.39 \pm 0.03$ fm, $|Q| = 42 \pm 2.5$ mb (1967SU1A). From results obtained for $E_e = 24.14$ to 97.19 MeV, $R_{\text{rms}} = 2.35 \pm 0.10$ fm (model independent), 2.29 ± 0.04 fm (shell model) (1969MO1J, 1972BU01). A study of the ratio of the electric charge scattering from ${}^6\text{Li}$ and from ${}^7\text{Li}$ as a function of (momentum transfer)² yields $\langle r^2 \rangle_6^{1/2} / \langle r^2 \rangle_7^{1/2} = 1.001 \pm 0.008$. The r.m.s. radius of the ground-state magnetization density distribution, $\langle r^2 \rangle_M^{1/2} = 2.98 \pm 0.05$ fm. From the ratio of

Table 7.4: Levels of ${}^7\text{Li}(e, e'){}^7\text{Li}^*$ ^a

E_x (MeV)	$J^\pi; T$	Γ_{γ_0} (eV)	Type	$\Gamma_{\gamma_0}/\Gamma_W$	Refs.
0.48	$\frac{1}{2}^-; \frac{1}{2}$	$(2.8 \pm 1.6) \times 10^{-7}$	E2	18	(1971VA20)
		$(6.30 \pm 0.31) \times 10^{-3}$	M1	2.8	(1971VA20)
4.63 ± 0.05	$\frac{7}{2}^-; \frac{1}{2}$		E2 ^e		(1963BE26, 1963BE53, 1968HU1C, 1969HU05)
6.6 ± 0.1 ^b	$\frac{5}{2}^-; \frac{1}{2}$		E2		(1968HU1C, 1969HU05)
7.5 ± 0.08	$\frac{5}{2}^-; \frac{1}{2}$	0.6 ± 0.3	E2		(1963BA19, 1963BE26)
		0.9 ± 0.4 ^f			(1964GR40) ^g
c					
11.25 ^d	$\frac{3}{2}^-; \frac{3}{2}$	1.3 ± 0.4	M1	0.043	(1967AR1A)

^a For a summary of $B(E2\uparrow)$ measurements, see Table 7.6 in (1966LA04) and ${}^7\text{Li}$, the ‘‘GENERAL’’ section.

^b $\Gamma_{\text{c.m.}} = 875_{-100}^{+200}$ keV (1968HU1C, 1969HU05).

^c The excitation of ${}^7\text{Li}^*(10.5, 12.5, 14.0)$ is reported by (1963BA19).

^d $\Gamma_{\text{cm}} = 200 \pm 100$ keV (1967AR1A).

^e Purely longitudinal (1968HU1C, 1969HU05).

^f $0.1 \rightarrow 0.5$ eV, from ${}^7\text{Li}(\gamma, n)$ (B.L. Berman, private communication).

^g From ${}^7\text{Li}(\gamma, n)$.

the transverse inelastic and elastic cross sections at 180° , $B(M1, \uparrow; 0.48) = 2.50 \pm 0.12 \mu_N^2$. The cross section of the longitudinal excitation of ${}^7\text{Li}^*(0.48)$ has been found from the scattering through angles of 90° to 150° , $B(C2, \uparrow; 0.48) = 7 \pm 4 \text{ fm}^4$. The harmonic oscillator length parameter of the $1p$ shell is found to be $a_{1p} = 1.90 \pm 0.03 \text{ fm}$ (1971VA20).

The magnetic form factor has been measured for $E_e = 70$ to 200 MeV. The ratio of the magnetic octupole moment to the dipole moment $\Omega/\mu = 2.30 \pm 0.50 \text{ fm}^2$ (1966RA29).

Inelastic scattering studies show peaks corresponding to ${}^7\text{Li}^*(0+0.48, 4.63, 6.68, 7.47, 11.25)$: see (1967AR1A, 1968HU1C, 1969HU05), (1966LA04) and Table 7.4. See also the review by (1972THZF). For reaction (b) see (1970WO10).

See also (1970WA1N), (1966GO1C, 1966GU1C, 1968GO1J) and (1966MU1A, 1967BO22, 1967EL1B, 1967KA1A, 1968BO1R, 1968KU1D, 1968KU1B, 1969HA1N, 1969KR16, 1969KU1C, 1969VI02, 1969WI21, 1971DRZW, 1973HI03; theor.).

19. (a) ${}^7\text{Li}(n, n'){}^7\text{Li}^*$

(b) ${}^7\text{Li}(n, nt){}^4\text{He}$ $Q_m = -2.4668$

Angular distributions have recently been measured at $E_n = 1.12$ to 2.30 MeV (1968KN1B; n_0, n_1), 3.35 and 4.83 MeV (1968HO03; n_0), 5.74 and 7.50 MeV (1968HO03; $n_0 + n_1$) and 14 MeV (1966RE1B; $n_0 + n_1, n_2$). See (1966LA04) for a listing of earlier references and (1970GA1A). At

$E_n = 14$ MeV no states other than ${}^7\text{Li}^*(0, 0.48, 4.63)$ are populated (1968HA1J): see, however, (1954AL24, 1966RE1B). See also (1966HU1B, 1967CO01) and (1969WA11; theor.). See also ${}^8\text{Li}$. For reaction (b) see (1972AN1Q) and ${}^5\text{He}$.

20. ${}^7\text{Li}(p, p'){}^7\text{Li}^*$

Angular distributions are reported at $E_p = 1.0$ to 2.0 MeV (1966BA1Q; p_0, p_1), 24.4 MeV (1967CR1E; p_0, p_1, p_2), 33.6 MeV (1970KU1D; p_0), 49.8 MeV (1971MA13, 1971MA44; p_0, p_1, p_2), 100 MeV (1966MA38, 1968LI1C; $p_0 + p_1, p_2$), 144 MeV (1972JA07; p_0), 152 MeV (1966RO1C; p_0), 155 MeV (1968GE04; p_0, p_1, p_2) and 185 MeV (1967JO1F; p_1). (1967CR1E) report that the p_2 group is strongly excited and that the angular distribution is consistent with the predictions of the collective model for an $l = 2$ transition. Earlier measurements are reported in (1966LA04).

Inelastic proton groups have been observed corresponding to ${}^7\text{Li}^*(0.48, 4.63, 6.68, 7.47)$: see (1952AJ38). At $E_p = 185$ MeV proton groups are observed to these states [$E_x = 4.62 \pm 0.04, 6.55 \pm 0.20, 7.5 \pm 0.2$ MeV: (1967JO1F)] as well as to states at $E_x = 5.5 \pm 0.3$ MeV ($\Gamma \approx 0.4$ MeV) (1965HA17: not seen by (1967JO1F)) and 9.6 ± 0.2 MeV (1965HA17), 9.4 ± 0.2 MeV (1967JO1F). The width of ${}^7\text{Li}^*(6.7)$ is ≈ 1 MeV (1965HA17). At $E_p = 50$ MeV (1968MA02) report ten states of ${}^7\text{Li}$ with $E_x < 13.6$ MeV.

$\tau_m(0.48) = 0.106 \pm 0.014$ psec (1966PA11: Doppler shift measurement), a value consistent with intermediate coupling with LS coupling predominating ($a/K < 3.6$): see also Table 7.4 in (1966LA04). Analysis of the 155 MeV data yields $B(E2\uparrow) = 10.5 \pm 2, 28 \pm 6$ and 4.5 ± 2.3 fm⁴ for ${}^7\text{Li}^*(0.48, 4.63, 6.68)$; $\Gamma(E2\downarrow) = 0.43, 0.025$ and 0.029 μeV (1965JA1A).

A comparison of σ_t for ${}^7\text{Li}(p, p'){}^7\text{Li}^*(0.48)$ and ${}^7\text{Li}(p, n){}^7\text{Be}^*(0.43)$ has been carried out for $E_p = 23$ to 52 MeV: the spin-flip, isospin-flip part of the effective interaction is approximately independent of energy while the pure central part appears to decrease with increasing energy (1967LO07). See also (1966MA1N, 1968GL1A, 1968NE1B, 1969MA1P, 1969NE1A, 1969TI02, 1969WA11, 1970KI1E, 1973KA04; theor.).

21. (a) ${}^7\text{Li}(p, 2p){}^6\text{He}$	$Q_m = -9.978$
(b) ${}^7\text{Li}(p, pn){}^6\text{Li}$	$Q_m = -7.2506$
(c) ${}^7\text{Li}(p, pd){}^5\text{He}$	$Q_m = -9.61$
(d) ${}^7\text{Li}(p, p\alpha){}^3\text{H}$	$Q_m = -2.4668$
(e) ${}^7\text{Li}(p, \alpha){}^4\text{He}$	$Q_m = 17.348$
(f) ${}^7\text{Li}(p, 2d){}^4\text{He}$	$Q_m = -6.500$

For reaction (a) see ${}^6\text{He}$. See also (1966LA04), (1973CO2B) and (1965BE1E, 1966JA1A, 1967EL1C, 1967JA1E, 1968JA1G, 1969KO1J; theor.). For reaction (b) see (1970TH1F). For

reaction (c) see reaction 20 in ${}^6\text{Li}$ and (1973CO2B, 1973KO1M). The momentum distribution of the α -particles in reaction (d) has the shape expected for the knockout of an $L = 1$ α -cluster from ${}^7\text{Li}$ (1970JA17). The reaction proceeds sequentially via ${}^7\text{Li}^*(4.63)$ (1967JO1C, 1970JA17) and via ${}^7\text{Li}^*(6.68)$ (1970JA17). See also (1969HO1K, 1971GA1J, 1973CO2B), (1972RA1E) and (1972JA23; theor.). For reaction (e) see ${}^8\text{Be}$. For reaction (f) see (1972FU07, 1973CO2B) and reaction 20 in ${}^6\text{Li}$.

22. ${}^7\text{Li}(d, d'){}^7\text{Li}^*$

Angular distributions have been measured at $E_d = 11.8$ MeV (1968LU02; d_0, d_1), 12 MeV (1971BI11; d_0), 14.7 MeV (1969MA13; d_0, d_1, d_2) and 28 MeV (1962SL02; d_0, d_2). See also (1970EL16; theor.) and (1966LA04).

23. ${}^7\text{Li}({}^3\text{He}, {}^3\text{He}'){}^7\text{Li}^*$

Angular distributions are reported at $E({}^3\text{He}) = 8.7$ and 9.7 MeV (1969MA1J; elastic), 11 MeV (1970SC23; elastic), 21, 24 and 27 MeV (1966VA1B, 1967BL1E: to ${}^7\text{Li}^*(0, 0.48, 4.63)$) and 25.2 MeV (1968BR1G: to ${}^7\text{Li}^*(0, 0.48, 4.63)$). See also (1967CO1J).

24. (a) ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}^*$

(b) ${}^7\text{Li}(\alpha, 2\alpha){}^3\text{H} \quad Q_m = -2.4668$

Angular distributions (reaction (a)) are reported at $E_\alpha = 3.6$ MeV (1972BO07; α_0, α_1), 12.0 to 18.0 MeV (1970BI1B, 1971BI12; α_0), 25 MeV (1968DO13; $\alpha_0 + \alpha_1, \alpha_2$) and 29.4 MeV (1968MA25, 1969MA13; $\alpha_0, \alpha_1, \alpha_2$). See also (1972BE1Z).

Reaction (b) has been studied at $E_\alpha = 23.6$ MeV (1968BE1Q), 25 MeV (1968DO13, 1969DO02, 1969DO03), 29.4 MeV (1968MA25), 50 MeV (1970LA14, 1973LA1Q), 55 MeV (1968PI04, 1969PI11, 1970PI1D), 64.3 MeV (1970JA17) and 104 MeV (1969VE1B). ${}^7\text{Li}^*(4.63)$ is strongly involved in the sequential decay: see (1968DO13, 1968MA25, 1970JA17, 1970LA14). The population of ${}^7\text{Li}^*(7.48)$ is reported by (1970LA14), and (1968MA25) suggests that ${}^7\text{Li}^*(6.68)$ is also involved in the sequential decay. At $E_\alpha = 55$ MeV, the effective number of α clusters, $N_{\text{eff}} = 4_{-2}^{+4}\%$ (1969PI11). See also (1969BA2C), (1969HO1K, 1971GA1J) and (1968BA1H, 1972AV04; theor.). See also (1966LA04).

25. ${}^7\text{Li}({}^7\text{Li}, {}^7\text{Li}){}^7\text{Li}$

The elastic angular distribution has been studied for $E(^7\text{Li}) = 4.0$ to 6.5 MeV (1966PI02).

26. $^7\text{Li}(^{10}\text{B}, ^{10}\text{B})^7\text{Li}$

See (1969RO1G) and ^{10}B .

27. (a) $^7\text{Li}(^{12}\text{C}, ^{12}\text{C})^7\text{Li}$

(b) $^7\text{Li}(^{13}\text{C}, ^{13}\text{C})^7\text{Li}$

The elastic scattering has been studied at $E(^7\text{Li}) = 20$ MeV (1969BE90) [on ^{13}C] and at 34 [^{13}C] and 36 [^{12}C] MeV (1973SC26). See also ^{12}C in (1975AJ02) and ^{13}C in (1976AJ04). The inelastic scattering angular distributions involving $^7\text{Li}_{0.48}^* + ^{12}\text{C}_{\text{g.s.}}$ and $^7\text{Li}_{0.48}^* + ^{12}\text{C}_{4.43}^*$ have been measured at $E(^7\text{Li}) = 36$ MeV (1973SC26).

28. $^7\text{Li}(^{16}\text{O}, ^{16}\text{O})^7\text{Li}$

See (1969BE90, 1971OR02).

29. $^7\text{Li}(^{20}\text{Ne}, ^{20}\text{Ne}')^7\text{Li}^*$

See (1966LA04).

30. $^7\text{Be}(\epsilon)^7\text{Li}$

$$Q_m = 0.862$$

$$Q_0 = 0.851 \pm 0.012 \text{ (1973MU19)}.$$

The decay proceeds to the ground and 0.48 MeV states. The branching ratio to $^7\text{Li}^*(0.48)$ is $10.32 \pm 0.16\%$ (1962TA11), $10.4 \pm 0.3\%$ (1972SZ02), $10.42 \pm 0.18\%$ (1973PO10). [See also Table 7.8 in (1966LA04)]. The weighted mean value of the half-life is 53.44 ± 0.09 days (1949SE20, 1953KR16, 1956BO36, 1970JO21). Both transitions are superallowed. $\log ft = 3.30$ and 3.54 for the decays to $^7\text{Li}^*(0, 0.48)$ respectively[‡].

The energy of the γ -ray is 477.57 ± 0.05 keV (1967BL03), 477.4 ± 0.2 keV (1965RO09), 477.593 ± 0.012 keV (1971HE20) [$E_x = 477.611 \pm 0.012$ keV]. A measurement of the bremsstrahlung

[‡] G. Fox and B. Zimmerman, private communication.

spectrum to ${}^7\text{Li}^*(0.48)$ measured in coincidence with the 478 keV γ -ray, leads to a transition energy of 395 ± 25 keV (1971LA03), 388 ± 8 keV (1972PE05). See also (1973MU19).

For discussions of astrophysical considerations, see (1969BA1U, 1969FO1D, 1969YI1B, 1971CA1B, 1972BA2M, 1972KO1A, 1973BA2C). See also (1967GE1A, 1969HE1M, 1973HE1M), (1972EM03) and (1965PR04, 1966EL08, 1966BA26, 1968FI02, 1969LE1D, 1969SU15, 1970DA21, 1970FA14, 1970KO41, 1973HA49, 1973MU12, 1973WI11; theor.).

$$31. {}^9\text{Be}(\gamma, d){}^7\text{Li} \quad Q_m = -16.6965$$

See ${}^9\text{Be}$ and (1955AJ61).

$$32. {}^9\text{Be}(n, t){}^7\text{Li} \quad Q_m = -10.4389$$

See ${}^{10}\text{Be}$ and (1966LA04).

$$33. {}^9\text{Be}(p, {}^3\text{He}){}^7\text{Li} \quad Q_m = -11.2027$$

At $E_p = 43.7$ angular distributions have been obtained for the ${}^3\text{He}$ particles corresponding to ${}^7\text{Li}^*(0, 0.48, 4.63, 7.47)$. The 7.47 MeV state is strongly excited while the mirror state in ${}^7\text{Be}$ is not appreciably populated in the mirror reaction (see reaction 17 in ${}^7\text{Be}$). The angular distribution indicates that the transition to ${}^7\text{Li}^*(7.47)$ involves both $L = 0$ and 2, with a somewhat dominant $L = 0$ character (1966CE05). Reanalysis of the data of (1965DE08) places the $J^\pi = \frac{3}{2}^-$; $T = \frac{3}{2}$ level at $E_x = 11.28 \pm 0.04$ MeV, $\Gamma = 260 \pm 50$ keV (1967MC14). See also (1969BA1Z, 1969IN1A).

$$34. {}^9\text{Be}(p, pd){}^7\text{Li} \quad Q_m = 16.6965$$

See (1966LA04).

$$35. \text{(a) } {}^9\text{Be}(d, \alpha){}^7\text{Li} \quad Q_m = 7.1511$$

$$\text{(b) } {}^9\text{Be}(d, t){}^4\text{He}{}^4\text{He} \quad Q_m = 4.684$$

$$Q_0 = 7.157 \pm 0.008 \text{ (1967SP09).}$$

Angular distributions have been measured recently at $E_d = 0.3$ to 1.0 MeV (1968BE1E; α_0 , α_1), 0.9 to 2.2 MeV (1971SA27; α_0 , α_1), 11.4 and 12.4 MeV (1966DO1A; α_0 , α_1). For older measurements see (1966LA04). A study at $E_d = 11$ MeV finds $\Gamma_{cm} = 93 \pm 25$ and 80 ± 20 keV, respectively, for ${}^7\text{Li}^*(4.63, 7.47)$. No evidence was observed for ${}^7\text{Li}^*(5.5, 8.6, 9.7, 12.5)$ or for the $T = \frac{3}{2}$ state ${}^7\text{Li}^*(11.25)$ (1966HA09). See also (1964MA57, 1966JA05) and (1966ME1E; theor.). In a kinematically complete study of reaction (b) at $E_d = 26.3$ MeV, ${}^7\text{Li}^*(4.6, 6.5 + 7.5, 9.4)$ are strongly excited. No sharp α -decaying states of ${}^7\text{Li}$ are observed with $10 < E_x < 25$ MeV. Parameters for ${}^7\text{Li}^*(9.6)$ are $E_x = 9.36 \pm 0.05$ MeV, $\Gamma = 0.8 \pm 0.2$ MeV (1973SO08). See also ${}^8\text{Be}$.

$$36. {}^9\text{Be}({}^3\text{He}, \alpha p){}^7\text{Li} \quad Q_m = 1.657$$

See (1967ST1D).

$$37. {}^9\text{Be}({}^6\text{Li}, 2\alpha){}^7\text{Li} \quad Q_m = 5.677$$

See (1966SA04, 1968JA08) and ${}^{11}\text{B}$ in (1975AJ02).

$$38. \text{(a) } {}^{10}\text{B}(n, \alpha){}^7\text{Li} \quad Q_m = 2.791$$

$$\text{(b) } {}^{10}\text{B}(n, t){}^4\text{He}{}^4\text{He} \quad Q_m = 0.3237$$

$$Q_0 = 2.8008 \pm 0.0076 \text{ (1967DE15).}$$

At $E_n = 14.4$ MeV angular distributions have been measured for the $n_0 + n_1$ and for the n_2 groups (1969AN25). The half-life of ${}^7\text{Li}^*(0.48)$ is 92 ± 11 fsec (1967CA02). See also (1966LA04), ${}^{11}\text{B}$ in (1975AJ02), (1969VA1F, 1970NE03, 1972SE1K) and (1972HA04; theor.). For reaction (b) see (1967VA12).

$$39. {}^{10}\text{B}(\alpha, {}^7\text{Be}){}^7\text{Li} \quad Q_m = -16.202$$

See reaction 19 in ${}^7\text{Be}$ (1969FO06).

$$40. {}^{11}\text{B}(\gamma, \alpha){}^7\text{Li} \quad Q_m = -8.666$$

See ${}^{11}\text{B}$ in (1975AJ02) and (1969MU10).

$$41. \text{}^{11}\text{B}(\text{p}, \text{p}\alpha)^7\text{Li} \quad Q_{\text{m}} = -8.666$$

See (1964BA1C).

$$42. \text{(a)} \text{}^{11}\text{B}(\alpha, 2\alpha)^7\text{Li} \quad Q_{\text{m}} = -8.666$$

$$\text{(b)} \text{}^{11}\text{B}(\alpha, \text{}^8\text{Be})^7\text{Li} \quad Q_{\text{m}} = -8.758$$

For reaction (a) see (1966GE12, 1969FU09). Angular distributions have been measured in reaction (b) at $E_{\alpha} = 28.4$ and 29.0 MeV for the transitions to ${}^8\text{Be}^*(0, 2.9)$ and ${}^7\text{Li}^*(0, 0.48)$ (1968KA24). At $E_{\alpha} = 65$ MeV ${}^7\text{Li}^*(0, 4.63)$ are strongly populated and ${}^7\text{Li}^*(7.47)$ is weakly excited. The intensity of the group to ${}^7\text{Li}^*(0.48)$ is $< 15\%$ of the group to ${}^7\text{Li}(0)$ (1973WO06). See also (1966GE12).

$$43. \text{}^{11}\text{B}(\text{d}, \text{}^6\text{Li})^7\text{Li} \quad Q_{\text{m}} = -7.192$$

At $E_{\text{d}} = 19.5$ MeV, angular distributions have been measured for the transitions to ${}^6\text{Li}(0)$ and ${}^7\text{Li}^*(0, 0.48)$ (1971GU07). At $E_{\text{d}} = 40$ MeV the cross section for the transition to ${}^6\text{Li}^*(3.56) + {}^7\text{Li}(0)$ is half that for ${}^6\text{He}(0) + {}^7\text{Be}(0)$ [to $\pm 10\%$] in agreement with isospin conservation (1972GO1P). See also (1972GA1E).

$$44. \text{}^{11}\text{B}(\text{}^{16}\text{O}, \text{}^{20}\text{Ne})^7\text{Li} \quad Q_{\text{m}} = -3.936$$

See (1968OK06).

$$45. \text{}^{12}\text{C}(\text{d}, \text{}^7\text{Be})^7\text{Li} \quad Q_{\text{m}} = -17.543$$

At $E_{\text{d}} = 39.8$ MeV, angular distributions have been measured for the transitions to ${}^7\text{Li}(0) + {}^7\text{Be}(0)$, ${}^7\text{Li}^*(0.48) + {}^7\text{Be}(0)$, ${}^7\text{Li}(0) + {}^7\text{Be}^*(0.43)$, and ${}^7\text{Li}^*(0.48) + {}^7\text{Be}^*(0.43)$. Asymmetries exceeding 20% are observed in the ratio of the cross sections to ${}^7\text{Li}(0)$ and ${}^7\text{Be}(0)$ (1971HO1K, 1971YO06). See also (1971SI28; theor.).

$$46. \text{}^{12}\text{C}(\text{}^3\text{He}, \text{}^8\text{B})^7\text{Li} \quad Q_{\text{m}} = -22.899$$

This reaction has been studied at $E({}^3\text{He}) = 40.7$ MeV (1971DE37).

47. $^{12}\text{C}(^6\text{Li}, ^{11}\text{C})^7\text{Li}$ $Q_m = -11.471$

At $E(^6\text{Li}) = 36$ MeV, angular distributions have been obtained for the transitions involving $^7\text{Li}_{\text{g.s.}} + ^{11}\text{C}_{\text{g.s.}}$ and $^7\text{Li}_{0.48}^* + ^{11}\text{C}_{\text{g.s.}}$ (1973SC26).

48. $^{13}\text{C}(\text{p}, ^7\text{Be})^7\text{Li}$ $Q_m = -20.264$

At $E_p = 45.0$ MeV angular distribution has been measured for the transition to $^7\text{Be}(0) + ^7\text{Li}(0)$ (1971BR07).

49. $^{13}\text{C}(\text{d}, ^8\text{Be})^7\text{Li}$ $Q_m = -3.589$

At $E_d = 14.6$ MeV, angular distributions are reported for the transitions to $^8\text{Be}(0)$ and $^7\text{Li}^*(0, 0.48)$ (1967DE03).

50. $^{13}\text{C}(^6\text{Li}, ^{12}\text{C})^7\text{Li}$ $Q_m = 2.304$

At $E(^6\text{Li}) = 34$ MeV angular distributions have been measured for the transitions involving $^7\text{Li}_{\text{g.s.}} + ^{12}\text{C}_{\text{g.s.}}$, $^7\text{Li}_{0.48}^* + ^{12}\text{C}_{\text{g.s.}}$, $^7\text{Li}_{\text{g.s.}} + ^{12}\text{C}_{4.4}^*$, and $^7\text{Li}_{0.48}^* + ^{12}\text{C}_{4.4}^*$ (1973SC26).

51. $^{14}\text{N}(\text{n}, 2\alpha)^7\text{Li}$ $Q_m = -8.823$

At $E_n = 14.1$ MeV, $^7\text{Li}^*(0, 0.48)$ are produced with about equal probability (1971SC16).

52. $^{16}\text{O}(\alpha, ^{13}\text{N})^7\text{Li}$ $Q_m = -22.566$

See (1972RU03).

53. $^{16}\text{O}(^6\text{Li}, ^{15}\text{O})^7\text{Li}$ $Q_m = -8.419$

At $E(^6\text{Li}) = 36$ MeV, angular distributions have been determined for the transitions involving $^7\text{Li}_{\text{g.s.}} + ^{15}\text{O}_{\text{g.s.}}$ and $^7\text{Li}_{0.48}^* + ^{15}\text{O}_{\text{g.s.}}$ (1973SC26).

54. (a) $^{17}\text{O}(\text{d}, ^{12}\text{C})^7\text{Li}$ $Q_{\text{m}} = -2.580$
(b) $^{18}\text{O}(\text{d}, ^{13}\text{C})^7\text{Li}$ $Q_{\text{m}} = -5.680$
(c) $^{19}\text{F}(\text{d}, ^{14}\text{N})^7\text{Li}$ $Q_{\text{m}} = -6.122$

At $E_{\text{d}} = 14.6$ to 15.0 MeV, angular distributions have been measured for the transitions to $^{12}\text{C}(0) + ^7\text{Li}^*(0, 0.48)$ [reaction (a)], $^{13}\text{C}(0) + ^7\text{Li}^*(0, 0.48)$ [reaction (b)] and $^{14}\text{N}(0) + ^7\text{Li}^*(0, 0.48)$ [reaction (c)] ([1967DE03](#)).

${}^7\text{Be}$
(Figs. 9 and 10)

GENERAL: (See also (1966LA04).)

Shell model: (1961KO1A, 1965VO1A, 1966BA26, 1966HA18, 1967FA1A, 1968GO01, 1969TA1H, 1971CO28, 1971NO02, 1972LE1L, 1973HA49).

Cluster model: (1965NE1B, 1968HA1G, 1971NO02, 1972HI16, 1972KU12, 1972LE1L).

Rotational and deformed models: (1965VO1A, 1966EL08).

Special levels: (1966BA26, 1966EL08, 1967FA1A, 1969HA1G, 1969HA1F, 1971CO28, 1971NO02, 1972BB26, 1973AS02, 1973FE1J).

Electromagnetic transitions: (1966BA26, 1966EL08, 1969HA1G, 1969HA1F, 1973AS02, 1973HA49).

Astrophysical questions: (1968BA2F, 1968HA1C, 1970BA1M, 1972KO1E, 1972PA1C, 1972UL1A, 1973LA19, 1973RA37, 1973SC1T).

Special reactions: (1965FU19, 1966GA15, 1966MI1C, 1967AU1B, 1967FU1E, 1967WI06, 1967WI20, 1968BE1F, 1968DI1B, 1968HU18, 1968MI1D, 1968RA34, 1968SH26, 1968YI01, 1969DI18, 1969HI1A, 1969YI1A, 1970BR13, 1970MA1E, 1970MO38, 1971AR02, 1971BA58, 1971BI22, 1971BR36, 1971DM01, 1971EP02, 1971HE24, 1971NO09, 1971ST30, 1972AM04, 1973ER1G, 1973HO11, 1973JO07, 1973LA19, 1973MI02, 1973VO1G).

Reactions involving pions: (1968BE1F).

Other topics: (1965BO1C, 1965VO1A, 1966DE1E, 1966HA18, 1966YO1B, 1967CA17, 1967FA1A, 1968BE1F, 1968GO01, 1969HE1N, 1969VI1C, 1970DE1P, 1971ZA1D, 1972AB14, 1972AN05, 1972BB26, 1972CA37, 1972LE1L, 1972PN1A, 1973JU2A, 1973RO1R).

Ground-state properties: (1965VO1A, 1966BA26, 1966EL08, 1968PE16, 1972LE1L, 1973MA1K).

1. ${}^7\text{Be}(\epsilon){}^7\text{Li}$ $Q_m = 0.8618$

The decay is complex: see ${}^7\text{Li}$.

2. ${}^4\text{He}({}^3\text{He}, \gamma){}^7\text{Be}$ $Q_m = 1.5864$

In the range $E_\alpha = 0.38$ to 5.80 MeV the cross section rises from 8×10^{-3} to $4 \mu\text{b}$ (1963PA12, 1969NA24). The capture proceeds mainly by E1, with both s- and d-waves contributing above

Table 7.5: Energy levels of ${}^7\text{Be}$

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	$\tau_{1/2} = 53.44 \pm 0.09$ d	ϵ	1, 2, 4, 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31
0.42920 ± 0.10	$\frac{1}{2}^-; \frac{1}{2}$	$\tau_m = 192 \pm 20$ fsec	γ	5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 22, 23, 24, 25, 28, 29, 30, 31
4.57 ± 50	$\frac{7}{2}^-; \frac{1}{2}$	$\Gamma = 175 \pm 7$ keV	${}^3\text{He}, \alpha$	3, 12, 13, 14, 15, 16, 17, 19
6.73 ± 100	$\frac{5}{2}^-; \frac{1}{2}$	1.2 MeV	p, ${}^3\text{He}, \alpha$	3, 9, 10, 16, 17
7.21 ± 60	$\frac{5}{2}^-; \frac{1}{2}$	0.5 MeV	p, ${}^3\text{He}, \alpha$	3, 6, 9, 10, 13, 17
9.27 ± 100	$\frac{7}{2}^-; \frac{1}{2}$		p, ${}^3\text{He}, \alpha$	3, 13
9.9	$\frac{3}{2}^-; \frac{1}{2}$	≈ 1.8 MeV	p, ${}^3\text{He}, \alpha$	3, 6
$\gtrsim 10.0$	$\frac{1}{2}^-; \frac{1}{2}$	broad	p, ${}^3\text{He}, \alpha$	3
11.01 ± 30	$\frac{3}{2}^-; \frac{3}{2}$	320 ± 30	p, ${}^3\text{He}, \alpha$	3, 6, 13, 16

$E_\alpha = 1$ MeV (1963TO06). The branching ratio γ_1/γ_0 [${}^7\text{Be}^*(0.43)/{}^7\text{Be}(0)$] is approximately constant at 37% for $E_\alpha = 0.57$ to 3.2 MeV (1963PA12, 1969NA24). The zero-energy intercept of the cross-section factor $S = 0.61 \pm 0.07$ keV \cdot b and $(dS/dE)_0 = -(5.8 \pm 0.3) \times 10^{-4}$ b using all of the data. If the low-energy data ($E_{\text{cm}} \leq 0.7$ MeV) is fitted using a direct capture calculation (1963TO06) the zero-energy intercept of the cross-section factor is $S = 0.51 \pm 0.05$ keV \cdot b and $(dS/dE)_0 = -(2.8 \pm 0.4) \times 10^{-4}$ b (1969NA24). A second-order (in energy) polynomial fit to the low-energy data ($E_{\text{cm}} \leq 0.8$ MeV) determines $S = 0.61 \pm 0.07$ keV \cdot b and $(dS/dE)_0 = -(5.8 \pm 0.3)$ b (1969NA24). Recent papers discussing the astrophysical implications of this reaction are (1967TO1B, 1968BA2E, 1971CA1B, 1972KA1B, 1973BA2C, 1973TR1E). See also (1966LA04).

3. (a) ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$

$$E_b = 1.5864$$

(b) ${}^4\text{He}({}^3\text{He}, \text{p}){}^6\text{Li}$

$$Q_m = -4.0200$$

Elastic scattering studies have previously been reported for $E({}^3\text{He}) = 2.5$ to 30 MeV and for $E_\alpha = 11$ to 41 MeV: See (1966LA04). More recent measurements have been made at $E({}^3\text{He}) = 1.72, 2.46, 2.98$ MeV (1971CH42), 5 to 18 MeV (1967SP10), 12.0 to 19.0 MeV (1967DU1B), 17.8

to 30.0 MeV (1970JA04), 18 to 70 MeV (1969BA2B), 19.5 to 51.1 MeV (1970BR42), 27.2 to 42.8 MeV (1969SC16) and at $E_\alpha = 5.94$ to 7.90 MeV (1968IV01), 42 MeV (1970VI01), 43.4 to 58.2 MeV (1969CA1E), 66 to 104 MeV (1971FE02, 1971FE03, 1973FE11) and 140 MeV (1972FR1J, 1972PU1C, 1972FR1K). Differential cross sections have been calculated by (1973KO1Q) for $E_{\text{cm}} < 44.5$ MeV using the one-channel resonating-group method, and including exchange terms and odd-even absorption. Polarization measurements have been carried out at $E(^3\text{He}) = 7.8$ to 13.0 MeV (1969AR07, 1971AR1K) and at 11.5 to 13.0 MeV (1970MC07), and at $E_\alpha = 4.33$ to 9.83 MeV (1972BO42), 7.5 to 18.5 MeV (1970HA1M, 1970HA1P, 1972HA64), 13.0 MeV (1971AR1K) and 98 MeV (1973FE11). The ratios of γ^2 to the Wigner limit are, respectively, $0.31_{-0.04}^{+0.09}$ and $0.29_{-0.07}^{+0.16}$ for ${}^7\text{Be}^*(0, 0.43)$ (1972BO42).

For $l \leq 4$, only f-wave phase shifts show resonance structure for $E(^3\text{He}) < 18$ MeV, corresponding to ${}^7\text{Be}^*(4.57, 6.73, 9.27)$: see Table 7.6 (1967SP10, 1968IV01). No structure corresponding to ${}^7\text{Be}^*(7.21)$ ($J^\pi = \frac{5}{2}^-$) is seen in the elastic data. The s-wave phase shift is somewhat greater than hard-sphere; the p-wave splitting agrees with (1964BA09, 1967SP10). The decay of ${}^7\text{Be}^*(9.27)$ ($J^\pi = \frac{7}{2}^-$) to ${}^6\text{Li}(0)$ requires f-shell configuration admixture. An estimate of the yield of ground state protons relative to those corresponding to ${}^6\text{Li}^*(2.19)$ yields $\gamma^2(p_0)/\gamma^2(p_1) = 16_{-10}^{+5}\%$ (1967SP10). At higher energies [$E(^3\text{He}) = 27.2$ to 42.8 MeV] (1969SC16) report that the s- and f-wave phase shifts fall appreciably below the predictions of resonating group calculations, while (1970BR42) see some indication of broad resonant structure at $E(^3\text{He}) \approx 34$ MeV, in rough qualitative agreement with such calculations. For inelastic scattering in reaction (a) see (1971HA21) and (1973FI04). The bremsstrahlung cross section at $E(^3\text{He}) = 7.4$ MeV is 12.6 ± 3.4 $\mu\text{b}/\text{sr}^2$ (1973FR17).

The differential reaction cross section for reaction (b) has been determined for $E(^3\text{He}) = 8$ to 18 MeV: resonances are observed corresponding to ${}^7\text{Be}^*(7.21, 9.27)$ in the p_0 yield and to ${}^7\text{Be}^*(9.27)$ in the p_1 yield: see Table 7.6 (1967SP10). A study of the gamma rays from ${}^6\text{Li}^*(3.56)$ (p_2) carried out at $E(^3\text{He}) = 13.8$ to 18.5 MeV shows the excitation of two $J^\pi = \frac{3}{2}^-$ states at $E_x \approx 10.0$ MeV ($T = \frac{1}{2}$) and 11.00 ± 0.05 MeV ($\Gamma = 400 \pm 50$ keV, $\theta_{p_2}^2 = 0.13 \pm 0.02$, $T = \frac{3}{2}$). The $T = \frac{3}{2}$ resonance is evidenced mainly through interference. There is also evidence for an extremely broad $J^\pi = \frac{1}{2}^-$ structure at $E_x \gtrsim 10$ MeV (1967HA07, 1967HA08: see also ${}^6\text{Li}(p, p){}^6\text{Li}$).

See also (1970LI06, 1972BI1G, 1973KO1R), (1966PH1A) and (1966BA26, 1966RA1B, 1966TH1C, 1967OK1A, 1968BR1H, 1968LE1K, 1969TA1G, 1970NE1F, 1971FU09, 1971KU22, 1971PL06, 1971TA23, 1972BR1Q, 1972CL1C, 1972NE17; theor.).

$$4. {}^4\text{He}(\alpha, n){}^7\text{Be} \quad Q_m = -18.9921$$

See ${}^8\text{Be}$.

$$5. {}^6\text{Li}(p, \gamma){}^7\text{Be} \quad Q_m = 5.6064$$

Table 7.6: ${}^7\text{Be}$ levels from ${}^3\text{He} + {}^4\text{He}$ ^a

E_x (MeV \pm keV)	J^π	l_α	LS term	R (fm)	θ_α^2 ^b	θ_p^2	$\theta_{p'}^2$	Refs.
4.57 ± 50 ^c	$\frac{7}{2}^-$	3	${}^2\text{F}_{7/2}$	4.0	0.70 ± 0.04			(1967SP10)
4.566				4.4	0.34			(1968IV01)
6.73 ± 100 ^c	$\frac{5}{2}^-$	3	${}^2\text{F}_{5/2}$	4.0	1.36 ± 0.13	0.000 ± 0.002		(1967SP10)
7.21 ± 60 ^c	$\frac{5}{2}^-$	3	${}^4\text{P}_{5/2}$	4.0	0.010 ± 0.001	0.26 ± 0.02		(1967SP10)
9.27 ± 100	$\frac{7}{2}^-$	3	${}^4\text{D}_{7/2}$	4.0	0.70 ± 0.26	$0.29^{+0.09}_{-0.18}$	1.8 ± 0.5	(1967SP10)
10.0 ^d	$\frac{3}{2}^-$	1	$({}^4\text{P}_{3/2})$					(1967HA07, 1967HA08)
≈ 10.0 ^e	$\frac{1}{2}^-$		$({}^4\text{P}_{1/2})$					(1967HA07, 1967HA08)
11.00 ± 50 ^f	$\frac{3}{2}^-$	1	$({}^2\text{P}_{3/2}, {}^2\text{D}_{3/2})$			0.13 ± 0.02 ^g		(1967HA07, 1967HA08)

^a Compare to Table 7.10 in (1966LA04).

^b $\gamma^2 / (\frac{3}{2} \hbar^2 / \mu a^2)$.

^c See also (1968LE1K).

^d $\Gamma = 1.8$ MeV.

^e Broad.

^f $\Gamma = 0.4 \pm 0.05$ MeV; $T = \frac{3}{2}$.

^g $\theta_p^{2''}$.

Gamma transitions are observed to the ground (γ_0) and to the 0.43 MeV (γ_1) states. The yield shows no evidence of resonance for $E_p = 0.2$ to 1.0 MeV and the branching ratio remains approximately constant at $(62 \pm 5)\%$ to the ground state, 38% to ${}^7\text{Be}^*(0.43)$, $< 4\%$ to ${}^7\text{Be}^*(4.57)$ (1955BA59, 1956WA03, 1969JO1K, 1969SW1C). At $E_p = 1.06$ MeV a resonance in the yield of 0.43 MeV γ -rays is reported by (1968WO1E), corresponding to ${}^7\text{Be}^*(6.52)$. See, however, (1963MC09). See also (1966LA1D) and (1966LA04).

6. (a) ${}^6\text{Li}(p, p){}^6\text{Li}$

$$E_b = 5.6064$$

(b) ${}^6\text{Li}(p, 2p){}^5\text{He}$

$$Q_m = -4.59$$

Measurements of elastic angular distributions have recently been reported by (1968ME25: $E_p = 14$ to 15.8 MeV). Earlier measurements are listed in (1966LA04). Two resonances are reported at $E_p = 1.84$ and 5 MeV in the elastic yield [${}^7\text{Be}^*(7.21, 9.9)$]. The parameters of the lower resonance are shown in Table 7.3 (1963MC09). The 5 MeV resonance has $\Gamma \approx 1.8$ MeV and appears to also be formed by p-waves: γ_p^2 is then 3 ± 2 MeV \cdot fm. A weak rise near $E_p = 8$ to 9 MeV may indicate a further level, ${}^7\text{Be}^* \approx 13$ MeV (1963HA53). Differential cross sections are reported by (1971BI11: 6.868 MeV) and by (1969LE08: 1.36 MeV). See also ${}^6\text{Li}$. Polarization measurements (elastic scattering) have been carried out at $E_p = 1.21$ to 3.22 MeV (1969PE22), 14.5 MeV (1965RO22), 49.8 MeV (1971MA13), 152 MeV (1966RO1C), and 155 MeV (1968GE04). A phase-shifty analysis for $E_p = 0.5$ to 5.6 MeV shows that only 2S , 4S and

4P are involved. The $^4P_{5/2}$ phase resonances at $E_p = 1.8$ MeV, and the broad resonance at 5 MeV can be reproduced equally well by either $^4P_{3/2}$ or $^4P_{1/2}$: tensor polarization measurements are necessary to distinguish between the two (1969PE22). See (1966LA04) for earlier results. An S -matrix analysis of the cross section of this reaction and of reactions 3 and 9 has been reported by (1966HU1C, 1968LE1K).

The reaction cross section for formation of $^6\text{Li}^*(2.19)$ has been measured for $E_p = 3.6$ to 9.40 MeV: a broad resonance indicates the presence of a state with $E_x \approx 10$ MeV, $\Gamma = 1.8$ MeV, $J^\pi = (\frac{3}{2}, \frac{5}{2})^-$; $T = \frac{1}{2}$ (1967HA07, 1967HA08). The cross section and terms of two $J^\pi = \frac{3}{2}^-$ states at $E_x \approx 10$ and 11 MeV: see reaction 3 (1967HA07, 1967HA08). The total cross section for formation of $^6\text{Li}^*(3.56)$ decreases slowly with energy for $E_p = 24.3$ to 46.4 MeV (1968AU06). Polarization measurements involving the p_1 and p_2 groups have been carried out $E_p = 49.8$ MeV (1971MA44). For the total scattering cross section at 1 GeV, see (1967IG1A).

See also (1967CA1G, 1968OL1B), (1971PL1C), (1966LE1E, 1969WA11, 1970LE02, 1972BR20, 1973LA26; theor.). For reaction (b) see ^6Li , ^5He and (1973NA1M).

$$7. \ ^6\text{Li}(p, n)^6\text{Be} \qquad Q_m = -5.070 \qquad E_b = 5.6064$$

The yield of neutrons increases approximately monotonically from threshold to $E_p = 14.3$ MeV (1964BA16). See also (1971BU1D). For polarization measurements at $E_p = 30$ and 50 MeV, see (1969RO20). See also (1969CL06, 1971JU05), (1970RA33; theor.) and ^6Be .

$$8. \ ^6\text{Li}(p, d)^5\text{Li} \qquad Q_m = -3.44 \qquad E_b = 5.6064$$

See ^5Li .

$$9. \ ^6\text{Li}(p, \alpha)^3\text{He} \qquad Q_m = 4.0200 \qquad E_b = 5.6064$$

Over the range $E_p = 25$ to 50 keV, the cross section rises from 0.8 to 72 μb : in the formula $\sigma \approx E^{-1}e^{-B/\sqrt{E}}$, $B = 90 \pm 6$ keV $^{1/2}$ (1967FI05). Cross section measurements for $E_p = 62$ to 188 keV show deviation from an s-wave Gamow plot above ≈ 130 keV (1966GE11). Using cross-section measurements at $E_p = 151$ and 317 keV, as well as the (1966GE11) data, (1971SP05) calculate $S(0) = 3.0$ MeV \cdot b. See also (1969AU1C).

At higher energies the cross section exhibits a broad, low maximum near $E_p = 1$ MeV and a pronounced resonance at $E_p = 1.85$ MeV (1951BA79, 1956MA91). No other structure is reported up to $E_p = 5.6$ MeV (1963JE03, 1964FA03). Measurements between $E_p = 0.4$ and 3.4 MeV show that the polarizations are generally large and positive. The $E_p = 1.9$ MeV resonance appears in A_1 and A_2 (1968BR18). S -matrix analysis of the cross section of this reaction and of reactions 3 and 6 are reported by (1966HU1C, 1968LE1K). Angular distributions are reported at $E_p = 151$

and 317 keV (1971SP05), at $E_p = 0.5$ to 1.82 MeV (1969JO1J), 3 to 45 MeV (1973SC1V) and 12, 14 and 16 MeV (1973WE07). Angular distributions at $E_p = 8$ to 18.5 MeV have been analyzed using a finite-range multi-interaction DWBA formalism. The analysis leads to reduced widths of 0.69 for $\alpha + d$ in a relative s-state, 0.04 for $\alpha + d$ in a relative d-state and 0.44 for ${}^3\text{He} + t$ in a relative s-state (1973WE07). Yield and cross section measurements are reported by (1968BE1P: 0.3 to 1.0 MeV) and (1969LE08: $E_p = 1.36$ MeV). See also (1974GO1V) and see (1966LA04) for earlier measurements.

Searches for excited states of ${}^3\text{He}$ have been unsuccessful: see (1968OL1B: $E_p = 20$ MeV), (1971BR12, 1972BU16: $E_p = 45$ MeV). See also (1970KO25, 1972BE1Y). See (1971BR12, 1973FI04) for discussions of the excited states of ${}^4\text{He}$.

See also (1967SP09, 1967VA1F) and (1969BE1M, 1969BO1G, 1972CL1C, 1972HU09: theor.).

10. ${}^6\text{Li}(d, n){}^7\text{Be}$ $Q_m = 3.3818$

Two neutron groups are observed, corresponding to ${}^7\text{Be}^*(0, 0.43)$. Angular distributions of the n_0 and n_1 groups have been measured at $E_d = 0.24$ to 3.5 MeV [see (1966LA04) and (1966SC26)]: $l_p = 1$, $J^\pi \leq \frac{5}{2}^-$ for both states. At $E_d = 12, 15$ and 17 MeV differential cross sections for the population of ${}^7\text{Be}(\text{g.s.} + 0.43)$ have been measured by (1970GA07).

The n - γ correlations are isotropic, indicating $J^\pi = \frac{1}{2}^-$ for ${}^7\text{Be}^*(0.43)$ (1956NE13): $E_\gamma = 428.9 \pm 2$ keV (1952TH24). Broad maxima are observed in the ratio of low-energy to high-energy neutrons at $E_d = 4.2$ and 5.1 MeV [${}^7\text{Be}^*(6.5, 7.2)$, $\Gamma_{\text{cm}} = 1.2$ and 0.5 MeV, respectively] (1957SL01).

11. ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$ $Q_m = 0.1126$

Angular distributions of the d_0 and d_1 groups to ${}^7\text{Be}^*(0, 0.43)$ have been measured at $E_d = 8, 10, 14$ and 18 MeV: all the distributions show an $l = 1$ maximum at small angles. The DWBA analysis leads to a ratio of spectroscopic factors S^*/S [for ${}^7\text{Be}^*(0.43)/{}^7\text{Be}(0)$] = 1.55, in fair agreement with other measurements (1968LU02). See also (1964MA57) and (1970JA1J: theor.).

12. ${}^6\text{Li}(\alpha, t){}^7\text{Be}$ $Q_m = -14.2082$

Angular distributions of triton groups have been reported at $E_\alpha = 40$ MeV (1965OG03: ${}^7\text{Be}^*(0 + 0.43, 4.57 + (5.0 \pm 0.3))$), 43 MeV (1967DE1K: ${}^7\text{Be}^*(0, 0.43)$) and 46 MeV (1969FO1C: ${}^7\text{Be}^*(0, 0.43, 4.57)$). See also (1967OG1A).

13. ${}^6\text{Li}(p, n){}^7\text{Be}$ $Q_m = -1.64422$

$E_{\text{thresh.}} = 1880.612 \pm 0.090$ keV (1970RO07); the recommended value for $E_{\text{thresh.}}$ based on this and on other experiments is 1880.59 ± 0.08 keV (1970RO07). See also (1966MA60, 1966RO09, 1967MA1D, 1973MA1V). The excitation energy of the first excited state is 429.20 ± 0.10 keV (1972BO02). The lifetime of ${}^7\text{Be}^*(0.43)$, $\tau_m = 1.92 \pm 0.2$ msec. The ratio of this lifetime to that of ${}^7\text{Li}^*(0.48)$ is 1.82 ± 0.20 . This value, and the values of τ_m are in agreement with intermediate coupling calculations assuming LS coupling predominates ($a/K < 3.6$) (1966PA11).

Angular distributions are reported at $E_p = 1.9$ to 2.36 MeV (1967BE61; n_0), 2.1 to 3.8 MeV (1971BU1D; n_0), 2.6 to 5.4 MeV (1972EL19; n_0), 3.2 to 5.4 MeV (1972EL19; n_1), at 17.5 MeV (1972AZ01; n_0, n_1, n_2) and at 30 and 50 MeV (1969CL06; $n_0 + n_1$). Above $E_p \approx 7$ MeV, neutrons corresponding to ${}^7\text{Be}^*(4.57)$ are seen (1959AJ81, 1960HI04, 1963BO06). At $E_p = 30$ to 50 MeV, neutron groups are observed to states at $E_x = 4.61 \pm 0.07, 7.21 \pm 0.06, 9.6 \pm 0.3, 11.3 \pm 0.2, 12.3 \pm 0.2, 13.24 \pm 0.15, 14.39 \pm 0.15, 15.3 \pm 0.2, 16.3 \pm 0.2, 18.3 \pm 0.2, 19.7 \pm 0.3$ and 20.5 ± 0.2 MeV (1965BA39). See also (1967LO07, 1969JU1A, 1970BO1U, 1971DA24, 1971WA1J), (1966PA04, 1969MA1P, 1969MA1G, 1968TH1H, 1971TH1K; theor.), (1966LA04) and ${}^8\text{Be}$.

14. ${}^7\text{Li}({}^3\text{He}, t){}^7\text{Be}$ $Q_m = -0.8804$

Angular distributions have been measured at $E({}^3\text{He}) = 3.0$ to 4.0 MeV (1969OR01; t_0, t_1), 8.7 and 9.7 MeV (1968MA1W; t_0, t_1), 14 MeV (1969NU1A; t_0, t_1) and 25.2 MeV (1968BR1G; t_0, t_1, t_2). The width of ${}^7\text{Be}^*(4.57)$, $\Gamma_{\text{cm}} = 175 \pm 7$ keV (1971PI06). See also (1967BL1E) and (1971WE1L, 1971WE1M; theor.).

15. ${}^7\text{Li}({}^6\text{Li}, {}^6\text{He}){}^7\text{Be}$ $Q_m = -4.372$

At $E({}^6\text{Li}) = 31.8$ MeV, the reaction is observed to ${}^7\text{Be}^*((0 + 0.43), 4.57)$ (1971CH1B).

16. ${}^9\text{Be}(p, t){}^7\text{Be}$ $Q_m = -12.0831$

Angular distributions of tritons have been measured at $E_p = 43.7$ MeV (1965DE08, 1966CE05, 1968BR23: ${}^7\text{Be}^*(0, 0.43, 4.57, 6.51, 11.01)$) and 46 MeV (1967VE01: ${}^7\text{Be}^*(0 + 0.43, 4.57, 6.51, 10.79)$). The 11 MeV state has $E_x = 11.01 \pm 0.04$ MeV (1968BR23), $\Gamma = 298 \pm 25$ keV, $J^\pi = \frac{3}{2}^-$; $T = \frac{3}{2}$ [the J^π ; T assignments are based on the similarity of the angular distribution to that in the $(p, {}^3\text{He})$ reaction to ${}^7\text{Li}^*(11.13)$] (1965DE08). See also (1967MC14), (1969BA1Z, 1969IN1A) and (1967BA1E; theor.).

17. ${}^{10}\text{B}(p, \alpha){}^7\text{Be}$ $Q_m = 1.1462$

Alpha groups corresponding to ${}^7\text{Be}^*(0, 0.43)$ have been studied by many observers: see (1952AJ38, 1959AJ76). Some reported values for the energy of the first excited state are: 431 ± 5 keV (1950VA01), 434.4 ± 4 keV (1951BR10), 429 ± 3 keV (1952CR30), 428.5 ± 1.8 keV (1952TH24). In addition the excitation of ${}^7\text{Li}^*(4.72 \pm 0.08, 6.27 \pm 0.10, 7.21 \pm 0.10, (14.6 \pm 0.3))$ is reported by (1955RE16). At $E_p = 18.6$ MeV ($\theta = 30^\circ$) (1968PA15) find that the intensity of a group to ${}^7\text{Be}^*(14.6)$ is $< 2\%$ that reported by (1955RE16). Angular distributions have been measured at $E_p = 2.8$ to 7.0 MeV (1964JE01). See also ${}^{11}\text{C}$ and (1966LA04).

$$18. {}^{10}\text{B}({}^3\text{He}, {}^6\text{Li}){}^7\text{Be} \quad Q_m = -2.874$$

At $E({}^3\text{He}) = 30.0$ MeV angular distributions have been obtained for the transitions to ${}^7\text{Be}^*(0, 0.43) + {}^6\text{Li}^*(0, 2.19)$: see (1970DE12, 1972OH01).

$$19. {}^{10}\text{B}(\alpha, {}^7\text{Li}){}^7\text{Be} \quad Q_m = -16.202$$

At $E_\alpha = 45.6$ MeV (1969FO06) have measured the angular distributions of the ${}^7\text{Li}$ and of the ${}^7\text{Be}$ ions, corresponding to the ground-state transitions. At a given angle the intensities of the two ions are the same, implying that the wave functions of the ground states of ${}^7\text{Li}$ and ${}^7\text{Be}$ are very similar (1969FO06). See also (1971RO1A, 1971SI28, 1973NA1N; theor.).

$$20. {}^{11}\text{B}(d, {}^6\text{He}){}^7\text{Be} \quad Q_m = -11.563$$

At $E_d = 40$ MeV the cross section of the ground-state reaction is twice (to $\pm 10\%$) that for the reaction to ${}^6\text{Li}^*(3.56) + {}^7\text{Li}(0)$, in agreement with isospin conservation (1972GO1P).

$$21. {}^{11}\text{B}({}^3\text{He}, {}^7\text{Li}){}^7\text{Be} \quad Q_m = -7.079$$

See (1971RO1A).

$$22. {}^{12}\text{C}(p, {}^6\text{Li}){}^7\text{Be} \quad Q_m = -22.569$$

At $E_p = 36.0, 40.7, 45.0, 50.0$ and 56.8 MeV angular distributions have been obtained for the transitions to ${}^7\text{Be}^*(0 + 0.43)$ (1971BR07, 1971HO25, 1971HO1K).

23. $^{12}\text{C}(\text{d}, ^7\text{Li})^7\text{Be}$ $Q_m = -17.543$

At $E_d = 39.8$ MeV angular distributions have been measured for the transitions $^7\text{Li}(0)+^7\text{Be}(0)$, $^7\text{Li}^*(0.43) + ^7\text{Be}(0)$, $^7\text{Li}(0) + ^7\text{Be}^*(0.43)$ and $^7\text{Li}^*(0.43) + ^7\text{Be}^*(0.43)$. The ratios of the $^7\text{Li}(0)$ and $^7\text{Be}(0)$ cross sections show asymmetries exceeding 20% (1971HO1K, 1971YO06). See also (1971SI28; theor.).

24. $^{12}\text{C}(^3\text{He}, ^8\text{Be})^7\text{Be}$ $Q_m = -5.780$

Angular distributions have been obtained at $E(^3\text{He}) = 25.5$ to 29 MeV (1972PI1A, 1973PI1D: $^7\text{Be}^*(0, 0.43)$), 28 and 30 MeV (1970DE12, 1973KL1B: $^7\text{Be}^*(0, 0.43) + ^8\text{Be}^*(0, 2.9)$), 35.6 MeV (1969NE1D, 1969ZE1A, 1970FO1D: $^7\text{Be}^*(0, 0.43)$) and 37 and 41 MeV (1967ZA1B: $^7\text{Be}^*(0 + 0.43)$). See also (1965EN01).

25. $^{12}\text{C}(\alpha, ^9\text{Be})^7\text{Be}$ $Q_m = -24.694$

At $E_\alpha = 42$ MeV, angular distributions have been measured to $^7\text{Be}^*(0, 0.43)+^9\text{Be}(0)$ (1972RU03).

26. $^{13}\text{C}(\text{p}, ^7\text{Li})^7\text{Be}$ $Q_m = -20.264$

At $E_p = 45.0$ MeV the angular distribution has been measured for the transition to $^7\text{Li}(0) + ^7\text{Be}(0)$ (1971BR07).

27. $^{16}\text{O}(\text{p}, ^{10}\text{B})^7\text{Be}$ $Q_m = -25.270$

See (1969HO1H).

28. $^{16}\text{O}(^3\text{He}, ^{12}\text{C})^7\text{Be}$ $Q_m = -5.575$

Angular distributions are reported at $E(^3\text{He}) = 25.5$ to 29 MeV (1972PI1A: to $^7\text{Be}^*(0, 0.43)$), 30 MeV (1970DE12, 1973KL1B: to $^{12}\text{C}^*(0, 4.4, 7.7, 9.6) + ^7\text{Be}^*(0, 0.43)$) and 41 MeV (1967ZA1B).

29. $^{16}\text{O}(\alpha, ^{13}\text{C})^7\text{Be}$ $Q_m = -21.207$

At $E_\alpha = 42$ MeV, angular distributions have been obtained for the transitions to $^7\text{Be}^*(0, 0.43) + ^{13}\text{C}(0)$ (1972RU03).

30. $^{19}\text{F}(\text{d}, ^{14}\text{C})^7\text{Be}$ $Q_m = -7.140$

The angular distribution to $^7\text{Be}^*(0 + 0.43) + ^{14}\text{C}(0)$ has been measured at $E_d = 14.9$ MeV (1967DE03).

31. (a) $^{19}\text{F}(^3\text{He}, ^{15}\text{N})^7\text{Be}$ $Q_m = -2.426$

(b) $^{20}\text{Ne}(^3\text{He}, ^{16}\text{O})^7\text{Be}$ $Q_m = -3.144$

(1970DE12, 1973KL1B) have studied, at $E(^3\text{He}) = 30$ MeV, the angular distributions to $^{15}\text{N}(0) + ^7\text{Be}^*(0, 0.43)$ and to $^{16}\text{O}^*(0, 6.06 + 6.13) + ^7\text{Be}^*(0, 0.43)$.

^7B

(Fig. 10)

1. $^{10}\text{B}(^3\text{He}, ^6\text{He})^7\text{B}$ $Q_m = -18.55$

A ^6He group corresponding to the unbound ground state of ^7B has been identified at $E(^3\text{He}) = 50$ MeV: $M - A (^7\text{B}) = 27.94 \pm 0.10$, $\Gamma = 1.4 \pm 0.2$ MeV. The isobaric quartet mass law would predict $M - A = 27.76 \pm 0.17$ MeV. ^7B is unbound with respect to $^6\text{Be} + \text{p}$ ($Q = 2.27$), $^5\text{Li} + 2\text{p}$ ($Q = 1.68$), $^4\text{He} + 3\text{p}$ ($Q = 3.65$). The expected single-particle width is $\Gamma = 0.64$ MeV: it is suggested that the two-proton and three-proton decays make an appreciable contribution to the width (1967MC14). See also (1960GO1B, 1968CE01, 1968ST1J, 1969GA1P, 1972CA37, 1972CE1A).

References

(Closed December 31, 1973)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1949SE20 E. Segre and C.E. Wiegand, Phys. Rev. 75 (1949) 39; Erratum Phys. Rev. 81 (1951) 284
- 1950VA01 D.M. Van Patter, A. Sperduto, E.N. Strait and W.W. Buechner, Phys. Rev. 79 (1950) 900
- 1951BA79 S. Bashkin and H.T. Richards, Phys. Rev. 84 (1951) 1124
- 1951BR10 A.B. Brown, C.W. Snyder, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 82 (1951) 159
- 1952AJ38 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 24 (1952) 321
- 1952CR30 D.S. Craig, D.J. Donahue and K.W. Jones, Phys. Rev. 88 (1952) 808
- 1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
- 1953KR16 J.J. Kraushaar, E.D. Wilson and K.T. Bainbridge, Phys. Rev. 90 (1953) 610
- 1954AL24 D.L. Allan, Nature 174 (1954) 267
- 1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684
- 1954GO1A Goldemberg and Katz, Can. J. Phys. 32 (1954) 49
- 1954JO17 C.H. Johnson, H.B. Willard and J.K. Bair, Phys. Rev. 96 (1954) 985
- 1954TI16 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. A67 (1954) 469
- 1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
- 1955BA59 S. Bashkin and R.R. Carlson, Phys. Rev. 97 (1955) 1245
- 1955RE16 J.B. Reynolds, Phys. Rev. 98 (1955) 1289
- 1956BO36 P. Bouchez, J. Tobailem, J. Robert, R. Muxart, R. Mellet and P. Daud, J. Phys. Rad. 17 (1956) 363
- 1956MA91 J.B. Marion, G. Weber and F.S. Mozer, Phys. Rev. 104 (1956) 1402
- 1956NE13 G.C. Neilson and J.B. Warren, Phys. Rev. 103 (1956) 1758
- 1956WA03 J.B. Warren, T.K. Alexander and G.B. Chadwick, Phys. Rev. 101 (1956) 242
- 1957BR97 C.P. Browne, Bull. Amer. Phys. Soc. 2 (1957) 350, N3
- 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809

- 1958BU38 W.E. Burcham, G.P. McCauley, D. Bredin, W.M. Gibson, D.J. Prowse and J. Rotblat, Nucl. Phys. 5 (1958) 141
- 1958RY77 T.W. Rybka and L. Katz, Phys. Rev. 110 (1958) 1123
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AJ81 F. Ajzenberg-Selove, C.F. Osgood and C.P. Baker, Phys. Rev. 116 (1959) 1521
- 1959BA46 A.J. Bame, Jr. and R.L. Cubitt, Phys. Rev. 114 (1959) 1580
- 1959GA08 F. Gabbard, R.H. Davis and T.W. Bonner, Phys. Rev. 114 (1959) 201
- 1959HO03 H.D. Holmgren and R.L. Johnston, Phys. Rev. 113 (1959) 1556
- 1959MU25 R.B. Murray and H.W. Schmitt, Phys. Rev. 115 (1959) 1707
- 1960GO1B Goldansky, Nucl. Phys. 19 (1960) 482
- 1960HA14 E.W. Hamburger and J.R. Cameron, Phys. Rev. 117 (1960) 781
- 1960HI04 K. Hisatake, Y. Ishizaki, A. Isoya, T. Nakamura, Y. Nakano, B. Saheki, Y. Saji and K. Yuasa, J. Phys. Soc. Jpn. 15 (1960) 741
- 1960HU08 D.J. Hughes, B.A. Magurno and M.K. Brussel, BNL 325, 2nd Edition, Suppl. Vol. 1 (1960)
- 1960MA32 R.D. Macfarlane and J.B. French, Rev. Mod. Phys. 32 (1960) 567
- 1961GR27 G.M. Griffiths, R.A. Morrow, P.J. Riley and J.B. Warren, Can. J. Phys. 39 (1961) 1397
- 1961HO21 H.D. Holmgren and L.M. Cameron, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 531
- 1961KO1A Koltun, Phys. Rev. 124 (1961) 1162
- 1961LA1A Lane, Langsdorf, Monahan and Elwyn, Ann. Phys. 12 (1961) 135
- 1962GR08 A.G. Gregory, T.R. Sherwood and E.W. Titterton, Nucl. Phys. 32 (1962) 543
- 1962SL02 R.J. Slobodrian, Phys. Rev. 125 (1962) 1003
- 1962TA11 J.G.V. Taylor and J.S. Merritt, Can. J. Phys. 40 (1962) 926
- 1963AS01 V.J. Ashby, H.C. Catron, M.D. Goldberg, R.W. Hill, J.M. Le Blanc, L.L. Newkirk, J.P. Stoering, C.J. Taylor and M.A. Williamson, Phys. Rev. 129 (1963) 1771
- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963BE26 M. Bernheim and G.R. Bishop, Phys. Lett. 5 (1963) 294
- 1963BE53 M. Bernheim and G.R. Bishop, J. Phys. (France) 24 (1963) 970
- 1963BO06 R.R. Borchers and C.H. Poppe, Phys. Rev. 129 (1963) 2679
- 1963HA53 W.D. Harrison and A.B. Whitehead, Phys. Rev. 132 (1963) 2607
- 1963JE03 J.M.F. Jeronymo, G.S. Mani and A. Sadeghi, Nucl. Phys. 43 (1963) 424

1963MC09 J.A. McCray, Phys. Rev. 130 (1963) 2034
 1963PA12 P.D. Parker and R.W. Kavanagh, Phys. Rev. 131 (1963) 2578
 1963TO06 T.A. Tombrello and P.D. Parker, Phys. Rev. 131 (1963) 2582
 1964BA09 A.C.L. Barnard, C.M. Jones and G.C. Phillips, Nucl. Phys. 50 (1964) 629
 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
 1964FA03 U. Fasoli, D. Toniolo and G. Zago, Phys. Lett. 8 (1964) 127
 1964FO1A Fowler and Vogl, Lect. in Theor. Phys., Vol. 6 (1964) 379
 1964GR40 L. Green and D.J. Donahue, Phys. Rev. 135 (1964) B701
 1964JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Nucl. Phys. 50 (1964) 516
 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed., W.H. Johnson, Jr. (1964) 305
 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Edition, Suppl. Vol. 1 (1964)
 1965BA39 C.J. Batty, E. Friedman, P.C. Rowe and J.B. Hunt, Phys. Lett. 19 (1965) 35
 1965BE1E Berggren, Ark. Fys. 30 (1965) 508
 1965BO1C Bodmer and Murphy, Nucl. Phys. 73 (1965) 664
 1965BU1C Burnett, Fowler and Hoyle, Geochim. Et Cosmochim. Acta 29 (1965) 1209
 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707
 1965DA06 P.J. Dallimore, K.S. Lam and H.H. Thies, Aust. J. Phys. 18 (1965) 389
 1965DE08 C. Detraz, J. Cerny and R.H. Pehl, Phys. Rev. Lett. 14 (1965) 708
 1965EN01 J.B.A. England and B.L. Reece, Nucl. Phys. 72 (1965) 449
 1965FU19 M. Furukawa, S. Kume and M. Ogawa, Nucl. Phys. 69 (1965) 362
 1965GI10 L. Gilly, M. Jean, R. Meunier, M. Spighel, J.P. Stroot and P. Duteil, Phys. Lett. 19 (1965) 335
 1965GR1C Gradsztajn, Ann. Phys. 10 (1965) 791
 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
 1965HA19 E. Hayward and T. Stovall, Nucl. Phys. 69 (1965) 241
 1965HU13 R.W. Huggins and J.H. Sanders, Proc. Phys. Soc. 86 (1965) 53
 1965JA1A Jacmart, Cahiers Phys. (France) 19 (1965) 1
 1965KU09 D. Kurath, Phys. Rev. 140 (1965) B1190
 1965LA1B Law, Nuovo Cim. 38 (1965) 807

1965LO1B Lodder and Jonker, Phys. Lett. 18 (1965) 310
 1965MO17 B.M. Morris and R.D. Present, Phys. Rev. 140 (1965) B1197
 1965NE1B Neudachin and Smirnov, At. Energy Rev. 3 (1965) 157
 1965OG03 A.A. Ogloblin and V.I. Chuev, Yad. Fiz. 2 (1965) 670; Sov. J. Nucl. Phys. 2 (1966) 480
 1965PR04 R.D. Present, Phys. Rev. 139 (1965) B300
 1965RI09 J. Rickards, Rev. Mex. Fis. 14 (1965) 241
 1965RO09 R.L. Robinson, P.H. Stelson, F.K. McGowan, J.L.C. Ford, Jr. and W.T. Milner, Nucl. Phys. 74 (1965) 281
 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96
 1965SC07 S. Schwarz, L.G. Stromberg and A. Bergstrom, Nucl. Phys. 63 (1965) 593
 1965VO1A Volkov, Nucl. Phys. 74 (1965) 33
 1965WA19 A. Wataghin, M. Scotto and G. Paoli, Nuovo Cim. B40 (1965) 441
 1965ZH1A Zhdanov, Kuzmin and Yakovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 239
 1966AG1A Agee and Rosen, LA-3538-MS (1966)
 1966AU1A Austin et al., Bull. Amer. Phys. Soc. 11 (1966) 10
 1966BA1Q Bardolle, Cabe, Chretien and Laurat, J. Phys. (Paris) 27 (1966) C1-96
 1966BA1V Barry, Bull. Amer. Phys. Soc. 11 (1966) 655
 1966BA1W Bazhanov, Komar and Kulikov, Dokl. Akad. Nauk SSSR 171 (1966) 549
 1966BA26 F.C. Barker, Nucl. Phys. 83 (1966) 418
 1966BE1E Beaumevieuille et al., J. Phys. (Paris) 27 (1966) C1-150
 1966BL1C Bluet, Fort and Leroy, Antwerp 1965 Neutron Conf. (North Holland, 1966) 566
 1966BR1M Bramblett et al., Bull. Amer. Phys. Soc. 11 (1966) 367
 1966BR25 G. Bruno, J. Decharge, A. Perrin, G. Surget and C. Thibault, J. Phys. (France) 27 (1966) 517
 1966CA1F Cade, Bull. Amer. Phys. Soc. 11 (1966) 11
 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, Phys. Rev. 152 (1966) 950
 1966DA1A D. Davies, H. Muirhead and J.N. Woulds, Nucl. Phys. 78 (1966) 663
 1966DE1E Detraz, J. Phys. (Paris) 27 (1966) C1-64
 1966DE1G Delorme and Ericson, J. Phys. (Paris) 27 (1966) C1-120
 1966DO1A Dolinov and Melikov, Vestnik Mosk. Univ. Fiz. Astron. (1966) 116
 1966DZ07 R.I. Dzhibuti, V.I. Mamasakhlisov and T.S. Macharadze, Yad. Fiz. 4 (1966) 52; Sov. J. Nucl. Phys. 4 (1967) 37

1966EL08 F. El-Batanoni and A.A. Kresnin, Nucl. Phys. 89 (1966) 577
 1966GA15 H. Gauvin, Compt. Rend. 263 (1966) 752
 1966GE11 W. Gemeinhardt, D. Kamke and C. von Rhoneck, Z. Phys. 197 (1966) 58
 1966GE12 L.B. Geesaman, H.B. Knowles and J.B. Mead, Nucl. Phys. 88 (1966) 177
 1966GO1C Goldemberg and Pratt, Rev. Mod. Phys. 38 (1966) 311
 1966GU1C Guth and Edge, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 238
 1966HA09 E.T. Hazzard, F. Ajzenberg-Selove and P.V. Hewka, Nucl. Phys. 75 (1966) 592; Erratum Nucl. Phys. 89 (1966) 706
 1966HA18 E.C. Halbert, Y.E. Kim and T.T.S. Kuo, Phys. Lett. 20 (1966) 657
 1966HU1B Humphrey and Finlay, Bull. Amer. Phys. Soc. 11 (1966) 831
 1966HU1C J. Humblet and A. Lejeune, Phys. Lett. 23 (1966) 561
 1966IS01 R. Isler, S. Marcus and R. Novick, Bull. Amer. Phys. Soc. 11 (1966) 62, CH13
 1966JA05 R. Jahr, K. Kayser, A. Kostka and J.P. Wurm, Nucl. Phys. 76 (1966) 79
 1966JA1A Jacob and Maris, Rev. Mod. Phys. 38 (1966) 121
 1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
 1966LA1D Langevin-Joliot, Z. Naturforsch. A21 (1966) 1737
 1966LE1E Lejeune, Bull. Soc. Roy. Sci. Liege (Belgium) 35 (1966) 566
 1966MA17 G.E. Manuzio, G. Ricco and M. Sanzone, Nuovo Cim. B42 (1966) 348
 1966MA1L Mahaux and Robaye, Antwerp 1965 Neutron Conf. (1966) 499
 1966MA1N Mahalanabis and Jackson, Proc. Phys. Soc. 87 (1966) 677
 1966MA38 S.K. Mark, P.M. Portner and R.B. Moore, Can. J. Phys. 44 (1966) 2961
 1966MA60 J.B. Marion, Rev. Mod. Phys. 38 (1966) 660
 1966ME1C Merchez et al., J. Phys. (Paris) 27 (1966) C1-61
 1966ME1E Melikov, Vestnik Mosk. Univ. Fiz. Astron. 6 (1966) 102
 1966MI1C T. Mikumo, R. Seki and Y. Tagishi, M. Furukawa and H. Yamaguchi, Phys. Lett. 23 (1966) 586
 1966MU1A Murray and Strachan, Proc. Phys. Soc. 87 (1966) 641
 1966PA04 D.W. Palmer, Nucl. Phys. 75 (1966) 529
 1966PA11 P. Paul, J.B. Thomas and S.S. Hanna, Phys. Rev. 147 (1966) 774
 1966PH1A Phillips, 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe (1966)
 1966PI02 L.L. Pinsonneault and J.M. Blair, Phys. Rev. 141 (1966) 961

1966RA1B Rahman, Khan and Sen Gupta, Nuovo Cim. 44 (1966) 36
 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246
 1966RE1B Regis et al., J. Phys. (Paris) 27 (1966) C1-84
 1966RO09 M.L. Roush, L.A. West, J.V. Mullendore, H.L. Fann and J.B. Marion, Phys. Lett. 23 (1966) 355
 1966RO1C Rolland et al., J. Phys. (Paris) 27 (1966) C1-126
 1966RO1J Robson, Proc. Phys. Soc. 88 (1966) 351
 1966RO1L Robaye et al., Antwerp 1965 Neutron Conf. (1966) 500
 1966SA04 B. Sahai, Phys. Rev. 142 (1966) 612
 1966SC1E Schuller, Z. Phys. 192 (1966) 173
 1966SC26 W.A. Schier, G. Michel and R.E. Benenson, Nucl. Phys. 88 (1966) 373
 1966ST09 M.F. Steuer and C.A. Kelsey, Nucl. Phys. 83 (1966) 401
 1966TH1C W.J. Thompson, Phys. Lett. 23 (1966) 132
 1966VA1B Van Bluemel and Brussel, Bull. Amer. Phys. Soc. 11 (1966) 10
 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97
 1966YO1B Yoccoz, J. Phys. (Paris) 27 (1966) C1-3
 1967AR1A Artus et al., Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 314
 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
 1967BA1E Bang, Zelemskaya, Magzumov and Neudachin, Sov. J. Nucl. Phys. 4 (1967) 688
 1967BE1F Beery, Harper, Stovall and Rosen, Los Alamos Sci. Lab. Rept. LA-3788 (1967)
 1967BE61 A. Bergstrom, S. Schwarz, L.G. Stromberg and L. Wallin, Ark. Fys. 34 (1967) 153
 1967BI1E Bing, Seltz, Gerardin and Magnac-Valette, Private Communication (1967)
 1967BL03 W.W. Black and R.L. Heath, Nucl. Phys. A90 (1967) 650
 1967BL1E Bluemel, Thesis, Univ. of Illinois (1967)
 1967BO1C M. Bouten, P. Van Leuven, H. Depuydt and L. Schotsmans, Nucl. Phys. A100 (1967) 90
 1967BO22 M. Bouten, M-C. Bouten and P. Van Leuven, Nucl. Phys. A102 (1967) 322
 1967CA02 A.L. Catz and S. Amiel, Nucl. Phys. A92 (1967) 222
 1967CA17 P. Camiz, Nuovo Cim. B51 (1967) 190
 1967CA1G Cabe, Bardolle and Laurat, Private Communication (1967)
 1967CO01 J.A. Cookson, D. Dandy and J.C. Hopkins, Nucl. Phys. A91 (1967) 273
 1967CO1J Coop, Poate and Titterton, Aust. J. Phys. 20 (1967) 609

1967CO1N Cox and Pontet, J. Nucl. Energy 21 (1967) 271
 1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1
 1967CO36 S.W. Cospers, J. Cerny and R.C. Gatti, Phys. Rev. 154 (1967) 1193
 1967CR1E Crawley and Austin, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 165
 1967DA1C Danzinger, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 81
 1967DE03 L.J. Denes and W.W. Daehnick, Phys. Rev. 154 (1967) 928
 1967DE11 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 5 (1967) 490; Sov. J. Nucl. Phys. 5 (1967) 344
 1967DE15 A.J. Deruytter and P. Pelfer, J. Nucl. Energy 21 (1967) 833
 1967DE1K Dehnhard and Siemssen, Bull. Amer. Phys. Soc. 12 (1967) 17
 1967DU1B F. Dunnill, T. J. Gray, H.T. Fortune and N.R. Fletcher, Nucl. Phys. A93 (1967) 201
 1967EL1B L.R.B. Elton and A. Swift, Nucl. Phys. A94 (1967) 52
 1967EL1C L.R. Elton and D.F. Jackson, Phys. Rev. 155 (1967) 1070
 1967FA1A W.M. Fairbairn, Nucl. Phys. A90 (1967) 135
 1967FI05 O. Fiedler and P. Kunze, Nucl. Phys. A96 (1967) 513
 1967FU1E Fulmer, Williams, Dell and Blumberg, Bull. Amer. Phys. Soc. 12 (1967) 499
 1967GE1A Genz, Renier, Ledingham and Fink, Bull. Amer. Phys. Soc. 12 (1967) 1123
 1967HA07 W.D. Harrison, Nucl. Phys. A92 (1967) 253
 1967HA08 W.D. Harrison, Nucl. Phys. A92 (1967) 260
 1967IG1A G.J. Igo, J.L. Friedes, H. Palevsky, R. Sutter, G. Bennett, W.D. Simpson, D.M. Corley and R.L. Stearns, Nucl. Phys. B3 (1967) 181
 1967JA1E B.K. Jain and D.F. Jackson, Nucl. Phys. A99 (1967) 113
 1967JO1C Joseph et al., Bull. Amer. Phys. Soc. 12 (1967) 465
 1967JO1F Johansson, Sundberg and Tibell, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 63
 1967KA1A Kabachnik and Grishanova, Sov. J. Nucl. Phys. 4 (1967) 583
 1967LO07 P.J. Locard, S.M. Austin and W. Benenson, Phys. Rev. Lett. 19 (1967) 1141
 1967MA1D Marion, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 497
 1967MC14 R.L. McGrath, J. Cerny and E. Norbeck, Phys. Rev. Lett. 19 (1967) 1442
 1967ME11 D.F. Measday and J.N. Palmieri, Phys. Rev. 161 (1967) 1071
 1967MI1A Mitler, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 59
 1967OG1A Ogloblin, Proc. Problem Symp. on Nucl. Phys., Tbilisi, 1967 (1967) 169
 1967OK1A Okai, Prog. Theor. Phys. 38 (1967) 1413

- 1967PA1G Pauling, Proc. Nat. Acad. Sci. U. S. 58 (1967) 2175
- 1967RA24 N.C. Rasmussen, V.J. Orphan and Y. Hukai, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 278
- 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
- 1967SH05 Y.Y. Sharon, Nucl. Phys. A99 (1967) 321
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SH1E Shevchenko, Proc. Int. Conf. Electromag. Inter. Dubna, 1967, Vol. 3 (1967) 206
- 1967SM1A I.L. Smith, J. Garvey, J.G. Rutherglen and G.R. Brookes, Nucl. Phys. B1 (1967) 483
- 1967SP09 A. Sperduto, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 657
- 1967SP10 R.J. Spiger and T.A. Tombrello, Phys. Rev. 163 (1967) 964
- 1967ST04 R.H. Stokes and P.G. Young, Phys. Rev. Lett. 18 (1967) 611
- 1967ST1D Stotland, Allen, Parker and Yerke, Bull. Amer. Phys. Soc. 12 (1967) 483
- 1967SU1A L.R. Suelzle, M.R. Yearian and H. Crannell, Phys. Rev. 162 (1967) 992; Erratum Phys. Rev. 168 (1968) 1414
- 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, Nucl. Instrum. Meth. 56 (1967) 325
- 1967TO1B Tombrello, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 195
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
- 1967VA1F Valkovic, Emerson, Jackson and Phillips, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 989
- 1967VE01 J.W. Verba, H. Willmes, R.F. Carlson, I. Slaus, J.R. Richardson and E.L. Petersen, Phys. Rev. 153 (1967) 1127
- 1967WI06 I.R. Williams and C.B. Fulmer, Phys. Rev. 154 (1967) 1005
- 1967WI20 I.R. Williams and C.B. Fulmer, Phys. Rev. 162 (1967) 1055
- 1967ZA1B Zafiratos and Slee, Bull. Amer. Phys. Soc. 12 (1967) 893
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968AU06 S.M. Austin, P.J. Locard, W. Benenson and G.M. Crawley, Phys. Rev. 176 (1968) 1227
- 1968BA1H V.V. Balashov and D.V. Meboniya, Nucl. Phys. A107 (1968) 369
- 1968BA2E J.N. Bahcall, N.A. Bahcall, W.A. Fowler and G. Shaviv, Phys. Lett. B26 (1968) 359
- 1968BA2F J.N. Bahcall, N.A. Bahcall and G. Shaviv, Phys. Rev. Lett. 20 (1968) 1209
- 1968BE1E Bertrand, Grenier and Pernet, Comm. A L'energie Atomique, Rept. CEA 3504 (1968)
- 1968BE1F H.W. Bertini, Phys. Rev. 171 (1968) 1261

- 1968BE1P Bertrand, Grenier and Pornet, France, Comm. A L'energie Atomique, Rept. CEA 3428 (1968)
- 1968BE1Q Becker, Bahr, Bilaniuk and Jahr, Bull. Amer. Phys. Soc. 13 (1968) 565
- 1968BO1R M. Bouten, M.-C. Bouten and P. Van Leuven, Nucl. Phys. A111 (1968) 385
- 1968BO1T P.E. Boynton, T.J. Devlin, J. Solomon and V. Perez-Mendez, Phys. Rev. 174 (1968) 1083
- 1968BO32 N.E. Booth, A. Beretvas, R.E.P. Davis, C. Dolnick, R.E. Hill, M. Raymond and D. Sherden, Nucl. Phys. A119 (1968) 233
- 1968BR18 L. Brown and C. Petitjean, Nucl. Phys. A117 (1968) 343
- 1968BR1G Brussel, Hoffswell, Noweir and Warshaw, Bull. Amer. Phys. Soc. 13 (1968) 83
- 1968BR1H R.E. Brown and Y.C. Tang, Phys. Rev. 176 (1968) 1235
- 1968BR23 H. Brunnader, J.C. Hardy and J. Cerny, Phys. Rev. 174 (1968) 1247
- 1968CE01 J. Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968DI1B Di Napoli et al., Nuovo Cim. B55 (1968) 95
- 1968DO13 V.K. Dolinov, Yad. Fiz. 8 (1968) 26; Sov. J. Nucl. Phys. 8 (1969) 15
- 1968EL06 S.K. el Samarai, Y.F. Smirnov and B.A. Yurev, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1709; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1575
- 1968FA1D Farrell and Pineo, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 153
- 1968FI02 R.W. Fink, Nucl. Phys. A110 (1968) 379
- 1968FOZY D.T. Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299, Vol. 1 (1968) 1
- 1968GE04 B. Geoffrion, N. Marty, M. Morlet, B. Tatischeff and A. Willis, Nucl. Phys. A116 (1968) 209
- 1968GI1D Gibbons, Neutron Cross Sections and Tech., NBS Special Pub. 299 (1968) 111
- 1968GL1A Glauber, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 41, 60
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GO1J Goldemberg, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 379
- 1968HA1C Hayakawa, Suppl. Prog. Theor. Phys. (1968) 156
- 1968HA1G Hackenbroich, Wildermuth and Wittern, in Tokyo (1968) 44
- 1968HA1J Harvey, Aust. J. Phys. 21 (1968) 103
- 1968HI1E Hibdon and Mooring, Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 159
- 1968HO03 J.C. Hopkins, D.M. Drake and H. Conde, Nucl. Phys. A107 (1968) 139
- 1968HU18 J. Hudis and S. Tanaka, Phys. Rev. 171 (1968) 1297

- 1968HU1C Hutcheon, Thesis, Univ. of Saskatchewan (1968)
- 1968IV01 M. Ivanovich, P.G. Young and G.G. Ohlsen, Nucl. Phys. A110 (1968) 441
- 1968JA08 R. Jambunathan and R.K. Hobbie, Phys. Rev. 172 (1968) 1065
- 1968JA1G Jackson, in Tokyo (1968) 105
- 1968KA24 S. Kakigi, N. Fujiwara, K. Fukunaga, D.-C. Nguyen, S. Yamashita and T. Yanabu, J. Phys. Soc. Jpn. 25 (1968) 1214
- 1968KA38 N.N. Kaushal, E.J. Winhold, P.F. Yergin, H.A. Medicus and R.H. Augustson, Phys. Rev. 175 (1968) 1330
- 1968KE03 P.W. Keaton, Jr., D.D. Armstrong and L.R. Veaser, Phys. Rev. Lett. 20 (1968) 1392
- 1968KN1B Knitter and Coppola, Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 827
- 1968KU1B Kudeyarov, Neudachin, Serebryakov and Smirnov, Sov. J. Nucl. Phys. 6 (1968) 876
- 1968KU1D Kurath, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 393
- 1968LE1K A. Lejeune, Nucl. Phys. A116 (1968) 72
- 1968LI1C Li and Mark, Can. J. Phys. 46 (1968) 2645
- 1968LO1A Locci and Picchi, Nuovo Cim. A57 (1968) 803
- 1968LU02 H. Ludecke, T. Wan-Tjin, H. Werner and J. Zimmerer, Nucl. Phys. A109 (1968) 676
- 1968MA02 G.S. Mani and A.D.B. Dix, Nucl. Phys. A106 (1968) 251
- 1968MA19 J.L. Matthews, W. Bertozzi, S. Kowalski, C.P. Sargent and W. Turchinets, Nucl. Phys. A112 (1968) 654
- 1968MA1W Mancuso, Knudson and Wolicki, Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968MA25 S. Matsuki, J. Phys. Soc. Jpn. 24 (1968) 1203
- 1968ME25 F. Merchez, R. Bouchez, R.A. Hoffswell and A.I. Yavin, J. Phys. (France) 29 (1968) 969
- 1968MI1D Mikumo et al., in Tokyo (1968) 256
- 1968NE1B Neumann, Bull. Amer. Phys. Soc. 13 (1968) 100
- 1968NO1A M.E. Nordberg, K.F. Kinsey and R.L. Burman, Phys. Rev. 165 (1968) 1096
- 1968OK06 Y. Okuma, J. Phys. Soc. Jpn. 25 (1968) 1
- 1968OL1B D.K. Olsen and R.E. Brown, Phys. Rev. 176 (1968) 1192
- 1968PA15 P.D. Parker, D.W. Mingay and J.C. Overley, Phys. Rev. 176 (1968) 1211
- 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
- 1968PE1B Pellegrini, Nuovo Cim. B54 (1968) 335

- 1968PI04 J.R. Pizzi, R. Bouche, M. Gaillard, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and C. Ruhla, *Phys. Lett.* B28 (1968) 32
- 1968RA1E Rawlins and Ricco, *Bull. Amer. Phys. Soc.* 13 (1968) 717
- 1968RA34 G.V.S. Rayudu, *J. Inorg. Nucl. Chem.* 30 (1968) 2311
- 1968SE1A Sergeev, Geradin, Wery and Magnac-Valette, *Compt. Rend.* 261 (1965) 291
- 1968SH26 D.W. Sheffey, I.R. Williams and C.B. Fulmer, *Phys. Rev.* 172 (1968) 1094
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, *Nucl. Phys.* A113 (1968) 395
- 1968ST1J Stokes and Young, in Tokyo (1968) 41
- 1968TH1H N. Thurlow, *Nucl. Phys.* A109 (1968) 471
- 1968TU1A Turkiewicz et al., *INR P* 849/1/PL (1968)
- 1968WI1B Wilkinson, *Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn.* 24 (1968) 469
- 1968WO1E Wood and Dunnam, *Bull. Amer. Phys. Soc.* 13 (1968) 1698
- 1968WO1F Wood and Steuer, *Bull. Amer. Phys. Soc.* 13 (1968) 236
- 1968YI01 F. Yiou, *Ann. Phys. (Paris)* 3 (1968) 169
- 1969AN1H Antufyev et al., *Ukr. Fiz. Zh.* 14 (1969) 248
- 1969AN20 Y.P. Antufev, Y.V. Zhebrovskii, L.Y. Kolesnikov, V.S. Kuzmenko, I.I. Miroshnichenko, A.L. Rubashkin and P.V. Sorokin, *Yad. Fiz.* 9 (1969) 680; *Sov. J. Nucl. Phys.* 9 (1969) 394
- 1969AN25 B. Antolkovic, J. Hudomalj, B. Janko, G. Paic and M. Turk, *Nucl. Phys.* A139 (1969) 10
- 1969AR07 D.D. Armstrong, L.L. Catlin, P.W. Keaton, Jr. and L.R. Veaser, *Phys. Rev. Lett.* 23 (1969) 135
- 1969AU1C J. Audouze and H. Reeves, *Astrophys. J.* 158 (1969) 419
- 1969BA1P Bacon et al., *Acta Cryst.* A25 (1969) 391
- 1969BA1U Bahcall and Moeller, *Astrophys. J.* 155 (1969) 511
- 1969BA1Z Barnes, *Nucl. Isospin, Proc. 1969 Asilomar Conf.* (1969) 179
- 1969BA2A J.N. Bahcall, *Phys. Rev. Lett.* 23 (1969) 251
- 1969BA2B Bacher et al., *Bull. Amer. Phys. Soc.* 14 (1969) 851
- 1969BA2C Bahr, Becker, Rausch and Zell, *Bochum Conf. STI/PUB/232 IAEA* (1969) 212
- 1969BE1M H. C. Benohr and K. Wildermuth, *Nucl. Phys.* A128 (1969) 1
- 1969BE90 K. Bethge, C.M. Fou and R.W. Zurmu, *Nucl. Phys.* A123 (1969) 521
- 1969BO1G S. Boffi and F.D. Pacati, *Nucl. Phys.* A129 (1969) 673

1969BU1C Burman and Nordberg, Bull. Amer. Phys. Soc. 14 (1969) 537
1969CA1E Cahill and Martens, Bull. Amer. Phys. Soc. 14 (1969) 553
1969CL06 A.S. Clough, C.J. Batty, B.E. Bonner, C. Tschalar, L.E. Williams and E. Friedman, Nucl. Phys. A137 (1969) 222
1969DE28 V.P. Denisov, A.P. Komar, L.A. Kulchitskii and I.Y. Chubukov, Yad. Fiz. 10 (1969) 1116; Sov. J. Nucl. Phys. 10 (1970) 635
1969DI18 V. di Napoli, Nucl. Instrum. Meth. 69 (1969) 155
1969DO02 V.K. Dolinov, Y.V. Melikov, A.F. Tulinov and O.V. Bormot, Nucl. Phys. A129 (1969) 577
1969DO03 V.K. Dolinov, D.V. Meboniya and A.F. Tulinov, Nucl. Phys. A129 (1969) 597
1969FO06 H.T. Fortune, A. Richter and B. Zeidman, Phys. Lett. B30 (1969) 175
1969FO1C Fortune, Dehnhard, Siemssen and Zeidman, Bull. Amer. Phys. Soc. 14 (1969) 487
1969FO1D Fowler, Contemp. Phys. 1 (1969) 359
1969FU09 N. Fujiwara, J. Phys. Soc. Jpn. 27 (1969) 1380
1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
1969GA18 J. Galin, B. Gatty, M. Lefort, J. Peter, X. Tarrago and R. Basile, Phys. Rev. 182 (1969) 1267
1969GA1P Garvey, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 703
1969GU03 M.R. Gunye, J. Law and R.K. Bhaduri, Nucl. Phys. A132 (1969) 225
1969HA1F Hanna, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 591
1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
1969HA1N Hansteen, Bochum Conf. STI/PUB/232 IAEA (1969) 45
1969HE1M Helmer, Greenwood and Gehrke, Bull. Amer. Phys. Soc. 14 (1969) 1243
1969HE1N E.M. Henley and C.E. Lacy, Phys. Rev. 184 (1969) 1228
1969HI1A Higgins, AD 703259, Naval Postgrad. School, Monterey, CA (1969)
1969HO1H Holman et al., Bull. Amer. Phys. Soc. 14 (1969) 1168
1969HO1K Holmgren, Bochum Conf. STI/PUB/232 IAEA (1969) 17
1969HO1M T.H. Ho and A.B. Volkov, Phys. Lett. B30 (1969) 303
1969HO39 T. Holtebekk and A.E. Monsrud, Phys. Norvegica 3 (1969) 215
1969HU05 R.M. Hutcheon and H.S. Caplan, Nucl. Phys. A127 (1969) 417
1969HU1E Hurst, Seeser and Schupp, Bull. Amer. Phys. Soc. 14 (1969) 190
1969IN1A Inglis, Nucl. Isospin, Proc. 1969 Asilomra Conf. (1969) 789
1969JO1J Johnston and Sargood, Contrib., Montreal (1969) 297

- 1969JO1K Johnston, Switkowski and Sargood, Contrib., Montreal (1969) 272
- 1969JU1A Jungerman et al., Bull. Amer. Phys. Soc. 14 (1969) 488
- 1969KO1J V.M. Kolybasov and N.Ya. Smorodinskaya, Nucl. Phys. A136 (1969) 165
- 1969KO30 T.I. Kopaleishvili and I.Z. Machabeli, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 69; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 65
- 1969KR16 J. Kruger and P. van Leuven, Nucl. Phys. A139 (1969) 418
- 1969KU1C Yu.A. Kudeyarov, I.V. Kurdyumov, V.G. Neudatchin and Yu.F. Smirnov, Nucl. Phys. A126 (1969) 36
- 1969LE08 G.M. Lerner and J.B. Marion, Nucl. Instrum. Meth. 69 (1969) 115
- 1969LE1D Leonardi and Rosa-Clot, Lett. Nuovo Cim. 1 (1969) 829
- 1969LE22 C.C. Lee, J. Korean Phys. Soc. 2 (1969) 1
- 1969LI1F Lindsay, Evans, Toews and Veit, Bull. Amer. Phys. Soc. 14 (1969) 851
- 1969MA13 S. Matsuki, S. Yamashita, K. Fukunaga, D.C. Nguyen, N. Fujiwara and T. Yanabu, J. Phys. Soc. Jpn. 26 (1969) 1344
- 1969MA1G Madsen, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 149
- 1969MA1J Mancuso, Knudson, Wolicki and Bluemel, Bull. Amer. Phys. Soc. 14 (1969) 487
- 1969MA1P Mahalanabis, Proc. Phys. Soc. A2 (1969) 66
- 1969ME1C Meyer, Amer. J. Phys. 37 (1969) 296
- 1969MI10 R.C. Minehart, L. Coulson, W.F. Grubb, III and K. Ziock, Phys. Rev. 177 (1969) 1455
- 1969MI1G R.C. Minehart, L. Coulson, W.F. Grubb, III and K. Ziock, Phys. Rev. 177 (1969) 1464
- 1969MO1E L. Moyer and D.S. Koltun, Phys. Rev. 182 (1969) 999
- 1969MO1J Monson, Rept. Naval Postgrad. School, Monterey (1969)
- 1969MU10 K.M. Murray, Phys. Rev. Lett. 23 (1969) 1461
- 1969MU1C Murray, Bull. Amer. Phys. Soc. 14 (1969) 1213
- 1969NA24 K. Nagatani, M.R. Dwarakanath and D. Ashery, Nucl. Phys. A128 (1969) 325
- 1969NE1A Neumann, Bull. Amer. Phys. Soc. 14 (1969) 1224
- 1969NE1D Neogy, Fortune, Scholz and Zeidman, Bull. Amer. Phys. Soc. 14 (1969) 1226
- 1969NU1A Nusslin, Werner and Zimmerer, Bochum Conf. STI/PUB/232 IAEA (1969) 229
- 1969OR01 H. Orihara, T. Nakagawa, H. Ueno, T. Tohei, T. Yamaya, Y. Nakagome, M. Baba and S. Morita, Nucl. Phys. A139 (1969) 226
- 1969PE22 C. Petitjean, L. Brown and R.G. Seyler, Nucl. Phys. A129 (1969) 209
- 1969PI11 J.R. Pizzi, M. Gaillard, P. Gaillard, A. Guichard, M. Gusakov, G. Reboulet and C. Ruhla, Nucl. Phys. A136 (1969) 496

- 1969PR04 G. Presser, R. Bass and K. Kruger, Nucl. Phys. A131 (1969) 679
- 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
- 1969RO20 L.P. Robertson, R.C. Hanna, K. Ramavataram, D.W. Devins, T.A. Hodges, Z.J. Moroz, S.J. Hoey and D.J. Plummer, Nucl. Phys. A134 (1969) 545
- 1969SC16 P. Schwandt, B.W. Ridley, S. Hayakawa, L. Put and J.J. Kraushaar, Phys. Lett. B30 (1969) 30
- 1969SM1A Smirnov, Bochum Conf. STI/PUB/232 IAEA (1969) 153
- 1969SU15 K. Sugimoto, Phys. Rev. 182 (1969) 1051
- 1969SW1C Sweeney, Bull. Amer. Phys. Soc. 14 (1969) 1007
- 1969TA1G Tang, Bochum Conf. STI/PUB/232 IAEA (1969) 109
- 1969TA1H Tartakovsky and Fursayev, Ukr. Fiz. Zh. 14 (1969) 895
- 1969TI02 G. Tibell, Phys. Lett. B28 (1969) 638
- 1969VA1C S. Varma and P. Goldhammer, Nucl. Phys. A125 (1969) 193
- 1969VA1F Valkovic et al., Bochum Conf. STI/PUB/232 IAEA (1969) 228
- 1969VE1B Velten, Kern. Karlsruhe, Rept. No. 3/69 24 (1969)
- 1969VI02 D. Vinciguerra and T. Stovall, Nucl. Phys. A132 (1969) 410
- 1969VI06 P. Villar, J. Casanova and J. Nalda, An. Fis. 65 (1969) 127
- 1969VI1C P. Vilain, G. Wilquet, J. Sacton, D.M. Harmsen, D.H. Davis, J.H. Wickens and W. Gajewski, Nucl. Phys. B13 (1969) 451
- 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
- 1969WI21 H.W. Wittern, Z. Phys. 229 (1969) 420
- 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
- 1969YI1A Yiou, Seide and Bernas, J. Geophys. Res. 74 (1969) 2447
- 1969YI1B Yiou and Guchan-Beck, J. Phys. (France) 30 (1969) 401
- 1969ZE1A Zeidman and Fortune, Bull. Amer. Phys. Soc. 14 (1969) 507
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1970AN05 Y.P. Antufev, V.I. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko, V.M. Sanin and P.V. Sorokin, Yad. Fiz. 11 (1970) 948; Sov. J. Nucl. Phys. 11 (1970) 527
- 1970AS1E Asami and Moxon, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki, IAEA (1970) 153
- 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
- 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119

1970BA1Q Barashenkov and Abdinov, Acta Phys. Pol. B1 (1970) 65
1970BI1B Bingham, Thesis, Florida State Univ. (1970)
1970BO1U Bonner, Gabbard and Morgan, Bull. Amer. Phys. Soc. 15 (1970) 85
1970BO1V Boynton, Thesis, Princeton Univ. (1970)
1970BR13 R.L. Brodzinski and N.A. Wogman, Phys. Rev. C1 (1970) 1955
1970BR42 R.E. Brown, E.E. Gross and A. van der Woude, Phys. Rev. Lett. 25 (1970) 1346
1970DA21 A. D'Andrea and M. Scalia, Nuovo Cim. A69 (1970) 702
1970DE12 C. Detraz, H.H. Duham and H. Hafner, Nucl. Phys. A147 (1970) 488
1970DE1H Deruytter, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki, IAEA (1970) 127
1970DE1P C. Detraz, Phys. Lett. B32 (1970) 575
1970EL16 M. El-Nadi, O. Zohni and H.M. Hussein, Ann. Phys. (Leipzig) 25 (1970) 1
1970FA14 A. Faessler, E. Huster, O. Krafft and F. Krahn, Z. Phys. 238 (1970) 352
1970FA15 D. Favart, F. Brouillard, L. Grenacs, P. Igo-Kemenes, P. Lipnik and P.C. Macq, Phys. Rev. Lett. 25 (1970) 1348
1970FI07 D. Fick, R. Kankowsky, K. Kilian and E. Salzborn, Phys. Rev. Lett. 24 (1970) 1503
1970FO1D Fortune and Zeidman, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 307
1970FO1E Fort, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki (1970) 253
1970FR1C Frahn, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 297
1970GA07 S. Gangadharan and R.L. Wolke, Phys. Rev. C1 (1970) 1333
1970GA1A Garber et al., BNL 400, 3rd Edition, Vol. 1 (1970)
1970HA1F Hayward, NBS Monograph 118 (1970)
1970HA1M Hardy, Thesis, Rice Univ. (1970)
1970HA1P D.M. Hardy, R.J. Spiger, S.D. Baker, Y.S. Chen and T.A. Tombrello, Phys. Lett. B31 (1970) 355
1970HO1J T.H. Ho and A.B. Volkov, Phys. Lett. B31 (1970) 259
1970JA04 C.G. Jacobs, Jr. and R.E. Brown, Phys. Rev. C1 (1970) 1615
1970JA17 M. Jain, P.G. Roos, H.G. Pugh and H.D. Holmgren, Nucl. Phys. A153 (1970) 49
1970JA1J Janouch and Mach, Nucl. Phys. A158 (1970) 193
1970JA23 B.K. Jain and B. Banerjee, Nuovo Cim. A69 (1970) 419
1970JO21 H.W. Johlige, D.C. Aumann and H.-J. Born, Phys. Rev. C2 (1970) 1616
1970KI1E Kinkaid and Gettys, Bull. Amer. Phys. Soc. 15 (1970) 162

- 1970KO25 V.I. Komarov, G.E. Kosarev and O.V. Savchenko, *Yad. Fiz.* 11 (1970) 711; *Sov. J. Nucl. Phys.* 11 (1970) 399
- 1970KO41 S.-I. Koyama, K. Takahashi and M. Yamada, *Prog. Theor. Phys.* 44 (1970) 663
- 1970KU1D Kull, Thesis, Michigan State Univ. (1970)
- 1970LA14 J.M. Lambert, P.A. Treado, L.A. Beach, R.B. Theus and E.L. Petersen, *Nucl. Phys.* A152 (1970) 516
- 1970LE02 A. Lejeune, *Nucl. Phys.* A141 (1970) 123
- 1970LI06 W.-K. Lin, F. Scheibling and R.W. Kavanagh, *Phys. Rev. C* 1 (1970) 816
- 1970LO1E M.A. Lodhi, *Phys. Rev. C* 1 (1970) 1732
- 1970MA1E Martens and Schweimer, *Z. Phys.* 233 (1970) 170
- 1970MC07 W.S. McEver, T.B. Clegg, J.M. Joyce, E.J. Ludwig and R.L. Walter, *Phys. Lett.* B31 (1970) 560
- 1970MC1A McPherson et al., *Bull. Amer. Phys. Soc.* 15 (1970) 162
- 1970ME1F Meadows and Whalen, *Nucl. Sci. Eng.* 40 (1970) 12
- 1970MO38 L.P. Moskaleva, G.A. Fedoseev and A.N. Khalemskii, *Yad. Fiz.* 12 (1970) 871; *Sov. J. Nucl. Phys.* 12 (1971) 472
- 1970NE03 D.O. Nellis, W.E. Tucker and I.L. Morgan, *Phys. Rev. C* 1 (1970) 847
- 1970NE1F Neudatchin and Wildermuth, *Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969* (1970) 327
- 1970PI1D Pizzi, Thesis, Univ. of Lyon (1970)
- 1970PO03 D.L. Powell, G.M. Crawley, B.V.N. Rao and B.A. Robson, *Nucl. Phys.* A147 (1970) 65
- 1970RA1H Ramachandran, Prabhakar and Rej, *Nucl. Phys.* B20 (1970) 369
- 1970RA33 K. Ramavataram and Q. Ho-Kim, *Nucl. Phys.* A156 (1970) 395
- 1970RO07 M.L. Roush, L.A. West and J.B. Marion, *Nucl. Phys.* A147 (1970) 235
- 1970SA14 M. Sanzone, G. Ricco, S. Costa and L. Ferrero, *Nucl. Phys.* A153 (1970) 401
- 1970SC23 G. Scheklinski, U. Strobusch and B. Goel, *Nucl. Phys.* A153 (1970) 97
- 1970SE1A Seeser, Hurst and Schupp. *Bull. Amer. Phys. Soc.* 15 (1970) 23
- 1970SE1D Seeser, Thesis, Univ. of Missouri (1970)
- 1970SO1A Sowerby, Patrick, Uttley and Diment, *J. Nucl. Energy* 24 (1970) 323
- 1970TH1F Theus et al., *Bull. Amer. Phys. Soc.* 15 (1970) 1695
- 1970TO09 S.G. Tonapetyan, O.G. Konovalov, A.I. Derebchinskii, A.A. Zybalov, V.M. Khvorostyan, N.V. Goncharov and V.A. Goldshtein, *Pisma Zh. Eksp. Teor. Fiz.* 11 (1970) 165; *JETP Lett.* 11 (1970) 101

- 1970VI01 J.S. Vincent and E.T. Boschitz, Nucl. Phys. A143 (1970) 121
- 1970WA1N Walcher, Private Communication (1970)
- 1970WO10 C.F. Wong, R.M. Hutcheon, Y.M. Shin and H.S. Caplan, Can. J. Phys. 48 (1970) 1917
- 1970ZO1A Zofka, Czech. J. Phys. 20 (1970) 926
- 1971AN04 Y.P. Antufev, V.L. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, Yad. Fiz. 13 (1971) 473; Sov. J. Nucl. Phys. 13 (1971) 265
- 1971AN15 Y.P. Antufev, V.L. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, Yad. Fiz. 14 (1971) 898; Sov. J. Nucl. Phys. 14 (1972) 502
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AR1K Armstrong, Keaton and Vesser, Polarization, Madison, 1970 (1971) 677, 680
- 1971BA2Y Baus-Baghdikian, Univ. libre Bruxelles, Bull. No. 45 (1971)
- 1971BA58 M. Barbier and S. Regnier, J. Inorg. Nucl. Chem. (GB) 33 (1971) 2720
- 1971BI11 H.G. Bingham, A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 265
- 1971BI12 H.G. Bingham, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A175 (1971) 374
- 1971BI22 R. Bimbot and H. Gauvin, Compt. Rend. B273 (1971) 1054
- 1971BR07 K.H. Bray, M. Jain, K.S. Jayaraman, G. LoBianco, W.T.H. Van Oers and Y.I. Wu, Nucl. Phys. A163 (1971) 649
- 1971BR12 K.H. Bray, S.N. Bunker, M. Jain, K.S. Jayaraman, G.A. Moss, W.T.H. van Oers, D.O. Wells and Y.I. Wu, Phys. Rev. C3 (1971) 1771
- 1971BR36 R.L. Brodzinski, L.A. Rancitelli, J.A. Cooper and N.A. Wogman, Phys. Rev. C4 (1971) 1250
- 1971BU1D Burke, Lunnon and Lefevre, Bull. Amer. Phys. Soc. 16 (1971) 829
- 1971CA01 P.J. Castleberry, L. Coulson, R.C. Minehart and K.O.H. Ziock, Phys. Lett. B34 (1971) 57
- 1971CA1B Cameron and Fowler, Astrophys. J. 164 (1971) 111
- 1971CA1J Castleberry, Thesis, Univ. of Virginia (1971)
- 1971CH1B Chuev et al., J. Phys. (Paris) 32 (1971) C6-167
- 1971CH1H D. Chlebowska, Phys. Lett. B35 (1971) 393
- 1971CH42 L.S. Chuang, Nucl. Phys. A174 (1971) 399
- 1971CO28 B.S. Cooper, J.B. Seaborn and S.A. Williams, Phys. Rev. C4 (1971) 1997
- 1971CU1B Cuzzocrea, Perillo and Notarrigo, Nuovo Cim. A4 (1971) 251

- 1971DA24 J.C. Davis, K.A. Weaver, D. Hilscher, H.H. Barschall and A.B. Smith, Phys. Rev. C4 (1971) 1061
- 1971DE2D Devons, Proc. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 619
- 1971DE37 C. Detraz, C.D. Zafiratos, C.E. Moss and C.S. Zaidins, Nucl. Phys. A177 (1971) 258
- 1971DM01 P.P. Dmitriev, N.N. Krasnov, G.A. Molin and M.V. Panarin, Atomn. Energ. (USSR) 31 (1971) 157; Sov. At. Energy 31 (1972) 876
- 1971DRZW T. Drake, Proc. Mont Tremblant Summer School Meeting on Dynamic Struct. of Nucl. States, 1971, Toronto, Canada (1972) 420
- 1971EP02 M. Epherre and C. Seide, Phys. Rev. C3 (1971) 2167; Erratum Phys. Rev. C4 (1971) 1494
- 1971FA09 J. Favier, T. Bressani, G. Charpak, L. Massonnet, W.E. Meyerhof and C. Zupancic, Nucl. Phys. A169 (1971) 540
- 1971FE02 W. Fetscher, K. Sattler, N.C. Schmeing, E. Seibt, C. Weddigen and E.J. Kanellopoulos, Phys. Lett. B34 (1971) 171
- 1971FE03 W. Fetscher, E. Seibt, C. Weddigen and E.J. Kanellopoulos, Phys. Lett. B35 (1971) 31
- 1971FO1P D.G. Foster and D.W. Glasgow, Phys. Rev. C3 (1971) 604
- 1971FO24 D.G. Foster and D.W. Glasgow, Phys. Rev. C3 (1971) 576
- 1971FU09 R.C. Fuller, Phys. Rev. C4 (1971) 1968
- 1971GA1J Gaillard, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971GO22 N.V. Goncharov, A.I. Derebchinskii, O.P. Konovalov, S.G. Tonapetyan and V.M. Khvorostyan, Yad. Fiz. 14 (1971) 31; Sov. J. Nucl. Phys. 14 (1972) 18
- 1971GR16 R.C. Grimm, I.E. McCarthy and R.G. Storer, Nucl. Phys. A166 (1971) 330
- 1971GU07 H.H. Gutbrod, H. Yoshida and R. Bock, Nucl. Phys. A165 (1971) 240
- 1971HA21 M.L. Halbert and A. van der Woude, Phys. Rev. Lett. 26 (1971) 1124
- 1971HE20 R.G. Helmer, R.C. Greenwood and R.J. Gehrke, Nucl. Instrum. Meth. 96 (1971) 173
- 1971HE24 H.H. Heckman, D.E. Greiner, P.J. Lindstrom and F.S. Bieser, Science 174 (1971) 1130
- 1971HO1K Holman, Thesis, Univ. of California, Los Angeles (1971)
- 1971HO25 A.B. Holman, J.-L. Perrenoud, J.C. Young, M.B. Epstein and B.T. Wright, Nucl. Phys. A174 (1971) 161
- 1971JU05 J.A. Jungerman, F.P. Brady, W.J. Knox, T. Montgomery, M.R. McGie, J.L. Romero and Y. Ishizaki, Nucl. Instrum. Meth. 94 (1971) 421

- 1971KA70 N.N. Kaushal, E.J. Winhold, R.H. Augustson, P.F. Yergin and H.A. Medicus, Nucl. Energy 25 (1971) 91
- 1971KO02 T.I. Kopaleishvili and I.Z. Machabeli, Nucl. Phys. A160 (1971) 204
- 1971KO24 K.F. Koral, E.A. Silverstein and P.R. Bevington, Nucl. Phys. A175 (1971) 156
- 1971KU22 L.M. Kuznetsova, V.I. Kukulin and V.G. Neudachin, Yad. Fiz. 13 (1971) 694; Sov. J. Nucl. Phys. 13 (1971) 394
- 1971LA03 H. Lancman and J. Lebowitz, Phys. Rev. C3 (1971) 465
- 1971MA13 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A165 (1971) 145
- 1971MA1Y Macklin, Hill and Allen, Nucl. Instrum. Meth. 96 (1971) 509
- 1971MA44 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A172 (1971) 166
- 1971NO02 J.L. Norton and P. Goldhammer, Nucl. Phys. A165 (1971) 33
- 1971NO09 V.I. Noga, Y.N. Ranyuk, P.V. Sorokin and V.A. Tkachenko, Ukr. Fiz. Zh. 16 (1971) 1850
- 1971OR02 J. Orloff and W.W. Daehnick, Phys. Rev. C3 (1971) 430
- 1971OV1A Overley, Sealock and Ehlers, Bull. Amer. Phys. Soc. 16 (1971) 16
- 1971PI06 C.J. Piluso, R.H. Spear, K.W. Carter, D.C. Kean and F.C. Barker, Aust. J. Phys. 24 (1971) 459
- 1971PL06 G.R. Plattner and A.D. Bacher, Phys. Lett. B36 (1971) 211
- 1971PL1C Plattner, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haerberli (1971) 107
- 1971PO1A Popic, Stepanic and Stanojevic, Fizika 4 (1971) 40
- 1971PR09 R. Prasad and D.C. Sarkar, Nuovo Cim. A3 (1971) 467
- 1971RE07 B. Remaud, Compt. Rend. B272 (1971) 390
- 1971RO1A Robson and Richter, Ann. Phys. 63 (1971) 261
- 1971SA27 A. Saganek, I. Sledzinska, A. Turos, Z. Wilhelmi and B. Zwiaglinski, Acta Phys. Pol. B2 (1971) 473
- 1971SC16 G. Schmidt, J. Mosner and J. Schintlmeister, Nucl. Phys. A173 (1971) 449
- 1971SH26 V.S. Shirley, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1255
- 1971SI28 M. Simonius, Phys. Lett. B37 (1971) 446
- 1971SP05 H. Spinka, T. Tombrello and H. Winkler, Nucl. Phys. A164 (1971) 1; Erratum Nucl. Phys. A196 (1972) 634
- 1971ST30 G.B. Stapleton and R.H. Thomas, Nucl. Phys. A175 (1971) 124; Erratum Nucl. Phys. A196 (1972) 635

- 1971TA23 Y.C. Tang and R.E. Brown, Phys. Rev. C4 (1971) 1979
- 1971TH1K Theobald, Migneco and Cervini, Nucl. Instrum. Meth. 95 (1971) 1
- 1971VA20 G.J.C. van Niftrik, L. Lapikas, H. de Vries and G. Box, Nucl. Phys. A174 (1971) 173
- 1971VO07 W. von Witsch, M. Ivanovich, V.A. Otte, D. Rendic and G.C. Phillips, Nucl. Phys. A172 (1971) 633
- 1971WA1J Walker and Weaver, Nucl. Instrum. Meth. 96 (1971) 93
- 1971WE1L Werby, Thesis, Florida State Univ. (1971)
- 1971WE1M Werby and Edwards, Bull. Amer. Phys. Soc. 16 (1971) 1186
- 1971YO06 J.C. Young, A.B. Holman, I. Slaus and T.A. Cahill, Phys. Lett. B37 (1971) 377
- 1971ZA1D D.A. Zaikin, Nucl. Phys. A170 (1971) 584
- 1972AB14 O.B. Abidinov and V.S. Barashenkov, Acta Phys. Pol. B3 (1972) 385
- 1972AM04 B.S. Amin, S. Biswas, D. Lal and B.L.K. Somayajulu, Nucl. Phys. A195 (1972) 311
- 1972AN05 R.K. Anderson, M.R. Wilson and P. Goldhammer, Phys. Rev. C6 (1972) 136
- 1972AN1Q Antolkovic, Few Particle Problems, UCLA, 1972 (1972) 695
- 1972AV04 G.V. Avakov, E.I. Dolinsky and V.V. Turovtsev, Nucl. Phys. A196 (1972) 529
- 1972AZ01 S.A. Azimov, U.R. Arifkhanov, M. Gulyamov, B.I. Islamov, T. Iskhakov, U.I. Faizul-
laev and E. Ergashov, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 173; Bull. Acad. Sci.
USSR Phys. Ser. 36 (1973) 162
- 1972BA2M Bahcall, Comments on Nucl. Part. Phys. 5 (1972) 59
- 1972BA77 A. Bamberger, G. Jansen, B. Povh, D. Schwalm and U. Smilansky, Nucl. Phys. A194
(1972) 193
- 1972BB26 F.C. Barker, Aust. J. Phys. 25 (1972) 341
- 1972BE1Y Beveridge et al., Helv. Phys. Acta 45 (1972) 943
- 1972BE1Z Becker and Beccaria, Communications, Proc. Aix-en-Provence Conf. 2 (1972) 45
- 1972BI1G Birchall, Frois, Roy and Slobodrian, Few Particle Problems, UCLA, 1972 (1972) 61
- 1972BO02 G.J. Bock, E.A. Samworth, J.W. Olness and E.K. Warburton, Phys. Rev. C5 (1972)
284
- 1972BO07 H. Bohlen, N. Marquardt, W. von Oertzen and P. Gorodetzky, Nucl. Phys. A179
(1972) 504
- 1972BO42 W.R. Boykin, S.D. Baker and D.M. Hardy, Nucl. Phys. A195 (1972) 241
- 1972BR1Q Brown, Tang and Thompson, Few Particle Problems, UCLA, 1972 (1972) 703
- 1972BR20 K.H. Bray, M. Jain, K.S. Jayaraman, G. Lobianco, G.A. Moss, W.T.H. van Oers and
D.O. Wells, Nucl. Phys. A189 (1972) 35
- 1972BU01 F.A. Bumiller, F.R. Buskirk, J.N. Dyer and W.A. Monson, Phys. Rev. C5 (1972) 391

- 1972BU16 S.N. Bunker, M. Jain, C.A. Miller, J.M. Nelson, P.J. Tivin and W.T.H. van Oers, *Can. J. Phys.* 50 (1972) 1295
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, *Nuovo Cim.* A12 (1972) 71
- 1972CE1A Cerny, *At. Masses Fund. Constants*, Teddington, 1971 (1972) 26
- 1972CL1A Clayton, *Encyclopedia of the Twentieth Century* (1972)
- 1972CL1C Clement et al., *Few Particle Problems*, UCLA, 1972 (1972) 953
- 1972DA21 A.V. Davydov and I.M. Narodetskii, *Pisma Zh. Eksp. Teor. Fiz.* 15 (1972) 741; *JETP Lett.* 15 (1972) 525
- 1972ED01 G. Eder, G. Winkler and P. Hille, *Z. Phys.* 253 (1972) 335
- 1972EL19 S.A. Elbakt, I.J. Van Heerden, W.J. McDonald and G.C. Neilson, *Nucl. Instrum. Meth.* 105 (1972) 519
- 1972EM03 G.T. Emery, *Ann. Rev. Nucl. Sci.* 22 (1972) 165
- 1972FR09 V. Franco, *Phys. Rev.* C6 (1972) 748
- 1972FR1J Frisbee, Pugh and Holmgren, *Bull. Amer. Phys. Soc.* 17 (1972) 586
- 1972FR1K Frisbee, Thesis, Univ. of Maryland (1972)
- 1972FU07 M. Furic, R.K. Cole, H.H. Forster, C.C. Kim, D.Y. Park, J. Rucker, H. Spitzer and C.N. Waddell, *Phys. Lett.* B39 (1972) 629
- 1972GA1E Garrett, *Symp. on Two-Nucleon Transfer and Pairing Excitations*, Argonne (1972) 232; CONF-720309
- 1972GA1L Gal, Soper and Dalitz, *Ann. Phys.* 72 (1972) 445
- 1972GO1P Goodman, Hensley, van der Woude and Raman, *Bull. Amer. Phys. Soc.* 17 (1972) 929
- 1972HA04 S.L. Hausladen, R.O. Lane and J.E. Monahan, *Phys. Rev.* C5 (1972) 277
- 1972HA06 O. Hausser, A.B. McDonald, T.K. Alexander, A.J. Ferguson and R.E. Warner, *Phys. Lett.* B38 (1972) 75
- 1972HA64 D.M. Hardy, R.J. Spiger, S.D. Baker, Y.S. Chen and T.A. Tombrello, *Nucl. Phys.* A195 (1972) 250
- 1972HI16 J. Hiura and R. Tamagaki, *Suppl. Prog. Theor. Phys.* 52 (1972) 25
- 1972HU09 R. Hub, D. Clement and K. Wildermuth, *Z. Phys.* 252 (1972) 324
- 1972HU1A W.T. Huang, C.A. Levinson and M.K. Banerjee, *Phys. Rev.* C5 (1972) 651
- 1972JA07 O.N. Jarvis, C. Whitehead and M. Shah, *Nucl. Phys.* A184 (1972) 615
- 1972JA23 A.K. Jain and N. Sarma, *Nucl. Phys.* A195 (1972) 566
- 1972KA1B Kavanagh, *Cosmology, Fusion and Other Matters*; Eds., F. Reines and G. Gamow, *Memorial Vol.* (1972) 169
- 1972KO1A Kocharov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 36 (1972) 2052

1972KO1E Koike, Prog. Theor. Phys. 48 (1972) 66
 1972KU12 K.-I. Kubo and M. Hirata, Nucl. Phys. A187 (1972) 186
 1972LA1F Lane, COO-1717-3 (1972)
 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353
 1972ME17 J.W. Meadows and J.F. Whalen, Nucl. Sci. Eng. 48 (1972) 221
 1972NE17 V.G. Neudatchin, V.I. Kukulin, A.N. Boyarkina and V.P. Korennoy, Lett. Nuovo Cim. 5 (1972) 834
 1972OH01 I.K. Oh, C.S. Zaidins, C.D. Zafiratos and S.I. Hayakawa, Nucl. Phys. A178 (1972) 497
 1972OP01 A.M.F. Op den Kamp and A.M.J. Spits, Nucl. Phys. A180 (1972) 569
 1972PA1C Parker, Astrophys. J. 175 (1972) 261
 1972PE05 B.I. Persson and S.E. Koonin, Phys. Rev. C5 (1972) 1443
 1972PI1A Pisano, Cobern and Parker, Bull. Amer. Phys. Soc. 17 (1972) 914
 1972PN1A Pniewski, Few Particle Problems, UCLA, 1972 (1972) 145
 1972PU1C Pugh, Frisbee and Holmgren, Communications, Proc. Aix-en-Provence Conf. 2 (1972) 152
 1972RA1E Radvanyi, J. Phys. (Paris) 33 (1972) C5-141
 1972RA30 G.M. Raisbeck, J. Lestringuez and F. Yiou, Phys. Rev. C6 (1972) 685
 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430.
 1972SE1K Sealock and Overley, Bull. Amer. Phys. Soc. 17 (1972) 900
 1972SM02 U. Smilansky, B. Povh and K. Traxel, Phys. Lett. B38 (1972) 293
 1972SW1A Swannack, Thesis, Carnegie Mellon Univ. (1972)
 1972SZ02 J. Szabo, J. Csikai and M. Varnagy, Nucl. Phys. A195 (1972) 527
 1972THZF H. Theissen, Springer Tracts in Mod. Phys.; Ed., Hohler, Vol. 65 (1972) 1
 1972UL1A Ulrich and Scalo, OAP-303 (1972)
 1972VO06 A.A. Vorobyov, D.M. Seliverstov, V.T. Grachev, I.A. Kondurov, A.M. Nikitin, N.N. Smirnov and Y.K. Zalite, Phys. Lett. B40 (1972) 102
 1972ZV1A Zverev, Simakhin and Dutov, Atomn. Energ. (USSR) 32 (1972) 39
 1973AH1A Ahrens et al., in Asilomar (1973) 23
 1973AR1L Arad and Ben-David, Rev. Mod. Phys. 45 (1973) 230
 1973AS02 A. Aswad, H.R. Kissener, H.U. Jager and R.A. Eramzhian, Nucl. Phys. A208 (1973) 61

1973AU1H Audouze and Truran, OAP-310 (1973)
 1973BA1Y Bamberger et al., Nucl. Phys. B60 (1973) 1
 1973BA2C Bahcall, Nucl. Instrum. Meth. 110 (1973) 381
 1973BA2G Baer and Crowe, in Asilomar (1973) 583
 1973BA2R Backenstoss et al., Contrib., Uppsala (1973) 146
 1973BA2V Backenstoss et al., in Munich, 1 (1973) 317
 1973BO1Y Bollen et al., Bull. Amer. Phys. Soc. 18 (1973) 1600
 1973BR1M Bramblett et al., Asilomar (1973) Paper 2B15S
 1973CL09 C.F. Clement, Nucl. Phys. A213 (1973) 469
 1973CO1N Costa, in Asilomar (1973) 1319
 1973CO1P Cohen, J. Phys. Soc. Jpn. Suppl. 34 (1973) 63
 1973CO2B Cole et al., Bull. Amer. Phys. Soc. 18 (1973) 1600
 1973DO13 P. Dougan and W. Stiefler, Z. Phys. 265 (1973) 1
 1973DO1F C.B. Dover and R.H. Lemmer, Phys. Rev. C7 (1973) 2312
 1973ER1G Erb et al., in Munich, 1 (1973) 547
 1973FE11 W. Fetscher, E. Seibt and C. Weddigen, Nucl. Phys. A216 (1973) 47
 1973FE1J Federsel, Schraner, Teufel and Wildermuth, in Munich, 1 (1973) 499
 1973FI04 S. Fiarman and W.E. Meyerhof, Nucl. Phys. A206 (1973) 1
 1973FR17 B. Frois, J. Birchall, C.R. Lamontagne, U. von Moellendorff, R. Roy and R.J. Slobodrian, Phys. Rev. C8 (1973) 2132
 1973GA16 S.N. Gardiner, J.L. Matthews and R.O. Owens, Phys. Lett. B46 (1973) 186
 1973GO2B Goulding, Stoler, Clement and Seagrave, Bull. Amer. Phys. Soc. 18 (1973) 538
 1973HA1V Hausser, J. Phys. Soc. Jpn. Suppl. 34 (1973) 135
 1973HA47 O. Hausser, A.B. McDonald, T.K. Alexander, A.J. Ferguson and R.E. Warner, Nucl. Phys. A212 (1973) 613
 1973HA49 P.S. Hauge and S. Maripuu, Phys. Rev. C8 (1973) 1609
 1973HE1M Hensley, Huizenga and Bassett, EOS Trans. Am. Geophys. Union 54 (1973) 505
 1973HI03 S. Hiramatsu, T. Kamae, H. Muramatsu, K. Nakamura, N. Izutsu and Y. Watase, Phys. Lett. B44 (1973) 50
 1973HO11 R. Holub, A.F. Zeller, G.R. Choppin, R.J. De Meijer and H.S. Plendl, Phys. Lett. B43 (1973) 375
 1973JO07 G.G. Jonsson and K. Lindgren, Phys. Scr. 7 (1973) 49

- 1973JU2A M. Juric, G. Bohm, J. Klabuhn, U. Kreckler, F. Wysotzki, G. Coremans-Bertrand, J. Sacton, G. Wilquet, T. Cantwell, F. Esmael et al., Nucl. Phys. B52 (1973) 1
- 1973KA04 V.V. Karapetyan, V.N. Mileev and N.N. Titarenko, Nucl. Phys. A203 (1973) 561
- 1973KL1B Klages and Duhm, in Munich, 1 (1973) 478
- 1973KO1D Kovar, Symp. on Heavy Ion Transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 59
- 1973KO1M Komarov et al., Contrib., Uppsala (1973) 179
- 1973KO1Q Koepke, Brown, Tang and Thompson, COO-1265-139
- 1973KO1R Koepke and Brown, Bull. Amer. Phys. Soc. 18 (1973) 1382
- 1973KU03 D. Kurath, Phys. Rev. C7 (1973) 1390
- 1973KU12 K.-I. Kubo, H.H. Duhm and N. Ueta, Phys. Lett. B45 (1973) 299
- 1973LA19 H. Laumer, S.M. Austin, L.M. Panggabean and C.N. Davids, Phys. Rev. C8 (1973) 483
- 1973LA1Q Lambert et al., Bull. Amer. Phys. Soc. 18 (1973) 1382
- 1973LA26 W. Laskar and B. Remaud, J. Phys. (France) 34 (1973) 783
- 1973LI02 R.H. Lindsay, W. Toews and J.J. Veit, Nucl. Phys. A199 (1973) 513
- 1973MA1K Maripuu, 5th Symp. Struct. Low-Medium Mass Nuclei (1973) 63
- 1973MA1V Mak, Jensen and Barnes, LAP-112 (1973)
- 1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
- 1973MI02 K. Miyano, J. Phys. Soc. Jpn. 34 (1973) 853
- 1973MU11 N.C. Mukhopadhyay, Phys. Lett. B45 (1973) 309
- 1973MU12 T. Mukoyama, Y. Isozumi, T. Kitahara and S. Shimizu, Phys. Rev. C8 (1973) 1308
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973MU19 M. Mutterer, Phys. Rev. C8 (1973) 2089
- 1973NA1M Nadejdin, Petrov and Satarov, Contrib., Uppsala (1973) 181
- 1973NA1N Nagatani, in Munich, 1 (1973) 561
- 1973NY04 E.M. Nyman, Nucl. Phys. A215 (1973) 397
- 1973OS1C Ost et al., in Munich, 1 (1973) 402
- 1973PE1E Pedroni et al., Contrib., Uppsala (1973) 75
- 1973PF02 K.O. Pfeiffer, E. Speth and K. Bethge, Nucl. Phys. A206 (1973) 545
- 1973PI1D Pisano, Bull. Amer. Phys. Soc. 18 (1973) 1384
- 1973PO10 W.P. Poenitz and A. Devolpi, Int. J. Appl. Radiat. Isotop. 24 (1973) 471
- 1973RA37 G.M. Raisbeck, J. Lestringuez and F. Yiou, Nature 244 (1973) 28

- 1973RE1G Reeves, Audouze, Fowler and Schramm, *Astrophys. J.* 179 (1973) 909
- 1973RO1R Robson, *Nucl. Phys.* A204 (1973) 523
- 1973SA1J Sackmann, Smith and Despain, OAP-324 (1973)
- 1973SC1T J.M. Scalo and R.K. Ulrich, *Astrophys. J.* 183 (1973) 151
- 1973SC1V Schenk et al., in Munich, 1 (1973) 450
- 1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, *Nucl. Phys.* A212 (1973) 573
- 1973SO08 M.A.A. Sonnemans, J.C. Waal and R. Van Dantzig, *Phys. Rev. Lett.* 31 (1973) 1359
- 1973SQ01 G.T.A. Squier, M.E. Cage, G.J. Pyle, A.S. Clough, G.K. Turner, B.W. Allardyce, C.J. Batty, D.J. Baugh, W.J. McDonald, R.A.J. Riddle et al., *Phys. Rev. Lett.* 31 (1973) 389
- 1973SU1C Sugimoto and Tanihata, *J. Phys. Soc, Jpn. Suppl.* 34 (1973) 245
- 1973TR1E Trimble and Reines, *Rev. Mod. Phys.* 45 (1973) 1
- 1973VO1G Vorobyov et al., in Munich, 1 (1973) 716
- 1973WE07 M.F. Werby, M.B. Greenfield, K.W. Kemper, D.L. McShan and S. Edwards, *Phys. Rev.* C8 (1973) 106
- 1973WI11 D.H. Wilkinson, *Nucl. Phys.* A209 (1973) 470
- 1973WO06 G.J. Wozniak, N.A. Jelley and J. Cerny, *Phys. Rev. Lett.* 31 (1973) 607
- 1974AU1A Audouze and Tinsley, *Astrophys. J.* 192 (1974) 487
- 1974BA1K Bartle, *Bull. Amer. Phys. Soc.* 19 (1974) 111
- 1974GO1V Gould, Nelson, Williams and Boyce, *Bull. Amer. Phys. Soc.* 19 (1974) 15
- 1975AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A248 (1975) 1
- 1976AJ04 F. Ajzenberg-Selove, *Nucl. Phys.* A268 (1976) 1

