

Energy Levels of Light Nuclei $A = 9$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A227* (1974), p. 1. This version of $A = 9$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 31, 1973)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 9$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [⁹He](#), [⁹Li](#), [⁹Be](#), [⁹B](#), [⁹C](#)

B. Tables of Recommended Level Energies:

[Table 9.1](#): Energy levels of ⁹Li

[Table 9.2](#): Energy levels of ⁹Be

[Table 9.9](#): Energy levels of ⁹B

[Table 9.12](#): Energy levels of ⁹C

C. [References](#)

D. Figures: [⁹Li](#), [⁹Be](#), [⁹B](#), [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

⁹He
(Not illustrated)

⁹He is predicted to be particle unstable: its calculated mass excess > 40.17 MeV (1970WA1G, 1972WA07), = 43.54 MeV (1972TH13). Particle instability with respect to ⁸He + n, ⁷He + 2n and ⁶He + 3n implies atomic mass excesses greater than 39.7, 42.25 and 41.812 MeV, respectively. See also (1968CE01). ⁹He has not been observed in a pion experiment [⁹Be(π^- , π^+)⁹He] (1965GI10) nor in the spontaneous fission of ²⁵²Cf (1967CO36).

⁹Li
(Figs. 15 and 18)

GENERAL:

Model calculations: (1966BA26).

Special reactions: (1965DO13, 1966GA15, 1966KL1C, 1967AU1B, 1967CA1J, 1967HA10, 1968DO20, 1972VO06, 1973KO1D, 1973MU12, 1973WI15).

Other topics: (1972CA37, 1972PN1A, 1973JU2A).

Ground state properties: (1966BA26, 1969JA1M).

Mass of ⁹Li: From the Q -value of ¹⁸O(⁷Li, ¹⁶O)⁹Li, the atomic mass excess of ⁹Li is 24.9654 ± 0.005 MeV (1969NE1E; prelim. results). (1971WA37) adopt 24.966 ± 0.005 MeV. We use the latter value.

1. ⁹Li(β^-)⁹Be $Q_m = 13.618$

The half-life of ⁹Li is 172 ± 3 msec [see (1966LA04)], 176 ± 1 msec (1965DO13), 177 ± 3 msec (1970CH07, 1970CH1T). We adopt $\tau_{1/2} = 176 \pm 2$ msec. See also (1968BO32). ⁹Li decays to ⁹Be*(0, 2.43, 2.78): see ⁹Be and Table 9.7 (1970CH07, 1970CH1T). See also (1963AL18, 1969MA11). Log ft values are listed in Table 9.7: the allowed nature of the transitions to ⁹Be*(0, 2.43, 2.78) with $J^\pi = \frac{3}{2}^-, \frac{5}{2}^-$ and ($\frac{1}{2}^-$) is evidence for $J^\pi = \frac{3}{2}$ for ⁹Li(0) (1970CH07).

E_{β^-} (max) have been measured by (1963AL18, 1963NE07, 1969KL08). Delayed neutrons are observed due to the decay of the neutron unbound states ⁹Be*(2.43, 2.78): see ⁹Be. See also (1966BA1A, 1966BA26, 1970DA21, 1971LI1H, 1971WI18, 1972WI28, 1972WI1C, 1973HA49, 1973TO14, 1973WI11; theor.).

2. ⁷Li(t, p)⁹Li $Q_m = -2.397$
 $Q_0 = -2.397 \pm 0.020$ (1964MI04).

Table 9.1: Energy levels of ${}^9\text{Li}$

| E_x (MeV \pm keV) | $J^\pi; T$ | $\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|--------------------------------|--|------------|------------------|
| g.s. | $(\frac{3}{2})^-; \frac{3}{2}$ | $\tau_{1/2} = 176 \pm 2$ msec | β^- | 1, 2, 3, 4, 8, 9 |
| 2.691 ± 5 | $(\frac{1}{2}^-)$ | | (γ) | 2 |
| 4.31 ± 30 | | $\Gamma = 250 \pm 30$ | | 2 |
| 5.38 ± 60 | | 600 ± 100 | | 2 |
| 6.41 ± 20 | | < 100 | | 2 |

Proton groups are observed to excited states at $E_x = 2.691 \pm 0.005$ MeV (1964MI04), 4.31 ± 0.03 , 5.38 ± 0.06 and 6.41 ± 0.02 MeV (1971YO04) [$\Gamma_{\text{cm}} = 250 \pm 30, 600 \pm 100, < 100$ keV, respectively]. Angular distributions are reported at $E_t = 11.3$ MeV (1964MI04; t_0) and 15 MeV (1971YO04; t_0, t_2, t_4). The angular distributions to ${}^9\text{Li}(0)$ are consistent with $J^\pi = \frac{3}{2}^-$ and the relative magnitude of the cross section to ${}^9\text{Li}^*(2.69)$ is consistent with $J^\pi = \frac{1}{2}^-$: see (1964MI04, 1971YO04). See also (1968HO1F; theor.) and (1969MA11, 1970CH07).

3. ${}^9\text{Be}(n, p){}^9\text{Li}$ $Q_m = -12.836$

See (1967ME11).

4. ${}^9\text{Be}(d, 2p){}^9\text{Li}$ $Q_m = -15.060$

See (1951GA30).

5. ${}^9\text{Be}(t, {}^3\text{He}){}^9\text{Li}$ $Q_m = -13.599$

Not reported.

6. (a) ${}^{11}\text{B}(\gamma, 2p){}^9\text{Li}$ $Q_m = -30.876$

(b) ${}^{11}\text{B}(p, 3p){}^9\text{Li}$ $Q_m = -30.876$

See (1966LA04).

7. $^{11}\text{B}(\text{n}, ^3\text{He})^9\text{Li}$ $Q_{\text{m}} = -23.158$

Not reported.

8. $^{12}\text{C}(\gamma, 3\text{p})^9\text{Li}$ $Q_{\text{m}} = -46.834$

See (1966LA04).

9. $^{18}\text{O}(^7\text{Li}, ^{16}\text{O})^9\text{Li}$ $Q_{\text{m}} = -6.103$

See “*Mass of ^9Li* ” in the GENERAL section here (1969NE1E). See also (1970CH07).

⁹Be
(Figs. 16 and 18)

GENERAL: (See also (1966LA04).)

Shell model: (1961KO1A, 1965CO25, 1965GR18, 1965VO1A, 1966AD06, 1966BA26, 1966HA18, 1966MA1P, 1966WI1E, 1967CO32, 1967ST1C, 1968GO01, 1969BO1V, 1969BO19, 1969BO33, 1969GU03, 1969VA1C, 1970CO1H, 1971CO28, 1971GR02, 1971NO02, 1972LE1L, 1973HA49, 1973KU03).

Alpha and cluster models: (1965NE1B, 1966HI1A, 1967TA1C, 1968KU1B, 1969BA1J, 1969NE1C, 1970BA1Q, 1971LE1N, 1971NO02, 1972AB19, 1972CH1N, 1972HI16, 1972IK1A, 1972LE1L, 1973KU03, 1973OK1B).

Collective and deformed models: (1965VO1A, 1966EL08, 1967BO1K, 1967BO34, 1973KU13, 1973SL02).

Special levels: (1966AD06, 1966BA26, 1966EL08, 1967CO32, 1967ST1C, 1968BO19, 1968GO01, 1969BO1V, 1969BO33, 1969GU03, 1969HA1G, 1970PE18, 1970TO1E, 1971CO28, 1971GR02, 1971LI30, 1971NO02, 1972BE1E, 1972CH1N).

Electromagnetic transitions: (1965CO25, 1966BA26, 1966EL08, 1967KU1E, 1968KU1D, 1969HA1G, 1969VA1C, 1971GR02, 1972AB19, 1972NA05, 1973HA49, 1973SL02).

Astrophysical questions: (1968HA1C, 1970BA1M, 1972CL1A, 1972KO1E, 1973AU1H, 1973LA19, 1973RA37, 1973RE1G, 1974AU1A).

Special reactions: (1968HA1C, 1968YI01, 1969AR13, 1969GA18, 1969YI1A, 1971AR02, 1972VO06, 1973KO1D, 1973KU03, 1973LA19, 1973WI15).

Muon capture: (1968BA2G, 1969WU1A, 1970FA15, 1971DE2D, 1972BU29, 1973MU11).

Pion capture and reactions: (1967ME1F, 1967MI1B, 1968BA2G, 1968ER1A, 1968NO1A, 1968WI1B, 1969BU1C, 1969CA1B, 1969CH1C, 1969KO1F, 1969MO1E, 1970BA1E, 1970BE1J, 1970CA1L, 1970ER1A, 1970GO28, 1971CA01, 1971CA1J, 1971FA09, 1971GO14, 1971MA1C, 1971RE1H, 1971SE02, 1972AB1H, 1972BE34, 1972BU1P, 1972HU1A, 1972MA1H, 1972SA10, 1972SE1F, 1973BA2R, 1973BA2V, 1973DI1H, 1973GA20, 1973HS1A, 1973HS1B, 1973JA1K, 1973NY04, 1973OS01, 1973PE1E, 1973SQ01, 1973UL1D).

Kaon reactions: (1973BA1Y).

Other topics: (1965CO25, 1965VO1A, 1966CH1B, 1966DO1C, 1966HA18, 1966HE1C, 1966WI1E, 1967BA12, 1967CA17, 1967CH1H, 1967MO1H, 1968KO1H, 1968GO01, 1969GU03, 1969LE1G, 1970CO1H, 1970GR33, 1970KA1K, 1970PA1D, 1970PE18, 1970SA05, 1971DA13, 1971ER1C, 1971TU04, 1972AN05, 1972CA37, 1972CH1P, 1972FR09, 1972HA57, 1972LE1L, 1972PN1A, 1972RA1J, 1972TA31, 1973BA1Y, 1973BE1N, 1973CL09, 1973JU2A, 1973KO1J, 1973KU03, 1973MA48, 1973RA1E).

Table 9.2: Energy levels of ${}^9\text{Be}$

| E_x (MeV \pm keV) | $J^\pi; T$ | $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|--------------------------|--------------------------------|---------------------------------|---------------------------|--|
| g.s. | $\frac{3}{2}^-; \frac{1}{2}$ | | stable | 2, 3, 4, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 |
| 1.680 ± 15 | $\frac{1}{2}^+; \frac{1}{2}$ | 210 ± 25 | γ, n | 4, 10, 11, 12, 15, 19, 20, 21, 23, 25, 26, 31, 36, 37, 42, 44 |
| 2.4294 ± 1.3 | $\frac{5}{2}^-; \frac{1}{2}$ | 1.03 ± 0.18 | γ, n, α | 4, 10, 11, 12, 14, 15, 19, 20, 21, 23, 24, 25, 26, 31, 34, 36, 37, 41, 42, 44 |
| 2.78 ± 120 | $\frac{1}{2}^-; \frac{1}{2}$ | 1080 ± 110 | n | 4, 10, 14 |
| 3.058 ± 12 | $\frac{5}{2}^+; \frac{1}{2}$ | 292 ± 15 | γ, n | 4, 10, 12, 15, 19, 20, 21, 23, 25, 26, 31, 36, 37, 41, 42, 44 |
| 4.704 ± 25 | $(\frac{3}{2})^+; \frac{1}{2}$ | 743 ± 55 | γ, n | 4, 10, 19, 21, 23, 25, 42 |
| 6.76 ± 60 | $\frac{7}{2}^-; \frac{1}{2}$ | 2000 ± 200 | γ, n | 10, 19, 20, 21, 23, 25, 26 |
| 7.94 ± 80 | | ≈ 1000 | γ | 19, 21 |
| 11.283 ± 24 | $\pi = -$ | 575 ± 50 | γ, n | 10, 19, 21, 26, 37 |
| 11.81 ± 20 | $T = \frac{1}{2}$ | 400 ± 30 | γ, n | 10, 12, 15, 41 |
| 13.79 ± 30 | $T = \frac{1}{2}$ | 590 ± 60 | γ | 10, 12, 19, 41 |
| 14.396 ± 5^a | $\frac{3}{2}^-; \frac{3}{2}$ | 0.33 ± 0.06 | γ, n, α | 10, 19, 21, 25, 37, 41 |
| 14.4 ± 300 | | ≈ 800 | | 21, 37 |
| 15.10 ± 50 | | | γ | 12, 19, 41 |
| 15.96 ± 30 | $T = \frac{1}{2}$ | ≈ 300 | γ | 19, 41 |
| 16.671 ± 8 | | 41 ± 4 | γ | 10, 19, 21, 37 |
| 16.977 ± 2 | $\frac{1}{2}^-; \frac{3}{2}$ | < 0.47 | γ, n, p, d | 4, 5, 6, 19 |
| 17.300 ± 12 | $(\frac{5}{2})^-$ | 195 | γ, n, p, d, α | 5, 6, 7, 19 |

Table 9.2: Energy levels of ${}^9\text{Be}$ (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|--------------------------|--------------------------------|---------------------------------|---------------------------|-----------------|
| 17.498 \pm 15 | $(\frac{3}{2}, \frac{5}{2})^+$ | 47 | γ, n, p, d, α | 5, 6, 7, 19, 21 |
| 18.02 \pm 50 | | | γ, n, p, d | 5, 6, 19 |
| 18.58 \pm 40 | | | γ, p, d, α | 5, 6, 19 |
| 19.10 \pm 30 | | 300 \pm 100 | γ, n, p, d, t | 1, 6, 15, 21 |
| 19.51 \pm 50 | | | γ, n, p, d | 6, 19 |
| (20.47 \pm 40) | | | γ, p, d | 6, 15 |
| 20.74 \pm 30 | | \approx 1000 | γ, p, t | 1, 15, 19 |
| (21.50 \pm 50) | | | γ, n | 15, 19 |
| (22.4 \pm 700) | | broad | | 21 |
| (23.9 \pm 100) | | | γ, n | 15 |

^a See also Table 9.6.

Ground state properties: (1965CO25, 1965GR18, 1965VO1A, 1966AD06, 1966BA26, 1966EL08, 1966MA1P, 1966WI1E, 1967SH05, 1967SH14, 1968DZ1A, 1968PE16, 1969AF1A, 1969BO19, 1969GU03, 1969HE1N, 1969JA1M, 1969VA1C, 1971AU1G, 1972FR09, 1972LE1L, 1973MA1K).

$\mu = -1.1776$ nm (1969FU11). See also (1971SH26);

$Q = 0.065_{-0.006}^{+0.009}$ b (1973BE19). See also (1967BL09, 1969FU11, 1971SH26).

1. (a) ${}^6\text{Li}(t, n){}^8\text{Be}$ $Q_m = 16.024$ $E_b = 17.6895$
- (b) ${}^6\text{Li}(t, p){}^8\text{Li}$ $Q_m = 0.801$
- (c) ${}^6\text{Li}(t, d){}^7\text{Li}$ $Q_m = 0.9930$
- (d) ${}^6\text{Li}(t, \alpha){}^5\text{He}$ $Q_m = 15.22$
- (e) ${}^6\text{Li}(t, n){}^4\text{He}{}^4\text{He}$ $Q_m = 16.116$

The 0° differential cross section for reaction (a) increases monotonically between $E_t = 0.10$ and 2.4 MeV (1960SE12, 1961VA43, 1962SE1A) except for a resonance at $E_t = 1.875$ MeV (${}^9\text{Be}^* = 18.938$). The excitation function for ${}^8\text{Li}$ (reaction (b)) increases monotonically for $E_t = 0.275$ to 1.000 MeV (1972CI05). In the range $E_t = 2.0$ to 6.8 MeV, a broad peak [$\Gamma \approx 1.3$ MeV]

is observed at $E_t = 4.57$ MeV [$E_x = 20.73$ MeV] (1973AB10). For reactions (c) and (d) see (1966LA04). For reaction (e) see (1966LA02, 1967BE13). See also ^5He , ^7Li and ^8Li .

$$2. \text{}^6\text{Li}(\alpha, \text{p})^9\text{Be} \quad Q_m = -2.1251$$

$$Q_0 = -2.1256 \pm 0.0012 \text{ (1967BR1B)}.$$

Angular distributions of ground state protons have been measured at $E_\alpha = 10.2, 11.5$ and 13.5 MeV (1960MA15), 13.6 and 14.7 MeV (1962KO13) and 30 MeV (1960KL03). See also (1966LA04).

$$3. \text{}^6\text{Li}(\text{}^6\text{Li}, \text{}^3\text{He})^9\text{Be} \quad Q_m = 1.895$$

See (1964KI02).

$$4. \text{}^7\text{Li}(\text{d}, \gamma)^9\text{Be} \quad Q_m = 16.6965$$

For $E_d = 0.1$ to 1.1 MeV, a resonance in the yield of capture γ -rays is observed at $E_d = 362 \pm 3$ keV (1965WO01), 361 ± 2 keV (1965IM01), corresponding to $^9\text{Be}^*(16.977)$ with $\Gamma_{\text{cm}} < 0.47$ keV. The small width of this state and its energy correspondence with $^9\text{Li}^*(2.69)$ argue for $T = \frac{3}{2}$ (1965WO01). The angular distribution of the γ -rays to $^9\text{Be}(0)$ is isotropic to within 7% (1965IM01). The branching ratios to $^9\text{Be}^*(0, 1.7, 2.4, 2.8, 3.1, 4.7)$ are $100/8.5 \pm 4.3/10.6 \pm 5.3/- / \leq 4.5/9.6 \pm 4.8$ (1965IM01), $100/11.8 \pm 0.6/3.3 \pm 0.7/13.3 \pm 4.2/- /12.9 \pm 1.3$ (1971SC19). The E_x and Γ of $^9\text{Be}^*(2.8, 4.7)$ are 2.82 and 1.7 MeV, and 4.64 and 0.95 MeV, respectively. The character of the decay suggests $(\frac{1}{2})^-$ for the second $T = \frac{3}{2}$ state [$^9\text{Be}^*(16.98)$] and is consistent with $J^\pi = (\frac{1}{2})^-$ for $^9\text{Be}^*(2.8)$ (1971SC19). See also (1968SN1A).

$$5. \text{ (a) } ^7\text{Li}(\text{d}, \text{n})^8\text{Be} \quad Q_m = 15.031 \quad E_b = 16.6965$$

$$\text{ (b) } ^7\text{Li}(\text{d}, \alpha)^5\text{He} \quad Q_m = 14.23$$

$$\text{ (c) } ^7\text{Li}(\text{d}, \text{n})^4\text{He}^4\text{He} \quad Q_m = 15.1233$$

The yield of neutrons has been measured for $E_d = 0.2$ to 4.8 MeV (1952BA64, 1957SL01, 1965IM01), 0.86 to 1.33 MeV by (1969NU1C), 3 to 8 MeV by (1967KE1F: yield of neutrons to $^8\text{Be}^*(16.6, 17.6, 18.1)$, at one MeV intervals) and 5 to 19 MeV by (1973WE19). Polarization measurements have been reported by (1971MO1R, 1973VO07: $E_d = 0.64$ MeV; n_0) and (1970TH08: $E_d = 2.5$ to 3.7 ; n_0, n_1). See also (1972SE09; theor.) and (1966LA04). Resonances in the yield of neutrons are observed at $E_d = 0.36, 0.68, 0.98$ and (1.8) MeV: see Table 9.3.

Table 9.3: Resonances in ${}^7\text{Li} + \text{d}$

| ${}^7\text{Li}(\text{d}, \text{p}){}^8\text{Li}$ | | ${}^7\text{Li}(\text{d}, \text{n}){}^8\text{Be}$ | ${}^7\text{Li}(\text{d}, \alpha){}^5\text{He}$ | | |
|--|-----------------------------|--|--|----------------------|---|
| E_{res} (keV) | Γ_{lab} (keV) | E_{res} (keV) | E_{res} (keV) | E_{x} (MeV) | J^{π} |
| 360 ± 3 ^a | < 2 | 360 ^d | | 16.976 | |
| 777 ± 12 ^b | 250 | 680 ^e | 750 ^f | 17.300 | $(\frac{3}{2})^{-}$ ^h |
| 1031 ± 15 ^b | 60 | 980 ^e | 1000 ^f | 17.498 | $(\frac{3}{2}, \frac{5}{2})^{+}$ ^h |
| 2000 ^c | | (1800) ^e | | 18.3 | |
| 2375 ± 50 ^{c,i} | | | 2500 ^g | 18.54 | |
| 3220 ± 50 ⁱ | 400 ± 100 | | | 19.20 | |
| ≈ 4800 ⁱ | | | | 20.4 | |

^a In ${}^7\text{Li}(\text{d}, \gamma){}^9\text{Be}$ $E_{\text{res}} = 361 \pm 2$ keV (1965IM01, 1965WO01); $\Gamma_{\text{n}_0}/\Gamma_{\gamma} \approx 1.5$, $\Gamma_{\alpha_0}/\Gamma_{\gamma} < 20$ (1965IM01) for ${}^9\text{Be}^*(16.98)$.

^b (1952BA64, 1954BA46, 1972SCIU).

^c (1956BE1A).

^d (1965IM01).

^e (1952BA64, 1957SL01).

^f (1963PA04, 1969DE31, 1971FR04).

^g (1963PA04): broad structure.

^h (1972DE44). See, however, (1971FR04, 1973HE26).


ⁱ (1973AB10).

The yield of α -particles has been measured for $E_{\text{d}} = 0.2$ to 0.3 MeV by (1964MA60), 0.6 to 2.0 MeV by (1971FR04), 0.7 to 1.6 MeV (1969DE31) and 0.7 to 3.0 MeV by (1963PA04). The excitation function for reaction (b) shows resonances at $E_{\text{d}} = 0.75$ and 1.00 MeV (1971FR04), 0.78 and 1.08 MeV (1969DE31), as well as a broad structure at $E_{\text{d}} = 2.5$ MeV (1963PA04). Also reported are α -particles from reaction (a) associated with ${}^8\text{Be}^*(11.4, 16.6, 16.9)$; the α -particles from the decay of ${}^8\text{Be}^*(11.4)$ seems to show resonance behavior at $E_{\text{d}} = 0.7, 1.0$ and 1.75 MeV. It is not clear whether the α -particles corresponding to ${}^8\text{Be}^*(16.6)$ show resonance at $E_{\text{d}} = 2.5$ MeV or whether the ${}^8\text{Be}^*(16.9)$ α -particles are appearing at this point (1963PA04).

A study of the B_{L} coefficients of the angular distributions of the α_0 group for $E_{\text{d}} = 0.45$ to 2.0 MeV and of the $\alpha - \alpha$ angular correlation coefficient, together with a re-analysis of the (d, d₀) data of (1964FO13), lead to assignments of $\frac{3}{2}^{-}$ and $(\frac{3}{2}, \frac{5}{2})^{+}$ for ${}^9\text{Be}^*(17.30, 17.50)$. The analysis also requires a $(\frac{3}{2}, \frac{5}{2})^{-}$ state at lower excitation energy and a $(\frac{3}{2}, \frac{5}{2})^{+}$ state at higher energy (1972DE44). See also (1973VO07) for an interpretation of results from reaction (a).

A kinematically complete study of reaction (c) at $E_{\text{d}} = 1.0$ MeV shows that the yield is dominated by sequential decay via ${}^8\text{Be}^*(2.9)$ and ${}^5\text{He}_{\text{g.s.}}$. There is evidence also for the involvement of ${}^9\text{Be}^*(17.50)$ [measurement of relative yield for $E_{\text{d}} = 0.9$ to 1.1 MeV]: $J = \frac{3}{2}$ is suggested

(1973HE26). See also ^5He , ^8Be and (1966AS04, 1967KE1F, 1967VA11, 1968WI1E, 1973HE06). See also (1966NU1B, 1966PO1D, 1967BE13, 1967WI1C, 1968DA1H, 1971DA2I, 1972BR1R, 1972SU1E, 1973DA1R).



The yield of p_0 measured for $E_d = 0.29$ to 0.78 MeV shows a single resonance with $E_d = 360 \pm 3$ keV, $\Gamma < 2$ keV, $\Gamma_p/\Gamma_\gamma \approx 0.5$ (1965IM01, 1965WO01): see also reaction 4.


The yield of ^8Li has been measured for $E_d = 0.4$ to 4 MeV (1952BA64, 1954BA46, 1956BE1A, 1960KA05), for $E_d = 0.62$ to 1.97 MeV by (1972SC1U) and for $E_d = 2.0$ to 7.0 MeV by (1973AB10): observed resonances are displayed in Table 9.3. The yield of 0.98 -MeV γ -rays [from $^7\text{Li}(d, p)^8\text{Li}^*$] rises monotonically from $E_d = 1.9$ to 3.3 MeV (1962CH14).

The total cross section at the $E_d = 0.77$ MeV resonance is 202 ± 9 mb (1972SC1U). Earlier values were 176 ± 15 mb (1960KA05) and 211 ± 15 mb (1966PA16). We adopt 205 ± 8 mb. This cross section is important since the $^7\text{Be}(p, \gamma)^8\text{B}$ data are normalized to it and the S -factor is of interest in relation to the solar neutrino problem: see (1968PA1M). See also ^8Li .



The upper limit for the relative partial width for elastic scattering at $E_d = 0.36$ MeV ($^9\text{Be}^* = 16.98$), $\Gamma_{d_0}/\Gamma_\gamma$ is 400 (1965IM01).

The elastic scattering, at $E_d = 0.4$ to 1.8 MeV, shows a marked increase in cross section for $E_d = 0.8$ to 1.0 MeV [perhaps related to $^9\text{Be}^*(17.30)$ and a conspicuous anomaly at $E_d = 1.0$ MeV, due to p -wave deuterons [$^9\text{Be}^*(17.50)$] (1964FO13) [and discussion in (1972DE44) and in reaction 5]. The elastic scattering cross section ($\theta_{\text{cm}} = 162^\circ$) decreases monotonically for $E_d = 10.0$ to 12.0 MeV (1971BI11). See also ^7Li .



The cross section for reaction (a) rises from threshold to 95 mb at $E_d = 2.4$ MeV and then more slowly to ≈ 165 mb at $E_d = 4.1$ MeV (1955MA20). The t_0 yield curve ($\theta_{\text{lab}} = 155^\circ$) decreases monotonically for $E_d = 10.0$ to 12.0 MeV (1971ZA07). See also ^6Li . For reaction (b) see ^6He .



Table 9.4: Excited states of ${}^9\text{Be}$ from ${}^7\text{Li}({}^3\text{He}, \text{p}){}^9\text{Be}$ ^a

| (1968CO07) | | (1965LY01, 1971AD01) | | (1968KR02) | |
|-----------------------------|---------------------------------|-----------------------------|---------------------------------|--------------------------|---------------------------------|
| E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) |
| 1.64 | | | | | |
| 2.429 ± 12 | ≤ 35 | | | 2.4292 ± 1.7 | < 8 |
| | | 2.9 ± 250 | 1000 ± 250 ^b | | |
| 3.031 ± 10 | 274 ± 15 | | | 3.076 ± 15 | 289 ± 22 |
| 4.57 ± 100 ^d | 800 ± 200 | | | 4.704 ± 25 | 743 ± 55 |
| 6.7 ± 100 | 1950 ± 250 ^e | | | | |
| 11.29 ± 30 | 620 ± 70 | | | | |
| 11.81 ± 20 | 400 ± 30 | | | | |
| 13.78 ± 30 | 590 ± 60 | | | | |
| | | 14.396 ± 5 ^f | < 5 ^c | | |
| 16.671 ± 8 | 41 ± 4 | | | | |

^a See also Table 9.5 in (1966LA04).

^b From γ -decay of ${}^9\text{Be}^*(14.39)$.

^c See also Table 9.5.

^d 4.65 MeV, $\Gamma = 900 \pm 250$ keV (1968CO08).

^e 2300 \pm 500 keV (1968CO08).

^f Based on Q_m .

See (1959AJ77, 1962SE1A) and ${}^{10}\text{Be}$.

10. (a) ${}^7\text{Li}({}^3\text{He}, \text{p}){}^9\text{Be}$ $Q_m = 11.2027$

(b) ${}^7\text{Li}({}^3\text{He}, \text{np}){}^8\text{Be}$ $Q_m = 9.538$

Observed proton groups are listed in Table 9.4 (1968CO07). See also Table 9.5 in (1966LA04) for a listing of the older work. Angular distributions have been measured for the protons to ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1)$ at $E({}^3\text{He}) = 0.90$ to 1.10 MeV (1971ST35: not to ${}^9\text{Be}^*(1.7)$), 2.2 to 3.2 MeV (1969SA04, 1972LI31) and at $E({}^3\text{He}) = 10$ MeV (1970DI12, 1970DI1F), and for the group corresponding to ${}^9\text{Be}^*(14.39)$ at $E({}^3\text{He}) = 10$ MeV (1971AD01). The characteristics of the neutron and γ -decays of ${}^9\text{Be}$ states are displayed in Tables 9.5 and 9.6 (1965GR08, 1965LY01, 1966CH20, 1968CO08, 1968KR02, 1971AD01, 1972AD04, 1972MC1E). See also (1964MA57, 1970LI1Q), (1969BA1Z), (1968SA1G, 1968TA1N, 1970LK1A, 1971WE1L; theor.), ${}^8\text{Be}$ and ${}^{10}\text{B}$.

Table 9.5: Neutron decay of ${}^9\text{Be}$ states

| ${}^9\text{Be}$ state (MeV) | l_n | Decay (in %) to | | θ^2 ^a (%) | Refs. |
|--------------------------------|-------|--------------------------|------------------------|--------------------------------|----------------------|
| | | ${}^8\text{Be}(0)$ | ${}^8\text{Be}^*(2.9)$ | | |
| 2.43 | 3 | 7.5 ± 1 ^b | | 2.1 ± 0.6 | (1966CH20) |
| | | 6.4 ± 1.2 | | | |
| 2.78 | 1 | mainly | | 0.48 ± 0.06 | (1970CH07, 1970CH1T) |
| 3.06 | 2 | 87 ± 13 | | 81 ± 13 | (1966CH20, 1968CO08) |
| 4.70 | 2 | 13 ± 4 | | 6.0 ± 0.4 | (1968CO08) |
| 6.76 | 3 | ≤ 2 | | ≤ 6 | (1968CO08) |
| | 1 | | 55 ± 14 | 37 ± 10 | (1968CO08) |
| 11.28 | 1 | ≤ 2 | | ≤ 0.1 | (1968CO08) |
| | 1 | | 14 ± 4 | 0.93 ± 0.28 | (1968CO08) |
| | 3 | | | 4.0 ± 1.2 | (1968CO08) |
| 11.81 | 1 | ≤ 3 | | ≤ 0.1 | (1968CO08) |
| | 1 | | 12 ± 4 | 0.48 ± 0.16 | (1968CO08) |
| | 3 | | | 1.8 ± 0.6 | (1968CO08) |
| 14.40 | | < 7 | | | (1972MC1E) |
| | | | 50 ± 12 | | (1972MC1E) |

^a Expressed in units of $\hbar^2/mR^2 = 2.47$ MeV (1968CO08, 1970CH07).

^b See also (1959MA34).

11. ${}^7\text{Li}(\alpha, d){}^9\text{Be}$ $Q_m = -7.1511$

At $E_\alpha = 30$ MeV angular distributions have been measured for the deuterons corresponding to ${}^9\text{Be}^*(0, 1.7, 2.4)$ (1972ME07). See also (1971BU1K; theor.) and (1966LA04).

12. ${}^7\text{Li}({}^6\text{Li}, \alpha){}^9\text{Be}$ $Q_m = 15.223$

At $E({}^7\text{Li}) = 2.9$ MeV α -particle groups are observed corresponding to ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1, 11.9 \pm 0.2 [\Gamma = 0.5 \pm 0.1 \text{ MeV}])$ (1964ME07). Angular distributions of the α_0 group have been measured at $E({}^7\text{Li}) = 3.78$ to 5.95 MeV (1967KI03). At $E({}^6\text{Li}) = 26.0$ MeV and $E({}^7\text{Li}) = 30.3$ MeV the excitation of ${}^9\text{Be}^*(0, 11.8, 13.8, 15.2, 17.8, 21.0)$ is reported (1971GL07). See also (1967CH34, 1968DA20, 1970OG1A) and (1966RO1E, 1966RO1F, 1966RO1H, 1969RO1G; theor.).

Table 9.6: Parameters of ${}^9\text{Be}^*(14.40)$ ^a

| | | Refs. |
|--|------------------------------|----------------------|
| E_x (MeV \pm keV) | 14.396 ± 5 | (1965LY01) |
| $J^\pi; T$ | $\frac{3}{2}^-; \frac{3}{2}$ | |
| Γ_{γ_0} (eV) | 6.9 ± 0.5 | See (1973BE19) |
| Γ_{γ_0}/Γ | 0.021 ± 0.004 | (1971AD01) |
| Γ (eV) | 329 ± 60 | (1971AD01, 1973BE19) |
| $\Gamma_{\gamma_{2.43}}/\Gamma_{\gamma_0}$ | 1.19 ± 0.16 | (1971AD01) |
| $\Gamma_{\gamma_{2.9}}/\Gamma_{\gamma_{2.43}}$ | 0.30 ± 0.04 | (1971AD01) |
| $\Gamma_{n_0}/\Gamma_{\gamma_0}$ | 6.4 ± 2.0 | (1972AD04) |
| $\Gamma_{n_{2.9}}/\Gamma_{\gamma_0}$ | 20.4 ± 4.6 | (1972AD04) |
| $\Gamma_{\alpha_0}/\Gamma_{\gamma_0}$ | 31.2 ± 9.8 | (1972AD04) |
| Γ_{n_0} (eV) | 44 ± 15 | (1972AD04, 1973BE19) |
| $\Gamma_{n_{2.9}}$ (eV) | 141 ± 35 | (1972AD04, 1973BE19) |
| Γ_{α_0} (eV) | 215 ± 60 | (1972AD04, 1973BE19) |

^a See also Table 9.5 and reaction 4 in ${}^9\text{B}$ for the parameters of the analog state: ${}^9\text{Be}^*(14.66)$. See also (1965LY01, 1965GR08).

13. ${}^7\text{Li}({}^{11}\text{B}, {}^8\text{Be}){}^9\text{Be}$ $Q_m = 7.286$

See (1970LK1A; theor.).

14. ${}^9\text{Li}(\beta^-){}^9\text{Be}$ $Q_m = 13.618$

${}^9\text{Li}$ decays by β^- emission with $\tau_{1/2} = 176 \pm 2$ msec to ${}^9\text{Be}^*(0, 2.43, 2.78)$: see ${}^9\text{Li}$ and Table 9.7. ${}^9\text{Be}^*(2.43, 2.78)$ are neutron unstable. The probability that ${}^9\text{Li}$ decays to these two states is 0.35. The branching ratio for the ${}^9\text{Be}^*(2.43) \rightarrow {}^8\text{Be}(0) + n$ decay is $(6.4 \pm 1.2)\%$. ${}^9\text{Be}^*(2.78)$ decays mainly to ${}^8\text{Be}(0) + n$. The assignment $J^\pi = \frac{1}{2}^-$ to ${}^9\text{Be}^*(2.78)$ derives from the allowed nature of the ${}^9\text{Li}$ decay, and the large value of θ_p^2 [0.48 ± 0.06] which is in agreement with the shell-model prediction that the $\frac{1}{2}^-$ state should decay mainly by p-wave neutron emission to ${}^8\text{Be}(0)$: if $J^\pi = \frac{3}{2}^-$ this decay branch should be small (1970CH07, 1970CH1T). See also (1963AL18, 1965SC17, 1973RO2F) and (1969MA11) for a general discussion of the problems in identifying the parameters of the low-lying states of ${}^9\text{Be}$.

Table 9.7: Branching parameters in ${}^9\text{Li}$ β -decay (1970CH07, 1970CH1T) ^a

| E_x in ${}^9\text{Be}$ (MeV) | $J^\pi; T$ | Branching ratio (%) | $\log ft$ |
|--------------------------------|------------------------------|-----------------------------------|------------------------|
| 0 | $\frac{3}{2}^-; \frac{1}{2}$ | $65.0^{+2.7}_{-2.4}$ | $5.12^{+0.01}_{-0.02}$ |
| 2.43 | $\frac{5}{2}^-; \frac{1}{2}$ | $32.0^{+2.7}_{-3.7}$ ^c | $5.00^{+0.04}_{-0.05}$ |
| 2.78 ± 0.12 ^b | $\frac{1}{2}^-; \frac{1}{2}$ | $3.0^{+2.7}_{-0.3}$ ^c | $5.97^{+0.05}_{-0.28}$ |

^a See also (1963AL18, 1969MA11).

^b $\Gamma_{\text{c.m.}} = 1.10 \pm 0.12$ MeV; $\theta_p^2 = 0.48 \pm 0.06$ (1970CH07).

^c See also (1973RO2F).

15. (a) ${}^9\text{Be}(\gamma, n){}^8\text{Be}$ $Q_m = -1.6651$
 (b) ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$ $Q_m = -2.46$
 (c) ${}^9\text{Be}(\gamma, n){}^4\text{He}{}^4\text{He}$ $Q_m = -1.5732$
 (d) ${}^9\text{Be}(\gamma, 2n){}^7\text{Be}$ $Q_m = -20.565$

The photoneutron cross section has been measured from threshold to 320 MeV: see Table 9.6 in (1966LA04). A sharp peak occurs 6 keV above threshold (1967BE49) [but see discussion in (1968BA1C)] with $\sigma_{\text{max}} = 1.6$ mb. The cross section then decreases slowly to 1.2 mb at $E_\gamma = 40$ keV (1967BE49). A satisfactory fit to the cross section is obtained with a one-level approximation of R -matrix theory (1968BA1C). Peaks in the cross section of (γ, xn) are reported corresponding to $E_x = 2.43 \pm 0.03$, 3.00 ± 0.03 , 19.00 ± 0.03 , 21.50 ± 0.05 and 23.9 ± 0.1 MeV, and there is some indication also of structure at $E_x = 9.1$, 10.8, 12.8 and 14.8 MeV (1973HU1G). See also (1966CO16, 1972TH12) and (1966LA04) for references to earlier work.

The total absorption cross section has been measured for $E_\gamma = 10$ to 155 MeV by (1972AH1B, 1973AH1A)[†]. The integrated cross section for $E_\gamma = 16$ to 29 MeV is 53 MeV · mb (1972AH1B). (1969DO09) report an integrated cross section of 156 ± 15 MeV · mb for $E_\gamma = 10$ to 29 MeV and resonant structure at $E_\gamma = 11.8$, (13.5), 14.8, (17.3), (19.5), 21.0, (23.0) and (25.0) MeV (1969DO09). Fine structure is reported at $E_\gamma = 20.47 \pm 0.04$ and 20.73 ± 0.04 MeV (1964TE04). See also (1965WY02).

See (1966LA04) for a discussion of the early evidence on ${}^9\text{Be}$ levels from reaction (a). See also (1966TH03, 1967GL1B, 1968AD09, 1968KA38, 1971KA70, 1972BU1R, 1972CR1E), (1967SH1E) and (1965BO1B, 1965MA1H, 1967BO1K, 1967BO34, 1969BO1U, 1968MA1Y, 1968MA1X, 1968PA1H, 1969AU05, 1970SA17, 1972TA31, 1973SL02; theor.).

16. (a) ${}^9\text{Be}(\gamma, p){}^8\text{Li}$ $Q_m = -16.888$
 (b) ${}^9\text{Be}(\gamma, np){}^7\text{Li}$ $Q_m = -18.921$

[†] We are deeply indebted to E.G. Fuller for his very helpful remarks on the ${}^9\text{Be} + \gamma$ processes.

The yield shows structure in the energy region corresponding to the ${}^9\text{Be}$ levels at 17–19 MeV (1962CL06) followed by the giant resonances at $E_\gamma \approx 23$ MeV (1962CL06: $\sigma = 2.64 \pm 0.30$ mb). (1966DE13) report structure attributed to eleven states of ${}^9\text{Be}$ with $18.2 < E_x < 32.2$ MeV. Integrated cross sections have been obtained for each of these resonances, and over different energy intervals for protons leading to ${}^8\text{Li}^*(0 + 0.98, 2.26 + 3.21, 9.0, 17.0)$ (1966DE13). Angular and energy distributions of photoprotons in various energy intervals have been studied by many groups: see (1966LA04) and (1966VO06, 1968AD09). See also (1965KO1B, 1969AN1H, 1971AN04, 1973DO13), (1967SH1E, 1973CO1N) and (1968MA1X; theor.).

$$17. \text{ (a) } {}^9\text{Be}(\gamma, \text{d}){}^7\text{Li} \quad Q_m = -16.6965$$

$$\text{ (b) } {}^9\text{Be}(\gamma, \text{t}){}^6\text{Li} \quad Q_m = -17.6895$$

The integrated cross sections are reported to be 1.0 ± 0.5 MeV · mb ($E_\gamma = 21 \rightarrow 33$ MeV) for reaction (a) to ${}^7\text{Li}^*(0 + 0.4)$ and 0.6 ± 0.3 MeV · mb ($E_\gamma = 25 \rightarrow 33$ MeV) for reaction (b) to ${}^6\text{Li}(0)$. The total integrated cross section for $[(\gamma, \text{p}) + (\gamma, \text{pn}) + (\gamma, \text{d}) + (\gamma, \text{t})]$ is given as 33 ± 3 MeV · mb by (1966DE13), who also report resonances in the (γ, d) and (γ, t) cross sections corresponding to ${}^9\text{Be}^*(26.0 \pm 0.2)$ and ${}^9\text{Be}^*(32.2 \pm 0.3)$, respectively. See also (1966LA04) and (1966VO06, 1968AD09, 1969AN1H, 1971AN04, 1971AN15) for reaction (a) and (1966VO06, 1972AN09) for reaction (b). See also (1968MA1X; theor.).

$$18. {}^9\text{Be}(\gamma, \gamma){}^9\text{Be}$$

See (1967LO1B, 1968SN1A, 1969MO1H).

$$19. \text{ (a) } {}^9\text{Be}(\text{e}, \text{e}){}^9\text{Be}$$

$$\text{ (b) } {}^9\text{Be}(\text{e}, \text{en}){}^8\text{Be} \quad Q_m = -1.6651$$

$$\text{ (c) } {}^9\text{Be}(\text{e}, \text{ep}){}^8\text{Li} \quad Q_m = -16.888$$

$$\text{ (d) } {}^9\text{Be}(\text{e}, \text{e}\alpha){}^5\text{He} \quad Q_m = -2.46$$

$\langle r^2 \rangle^{1/2} = 2.46 \pm 0.11$ fm, $Q = 6.5_{-0.6}^{+0.9}$ fm², $b = 1.5_{-0.2}^{+0.3}$ fm [b = oscillator parameter] (1973BE19);

$\langle r^2 \rangle^{1/2} = 2.519 \pm 0.012$ fm, $Q = 6.4 \pm 2.4$ fm² ((1972JA10) and K. de Jager, private communication to J.C. Bergstrom);

$\langle r^2 \rangle^{1/2} = 2.43 \pm 0.08$ fm, $Q = 3$ fm² (1969BE21).

See also (1966AF1A, 1967BE26, 1967BE1P, 1973SL02). (1973LA1T) report $b = 1.80 \pm 0.03$ fm from magnetic scattering.

Table 9.8: Levels of ${}^9\text{Be}$ from ${}^9\text{Be}(e, e'){}^9\text{Be}^*$

| E_x in ${}^9\text{Be}$ (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | Transition | J^π | Γ_{γ_0} (eV) | Refs. |
|---|---------------------------------|------------|-----------------|----------------------------------|-------------------------|
| 1.78 ± 30^i | 150 ± 50 | E1 | $\frac{1}{2}^+$ | 0.3 ± 0.12 | (1968CL08) |
| | | | | 4.5 ± 0.6 | (1963NG01, 1965NG1A) |
| 2.44 ± 20^i | < 30 | M1 | $\frac{5}{2}^-$ | 0.13 ± 0.03 | (1960BA47) |
| | | | | 0.12 ± 0.02 | (1962ED02) |
| | | | | 0.13 ± 0.015 | (1968VA05) |
| | | | | 0.089 ± 0.010 | (1968CL08) |
| | | | | 0.12 ± 0.02 | “Best” |
| | | E2 | | $(1.89 \pm 0.14) \times 10^{-3}$ | (1968CL08) |
| 3.04 ± 20^i | 450 ± 150 | E1 | $\pi = +$ | 0.45 ± 0.35^e | (1968CL08) |
| | | M1 | $\pi = -$ | $(8.8 \pm 4.4) \times 10^{-2}^e$ | (1968CL08) |
| 4.7 ± 200 | 700 ± 300 | E(1) | | 2.4 ± 1.2^e | (1968CL08) |
| | | | | 0.3^e | (1968VA05) ^h |
| 6.4 ± 100^i | 2000 ± 500 | E2 | $\frac{7}{2}^-$ | 0.109 ± 0.005 | (1963NG01, 1965NG1A) |
| | 1100 ± 300 | | | 0.082 ± 0.035 | (1968CL08) |
| 8.0 ± 200 | | M1 | | 3.5 ± 1.0^e | (1968VA05) |
| 9.1 ± 200 | | M1 | | 1.9 ± 0.5^e | (1968VA05) ^h |
| 10.2 ± 200 | | M1 | | 1.7 ± 0.5^e | (1968VA05) ^h |
| 11.2 ± 200 | | M1 | | 5.6 ± 1.1^e | (1968VA05) ^h |
| 13.84 ± 50^a | | | | | (1973BE19) |
| 14.388 ± 15 | < 70 | M1 | $\frac{3}{2}^-$ | 6.2 ± 0.6 | (1973BE19) |
| | | | | 10.5 ± 1.5^j | (1966CL01) |
| | | | | 18 ± 9 | (1962ED02) |

Table 9.8: Levels of ${}^9\text{Be}$ from ${}^9\text{Be}(e, e'){}^9\text{Be}^*$ (continued)

| E_x in ${}^9\text{Be}$ (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | Transition | J^π | Γ_{γ_0} (eV) | Refs. |
|---|---------------------------------|-----------------|----------------------|--------------------------------|--------------------------------|
| | | | | 8 ± 2 | (1968VA05) ^h |
| | | | | 6.9 ± 0.5 ^f | “Best”: see (1973BE19) |
| 15.10 ± 50 ^a | | | | | (1973BE19) |
| 15.97 ± 30 ^a | ≈ 300 | | | 3.7 ± 0.8 ^{e,k} | (1966CL01, 1967AR1A, 1973BE19) |
| 16.631 ± 15 ^b | < 70 | M2 ^d | $\leq \frac{7}{2}^+$ | 0.30 ± 0.08 ^{e,k} | (1966CL01) |
| | | | | 0.26 ± 0.02 ^e | (1973BE19) |
| | | M1 | $\leq \frac{5}{2}^-$ | 2.0 ± 0.5 ^e | (1973BE19) |
| 16.961 ± 15 ^b | < 70 | M1 | $\frac{1}{2}^-$ | 18.8 ± 1.8 ^j | (1966CL01) |
| | | | | 11.5 ± 1.4 | (1973BE19) |
| 17.28 | | } M1 | $\leq \frac{5}{2}^-$ | 7.3 ± 1.3 ^e | (1973BE19) |
| 17.480 ± 20 | ≈ 100 | | } M2 ^d | $\leq \frac{7}{2}^+$ | 0.7 ± 0.2 ^{e,k} |
| | | | | 0.42 ± 0.10 ^e | See (1973BE19) ^g |
| | | | | 0.40 ± 0.03 ^e | (1973BE19) |
| 18.02 ± 50 ^a | | | | | (1973BE19) |
| 18.62 ± 50 ^{a,b} | | | | | (1973BE19) |
| 19.51 ± 50 ^a | | | | | (1973BE19) |
| 20.76 ± 50 ^{a,b} | | | | | (1973BE19) |
| ^c | | | | | |

^a Weak transition (1973BE19).

^b See also (1968VA05).

^c Higher states reached by M1 transitions are reported at 21.6 ± 0.2 , 22.5 ± 0.2 , 24.4 ± 0.2 and 25.7 ± 0.2 MeV (1968VA05).

^d Or pure spin-flip E1.

^e $g\Gamma_{\gamma_0}$, where $g = (2J_f + 1)/(2J_i + 1)$.

^f This value is calculated by (1973BE19): it is the weighted mean of 6.2 ± 0.6 eV, 8.1 ± 0.8 eV (an unpublished correction of (1966CL01): see (1972THZF)) and 6.7 ± 1.4 eV (an unpublished value by H.S. Caplan *et al*). We are grateful to Prof. J.C. Bergstrom for his comments.

^g Unpublished corrected value of (1966CL01)'s result: see (1973BE19).

^h And G.J. Vanpraet, private communication. All values for the cross sections listed in Table 1 of (1968VA05) for states with $E_x > 8$ MeV should be reduced by a factor of 3.13.

ⁱ See (1973SL02): $E_x = 1.79 \pm 0.06$ for ${}^9\text{Be}^*(1.7)$.

^j (1972THZF) list revised values for $\Gamma_{\gamma_0} = 8.1 \pm 0.8$ and 8.6 ± 0.9 eV for ${}^8\text{Be}^*(14.40, 16.96)$.

^k See also (1972THZF).

The elastic scattering of electrons has been studied at E_e up to 700 MeV: see (1966LA04), the references listed above, and (1965VA1G, 1966RA29, 1969BE50). Magnetic elastic scattering at $\theta = 180^\circ$ gives indication of both M1 and M3 contributions [(1965GR18, 1966RA29): see (1968KU1D)].

Inelastic scattering reveals a number of levels: Table 9.8 displays the parameters of these states (1960BA47, 1962ED02, 1963NG01, 1965NG1A, 1966CL01, 1967AR1A, 1968CL08, 1968VA05, 1973BE19, 1973SL02). See also (1966LA04) for a discussion of the earlier work.

Electron bremsstrahlung has been measured at $E_e = 1.0$ and 2.0 MeV by (1968RE11). See also (1966PE1E), (1968GO1J) and (1965GR1E, 1965NE1B, 1966DE1K, 1966KU1C, 1967KA1A, 1967ST1G, 1967WA1E, 1967WA1F, 1968JA1D, 1968KU1B, 1969BO1V, 1969BO19, 1969VI02, 1970BE1G, 1970TI1C, 1971DRZW, 1971GO14, 1972BL12, 1972BO01, 1973OK1B; theor.).

For reaction (b) see (1970AL1M; theor.). For reaction (c) see (1968AM1A, 1968BO46, 1970HI1F, 1970RE1E) and (1965AL1F, 1973HI03; theor.). For reaction (d) see (1973JU1E).

20. (a) ${}^9\text{Be}(n, n){}^9\text{Be}$

(b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$ $Q_m = -1.6651$

The neutron spectrum at $E_n = 3.7$ MeV is consistent with the excitation of ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1)$ with subsequent neutron decay of the two higher states (1957HU14, 1958WA05). About $\frac{1}{2}$ of the inelastic processes involve ${}^9\text{Be}^*(2.4)$ for $E_n = 2.6$ to 6.0 MeV; that level decays only $12 \pm 5\%$ via ${}^8\text{Be}(0) + n$ (1959MA34). See Table 9.5. At $E_n = 14$ MeV ${}^9\text{Be}^*(6.8)$ appears to be excited (1963JE05). Angular distributions have been measured at $E_n = 14$ MeV (1958AN32, 1958NA09, 1968RO1H, 1969RO1F; n_0, n_2). See also ${}^{10}\text{Be}$, (1966LA04) and (1965BO19, 1965FR1B, 1965GO1E, 1966AM1A, 1966BO1C, 1966BO1F, 1967BO1F, 1970DR11) and (1969WA11; theor.).

21. ${}^9\text{Be}(p, p){}^9\text{Be}$

The elastic scattering has been studied at many energies for $E_p = 5$ to 725 MeV: see (1966LA04). More recent angular distribution measurements have been carried out at $E_p = 2.009$ to 2.099 MeV (1970SI12; p_0), 6.36 and 6.48 MeV (1971VA34; p_1), 13.0, 14.0, 15.0, 21.35 and 30.3 MeV (1973VO02; p_0, p_2), 17.0, 21.0, 25.0, 29.1 (1973MO01; p_0, p_2 and p to ${}^9\text{Be}^*(3.1)$), 33.6 MeV (1970KU1D; p_0), 46 MeV (1967SA13, 1967VE01; p_0, p_2 and p to ${}^9\text{Be}^*(11.3, 14.4, 16.7, 17.5)$), 49.75 MeV (1971MA13, 1971MA44; p_0, p_2 and p to ${}^9\text{Be}^*(6.4)$), and 100 MeV (1966MA38, 1968LI1C; p_0, p_2 and p to ${}^9\text{Be}^*(4.7, 6.8)$). The elastic angular distributions show pronounced diffraction maxima characteristic of the optical model. See (1973MO01) for a discussion of optical model parameters. (1973VO02), in coupled channels analysis, find that a quadrupole-deformed optical model potential with a deformation parameter $\beta = 1.1$ [see also (1967SA13)] provides an improved description of the elastic data and a good fit to the p_2 data [to ${}^9\text{Be}^*(2.4)$].

The structure corresponding to ${}^9\text{Be}^*(1.7)$ is asymmetric: the line shape peaks 25_{-11}^{+15} keV above the threshold for ${}^8\text{Be} + n$ (1970TU06): see also the discussions in (1960SP08, 1971JE03) and in (1966LA04). The energy of ${}^9\text{Be}^*(2.4)$ is given as 2433 ± 5 (1951BR72), 2434 ± 5 (1956BO18), 2432 ± 4 (1955GO48), 2430 ± 5 keV (1960SP08). ${}^9\text{Be}^*(3.1)$ has a width of 250 ± 5 keV (1960SP08): $E_x = 3.03 \pm 0.03$ MeV (1956BO18), 3.04 ± 0.05 MeV (1960SP08), $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+$. Higher states are observed at $E_x = 4.8 \pm 0.2, 6.76 \pm 0.06$ [$J^\pi = \frac{1}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$ (but see below), $\Gamma = 1.2 \pm 0.2$ MeV], 7.94 ± 0.08 ($\Gamma \approx 1$ MeV), 11.3 ± 0.2 MeV ($\Gamma \approx 1$ MeV), 14.4 ± 0.3 ($\Gamma \approx 1$ MeV), $16.7 \pm 0.3, 17.4 \pm 0.3, 19.0 \pm 0.4, 21.1 \pm 0.5$ and 22.4 ± 0.7 MeV [the five highest states are all broad] (1956BE14, 1965HA17). See also Table 9.8 in (1966LA04). (1965JA1A) reports for ${}^9\text{Be}^*(2.4, 6.8)$ $B(E2\uparrow) = 49 \pm 6$ and 24 ± 4 fm⁴ and $\Gamma(E2\downarrow) = 0.0025$ and 0.10 eV, respectively. The strong population of ${}^9\text{Be}^*(2.4, 6.8)$ in this reaction is consistent with the assumption that they have $J^\pi = \frac{5}{2}^-$ and $\frac{7}{2}^-$, respectively, and are members of the ground state $K = \frac{3}{2}^-$ band: see discussion in (1966LA04). See also (1966MA38).

See also (1965HU10, 1971HU1A, 1971SC1N) and (1968GL1A, 1968NE1A, 1968NE1B, 1968SE1B, 1969NE1A, 1969NE08, 1969WA11, 1970MA04, 1970MA38, 1971BA87, 1971IN05, 1971RA36, 1972SO03, 1973GU08, 1973HU05; theor.).

| | |
|---|------------------|
| 22. (a) ${}^9\text{Be}(p, 2p){}^8\text{Li}$ | $Q_m = -16.888$ |
| (b) ${}^9\text{Be}(p, pd){}^7\text{Li}$ | $Q_m = -16.6965$ |
| (c) ${}^9\text{Be}(p, p\alpha){}^5\text{He}$ | $Q_m = -2.46$ |
| (d) ${}^9\text{Be}(p, pn){}^8\text{Be}$ | $Q_m = -1.6651$ |
| (e) ${}^9\text{Be}(p, p^3\text{He}){}^6\text{He}$ | $Q_m = -21.181$ |

The summed proton spectrum (reaction (a)) shows two peaks with $Q = -16.4 \pm 0.3$ and $Q = -25.4 \pm 0.5$ MeV, corresponding to removal of a p-proton and an s-proton respectively, and a third peak of uncertain assignment with $Q = -32.3 \pm 0.6$ MeV (probably due to unresolved states) (1966TY01: $E_p = 460$ MeV). See also (1966LA04), (1966WA12, 1968PE1A), (1965BE1E, 1966JA09, 1966JA1A, 1967BE1Q, 1967JA1E, KO67Q, 1968JA1G, 1969KO1J; theor.) and ${}^8\text{Li}$. For reaction (b) see (1966LA04) and (1973KO1M). Reaction (c) is characterized by a sequential mechanism at low energies and by a quasi elastic scattering mechanism at higher energies. At $E_p = 26.0, 35.0, 46.8$ and 57 MeV the data are consistent with quasi free scattering of the incident protons by zero-momentum α -particle clusters in ${}^9\text{Be}$: the probability of finding such clusters at these four energies is determined to be $0.166, 0.099, 0.124$ and 0.09 , respectively (1968RO19, 1970QU1C, 1972QU01). See also (1967TA1C, 1969YA1B, 1970GO12, 1973WO1E), (1969HO1K, 1971GA1J) and (1973HO30; theor.).

For reaction (d), see (1969WI1F, 1970TH1F, 1971HU1A, 1973MI1J) and (1967BE1Q; theor.). For reaction (e) see (1969KO1G). See also (1968KO1E, 1969ED01).

23. ${}^9\text{Be}(d, d){}^9\text{Be}$

Elastic scattering has been studied at many energies in the range $E_d = 7.8$ to 27.7 MeV: see (1966LA04). Elastic angular distributions are also reported for $E_d = 1.1$ to 2.5 MeV (1968MA1H), 4.5 to 6.0 MeV (1970PO03), 5.00 to 7.00 MeV (1971DJ02), 11.8 MeV (1967FI07; also d_2), 12.8 MeV (1973VA08; d_1), 13.6 MeV (1968VE11, 1968VE1C, 1970VE06, 1972MA47), 14.35 MeV (1966NG01; also d_2), 15.0 MeV (1969AR1B), 15.8 MeV (1966CO24; also d_2) and 410 MeV (1960BU25; forward angles). For optical model parameters see discussions in (1966CO24, 1967FI07).

Inelastic deuteron groups have been observed to ${}^9\text{Be}^*(1.7, 2.4, 3.1, 4.7, 6.8)$: see (1966LA04). (1968KR02) report $E_x = 2431.9 \pm 7.0$ keV and 3040 ± 15 keV [$\Gamma = 294 \pm 20$ keV]. See also (1963ST1A, 1969VE09, 1970EL16, 1973ZW1A; theor.) and ${}^{11}\text{B}$.

24. (a) ${}^9\text{Be}(t, t){}^9\text{Be}$

(b) ${}^9\text{Be}(t, nt){}^8\text{Be}$ $Q_m = -1.6651$

The angular distribution of elastically scattered tritons has been measured at $E_t = 2.10$ MeV (1969HE08, 1970CO04). Reaction (b) at $E_t = 4.2$ and 4.6 MeV proceeds via ${}^9\text{Be}^*(2.4)$ (1967SE11).

25. ${}^9\text{Be}({}^3\text{He}, {}^3\text{He}){}^9\text{Be}$

Angular distributions of elastically scattered ${}^3\text{He}$ particles have been obtained at $E({}^3\text{He}) = 4, 6, 8, 10, 15$ and 18 MeV (1969PA11), 6.0 and 8.0 MeV (1967EA01), $13.2, 20.4, 22.2$ and 27.0 MeV (1972BU30), 22.7 and 32.3 MeV (1965AR1E), and 217 MeV (1973WI07). See also (1970BA1P). For optical model parameters see discussions in (1972BU30, 1973WI07). At $E({}^3\text{He}) = 39.8$ MeV inelastic ${}^3\text{He}$ groups are observed to ${}^9\text{Be}^*(1.7, 2.4, 3.1, 4.7, 6.8, 14.4)$ (1968BA1E, 1969BA06). See also (1968HO1C, 1968PA1P, 1969HO27; theor.), (1966LA04) and reaction 28 in ${}^6\text{Li}$.

26. (a) ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$

(b) ${}^9\text{Be}(\alpha, 2\alpha){}^5\text{He}$ $Q_m = -2.46$

(c) ${}^9\text{Be}(\alpha, \alpha n){}^8\text{Be}$ $Q_m = -1.6651$

Elastic scattering has been studied at many energies for $E_\alpha = 9.5$ to 48 MeV: see (1966LA04). Recent measurements of the angular distributions of α_0 are reported at $E_\alpha = 8.76, 9.27$ and 10.13 MeV (1967BR39), 28.5 MeV (1967FU08; also α_2 , and partial angular distributions for α_1 and α_3) and 104 MeV (1969HA14, 1970HA1G, 1972DE01, 1972DE02). See (1972DE01, 1972DE02) for a discussion of optical model parameters. The structure $({}^5\text{He} + \alpha)$ for ${}^9\text{Be}$ is found to be much more probable than $({}^6\text{Li} + t)$: the ratio of the spectroscopic factors is about 30 (1972DE01, 1972DE02).

Inelastic groups have been observed to ${}^9\text{Be}^*(1.7, 2.4, 3.1, 6.8, 11.3)$: see (1966LA04). The angular distribution of the group corresponding to ${}^9\text{Be}^*(1.7)$ is consistent with $J^\pi = \frac{1}{2}^+$ (1964LU02: $E_\alpha = 18.4$ MeV). The angular distributions for α_2 are consistent with $l = 2$ ($J^\pi = (\frac{1}{2}, \frac{5}{2}, \frac{7}{2})^-$) (1958SU14: $E_\alpha = 48$ MeV) and (1967FU08: $E_\alpha = 28.5$ MeV). Analysis based on the rotational model leads to a deformation coefficient $\beta_2 = 0.46$ (1959BL31), 0.34 ± 0.01 (1964GR39). See also (1966GR1E, 1966GR1F, 1968LI1E, 1970WO1B, 1972DM01, 1972RA34).

The summed α -spectra from reaction (b) show a peak corresponding to quasi-elastic scattering leaving ${}^5\text{He}$ in the ground state. The angular distribution peaks at the angle corresponding to a zero-momentum α -cluster. The probability of formation of such clusters is $7_{-5}^{+13}\%$ (1969PI11, 1970PI1D: $E_\alpha = 55$ MeV). See also (1965YA02, 1968YA02, 1969DO02, 1969DO03) and ${}^5\text{He}$. The reaction cross section has been measured at $E_\alpha = 42.8$ and 49.2 MeV by (1971GU15).

See also (1965HI1B, 1965KU1B, 1966HI1A, 1967ME1C, 1968BA1H, 1968TA1K, 1969BA1J, 1970MI12, 1972AV04, 1972HI16; theor.) and (1971GA1J).

A study of continuum neutrons at $E_\alpha = 4.9$ to 6.4 MeV (reaction (c)) indicates that sequential decay takes place via ${}^9\text{Be}^*(1.7, 2.4, 3.0)$ (1972OB01). See also (1971GE09, 1973GE1J, 1973WE03).

27. (a) ${}^9\text{Be}({}^6\text{Li}, {}^6\text{Li}){}^9\text{Be}$
 (b) ${}^9\text{Be}({}^7\text{Li}, {}^7\text{Li}){}^9\text{Be}$

Elastic angular distributions have been measured at $E({}^6\text{Li}) = 24$ MeV (1968DA20) and $E({}^7\text{Li}) = 24$ MeV (1972WE08). See also (1970LK1A; theor.) for reaction (b).

28. ${}^9\text{Be}({}^{12}\text{C}, {}^{12}\text{C}){}^9\text{Be}$

Elastic scattering angular distributions have been obtained at $E({}^{12}\text{C}) = 12, 15, 18$ and 21 MeV. The neutron spectroscopic factor for ${}^9\text{Be}$ is 0.84 (1970BA49).

29. ${}^9\text{Be}({}^{14}\text{N}, {}^{14}\text{N}){}^9\text{Be}$

Elastic angular distributions have been measured at $E({}^{14}\text{N}) = 25$ MeV (1966OE1A) and 27.3 MeV (1959HA28). See also (1966LA04), (1969BR1D) and (1965BO37).

30. (a) ${}^9\text{Be}({}^{16}\text{O}, {}^{16}\text{O}){}^9\text{Be}$
 (b) ${}^9\text{Be}({}^{18}\text{O}, {}^{18}\text{O}){}^9\text{Be}$

Elastic angular distributions have been reported at $E(^{16}\text{O}) = 15, 18, 21.5$ and 25 MeV (1970BA49) and 30 MeV (1969KR03), and at $E(^{18}\text{O}) = 12.1, 16$ and 20 MeV (1971KN05).

$$31. \text{}^{10}\text{Be}(\text{d}, \text{t})^9\text{Be} \quad Q_{\text{m}} = -0.5544$$

Forward angular distributions have been obtained at $E_{\text{d}} = 15.0$ MeV for the tritons to $^9\text{Be}^*(0, 1.7, 2.4, 3.1)$. The ground state transition is well fitted by $l = 1$. The transition to $^9\text{Be}^*(1.7)$ [$\Gamma \approx 165 \pm 25$ keV] is consistent with $J^\pi = \frac{1}{2}^+$, that to $^9\text{Be}^*(2.4)$ is quite well fitted with $l = 3$ [$J^\pi = \frac{5}{2}^-$], and that to $^9\text{Be}^*(3.1)$ [$\Gamma = 280 \pm 25$ keV] is consistent with $l = 2$. No other narrow states are seen up to $E_{\text{x}} = 5.5$ MeV (1970AU02).

$$32. \text{}^{10}\text{B}(\gamma, \text{p})^9\text{Be} \quad Q_{\text{m}} = -6.5853$$

See ^{10}B .

$$33. \text{}^{10}\text{B}(\text{e}, \text{ep})^9\text{Be} \quad Q_{\text{m}} = -6.5853$$

See (1969BA1F; theor.) and ^{10}B .

$$34. \text{}^{10}\text{B}(\text{n}, \text{d})^9\text{Be} \quad Q_{\text{m}} = -4.3607$$

A good fit to the angular distributions of the deuterons to $^9\text{Be}^*(0, 2.4)$ has been obtained at $E_{\text{n}} = 14.4$ MeV, using DWBA. The spectroscopic factors are in close agreement with shell-model predictions (1965VA05). See also (1966GO1D), (1966WE1B, 1971MI12), (1966LA04) and ^{11}B in (1975AJ02).

$$35. \text{}^{10}\text{B}(\text{p}, 2\text{p})^9\text{Be} \quad Q_{\text{m}} = -6.5853$$

The summed proton spectrum at $E_{\text{p}} = 460$ MeV yields $Q = -6.7 \pm 0.5, -11.9 \pm 0.5, -17.1 \pm 0.6$ (all $l \neq 0$) and $Q = -30.5 \pm 0.6$ MeV ($l = 0$) (1966TY01). See also (1965BE1E, 1966JA09, 1966JA1A, 1967JA1E, 1967KO1B, 1968JA1G; theor.), ^{10}B and (1966LA04).

$$36. \text{}^{10}\text{B}(\text{d}, \text{}^3\text{He})^9\text{Be} \quad Q_{\text{m}} = -1.0916$$

Angular distributions of the ^3He groups corresponding to $^9\text{Be}^*(0, 2.4)$ have been measured at $E_d = 11.8$ MeV (1966BA21, 1967FI07: $S_{\text{exp}} = 0.76$ and 0.66 , respectively) and at 28 MeV [(1971IN1C); (1968GA13: ground state only)]. The data are very similar, as predicted, to those obtained in the mirror reaction: see reaction 12 in ^9B . See also (1965SY02, 1971WA1K, 1972WA1M).

$$37. \text{}^{10}\text{B}(t, \alpha)^9\text{Be} \quad Q_m = 13.229$$

At $E_t = 12.9$ MeV α groups are observed to the ground state of ^9Be and to excited states at $E_x = 1.75 \pm 0.03$, 2.43 , 3.02 ± 0.04 [$\Gamma = 320 \pm 60$ keV], 11.27 ± 0.04 [$\Gamma = 530 \pm 70$ keV], (14.4) [$\Gamma \approx 800$ keV], 14.39 and 16.67 MeV. The $T = \frac{3}{2}$ state $^9\text{Be}^*(14.39)$ is very weakly populated [$\approx 5\%$ of intensity of α_2]. The angular distribution of the α_2 group shows sharp forward and backward peaking. The α_0 group is not peaked in the backward direction (1968AJ01). A study at $E_t = 1.0$ to 3.2 MeV finds $E_x = 1.750 \pm 0.025$ MeV, $\Gamma = 220 \pm 8$ keV. The angular distribution of the corresponding α -group has been determined at $E_t = 2.5$ MeV (1971GE09).

$$38. \text{}^{10}\text{B}(^{14}\text{N}, ^{15}\text{O})^9\text{Be} \quad Q_m = 0.707$$

The ground state angular distribution has been measured at $E(^{14}\text{N}) = 27.5$ MeV (1962NE01).

$$39. \text{}^{10}\text{B}(^{16}\text{O}, ^{17}\text{F})^9\text{Be} \quad Q_m = -5.985$$

See (1968OK06).

$$40. \text{}^{11}\text{B}(n, t)^9\text{Be} \quad Q_m = -9.5591$$

The angular distribution of the ground state group has been measured at $E_n = 14.4$ MeV (1970MI14).

$$41. (a) \text{}^{11}\text{B}(p, ^3\text{He})^9\text{Be} \quad Q_m = -10.3229$$

$$(b) \text{}^{11}\text{B}(p, \text{pd})^9\text{Be} \quad Q_m = -15.8167$$

At $E_p = 45$ MeV angular distributions are reported for the ^3He ions corresponding to $^9\text{Be}^*(0, 2.4, 11.8, 13.8, 14.39 [T = \frac{3}{2}], 15.96 \pm 0.04 [T = \frac{1}{2}])$. In addition one or more states may be located at $^9\text{Be}^*(15.13)$. It is suggested that $^9\text{Be}^*(11.8, 13.8, 15.96)$ are the $J^\pi = \frac{3}{2}^-; T = \frac{1}{2}$ analogs to $^9\text{B}^*(12.06, 14.01, 16.02)$ (1971HA10). Angular distributions are also reported at $E_p = 40$ MeV (1971KA21; α_0, α_2). The intensity of the group to $^9\text{Be}^*(3.1)$ is $\approx 1\%$ of the ground state group (1971KA21). For reaction (b) see (1964BA1C).

42. (a) $^{11}\text{B}(d, \alpha)^9\text{Be}$ $Q_m = 8.0309$
 (b) $^{11}\text{B}(d, n\alpha)^4\text{He}^4\text{He}$ $Q_m = 6.4577$
 $Q_0 = 8.0297 \pm 0.0028$ (1967OD01). See also
 (1964MA57, 1967SP09).

Alpha groups are reported corresponding to $^9\text{Be}^*(0, 1.7, 2.4, 3.1)$. The width of $^9\text{Be}^*(1.7)$: $\Gamma_{\text{cm}} = 224 \pm 25$ keV (1958KA31, 1966PU02). The energy of $^9\text{Be}^*(2.4)$ is 2422 ± 5 keV (1951VA08), 2431 ± 6 keV (1954EL10), 2424 ± 5 keV (1956BO18). The $\frac{5}{2}^+$ state is at $E_x = 3.02 \pm 0.03$ MeV (1955LE36), 3.05 ± 0.03 MeV (1956BO18); $\Gamma_{\text{cm}} = 257 \pm 25$ keV (1958KA31, 1966PU02).

Angular distributions are reported at $E_d = 0.39$ to 0.7 MeV (1965SA15; α_0, α_2), 0.8 to 2.5 MeV (1968CO31; α_0, α_2) and at 12.6 MeV (1966DR04; α_0, α_2). See also (1966ME1E; theor.).

The ratio of the γ -decay width to the total width, Γ_γ/Γ , of $^9\text{Be}^*(2.4)$ is $(1.16 \pm 0.14) \times 10^{-4}$. Since Γ_γ is known from (e, e') [see Table 9.8], $\Gamma = 1.03 \pm 0.18$ keV. For $^9\text{Be}^*(1.7)$, $\Gamma_\gamma/\Gamma \leq 2.4 \times 10^{-5}$ (1966PU02).

Reaction (b), at $E_d = 10.4$ and 12.0 MeV, proceeds via $^9\text{Be}^*(2.4)$ and to some extent via $^9\text{Be}^*(3.1, 4.7)$ and possibly some higher excited states. The dominant decay of $^9\text{Be}^*(2.4)$ is to $^5\text{He}(0) + \alpha$ while $^9\text{Be}^*(3.1, 4.7)$ decay to $^9\text{Be}(0) + n$ (1971RE19). It should be noted, however, that the peaks corresponding to $^9\text{Be}^*(3.1)$ have a FWHM of ≈ 1 MeV, which may imply that $^9\text{Be}^*(2.8)$ is involved (1971RE19). See also ^8Be , ^{13}C in (1976AJ04) and (1966LA04).

43. $^{11}\text{B}(^{16}\text{O}, ^{18}\text{F})^9\text{Be}$ $Q_m = -8.290$

See (1968OK06).

44. (a) $^{12}\text{C}(n, \alpha)^9\text{Be}$ $Q_m = -5.7016$
 (b) $^{12}\text{C}(n, n\alpha)^4\text{He}^4\text{He}$ $Q_m = -7.2748$

Angular distributions of the α_0 group have been measured at $E_n = 14.1$ MeV (1969HS02, 1969KI02), 13.9 and 15.6 MeV (1968BR21), and 14.8 to 18.8 MeV (1971SA31). (1968BR21,

1969HS02) also report the population of ${}^9\text{Be}^*(1.7, 2.4, 3.1)$. Reaction (b) at $E_n = 13$ to 18 MeV involves ${}^9\text{Be}^*(2.4)$ (1966MO05). See also (1965MO09, 1966MI1D, 1968BE1J, 1969KA1D, 1969LO1D, 1971DO1K, 1971FA04), (1966CI1A, 1966LA04), (1967EL1D, 1968CH1J; theor.), ${}^{12}\text{C}$ in (1975AJ02), and ${}^{13}\text{C}$ in (1976AJ04).

45. ${}^{12}\text{C}(\alpha, {}^7\text{Be}){}^9\text{Be}$ $Q_m = -24.694$

Angular distributions have been obtained for the transitions to ${}^7\text{Be}^*(0, 0.43) + {}^9\text{Be}(0)$ (1972RU03; 42 MeV). See also (1971LE33).

46. ${}^{13}\text{C}(\gamma, \alpha){}^9\text{Be}$ $Q_m = -10.6480$

See ${}^{13}\text{C}$ in (1976AJ04).

47. ${}^{13}\text{C}(p, p\alpha){}^9\text{Be}$ $Q_m = -10.6480$

See (1971BR07).

48. ${}^{13}\text{C}(d, {}^6\text{Li}){}^9\text{Be}$ $Q_m = -9.174$

A partial angular distribution involving the transition to ${}^6\text{Li}(0) + {}^9\text{Be}(0)$ has been measured at $E_d = 14.6$ MeV (1966DE09).

49. ${}^{16}\text{O}(\alpha, {}^{11}\text{C}){}^9\text{Be}$ $Q_m = -15.387$

The angular distribution involving the transition to ${}^{11}\text{C}(0) + {}^9\text{Be}(0)$ has been measured at $E_\alpha = 42$ MeV (1972RU03).

⁹B

(Figs. 17 and 18)

GENERAL: (See also (1966LA04).)

Model calculations: (1966BA26, 1966EL08, 1967ST1C, 1971CO28, 1972LE1L, 1973HA49).

Special levels: (1966BA26, 1966EL08, 1967BA59, 1967ST1C, 1969HA1G, 1970TO1E, 1971CO28, 1971LI30, 1972BE1E).

Astrophysical questions: (1970BA1M).

Other topics: (1967CA17, 1967CH1H, 1970SA05, 1972AN05, 1972HA57, 1972CA37, 1972LE1L, 1972PN1A, 1973JU2A).

Ground state properties: (1966BA26, 1966EL08, 1968PE16, 1969HE1N, 1969JA1M, 1969LE1D, 1971AU1G, 1972LE1L).

| | | |
|--|----------------|----------------|
| 1. (a) ${}^6\text{Li}({}^3\text{He}, n){}^8\text{B}$ | $Q_m = -1.975$ | $E_b = 16.604$ |
| (b) ${}^6\text{Li}({}^3\text{He}, p){}^8\text{Be}$ | $Q_m = 16.788$ | |
| (c) ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$ | $Q_m = 0.1126$ | |
| (d) ${}^6\text{Li}({}^3\text{He}, t){}^6\text{Be}$ | $Q_m = -4.306$ | |
| (e) ${}^6\text{Li}({}^3\text{He}, {}^3\text{He}){}^6\text{Li}$ | | |

The total cross section for reaction (a) has been measured from threshold to $E({}^3\text{He}) = 3.5$ MeV: it increases monotonically reaching the value 4.3 mb at 3.5 MeV (1973MCZW). No structure is observed in the n_0 excitation curve for $E({}^3\text{He}) = 4.0$ to 5.7 MeV (1967VA24). The yield of ${}^8\text{B}$ has also been measured for $E({}^3\text{He}) = 8.9$ to 26.5 MeV (1973MA24). See also (1966FA1A) and ${}^8\text{B}$.

The excitation functions for protons leading to ${}^8\text{Be}^*(0, 2.9)$ [p_0, p_1] have been measured for $E({}^3\text{He}) = 0.9$ to 17 MeV (reaction (b)). Resonances are reported at $E({}^3\text{He}) = 1.6$ MeV ($\Gamma = 0.25$ MeV) [1.68 MeV: (1969VI05). $J^\pi = \frac{3}{2}^-$ or $\frac{5}{2}^-$] and 3.0 MeV ($\Gamma = 1.5$ MeV) (1956SC01). Above 5 MeV, the p_0 yield at 0° increases monotonically with energy to $E({}^3\text{He}) = 17$ MeV (1965FL03). Polarization measurements are reported at $E({}^3\text{He}) = 1.4$ to 2.0 MeV (1966SI1C, 1971SI1J; p_0, p_1). See also (1970GA1G) and ${}^8\text{Be}$.

The yields of 0.43 and 0.48 MeV γ -rays (reaction (c)), measured for $E({}^3\text{He}) = 0.5$ to 1.3 MeV, are reported to show the excitation of ${}^9\text{B}^*(17.20 \pm 0.02)$ with $\Gamma = 110 \pm 30$ keV, $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+$; $T = \frac{1}{2}$ (1970AL25). See also ${}^7\text{Be}$.

Excitation functions for ground state tritons (reaction (d)) have been measured for $E({}^3\text{He}) = 10$ to 16 MeV (1969NU1A) and 23.3 to 25.4 MeV (1972GI07). See also ${}^6\text{Be}$. Differential cross sections have been measured at several angles for the ${}^3\text{He}$ groups to ${}^6\text{Li}^*(0, 2.19)$ (reaction (e)) for $E({}^3\text{He}) = 23.3$ to 25.4 MeV (1972GI07). See also ${}^6\text{Li}$.

Table 9.9: Energy levels of ${}^9\text{B}$

| E_x (MeV \pm keV) | $J^\pi; T$ | $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|---|------------------------------|-----------------------|---|
| g.s. | $\frac{3}{2}^-; \frac{1}{2}$ | 0.54 ± 0.21 | p, α | 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 |
| (1.6) | | ≈ 700 | (p, α) | 13 |
| 2.361 ± 5 | $\frac{5}{2}^-; \frac{1}{2}$ | 81 ± 5 | α | 2, 4, 6, 7, 8, 11, 12, 13, 15, 16 |
| 2.788 ± 30 | $(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$ | 550 ± 40 | p | 4, 6, 11, 13, 16 |
| (4.8 ± 100) | | 1000 ± 200 | | 4, 9 |
| 6.97 ± 60 | $\frac{7}{2}^-; \frac{1}{2}$ | 2000 ± 200 | p | 4, 6, 9, 11, 15, 16 |
| 11.75 ± 100 | $(\frac{7}{2})^-; \frac{1}{2}$ | 800 ± 50 | p | 9, 11, 13 |
| 12.06 ± 60 | $;$ $\frac{1}{2}$ | 800 ± 200 | p | 4, 9, 15 |
| 14.01 ± 70 | $;$ $\frac{1}{2}$ | 390 ± 110 | | 4, 15 |
| 14.659 ± 5 | $\frac{3}{2}^-; \frac{3}{2}$ | $0.26^{+0.09}_{-0.12}$ | γ , p | 4, 7, 15 |
| 14.7 ± 200 | $(\frac{5}{2})^-; \frac{1}{2}$ | 1350 ± 200 | | 11 |
| 15.29 ± 40 | $;$ $\frac{1}{2}$ | | | 15 |
| 15.58 ± 40 | $;$ $\frac{1}{2}$ | | | 15 |
| 16.024 ± 25 | $;$ $(\frac{1}{2})$ | 180 ± 16 | | 4, 15 |
| 17.190 ± 25 | | 120 ± 40 | p, d, ${}^3\text{He}$ | 1, 4, 5, 15 |
| 17.637 ± 10 | | 71 ± 8 | p, d, ${}^3\text{He}$ | 1, 4, 5, 15 |
| (18.6) | | 1000 | p, ${}^3\text{He}$ | 1, 11 |

 2. ${}^6\text{Li}(\alpha, n){}^9\text{B}$

$$Q_m = -3.975$$

Angular distributions at $E_\alpha = 8.0, 10.0, 12.0$ and 14.0 MeV all display strong forward peaking (1963ME08). At $E_\alpha = 14.4$ MeV, neutron groups are observed to ${}^9\text{Be}^*(0, 2.4)$: the upper limit of the cross section to a state at ≈ 1.7 MeV is $100 \mu\text{b/sr}$ or < 0.1 of the ground state group (1964BA29). See also ${}^{10}\text{B}$.

 3. ${}^6\text{Li}({}^6\text{Li}, t){}^9\text{B}$

$$Q_m = 0.809$$

Angular distributions of the t_0 group have been measured for $E({}^6\text{Li}) = 4.0$ to 5.5 MeV and at 7.35 and 9.0 MeV. No evidence was observed for a group corresponding to ${}^9\text{B}^*(1.6)$ (1966KI09).

4. (a) ${}^7\text{Li}({}^3\text{He}, n){}^9\text{B}$ $Q_m = 9.353$
 (b) ${}^7\text{Li}({}^3\text{He}, np){}^8\text{Be}$ $Q_m = 9.538$

For $E({}^3\text{He})$ to 12.5 MeV this reaction populates ${}^9\text{B}^*(0, 2.4, 2.8, (7.0))$ (1963DU12) and levels at 12.06 ± 0.06 [0.8 ± 0.2], 14.01 ± 0.07 [0.39 ± 0.11], 14.670 ± 0.016 [< 0.045], 16.024 ± 0.025 [0.180 ± 0.016], 17.19 and 17.63 MeV [widths in brackets] (1964DI1A, 1965DI03). (1967BA59) report $Q = (-5306 \pm 5)$ keV and therefore $E_x = 14.659 \pm 0.005$ MeV (based on Q_m).

${}^9\text{B}^*(14.66)$ is the first $T = \frac{3}{2}$ state in ${}^9\text{B}$ (1965DI03). It γ -decays to ${}^9\text{B}^*(0, 2.4, 2.8)$ with branching ratios which appear to be similar to those for the analog state in ${}^9\text{Be}$: see Table 9.6. Assuming that the Γ_γ for both $T = \frac{3}{2}$ states in ${}^9\text{Be}$, ${}^9\text{B}$ are the same, the total Γ for ${}^9\text{B}^*(14.66)$ is then $\approx 80\%$ of that for ${}^9\text{Be}^*(14.40)$: 260_{-120}^{+90} eV (1971AD01) [corrected for revised value of Γ of ${}^9\text{Be}^*(14.40)$]. The ratio $\Gamma_{p_0}/\Gamma_{\gamma_0} < 1.5$, $\Gamma_{p_1}/\Gamma_{\gamma_0} = 13.9 \pm 2.1$ [Γ_{p_0} is the width for decay of ${}^9\text{B}^*(14.66)$ to the ground state of ${}^8\text{Be}$; Γ_{p_1} is that for decay to ${}^8\text{Be}^*(2.9)$]. Assuming Γ_{γ_0} for ${}^9\text{B}^*(14.66) = 6.9 \pm 0.5$ eV [the value for the analog state in ${}^9\text{Be}$: see Table 9.6], $\Gamma_{p_0} < 10$ eV, $\Gamma_{p_1} = 96 \pm 20$ eV (1972AD04). See also (1971AD1C).

Angular distributions have been reported for $E({}^3\text{He}) = 1.56$ to 5.27 MeV (1966DI04; n_0) and 3.1 MeV (1970GU08; n_0). (1970GU08) find that a structure in the neutron spectrum which might correspond to ${}^9\text{B}^*(1.5)$ is really due to sequential decay of ${}^9\text{Be}^*(11.8)$ to ${}^8\text{Be}(0)$. They also report a ${}^9\text{B}$ state with $E_x = 4.8 \pm 0.1$ MeV, $\Gamma = 1.0 \pm 0.2$ MeV. See also (1970LI1Q), (1969BA1Z) and ${}^{10}\text{B}$.

5. ${}^7\text{Be}(d, p){}^8\text{Be}$ $Q_m = 16.676$ $E_b = 16.491$

For $E_d = 0.75$ to 1.70 MeV, resonances in the yields of protons are observed at $E_d = 0.900 \pm 0.025$ MeV ($p_0, p_{2.4}$) and 1.475 ± 0.010 MeV ($p_{2.4}$ only) with $\Gamma_{\text{cm}} = 120 \pm 40$ and 71 ± 8 keV, respectively [${}^9\text{B}^* = 17.19$ and 17.64 MeV] (1960KA17). See (1972PA1C) for astrophysical considerations.

6. ${}^9\text{Be}(p, n){}^9\text{B}$ $Q_m = -1.8498$

A high resolution experiment at $E_p = 20$ MeV shows the population of ${}^9\text{B}^*(0, 2.4)$ and is consistent with the excitation of ${}^9\text{B}^*(2.8)$ [$\Gamma \approx 0.3$ MeV] and ${}^9\text{B}^*(7.0)$ [$\Gamma > 1$ MeV]. No other states are excited for $E_x < 7.1$ MeV (1970AN07). (1972AR22) report $E_x = 3.09 \pm 0.10$ MeV. Additional states have been reported by a number of groups: see (1966LA04) for earlier references and (1967SL04, 1970CL01). The width of the ground state is 540 ± 210 eV (1964TE01).

Angular distributions have been measured at $E_p = 3.5$ to 10.9 MeV (1965WA04), 6.8 MeV (1967DR08), 8 to 14 MeV (1960SA03), 18.5 MeV (1964AN1B) and 30.3 and 49.3 MeV (1970CL01). See also (1967BO1D, 1969JU1A, 1969VE02, 1970WI1B, 1971BE46, 1971CA1F, 1972CA1Q, 1973WA28), (1966LA04) and (1966PA1H, 1968TH1H; theor.).

7. ${}^9\text{Be}({}^3\text{He}, t){}^9\text{B}$ $Q_m = -1.0860$

Angular distributions have been reported at $E({}^3\text{He}) = 3.0$ to 3.8 MeV (1969OR01; t_0), 5.0 to 9.0 MeV (1967EA01; t_0), 5.7 MeV (1959HI69; t_0), 10 MeV (1967CR04; $t_0, t_{2.4}$), 20 to 27.8 MeV (1969OP1A, 1970OP1B; $t_0, t_{2.4}$) and 25 MeV (1960WE04; $t_0, t_{2.4}$). At $E({}^3\text{He}) = 39.8$ MeV ${}^9\text{Be}(0)$ is very strongly excited and ${}^9\text{B}^*(2.4, 14.7)$ are also observed. There is some indication that other known ${}^9\text{B}$ states are also populated (1969BA06). See also ${}^{12}\text{C}$ in (1975AJ02) and (1970CA28).

8. ${}^9\text{Be}({}^6\text{Li}, {}^6\text{He}){}^9\text{B}$ $Q_m = -4.577$

At $E({}^6\text{Li}) = 30.8$ and 31.8 MeV the ground state of ${}^9\text{B}$ is strongly excited. ${}^9\text{B}^*(2.4)$ is also observed (1970CH19, 1971CH1B). A partial angular distribution for the ground state transition is reported at the higher energy by (1971CH1B).

9. ${}^9\text{C}(\beta^+){}^9\text{B}^* \rightarrow {}^9\text{Be} + \text{p}$ $Q_m = 16.677$
 $\rightarrow {}^5\text{Li} + \alpha$ $Q_m = 14.80$

Several groups of delayed protons are observed indicating the involvement of a number of ${}^9\text{B}$ states: see Table 9.10 (1972ES05). It is not possible to determine ft values since some of the ${}^9\text{B}$ states involved in the ${}^9\text{C}$ decay may decay via ${}^5\text{Li} + \alpha$: see (1972ES05).

10. ${}^{10}\text{B}(\gamma, n){}^9\text{B}$ $Q_m = -8.435$

See ${}^{10}\text{B}$.

11. ${}^{10}\text{B}(\text{p}, \text{d}){}^9\text{B}$ $Q_m = -6.211$

At $E_p = 33.6$ MeV (1968KU04, 1970KU1D) and 155.6 MeV (1969BA05) deuteron groups are observed to ${}^9\text{B}^*(0, 2.4, 7.0, 11.7, 14.7)$. All have angular distributions characteristic of $l_n = 1$ and therefore odd parity and $J = (\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2})$: see Table 9.11. There is some evidence also for the population of ${}^9\text{B}^*(2.8, 18.4)$ (1968KU04, 1969BA05). Angular distributions have also been reported at $E_p = 49.5$ MeV (1970SQ01; $d_0, d_{2.4}, d_{11.7}$). See also (1968KR02), (1966LA04) and (1969TO1A; theor.).

Table 9.10: Delayed protons following the β^+ decay of ${}^9\text{C}$ ^a

| E_p (c.m.) (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | Corresponding state in ${}^9\text{B}$ (MeV) | |
|-------------------------------|------------------------------|--|---------------------------------------|
| | | if decay is to ${}^8\text{Be}_{\text{g.s.}}$ | if decay is to ${}^8\text{Be}^*(2.9)$ |
| 3.45 ± 0.25 | 200 ± 100 | 3.26 ± 0.25 ^d | ^c |
| (4.2 ± 0.3) | 1000 ± 200 | 4.0 ± 0.3 | 6.9 ± 0.3 |
| (5.0 ± 0.2) | 400 ± 200 | 4.8 ± 0.2 | ^c |
| 6.10 ± 0.10 | 400 ± 100 | 5.91 ± 0.10 | ^c |
| 9.28 ± 0.24 ^b | 1800 ± 200 | 9.09 ± 0.24 | 11.99 ± 0.24 |
| 12.30 ± 0.10 ^b | 450 ± 100 | 12.11 ± 0.10 | ^c |

^a (1972ES05). See also (1965HA09).

^b Ratio of the intensities $I_{9.28}/I_{12.30} = 1.2 \pm 0.2$.

^c The relatively narrow width of the proton group does not permit this option.

^d By analogy with the ${}^9\text{Li}$ decay, this decay may involve a $J^\pi = \frac{1}{2}^-$ analog of ${}^9\text{Be}^*(2.78)$. Such a state in ${}^9\text{B}$ has not been reported in any other reaction.

Table 9.11: Levels of ${}^9\text{B}$ from ${}^{10}\text{B}(p, d){}^9\text{B}$

| (1969BA05) ^a | | | (1968KU04) ^b | | | |
|-----------------------------|-------|---------------------------------|------------------------------|----------------------------|-------|----------------------|
| E_x (MeV) | l_n | F_{exp}^2 ^c | E_x (MeV) | Γ_{cm} (MeV) | l_n | J^π ^d |
| 0 | 1 | 0.44 | 0 | | 1 | $\frac{3}{2}^-$ |
| 2.4 ± 0.1 | 1 | 0.60 | 2.35 ± 0.02 | | 1 | $\frac{5}{2}^-$ |
| | | | (2.8) ^g | | | |
| 7.1 ± 0.2 ^e | 1 | 0.52 | 7.1 ± 0.2 | 1.95 ± 0.2 | 1 | $\frac{7}{2}^-$ |
| 11.5 ± 0.2 | 1 | 1.12 | 11.75 ± 0.1 ^h | 0.80 ± 0.05 | 1 | $(\frac{7}{2})^-$ |
| 14.9 ± 0.3 ^f | 1 | 0.32 | 14.6 ± 0.2 ^g | 1.35 ± 0.2 | (1) | $(\frac{5}{2})^-$ |
| (18.4) | | | | | | |

^a (1969BA05): $E_p = 155.6$ MeV.

^b (1968KU04, 1970KU1D): $E_p = 33.6$ MeV.

^c Spectroscopic factor.

^d J from best fit to theoretical spectroscopic factor.

^e $\Gamma = 2.4 \pm 0.2$ MeV.

^f $T = \frac{1}{2}$.

^g Weak group.

^h 11.66 ± 0.10 MeV (1970SQ01).

$$12. \text{}^{10}\text{B}(\text{d}, \text{t})\text{}^9\text{B} \quad Q_{\text{m}} = -2.178$$

$$Q_0 = -2.189 \pm 0.010 \text{ (1967SP09)}.$$

Angular distributions have been measured at $E_{\text{d}} = 11.8$ MeV (1966BA21, 1967FI07; t_0 , $t_{2.4}$) [S for ${}^9\text{B}^*(0, 2.4) = 0.80$ and 0.64 , respectively], 13.5 MeV (1964FU15; t_0) and 28 MeV [(1968GA13; t_0); (1971IN1C; t_0 , $t_{2.4}$ (partial))]. See also (1971WA1K, 1972WA1M) and (1966BA1X, 1971BO50; theor.).

$$13. \text{ (a) } {}^{10}\text{B}({}^3\text{He}, \alpha){}^9\text{B} \quad Q_{\text{m}} = 12.143$$

$$\text{ (b) } {}^{10}\text{B}({}^3\text{He}, \alpha\text{p}){}^8\text{Be} \quad Q_{\text{m}} = 12.328$$

Alpha particle spectra show the excitation of ${}^9\text{B}^*(0, 2.4, 2.8, 11.8)$: see (1966LA04). Measurements by (1968KR02) determine $E_{\text{x}} = 2.361 \pm 0.005$ and 2.788 ± 0.030 MeV, $\Gamma = 81 \pm 5$ and 548 ± 40 keV, respectively [see Table 9.11 in (1966LA04) for other values]. There is some evidence for a state with $E_{\text{x}} \approx 1.6$ MeV, $\Gamma \approx 0.7$ MeV, but it is not conclusive, in agreement with the older work [see (1960SP08, 1962BA1C)]. No evidence is found for any narrow levels in ${}^9\text{B}$ with $\Gamma \leq 100$ keV and $4 < E_{\text{x}} < 7$ MeV: the upper limit to the intensity of the corresponding α -group is 1% of the intensity of the group to ${}^9\text{B}^*(2.4)$ (1968KR02). Angular distributions have been determined at $E({}^3\text{He}) = 5.5$ MeV (1966CA02; α_0) and 33.7 MeV (1971SQ03; α_0 , $\alpha_{2.4}$, $\alpha_{11.8}$). DWBA does not seem to give a good description of the transition to ${}^9\text{B}^*(11.8)$ (1971SQ03).

In reaction (b) study of the decays of ${}^9\text{B}^*(2.4, 2.8)$ shows that ${}^9\text{B}^*(2.4)$ decays $< 0.5\%$ by proton emission to ${}^8\text{Be}(0)$ [$\theta_{\text{f}}^2 < 5.1 \times 10^{-3}$] [it decays to ${}^5\text{Li}(0)$ by α -emission] while the second state, $E_{\text{x}} = 2.71 \pm 0.03$ MeV [$\Gamma = 0.71 \pm 0.06$ MeV] decays almost 100% by that channel [$\theta_{\text{d}}^2 = 0.74$] (1966WI08). (1966WA16) find $\Gamma = 1.1 \pm 0.2$ MeV for ${}^9\text{B}^*(2.8)$ and suggest $J = \frac{1}{2}$ for this state [see, however, (1960SP08)]. No other excited states of ${}^9\text{B}$ with $3.5 < E_{\text{x}} < 9.5$ MeV decay by proton emission to ${}^8\text{Be}(0)$ (1968KR02). See also (1970BE1F, 1971FO1E) and (1968TA1M; theor.). See also ${}^8\text{Be}$, and ${}^{13}\text{N}$ in (1976AJ04).

$$14. \text{}^{10}\text{B}({}^{16}\text{O}, \text{}^{17}\text{O}){}^9\text{B} \quad Q_{\text{m}} = -4.293$$

See (1968OK06).

$$15. \text{}^{11}\text{B}(\text{p}, \text{t}){}^9\text{B} \quad Q_{\text{m}} = -11.409$$

At $E_{\text{p}} = 45$ MeV angular distributions have been obtained for the triton groups to ${}^9\text{B}^*(0, 2.36, 12.06, 14.01, 14.66, 16.02)$. In addition the spectra show some indication of the groups corresponding to ${}^9\text{B}^*(7.0, 17.19, 17.63)$. New $T = \frac{1}{2}$ states are reported at $E_{\text{x}} = 15.29 \pm 0.04$ and 15.58 ± 0.04 MeV (1971HA10). See also (1971KA04; theor.) and reaction 41 in ${}^9\text{Be}$.

16. (a) $^{12}\text{C}(p, \alpha)^9\text{B}$ $Q_m = -7.551$
 (b) $^{12}\text{C}(p, p)^4\text{He}^4\text{He}^4\text{He}$ $Q_m = -7.2748$

Angular distributions of ground state α -particles have been measured at $E_p = 14.0$ to 17.2 MeV ([1969KO1D](#)), 19.1 to 44 MeV ([1967AC01](#), [1969GA03](#), [1970GU06](#), [1971GU23](#)), 38.5 , 41.6 and 44.5 MeV [also partial back-angle measurements at $E_p = 30.5$, 32.0 , 33.0 , 34.5 and 39.5 MeV] ([1966CR05](#), [1967CR05](#), [1968LI1E](#)) and at $E_p = 54.1$ MeV ([1972MA21](#)). Alpha groups are also observed to $^9\text{B}^*(2.3, 2.9 \pm 0.2, 6.97 \pm 0.06)$: see ([1955RE16](#), [1962MA40](#), [1964BA29](#), [1972MA21](#)). See also ([1966LA04](#)). The angular distribution to $^9\text{B}^*(6.97)$ is consistent with $J^\pi = \frac{7}{2}^-$; $\Gamma \approx 2$ MeV ([1972MA21](#): $E_p = 54.1$ MeV). For reaction (b) see ([1972MA62](#)) and ([1966LA04](#)). See also ^{12}C in ([1975AJ02](#)) and ^{13}N in ([1976AJ04](#)).

17. $^{12}\text{C}(^3\text{He}, ^6\text{Li})^9\text{B}$ $Q_m = -11.571$

Angular distributions of ^6Li ions have been obtained at $E(^3\text{He}) = 28$ MeV ([1971KL1E](#)), 30.0 and 40.7 MeV ([1972OH01](#)) and 35.7 MeV ([1969ZE1A](#), [1970FO1D](#)).

${}^9\text{C}$
(Figs. 17 and 18)


GENERAL: (See also (1966LA04).)

Model calculations: (1966BA26).

Other topics: (1966BA26, 1966MC1C, 1972AN05, 1972CA37, 1973LA19).

Ground state properties, including theoretical mass predictions: (1965GO1D, 1966BA26, 1966GO1B, 1966KE16, 1969GA1P, 1969JA1M, 1972CE1A, 1973HA77).

Mass of ${}^9\text{C}$: From the threshold energy of ${}^7\text{Be}({}^3\text{He}, \text{n}){}^9\text{C}$ (1971MO01) the atomic mass excess of ${}^9\text{C}$ is 28.908 ± 0.004 MeV. This value is in good agreement with that obtained from the Q -value of ${}^{12}\text{C}({}^3\text{He}, {}^6\text{He}){}^9\text{C}$ [28.912 ± 0.009 MeV]. See also (1967BA59).



The half-life of ${}^9\text{C}$ is 126.5 ± 1.0 msec (1971HA05, 1972ES05), 126.5 ± 2 msec (1971MO01). Several groups of delayed protons are observed indicating the involvement of a number of ${}^9\text{B}$ states: see Table 9.10 (1972ES05). See also (1965HA09). See also (1966BA26, 1971WI18, 1972WI28, 1972WI1C, 1973TO14; theor.).



$E_{\text{thresh.}} = 8980 \pm 5$ keV (1971MO01). See also (1967BA59).



See (1965RO1G).



See (1965HA09).



Table 9.12: Energy levels of ${}^9\text{C}$

| E_x (MeV) | $J^\pi; T$ | $\tau_{1/2}$ (msec) | Decay | Reactions |
|-------------|--------------------------------|---------------------|-----------|------------------|
| g.s. | $(\frac{3}{2}^-); \frac{3}{2}$ | 126.5 ± 0.9 | β^+ | 1, 2, 3, 4, 5, 6 |

See (1965HA09).

6. ${}^{12}\text{C}({}^3\text{He}, {}^6\text{He}){}^9\text{C}$ $Q_m = -31.574$

$Q_0 = -31.578 \pm 0.008$ MeV (1970TR05, 1970TR1F, 1971TR03). See also (1964CE04).

References

(Closed December 31, 1973)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1951BR72 C.P. Browne, R.M. Williamson, D.S. Craig and D.J. Donahue, Phys. Rev. 83 (1951) 179
- 1951GA30 W.L. Gardner, N. Knable and B.J. Moyer, Phys. Rev. 83 (1951) 1054
- 1951VA08 D.M. Van Patter, A. Sperduto, K. Huang, E.N. Strait and W.W. Buechner, Phys. Rev. 81 (1951) 233
- 1952BA64 L.M. Baggett and S.J. Bame, Jr., Phys. Rev. 85 (1952) 434
- 1954BA46 S. Bashkin, Phys. Rev. 95 (1954) 1012
- 1954EL10 R.B. Elliott and D.J. Livesey, Proc. Roy. Soc. A224 (1954) 129
- 1955GO48 C.R. Gossett, G.C. Phillips, J.P. Schiffer and P.M. Windham, Phys. Rev. 100 (1955) 203
- 1955LE36 L.L. Lee, Jr. and D.R. Inglis, Phys. Rev. 99 (1955) 96
- 1955MA20 R.L. Macklin and H.E. Banta, Phys. Rev. 97 (1955) 753
- 1955RE16 J.B. Reynolds, Phys. Rev. 98 (1955) 1289
- 1956BE14 J. Benveniste, R.G. Finke and E.A. Martinelli, Phys. Rev. 101 (1956) 655
- 1956BE1A Bezrukov, Panov and Timoshuk, Sov. J. Nucl. Energy 4 (1956) 609
- 1956BO18 C.K. Bockelman, A. Leveque and W.W. Buechner, Phys. Rev. 104 (1956) 456
- 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., Phys. Rev. 104 (1956) 1064
- 1957HU14 P. Huber and R. Wagner, Helv. Phys. Acta 30 (1957) 257
- 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809
- 1958AN32 J.D. Anderson, C.C. Gardner, J.W. McClure, M.P. Nakada and C. Wong, Phys. Rev. 111 (1958) 572
- 1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503
- 1958NA09 M.P. Nakada, J.D. Anderson, C.C. Gardner and C. Wong, Phys. Rev. 110 (1958) 1439
- 1958SU14 R.G. Summers-Gill, Phys. Rev. 109 (1958) 1591
- 1958WA05 R. Wagner and P. Huber, Helv. Phys. Acta 31 (1958) 89

- 1959AJ77 F. Ajzenberg-Selove, N. Jarmie and E. Haddad, Bull. Amer. Phys. Soc. 4 (1959) 258, NA9
- 1959BL31 J.S. Blair, Phys. Rev. 115 (1959) 928
- 1959HA28 M.L. Halbert and A. Zucker, Phys. Rev. 115 (1959) 1635
- 1959HI69 S. Hinds and R. Middleton, Proc. Phys. Soc. A74 (1959) 196
- 1959MA34 J.B. Marion, J.S. Levin and L. Cranberg, Phys. Rev. 114 (1959) 1584
- 1960BA47 W.C. Barber, F. Berthold, G. Fricke and F.E. Gudden, Phys. Rev. 120 (1960) 2081
- 1960BU25 J. Button and R. Mermod, Phys. Rev. 118 (1960) 1333
- 1960KA05 R.W. Kavanagh, Nucl. Phys. 15 (1960) 411
- 1960KA17 R.W. Kavanagh, Nucl. Phys. 18 (1960) 492
- 1960KL03 P.R. Klein, N. Cindro, L.W. Swenson and N.S. Wall, Nucl. Phys. 16 (1960) 374
- 1960MA15 K.V. Makariunas and S.V. Starodubtsev, Zh. Eksp. Teor. Fiz. 38 (1960) 372; Sov. Phys. JETP 11 (1960) 271
- 1960SA03 Y. Saji, J. Phys. Soc. Jpn. 15 (1960) 367
- 1960SE12 R. Seltz and D. Magnac-Valette, Compt. Rend. 251 (1960) 2006
- 1960SP08 R.R. Spencer, G.C. Phillips and T.E. Young, Nucl. Phys. 21 (1960) 310
- 1960WE04 H.E. Wegner and W.S. Hall, Phys. Rev. 119 (1960) 1654
- 1961KO1A Koltun, Phys. Rev. 124 (1961) 1162
- 1961VA43 A.K. Val-ter, P.I. Vatsset, L.Y. Kolesnikov, S.G. Tonapetyan, K.K. Chernyavskii and A.I. Shpetnyi, Atomn. Energ. (USSR) 10 (1961) 577; Sov. J. At. Energy 10 (1962) 574; Correction At. Energy 123 (2018) 432
- 1962BA1C Barker and Treacy, Nucl. Phys. 38 (1962) 33
- 1962CH14 L.F. Chase, Jr., R.G. Johnson, F.J. Vaughn and E.K. Warburton, Phys. Rev. 127 (1962) 859
- 1962CL06 F.M. Clikeman, A.J. Bureau and M.G. Stewart, Phys. Rev. 126 (1962) 1822
- 1962ED02 R.D. Edge and G.A. Peterson, Phys. Rev. 128 (1962) 2750
- 1962KO13 M.P. Konstantinova, E.V. Myakinin, A.M. Petrov and A.N. Ronsnov, Zh. Eksp. Teor. Fiz. 43 (1962) 388; Sov. Phys. JETP 16 (1963) 278
- 1962MA40 D.R. Maxson, Phys. Rev. 128 (1962) 1321
- 1962NE01 E. Newman, Phys. Rev. 125 (1962) 600
- 1962SE1A Serov and Guzhovskii, Atomn. Energ. (USSR) 12 (1962) 5
- 1963AL18 D.E. Alburger, Phys. Rev. 132 (1963) 328
- 1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336

1963JE05 H. Jeremie, Nucl. Phys. 47 (1963) 225
 1963ME08 M.K. Mehta, W.E. Hunt, H.S. Plendl and R.H. Davis, Nucl. Phys. 48 (1963) 90
 1963NE07 B.M.K. Nefkens, Phys. Rev. Lett. 10 (1963) 243
 1963NG01 H. Nguyen Ngoc, N. Hors and J. Perez y Jorba, Nucl. Phys. 42 (1963) 62
 1963PA04 P. Paul and D. Kohler, Phys. Rev. 129 (1963) 2698
 1963ST1A Strzalkowski, Nukleonika 8 (1963) 301
 1964AN1B Anderson, Wong, McClure and Walker, Phys. Rev. 136 (1964) B118
 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
 1964BA29 R.W. Bauer, J.D. Anderson and C. Wong, Nucl. Phys. 56 (1964) 117
 1964CE04 J. Cerny, R.H. Pehl, F.S. Goulding and D.A. Landis, Phys. Rev. Lett. 13 (1964) 726
 1964DI1A Dietrich, Thesis, CalTech (1964)
 1964FO13 J.L.C. Ford, Jr., Phys. Rev. 136 (1964) B953
 1964FU15 R. Fulle, D. Netzband and K. Schlott, Nucl. Phys. 56 (1964) 512
 1964GR39 G. Gregoire and P.C. Macq, Phys. Lett. 8 (1964) 328
 1964KI02 K.G. Kibler and R.R. Carlson, Bull. Amer. Phys. Soc. 9 (1964) 406, CA1
 1964LU02 B.T. Lucas, S.W. Cospers and O.E. Johnson, Phys. Rev. 133 (1964) B963
 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba,
 Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed., W.H. Johnson, Jr.
 (1964) 305
 1964MA60 M. Manalis and J.E. Henkel, Phys. Rev. 136 (1964) B1741
 1964ME07 R.A. Mendelson, Jr., E. Norbeck and R.R. Carlson, Phys. Rev. 135 (1964) B1319
 1964MI04 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 50
 1964TE01 E. Teranishi and B. Furubayashi, Phys. Lett. 9 (1964) 157
 1964TE04 G. Tessler and W.E. Stephens, Phys. Rev. 135 (1964) B129
 1965AL1F Almadi, Venuti, Cortellessa, Fronterotta, Reale and Salvadori, RC Accad. Naz. Lincei
 38 (1965) 499
 1965AR1E Artemov et al, Yad. Fiz. 1 (1965) 629; Sov. J. Nucl. Phys. 1 (1965) 450
 1965BE1E Berggren, Ark. Fys. 30 (1965) 508
 1965BO19 R. Bouchez, C. Perrin, A. Giorni and R. Darves-Blanc, Compt. Rend. 261 (1965)
 1269
 1965BO1B Boffi, Sawicki and Scacciatelli, Nuovo Cim. B40 (1965) 1
 1965BO37 R. Bock, H.H. Duhm, M. Grosse-Schulte and R. Rudel, Nucl. Phys. 70 (1965) 481
 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707

- 1965DI03 F.S. Dietrich, Nucl. Phys. 69 (1965) 49
- 1965DO13 I. Dostrovsky, R. Davis, Jr., A.M. Poskanzer and P.L. Reeder, Phys. Rev. 139 (1965) B1513
- 1965FL03 N.R. Fletcher, J.D. Marshall and R.H. Davis, Nucl. Phys. 70 (1965) 471
- 1965FR1B Frasca, Finlay, Koshel and Cassola, Bull. Amer. Phys. Soc. 10 (1965) 1126
- 1965GI10 L. Gilly, M. Jean, R. Meunier, M. Spighel, J.P. Stroot and P. Duteil, Phys. Lett. 19 (1965) 335
- 1965GO1D Goldanskii, Usp. Fiz. Nauk 87 (1965) 255
- 1965GO1E Gondrand, CEA 2734 (1965)
- 1965GR08 G.M. Griffiths, Nucl. Phys. 65 (1965) 647
- 1965GR18 T.A. Griffy and D.U.L. Yu, Phys. Rev. 139 (1965) B880
- 1965GR1E Grishanova and Kabatchnik, Yad. Fiz. 2 (1965) 232
- 1965HA09 J.C. Hardy, R.I. Verrall, R. Barton and R.E. Bell, Phys. Rev. Lett. 14 (1965) 376
- 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
- 1965HI1B Hiura and Shimodaya, Prog. Theor. Phys. 34 (1965) 861
- 1965HU10 H. Hulubei, M. Ivascu, A. Berinde, I. Neamu, N. Scintei, I. Francz, N. Martalogu and E. Marincu, Rev. Roum. Phys. 10 (1965) 403
- 1965IM01 W.L. Imhof, L.F. Chase, Jr. and D.B. Fossan, Phys. Rev. 139 (1965) B904
- 1965JA1A Jacmart, Cahiers Phys. (France) 19 (1965) 1
- 1965KO1B Komar and Makhnovskii, Dokl. Akad. Nauk SSSR 160 (1965) 1300; Sov. Phys. Dokl. 10 (1965) 150
- 1965KU1B Kudo, Prog. Theor. Phys. 34 (1965) 942
- 1965LY01 B. Lynch, G.M. Griffiths and T. Lauritsen, Nucl. Phys. 65 (1965) 641
- 1965MA1H Mahaux, Nucl. Phys. 71 (1965) 241
- 1965MO09 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. 64 (1965) 169
- 1965NE1B Neudachin and Smirnov, At. Energy Rev. 3 (1965) 3-157
- 1965NG1A Nguyen Ngoc, Ann. Phys. (France) 10 (1965) 315
- 1965RO1G Rose and Fisher, Private Communication (1965)
- 1965SA15 D.G. Sargood and G.D. Putt, Aust. J. Phys. 18 (1965) 491
- 1965SC17 R. Schoneberg, H. Hansen, H. Weigmann and A. Flammersfeld, Z. Phys. 188 (1965) 38
- 1965SY02 G.D. Symons, Phys. Lett. 18 (1965) 142

- 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331
- 1965VA1G Van Praet and Kossanyi-Demay, Nuovo Cim. 39 (1965) 388
- 1965VO1A Volkov, Nucl. Phys. 74 (1965) 33
- 1965WA04 B.D. Walker, C. Wong, J.D. Anderson and J.W. McClure, Phys. Rev. 137 (1964) B1504
- 1965WO01 J.B. Woods and D.H. Wilkinson, Nucl. Phys. 61 (1965) 661
- 1965WY02 J.M. Wyckoff, B. Ziegler, H.W. Koch and R. Uhlig, Phys. Rev. 137 (1965) B576
- 1965YA02 T. Yanabu, S. Yamashita, K. Takimoto and K. Ogino, J. Phys. Soc. Jpn. 20 (1965) 1303
- 1966AD06 C. Adler, T. Corcoran and C. Mast, Nucl. Phys. 88 (1966) 145
- 1966AF1A Afanasyev et al, Bull. Acad. Sci. USSR Phys. 30 (1966) 376
- 1966AM1A Amster and Perkins, Nucl. Sci. Eng. 25 (1966) 440
- 1966AS04 P.A. Assimakopoulos, N.H. Gangas and S. Kossionides, Nucl. Phys. 81 (1966) 305
- 1966BA1A Bahcall, Nucl. Phys. 75 (1966) 10
- 1966BA1X Barz et al, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Eds., J. Schintlmeister; ZFK-122 (1966) 222
- 1966BA21 K. Bahr, W. Fitz, R. Jahr and R. Santo, Phys. Lett. 21 (1966) 686
- 1966BA26 F.C. Barker, Nucl. Phys. 83 (1966) 418
- 1966BO1C Bouchez et al, J. Phys. (Paris) 27 (1966) C1-79
- 1966BO1F Bouchez, Antwerp 1965 Neutron Conf. (1966) 566
- 1966CA02 R.T. Carpenter and H.R. Hiddleston, Phys. Lett. 20 (1966) 286
- 1966CH1B Cherdantsev and Gamalya, Izv. Vyss. Uch. Zaved. Fiz. 5 (1966) 24
- 1966CH20 P.R. Christensen and C.L. Cocke, Nucl. Phys. 89 (1966) 656
- 1966CI1A Cindro, Revs. Mod. Phys. 38 (1966) 391
- 1966CL01 H.G. Clerc, K.J. Wetzel and E. Spamer, Phys. Lett. 20 (1966) 667
- 1966CO16 S. Costa, L. Pasqualini, G. Piragino and L. Roasio, Nuovo Cim. B42 (1966) 306
- 1966CO24 A.A. Cowley, G. Heymann, R.L. Keizer and M.J. Scott, Nucl. Phys. 86 (1966) 363
- 1966CR05 R.M. Craig, B. Hird, C.J. Kost and T.Y. Li, Phys. Lett. 21 (1966) 177
- 1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097
- 1966DE13 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 3 (1966) 268; Sov. J. Nucl. Phys. 3 (1966) 192
- 1966DE1K De Forest and Walecka, Adv. Phys. 15 (1966) 1, 491

- 1966DI04 G.U. Din and J.L. Weil, Nucl. Phys. 86 (1966) 509
- 1966DO1C Donau, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122 (1966) 20
- 1966DR04 Y.G. Dragunov, Y.V. Melikov and A.F. Tulinov, Yad. Fiz. 4 (1966) 314; Sov. J. Nucl. Phys. 4 (1967) 225
- 1966EL08 F. El-Batanoni and A.A. Kresnin, Nucl. Phys. 89 (1966) 577
- 1966FA1A Fasoli, Toniolo and Zago, Antwerp 1965 Neutron Conf. (1966) 496
- 1966GA15 H. Gauvin, Compt. Rend. 263 (1966) 752
- 1966GO1B Goldanskii, Ann. Rev. Nucl. Sci. 16 (1966) 1
- 1966GO1D Gotz, Mittenbacher, Prager and Muhle, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122 (1966) 201
- 1966GR1E Gregoire, Ann. Soc. Sci. (Brussels) 80 (1966) 206
- 1966GR1F Gregoire and Macq, J. Phys. (Paris) 27 (1966) C1-136
- 1966HA18 E.C. Halbert, Y.E. Kim and T.T.S. Kuo, Phys. Lett. 20 (1966) 657
- 1966HE1C R.C. Herndon and Y.C. Tang, Phys. Rev. 149 (1966) 735
- 1966HI1A Hiura and Shimodaya, Prog. Theor. Phys. Jpn. 36 (1966) 977
- 1966JA09 D.F. Jackson, Nuovo Cim. B41 (1966) 86
- 1966JA1A Jacob and Maris, Revs. Mod. Phys. 38 (1966) 121
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932
- 1966KL1C Klapisch and Bernas, Nucl. Instrum. Meth. 38 (1966) 291
- 1966KU1C Kudiyarov, Neudachin and Smirnov. Bull. Acad. Sci. USSR Phys. 30 (1966) 240
- 1966LA02 J.P. Laugier, L. Marquez, N. Saunier and J. Rey, Nucl. Phys. 75 (1966) 418
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966MA1P MacFarlane, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 411
- 1966MA38 S.K. Mark, P.M. Portner and R.B. Moore, Can. J. Phys. 44 (1966) 2961
- 1966MC1C McPherson, F.S.U. Isobaric Spin Conf. (1966) 162
- 1966ME1E Melikov, Vestnik Mosk. Univ. Fiz. Astron. 6 (1966) 102
- 1966MI1D Mittenbacher, Gotz and Prager, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122 (1966) 293
- 1966MO05 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. 75 (1966) 113
- 1966NG01 D.-C. Nguyen, J. Phys. Soc. Jpn. 21 (1966) 2462; Erratum J. Phys. Soc. Jpn. 22 (1967) 684

- 1966NU1B Nussbaum, *Helv. Phys. Acta* 39 (1966) 218
- 1966OE1A Oertzen, Bock and Grosse-Schulte, *Z. Naturforsch.* 21A (1966) 946
- 1966PA16 P.D. Parker, *Phys. Rev.* 150 (1966) 851
- 1966PA1H Park, *Bull. Amer. Phys. Soc.* 11 (1966) 627
- 1966PE1E Peterson, Ziegler and Cole, *Bull. Amer. Phys. Soc.* 11 (1966) 120
- 1966PO1D Potenza, *Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch* (1966) 584
- 1966PU02 P. Purdom, Jr., P.A. Seeger and R.W. Kavanagh, *Nucl. Phys.* 83 (1966) 513
- 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, *Phys. Rev.* 144 (1966) 859; Erratum *Phys. Rev.* 148 (1966) 1246
- 1966RO1E Rotter and Zhusupov, *Ann. Phys.* 17 (1966) 57
- 1966RO1F Rotter, *Ann. Phys.* 17 (1966) 247
- 1966RO1H Rotter, *Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122* (1966) 244
- 1966SI1C Simons, *Bull. Amer. Phys. Soc.* 11 (1966) 301
- 1966TH03 M.N. Thompson and J.M. Taylor, *Nucl. Phys.* 76 (1966) 377
- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, *Nucl. Phys.* 79 (1966) 321; Erratum *Nucl. Phys.* A119 (1968) 692
- 1966VO06 Y.M. Volkov, A.P. Komar and V.P. Chizhov, *Yad. Fiz.* 3 (1966) 277; *Sov. J. Nucl. Phys.* 3 (1966) 198
- 1966WA12 N.S. Wall and P.R. Roos, *Phys. Rev.* 150 (1966) 811; Erratum *Phys. Rev.* 159 (1967) 1062
- 1966WA16 M.A. Waggoner, J.E. Etter, H.D. Holmgren and C. Moazed, *Nucl. Phys.* 88 (1966) 81
- 1966WE1B Weinberg, *Antwerp 1965 Neutron Conf.* (1966) 37
- 1966WI08 D.H. Wilkinson, J.T. Sample and D.E. Alburger, *Phys. Rev.* 146 (1966) 662
- 1966WI1E D.H. Wilkinson and M.E. Mafethe, *Nucl. Phys.* 85 (1966) 97
- 1967AC01 E. Acerbi, M. Castiglioni, G. Dutto, I. Iori, A. Luccio, S. Micheletti, N. Molho, M. Pignanelli, F. Resmini, G. Strini et al, *Suppt. Nuovo Cim.* 5 (1967) 1252
- 1967AR1A Artus et al, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (1967) 314
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, *Nucl. Phys.* A97 (1967) 144
- 1967BA12 G. Baron and H. Rouhaninejad, *J. Phys. (France)* 28 (1967) 142
- 1967BA59 C.A. Barnes, E.G. Adelberger, D.C. Hensley and A.B. MacDonald, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (1967) 261
- 1967BE13 M. Berrada, J.-P. Laugier, C. Lemeille, N. Saunier and L. Marquez, *J. Phys. (France)* 28 (1967) 135

1967BE1P H.A. Bentz, R. Engfer and W. Buhring, Nucl. Phys. A101 (1967) 527
 1967BE1Q H.W. Bertini, Phys. Rev. 162 (1967) 976
 1967BE26 M. Bernheim, T. Stovall and D. Vinciguerra, Nucl. Phys. A97 (1967) 488
 1967BE49 B.L. Berman, R.L. Van Hemert and C.D. Bowman, Phys. Rev. 163 (1967) 958; Erratum Phys. Rev. 168 (1968) 1414
 1967BL09 A.G. Blachman and A. Lurio, Phys. Rev. 153 (1967) 164
 1967BO1D Bodart and Deconninck, Ann. Soc. Sci. (Brussels) 81 (1967) 137
 1967BO1F Bouchez et al, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 287
 1967BO1K Boffi, Pacati and Sawicki, Nuovo Cim. B52 (1967) 210
 1967BO34 S. Boffi, F.D. Pacati and J. Sawicki, Nuovo Cim. B52 (1967) 244
 1967BR1B Browne and O'Donnell, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 508
 1967BR39 F.P. Brady, J.A. Jungerman and J.C. Young, Nucl. Phys. A98 (1967) 241
 1967CA17 P. Camiz, Nuovo Cim. B51 (1967) 190
 1967CA1J Cabrespine, Gauvin, Lefort and Sauvage, Ark. Fys. 36 (1967) 463
 1967CH1H D. Chlebowska, J. Wrzecionko and M. Zielinska-Pfabe, Phys. Lett. B25 (1967) 309
 1967CH34 V.I. Chuev, V.V. Davidov, A.A. Ogloblin and S.B. Sakuta, Ark. Fys. 36 (1967) 263
 1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1
 1967CO36 S.W. Cospser, J. Cerny and R.C. Gatti, Phys. Rev. 154 (1967) 1193
 1967CR04 M.A. Crosby and J.C. Legg, Nucl. Phys. A95 (1967) 639
 1967CR05 R.M. Craig, B. Hird, C.J. Kost and T.Y. Li, Nucl. Phys. A96 (1967) 367
 1967DR08 I.P. Dryapachenko, V.A. Kornilov, O.F. Nemetz and V.A. Pilipchenko, Yad. Fiz. 6 (1967) 440; Sov. J. Nucl. Phys. 6 (1968) 321
 1967EA01 L.G. Earwaker, Nucl. Phys. A90 (1967) 56
 1967EL1D El-Nadi, Ismail and Rihan, Nuovo Cim. B50 (1967) 224
 1967FI07 W. Fitz, R. Jahr and R. Santo, Nucl. Phys. A101 (1967) 449
 1967FU08 K. Fukunaga, H. Nakamura and N. Fujiwara, J. Phys. Soc. Jpn. 23 (1967) 911
 1967GL1B Glavina et al, Bull. Amer. Phys. Soc. 12 (1967) 651
 1967HA10 K.B. Haque and J.G. Valatin, Nucl. Phys. A95 (1967) 97
 1967JA1E B.K. Jain and D.F. Jackson, Nucl. Phys. A99 (1967) 113
 1967KA1A Kabachnik and Grishanova, Sov. J. Nucl. Phys. 4 (1967) 583
 1967KE1F Kerr, Thesis, Univ. of Wisconsin (1967)
 1967KI03 K.G. Kibler, Phys. Rev. 155 (1967) 1110

1967KO1B Kolybasov and Smorodinskaya, *Yad. Fiz.* 5 (1967) 777
1967KU1E D. Kurath and R.D. Lawson, *Phys. Rev.* 161 (1967) 915
1967LO1B Loiseaux, Maison and Langevin, *J. Phys. (France)* 28 (1967) 11
1967ME11 D.F. Measday and J.N. Palmieri, *Phys. Rev.* 161 (1967) 1071
1967ME1C Meboniya, *Bull. Moscow Univ., Phys. Astron.* 1 (1967) 114
1967ME1F Meyerhcf, *Proc. Problem Symp. on Nucl. Phys., Tbilisi, 1967* (1967) 350
1967MI1B Miller, Thesis, Princeton Univ. (1967)
1967MO1H S.K. Monga, *Phys. Rev.* 160 (1967) 846
1967OD01 F.H. O'Donnell and C.P. Browne, *Phys. Rev.* 158 (1967) 957
1967SA13 G.R. Satchler, *Nucl. Phys. A100* (1967) 497
1967SE11 V.I. Serov, S.N. Abramovich, V.A. Pereshivkin and L.A. Morkin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 31 (1967) 333; *Bull. Acad. Nauk USSR Phys. Ser.* 31 (1968) 317
1967SH05 Y.Y. Sharon, *Nucl. Phys. A99* (1967) 321
1967SH14 V.S. Shirley, UCRL-17990 (1967)
1967SH1E Shevchenko, *Proc. Int. Conf. Electromag. Inter. Dubna, 1967, Vol. 3* (1967) 206
1967SL04 R.J. Slobodrian, H. Bichsel, J.S.C. McKee and W.F. Tivol, *Phys. Rev. Lett.* 19 (1967) 595
1967SP09 A. Sperduto, *Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967* (1967) 657
1967ST1C Stephenson, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (1967) 906
1967ST1G T. Stovall, *Nucl. Phys. A102* (1967) 554
1967TA1C Takimoto, *Mem. Coll. Sci. Univ. Kyoto A31* (1967) 267
1967VA11 V. Valkovic, W.R. Jackson, Y.S. Chen, S.T. Emerson and G.C. Phillips, *Nucl. Phys. A96* (1967) 241
1967VA24 P. van der Merwe, W.R. McMurray and I.J. van Heerden, *Nucl. Phys. A103* (1967) 474
1967VE01 J.W. Verba, H. Willmes, R.F. Carlson, I. Slaus, J.R. Richardson and E.L. Petersen, *Phys. Rev.* 153 (1967) 1127
1967WA1E Walecka, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (1967) 289
1967WA1F Walecka, *Proc. Int. Conf. Electromag. Inter. Dubna, 1967, Vol. 3* (1967) 28
1967WI1C Wise, Knowles and Bunch, *Bull. Amer. Phys. Soc.* 12 (1967) 633
1968AD09 J.-O. Adler and B. Forkman, *Ark. Fys.* 38 (1968) 389
1968AJ01 F. Ajzenberg-Selove, R.D. Wardaski and R. Middleton, *Nucl. Phys. A116* (1968) 481
1968AM1A Amaldi et al, in Tokyo (1968) 369

- 1968BA1C Barker and Fitzpatrick, Aust. J. Phys. 21 (1968) 415
- 1968BA1E Ball, UCRL-18263 (1968)
- 1968BA1H V.V. Balashov and D.V. Meboniya, Nucl. Phys. A107 (1968) 369
- 1968BA2G Backenstoss et al, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968BE1J Berkowitz et al, Bull. Amer. Phys. Soc. 13 (1968) 607
- 1968BO19 M. Bouten, M.C. Bouten, H. Depuydt and L. Schotsmans, Phys. Lett. B27 (1968) 61
- 1968BO32 N.E. Booth, A. Beretvas, R.E.P. Davis, C. Dolnick, R.E. Hill, M. Raymond and D. Sherden, Nucl. Phys. A119 (1968) 233
- 1968BO46 V.F. Borzhkovskii, A.S. Cherkasov, N.G. Afanasev, I.A. Grishaev and I.I. Zalyubovskii, Yad. Fiz. 7 (1968) 261; Sov. J. Nucl. Phys. 7 (1968) 181
- 1968BR21 M. Brendle, M. Morike, G. Staudt and G. Steidle, Z. Naturforsch. A23 (1968) 1229
- 1968CE01 J. Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968CH1J Chatterjee and De, Indian J. Phys. 42 (1968) 486
- 1968CL08 H.-G. Clerc, K.J. Wetzel and E. Spamer, Nucl. Phys. A120 (1968) 441
- 1968CO07 C.L. Cocke, Nucl. Phys. A110 (1968) 321
- 1968CO08 C.L. Cocke and P.R. Christensen, Nucl. Phys. A111 (1968) 623
- 1968CO31 M.N.H. Comsan, M.A. Farouk, A.A. El-Kamhawy, M.S.M. El-Tahawy and A.N. Lvov, Atomkernenergie 13 (1968) 415
- 1968DA1H Daruga et al, Atomn. Energ. (USSR) 24 (1968) 66
- 1968DA20 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Yad. Fiz. 7 (1968) 758; Sov. J. Nucl. Phys. 7 (1968) 463
- 1968DO20 I. Dostrovsky, H. Gauvin and M. Lefort, Phys. Rev. 169 (1968) 836
- 1968DZ1A Dzhibuti et al, Krupennikova and Mamasakhlisov, Sov. J. Nucl. Phys. 7 (1968) 36
- 1968ER1A Ericson, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 103
- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161
- 1968GL1A Glauber, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 41, 60
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GO1J Goldemberg, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 379
- 1968HA1C Hayakawa, Suppl. Prog. Theor. Phys. (1968) 156
- 1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, 1967 (1968) 41
- 1968HO1F Honda, Mori and Yoshida, in Tokyo (1968) 260

1968JA1D Jackson, Adv. Phys. 17 (1968) 481
 1968JA1G Jackson, in Tokyo (1968) 105
 1968KA38 N.N. Kaushal, E.J. Winhold, P.F. Yergin, H.A. Medicus and R.H. Augustson, Phys. Rev. 175 (1968) 1330
 1968KO1E Komarov, Kosarev and Savchenko, Joint Inst. Nucl. Res., Lab. Nucl. Problems, USSR, Rept. P1 4227 (1968)
 1968KO1H Konshin et al, Sov. J. Nucl. Phys. 6 (1968) 363
 1968KR02 J.J. Kroepfl and C.P. Browne, Nucl. Phys. A108 (1968) 289
 1968KU04 L.A. Kull and E. Kashy, Phys. Rev. 167 (1968) 963
 1968KU1B Kudeyarov, Neudachin, Serebryakov and Smirnov, Sov. J. Nucl. Phys. 6 (1968) 876
 1968KU1D Kurath, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 393
 1968LI1C Li and Mark, Can. J. Phys. 46 (1968) 2645
 1968LI1E T.Y. Li and B. Hird, Phys. Rev. 174 (1968) 1130
 1968MA1H Machali et al, Atomkernenergie 13 (1968) 29
 1968MA1X Majling, Kukulín and Smirnov, Czech. J. Phys. B18 (1968) 1560
 1968MA1Y L. Majling, V.I. Kukulín and Yu.F. Smirnov, Phys. Lett. B27 (1968) 487
 1968NE1A NeuV.G. Neudatchin, N.S. Zelenskaya, E.Zh. Magzumov and E.A. Romanovsky, Phys. Lett. B27 (1968) 490
 1968NE1B Neumann, Bull. Amer. Phys. Soc. 13 (1968) 100
 1968NO1A M.E. Nordberg, K.F. Kinsey and R.L. Burman, Phys. Rev. 165 (1968) 1096
 1968OK06 Y. Okuma, J. Phys. Soc. Jpn. 25 (1968) 1
 1968PA1H Parthasaradhi, Current Sci. (India) 37 (1968) 101
 1968PA1M Parker, Astrophys. J. 153 (1968) L85
 1968PA1P Park, Can. J. Phys. 46 (1968) 1967
 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
 1968PE1A R.W. Peelle, T.A. Love, N.W. Hill and R.T. Santoro, Phys. Rev. 167 (1968) 981
 1968RE11 D.H. Rester, Nucl. Phys. A118 (1968) 129
 1968RO19 P.G. Roos, H.G. Pugh, M. Jain, H.D. Holmgren, M. Epstein and C.A. Ludemann, Phys. Rev. 176 (1968) 1246
 1968RO1H Roturier, NP 17794 (1968)
 1968SA1G Sanada and Takeuchi, in Tokyo (1968) 255

- 1968SE1B Segel, Watson and Singh, Bull. Amer. Phys. Soc. 13 (1968) 115
- 1968SN1A Snover, Paul, Healey and Hanna, Bull. Amer. Phys. Soc. 13 (1968) 882
- 1968TA1K Tanaka, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 51
- 1968TA1M Taffara and Vanzani, Nuovo Cim. B56 (1968) 166
- 1968TA1N Takeuchi and Sanada, TUENS-4 (1968)
- 1968TH1H N. Thurlow, Nucl. Phys. A109 (1968) 471
- 1968VA05 G.J. Vanpraet and W.C. Barber, Z. Phys. 211 (1968) 213
- 1968VE11 A.N. Vereshchagin, I.N. Korostova, L.S. Sokolov, V.V. Tokarevskii and I.P. Chernov, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 623; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 573
- 1968VE1C A.N. Vereshchagin et al, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1956
- 1968WI1B Wilkinson, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 469
- 1968WI1E Wise, Thesis, Washington State Univ. (1968)
- 1968YA02 T. Yanabu, S. Yamashita, K. Hosono, S. Matsuki, T. Tanabe, K. Takimoto, Y. Okuma, K. Ogino, S. Okumura and R. Ishiwari, J. Phys. Soc. Jpn. 24 (1968) 667
- 1968YI01 F. Yiou, Ann. Phys. (Paris) 3 (1968) 169
- 1969AF1A Afanasev et al, Sov. J. Nucl. Phys. 8 (1969) 646
- 1969AN1H Antufyev et al, Ukr. Fiz. Zh. 14 (1969) 248
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969AR1B Armstrong, Beery, Keaton and Veaser, LA 4177 (1969)
- 1969AU05 T. Aurdal, Z. Naturforsch. A24 (1969) 1188
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969BA1F V.V. Balashov, N.M. Kabachnik and V.I. Markov, Nucl. Phys. A129 (1969) 369
- 1969BA1J Balashov, Bochum Conf. STI/PUB/232 IAEA (1969) 59
- 1969BA1Z Barnes, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 179
- 1969BE21 H.A. Bentz, Z. Naturforsch. A24 (1969) 858
- 1969BE50 M. Bernheim, R. Riskalla, T. Stovall and D. Vinciguerra, Phys. Lett. B30 (1969) 412
- 1969BO19 M. Bouten, M.-C. Bouten and H. Depuydt, Nucl. Phys. A131 (1969) 385
- 1969BO1U Boffi, Nuovo Cim. B57 (1969) 247

- 1969BO1V M. Bouten, M.-C. Bouten, H. Depuydt and L. Schotsmans, Nucl. Phys. A127 (1969) 177
- 1969BO33 M. Bouten, Physica 44 (1969) 617
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 242
- 1969BU1C Burman and Nordberg, Bull. Amer. Phys. Soc. 14 (1969) 537
- 1969CA1B F. Calligaris, C. Cernigoi, I. Gabrielli and F. Pellegrini, Nucl. Phys. A126 (1969) 209
- 1969CH1C D.T. Chivers, E.M. Rimmer, B.W. Allardyce, R.C. Witcomb, J.J. Domingo and N.W. Tanner, Nucl. Phys. A126 (1969) 129
- 1969DE31 J.M. Delbrouck-Habaru, P.D. Dumont, M. Huez, G. Robaye and L. Winand, Bull. Soc. Roy. Sci. Liege 38 (1969) 240
- 1969DO02 V.K. Dolinov, Y.V. Melikov, A.F. Tulinov and O.V. Bormot, Nucl. Phys. A129 (1969) 577
- 1969DO03 V.K. Dolinov, D.V. Meboniya and A.F. Tulinov, Nucl. Phys. A129 (1969) 597
- 1969DO09 B.S. Dolbilkin, A.I. Isakov, V.I. Korin, L.E. Lazareva, N.V. Linkova and F.A. Nikolaev, Yad. Fiz. 9 (1969) 914; Sov. J. Nucl. Phys. 9 (1969) 534
- 1969ED01 R.D. Edge, D.H. Tompkins and J.W. Glenn, Phys. Rev. 183 (1969) 849
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA03 G. Gambarini, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Tagliaferri, Nucl. Phys. A126 (1969) 562
- 1969GA18 J. Galin, B. Gatty, M. Lefort, J. Peter, X. Tarrago and R. Basile, Phys. Rev. 182 (1969) 1267
- 1969GA1P Garvey, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 703
- 1969GU03 M.R. Gunye, J. Law and R.K. Bhaduri, Nucl. Phys. A132 (1969) 225
- 1969HA14 G. Hauser, R. Lohken, H. Rebel, G. Schatz, G.W. Schweimer and J. Specht, Nucl. Phys. A128 (1969) 81
- 1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
- 1969HE1N E.M. Henley and C.E. Lacy, Phys. Rev. 184 (1969) 1228
- 1969HO1K Holmgren, Bochum Conf. STI/PUB/232 IAEA (1969) 17
- 1969HO27 T. Honda, Nucl. Phys. A136 (1969) 183
- 1969HS02 Y.-C. Hsu, C.-Y. Huang, S.-Y. Lin, Y.-C. Hsu and M.-C. Chou, Chin. J. Phys. (Taiwan) 7 (1969) 1
- 1969JA1M J. Janecke, Nucl. Phys. A128 (1969) 632

- 1969JU1A Jungerman et al, Bull. Amer. Phys. Soc. 14 (1969) 488
- 1969KA1D Karge, Wiss. Z. Friedrich. Schiller Univ. Jena. Math. Naturwiss. Reihe (Germany) 18 (1969) 51
- 1969KI02 H. Kitazawa and N. Yamamuro, J. Phys. Soc. Jpn. 26 (1969) 600
- 1969KL08 R. Klapisch, C. Thibault-Philippe, C. Detraz, J. Chaumont, R. Bernas and E. Beck, Phys. Rev. Lett. 23 (1969) 652
- 1969KO1D Konig and Staudi, Bochum Conf. STI/PUB/232 IAEA (1969) 216
- 1969KO1F V.M. Kolybasov and N.Ya. Smorodinskaya, Phys. Lett. B30 (1969) 11
- 1969KO1G Komarov, Kosarev and Savchenko, Joint Inst. Nucl. Res., Lab. Nucl. Problems, USSR, Rept. P1 4373 (1969)
- 1969KO1J V.M. Kolybasov and N.Ya. Smorodinskaya, Nucl. Phys. A136 (1969) 165
- 1969KR03 E. Krubasik, H. Voit, E. Blatt, H.-D. Helb and G. Ischenko, Z. Phys. 219 (1969) 185
- 1969LE1D Leonardi and Rosa-Clot, Lett. Nuovo Cim. 1 (1969) 829
- 1969LE1G Leung and Park, Bull. Amer. Phys. Soc. 14 (1969) 511
- 1969LO1D Loude, Perroud and Sellum, Helv. Phys. Acta 42 (1969) 905
- 1969MA11 B.E.F. Macefield, B. Wakefield and D.H. Wilkinson, Nucl. Phys. A131 (1969) 250
- 1969MO1E L. Moyer and D.S. Koltun, Phys. Rev. 182 (1969) 999
- 1969MO1H R. Moreh, D. Salzmann and Y. Wand, Phys. Lett. B30 (1969) 536
- 1969NE08 V.G. Neudachin, E.Z. Magzumov and E.A. Romanovskii, Yad. Fiz. 9 (1969) 94; Sov. J. Nucl. Phys. 9 (1969) 56
- 1969NE1A Neumann, Bull. Amer. Phys. Soc. 14 (1969) 1224
- 1969NE1C Neudatchin, Bochum Conf. STI/PUB/232 IAEA (1969) 35
- 1969NE1E Nettles, Hensley and Tombrello, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 819
- 1969NU1A Nusslin, Werner and Zimmerer, Bochum Conf. STI/PUB/232 IAEA (1969) 229
- 1969NU1C Nussbaum, Helv. Phys. Acta 42 (1969) 361
- 1969OP1A Opelka, Brussel, Hoffswell and Yavin, Bull. Amer. Phys. Soc. 14 (1969) 506
- 1969OR01 H. Orihara, T. Nakagawa, H. Ueno, T. Tohei, T. Yamaya, Y. Nakagome, M. Baba and S. Morita, Nucl. Phys. A139 (1969) 226
- 1969PA11 J.Y. Park, J.L. Duggan, P.D. Miller, M.M. Duncan and R.L. Dangle, Nucl. Phys. A134 (1969) 277
- 1969PI11 J.R. Pizzi, M. Gaillard, P. Gaillard, A. Guichard, M. Gusakov, G. Reboulet and C. Ruhla, Nucl. Phys. A136 (1969) 496
- 1969RO1F Roturier, Ann. Phys. 4 (1969) 289

- 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
- 1969SA04 J. Sanada, Y.C. Liu, Y. Sugiyama and O. Mikoshiba, J. Phys. Soc. Jpn. 26 (1969) 853
- 1969TO1A I.S. Towner, Nucl. Phys. A126 (1969) 97
- 1969VA1C S. Varma and P. Goldhammer, Nucl. Phys. A125 (1969) 193
- 1969VE02 V.V. Verbinski and W.R. Burrus, Phys. Rev. 177 (1969) 1671
- 1969VE09 A.N. Vereshchagin, I.N. Simonov, K.O. Terenetskii, V.V. Tokarevskii and I.P. Chernov, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 2064; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1880
- 1969VI02 D. Vinciguerra and T. Stovall, Nucl. Phys. A132 (1969) 410
- 1969VI05 B. Vignon, E. Ligeon and J.P. Longequeue, J. Phys. (France) 30 (1969) 913
- 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
- 1969WI1F Wilson, Sandler, Otte and Phillips, Bull. Amer. Phys. Soc. 14 (1969) 37
- 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
- 1969YA1B Yamashita et al, J. Phys. Soc. Jpn. 26 (1969) 1078
- 1969YI1A Yiou, Seide and Bernas, J. Geophys. Res. 74 (1969) 2447
- 1969ZE1A Zeidman and Fortune, Bull. Amer. Phys. Soc. 14 (1969) 507
- 1970AL1M Alsmiller, Gabriel and Guthrie, Nucl. Sci. Eng. 40 (1970) 365
- 1970AL25 M.R. Aleksic, R.V. Popic, D.M. Stanojevic and B.Z. Stepancic, Fizika 2 (1970) 113
- 1970AN07 J.D. Anderson, C. Wong, B.A. Pohl and J.W. McClure, Phys. Rev. C2 (1970) 319
- 1970AU02 D.L. Auton, Nucl. Phys. A157 (1970) 305
- 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
- 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119
- 1970BA1P Barendoltz, Thesis, Univ. of Illinois (1970)
- 1970BA1Q Barashenkov and Abdinov, Acta Phys. Pol. B1 (1970) 65
- 1970BA49 P.H. Barker, A. Huber, H. Knoth, U. Matter, A. Gobbi and P. Marmier, Nucl. Phys. A155 (1970) 401
- 1970BE1F Bennett, Thesis, Washington State Univ. (1970)
- 1970BE1G Beck, Grunbaum and Tomaselli, Proc. 3rd Int. Conf. on High Energy Phys. and Nucl. Struct., New York, 1969 (1970) 63
- 1970BE1J H.W. Bertini, Phys. Rev. C1 (1970) 423
- 1970CA1L Calligaris, Cernigoi, Gabrielli and Pellegrini, Proc. 3rd Int. Conf. on High Energy Phys. and Nucl. Struct., New York, 1969 (1970) 367
- 1970CA28 J. Catala, A. Garcia, V. Martinez and F. Senent, An. Fis. 66 (1970) 297

1970CH07 Y.S. Chen, T.A. Tombrello and R.W. Kavanagh, Nucl. Phys. A146 (1970) 136

1970CH19 V.I. Chuev, V.V. Davydov, V.I. Manko, B.G. Novatsky, S.B. Sakuta and D.N. Stepanov, Phys. Lett. B31 (1970) 624

1970CH1T Chen, Thesis, CalTech (1970)

1970CL01 A.S. Clough, C.J. Batty, B.E. Bonner and L.E. Williams, Nucl. Phys. A143 (1970) 385

1970CO04 L. Cohen and G.H. Herling, Nucl. Phys. A141 (1970) 595

1970CO1H Cohen and Kurath, Nucl. Phys. A141 (1970) 145

1970DA21 A. D'Andrea and M. Scalia, Nuovo Cim. A69 (1970) 702

1970DI12 R.L. Dixon and R.D. Edge, Nucl. Phys. A156 (1970) 33

1970DI1F Dixon, Thesis, Univ. of South Carolina (1970)

1970DR11 D.M. Drake, J.C. Hopkins, C.S. Young and H. Conde, Nucl. Sci. Eng. 40 (1970) 294

1970EL16 M. El-Nadi, O. Zohni and H.M. Hussein, Ann. Phys. (Leipzig) 25 (1970) 1

1970ER1A Ericson and Locher, Nucl. Phys. A148 (1970) 1

1970FA15 D. Favart, F. Brouillard, L. Grenacs, P. Igo-Kemenes, P. Lipnik and P.C. Macq, Phys. Rev. Lett. 25 (1970) 1348

1970FO1D Fortune and Zeidman, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 307

1970GA1G Gagne, Lambert and Treado, Bull. Amer. Phys. Soc. 15 (1970) 1695

1970GO12 B. Gottschalk and S.L. Kannenberg, Phys. Rev. C2 (1970) 24

1970GO28 Y.M. Goryachev, V.P. Kanavets, I.I. Levintov, B.V. Morozov, N.A. Nikiforov and A.S. Starostin, Yad. Fiz. 11 (1970) 629; Sov. J. Nucl. Phys. 11 (1970) 353

1970GR33 R. Grubman and T. Witten, Jr., Nucl. Phys. A158 (1970) 289

1970GU06 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Tagliaferri, Nuovo Cim. A67 (1970) 407

1970GU08 K. Gul, B.H. Armitage and B.W. Hooton, Nucl. Phys. A153 (1970) 390

1970HA1G Habs et al, Kernforsch. Karlsruhe, Ext. Bericht 18-70 2 (1970)

1970HI1F Hiramatsu et al, Univ. Tokyo, Inst. Nucl. Study, Rept. INS J 122 (1970)

1970KA1K Kadenskii et al, Sov. J. Nucl. Phys. 10 (1970) 422

1970KU1D Kull, Thesis, Michigan State Univ. (1970)

1970LI1Q Lin, Thesis, Univ. of Georgia (1970)

1970LK1A Lkhagva and Rotter, Sov. J. Nucl. Phys. 11 (1970) 576

1970MA04 E.Z. Magzumov and V.G. Neudatchin, Phys. Lett. B31 (1970) 106

- 1970MA38 E.Z. Magzumov, V.G. Neudachin and M.S. Belkin, *Yad. Fiz.* 11 (1970) 589; *Sov. J. Nucl. Phys.* 11 (1970) 331
- 1970MI12 B. Mithra and R. Laverriere, *Nucl. Phys.* A155 (1970) 535
- 1970MI14 D. Miljanic, V. Valkovic, D. Rendic and M. Furic, *Nucl. Phys.* A156 (1970) 193
- 1970OG1A Ogloblin, *Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969* (1970) 231
- 1970OP1B Opelka, Thesis, Univ. of Illinois (1970)
- 1970PA1D Park and Rickett, *Bull. Amer. Phys. Soc.* 15 (1970) 1683
- 1970PE18 A.K. Petrauskas and L.Y. Sabalyauskas, *Yad. Fiz.* 12 (1970) 492; *Sov. J. Nucl. Phys.* 12 (1970) 269
- 1970PI1D Pizzi, Thesis, Univ. of Lyon (1970)
- 1970PO03 D.L. Powell, G.M. Crawley, B.V.N. Rao and B.A. Robson, *Nucl. Phys.* A147 (1970) 65
- 1970QU1C Quinn, Thesis, Univ. of California, Los Angeles (1970)
- 1970RE1E Rester, Dance and Derrickson, *J. Appl. Phys.* 41 (1970) 2682
- 1970SA05 J. Sanada, *J. Phys. Soc. Jpn.* 28 (1970) 524
- 1970SA17 A. Salyers, *Phys. Rev.* C2 (1970) 1653
- 1970SI12 V.S. Siksin, V.N. Domoratskii, L.V. Donetskov, A.N. Voronin, M.M. Novikov, A.N. Kudryavtsev, G.D. Batyshchev, S.K. Barvin and V.F. Tseshkovskaya, *Yad. Fiz.* 12 (1970) 15; *Sov. J. Nucl. Phys.* 12 (1971) 9
- 1970SQ01 G.T.A. Squier, A.R. Johnston, E.W. Spiers, S.A. Harbison and N.M. Stewart, *Nucl. Phys.* A141 (1970) 158
- 1970TH08 R.S. Thomason, G. Spalek and R.L. Walter, *Nucl. Phys.* A155 (1970) 659
- 1970TH1F Theus et al, *Bull. Amer. Phys. Soc.* 15 (1970) 1695
- 1970TI1C Tishchenko and Kresnin, *Sov. J. Nucl. Phys.* 10 (1970) 442
- 1970TO1E Tomaselli, *Z. Phys.* 233 (1970) 240
- 1970TR05 G.F. Trentelman, B.M. Preedom and E. Kashy, *Phys. Rev. Lett.* 25 (1970) 530
- 1970TR1F Trentelman, Thesis, Michigan State Univ. (1970)
- 1970TU06 S.N. Tucker, P.B. Treacy and V.V. Komarov, *Aust. J. Phys.* 23 (1970) 651
- 1970VE06 A.N. Vereshchagin, I.N. Korostova and I.P. Chernov, *Izv. Vys. Ucheb. Zav. Fiz.* 8 (1970) 105; *Sov. Phys. J.* 13 (1973) 1071
- 1970WA1G Walker and Stokes, LA-DC 11224 (1970)
- 1970WI1B Williams et al, *Bull. Amer. Phys. Soc.* 15 (1970) 62
- 1970WO1B Wolter and Ober, *Bull. Amer. Phys. Soc.* 15 (1970) 126
- 1971AD01 J.C. Adloff, K.H. Souw and C.L. Cocke, *Phys. Rev.* C3 (1971) 1808

- 1971AD1C Adelberger et al, Bull. Amer. Phys. Soc. 16 (1971) 829
- 1971AN04 Y.P. Antufev, V.L. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, Yad. Fiz. 13 (1971) 473; Sov. J. Nucl. Phys. 13 (1971) 265
- 1971AN15 Y.P. Antufev, V.L. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, Yad. Fiz. 14 (1971) 898; Sov. J. Nucl. Phys. 14 (1972) 502
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AU1G N. Auerbach, A. Lev and E. Kashy, Phys. Lett. B36 (1971) 453
- 1971BA87 C.J. Batty, Nucl. Phys. A178 (1971) 17
- 1971BE46 R.F. Bentley, J.D. Carlson, D.A. Lind, R.B. Perkins and C.D. Zafiratos, Phys. Rev. Lett. 27 (1971) 1081
- 1971BI11 H.G. Bingham, A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 265
- 1971BO50 I. Borbely, Phys. Lett. B37 (1971) 243
- 1971BR07 K.H. Bray, M. Jain, K.S. Jayaraman, G. LoBianco, W.T.H. Van Oers and Y.I. Wu, Nucl. Phys. A163 (1971) 649
- 1971BU1K Burymov et al, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 159
- 1971CA01 P.J. Castleberry, L. Coulson, R.C. Minehart and K.O.H. Ziock, Phys. Lett. B34 (1971) 57
- 1971CA1F Carlson, Lind and Zafiratos, Bull. Amer. Phys. Soc. 16 (1971) 1173
- 1971CA1J Castleberry, Thesis, Univ. of Virginia (1971)
- 1971CH1B Chuev et al, J. Phys. (Paris) 32 (1971) C6-167
- 1971CO28 B.S. Cooper, J.B. Seaborn and S.A. Williams, Phys. Rev. C4 (1971) 1997
- 1971DA13 S. Das Gupta, J.C. Hocquenghem and B. Giraud, Nucl. Phys. A168 (1971) 625
- 1971DA21 V.K. Daruga and N.N. Krasnov, Atomn. Energ. (USSR) 30 (1971) 399
- 1971DE2D Devons, Proc. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 619
- 1971DJ02 A. Djaloeis, H. Cords and J. Nurzynski, Nucl. Phys. A163 (1971) 131
- 1971DO1K Dolenc and Antolkovic, Fizika 4 (1971) 37
- 1971DRZW T. Drake, Proc. 1971 Mont Tremblant Summer School Meeting on Dynamic Struct. of Nucl. States, Toronto, Canada, 1971 (1972) 420
- 1971ER1C Ericson, Ann. Phys. 63 (1971) 562
- 1971FA04 H.-U. Fabian, Z. Naturforsch. A26 (1971) 317

- 1971FA09 J. Favier, T. Bressani, G. Charpak, L. Massonnet, W.E. Meyerhof and C. Zupancic, Nucl. Phys. A169 (1971) 540
- 1971FO1E Fou and Gagne, Bull. Amer. Phys. Soc. 16 (1971) 488
- 1971FR04 E. Friedland and I. Venter, Z. Phys. 243 (1971) 126
- 1971GA1J Gaillard, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971GE09 C. Gerardin, R. Seltz and D. Magnac-Valette, Nucl. Phys. A169 (1971) 521
- 1971GL07 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, Yad. Fiz. 13 (1971) 277; Sov. J. Nucl.Phys. 13 (1971) 154
- 1971GO14 N.F. Golovanova and V.G. Neudachin, Yad. Fiz. 13 (1971) 1248; Sov. J. Nucl. Phys. 13 (1971) 718
- 1971GR02 L. Grunbaum and M. Tomaselli, Nucl. Phys. A160 (1971) 437
- 1971GU15 A. Guichard, M. Chevallier, P. Gaillard, J-Y. Grossiord, M. Gusakow, J-R. Pizzi and C. Ruhla, Phys. Rev. C4 (1971) 700
- 1971GU23 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Semenescu, Phys. Rev. C4 (1971) 1092
- 1971HA05 J.C. Hardy, J.E. Esterl, R.G. Sextro and J. Cerny, Phys. Rev. C3 (1971) 700
- 1971HA10 J.C. Hardy, J.M. Loiseaux, J. Cerny and G.T. Garvey, Nucl. Phys. A162 (1971) 552
- 1971HU1A Hungerford et al, Bull. Amer. Phys. Soc. 16 (1971) 488
- 1971IN05 A. Ingemarsson and G. Tibell, Phys. Scr. 4 (1971) 235
- 1971IN1C Ingalls, Thesis, Princeton Univ. (1971)
- 1971JE03 H. Jeremie, Can. J. Phys. 49 (1971) 2085
- 1971KA04 S. Kahana and D. Kurath, Phys. Rev. C3 (1971) 543
- 1971KA21 D.G. Kamke and C.D. Goodman, Nucl. Phys. A172 (1971) 555
- 1971KA70 N.N. Kaushal, E.J. Winhold, R.H. Augustson, P.F. Yergin and H.A. Medicus, Nucl. Energy 25 (1971) 91
- 1971KL1E Klages et al, J. Phys. (Paris) 32 (1971) C6-209
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971LE1N Leclercq-Villain, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971LE33 J. Lestringuez, G.M. Raisbeck, F. Yiou and R. Bernas, Phys. Lett. B36 (1971) 331
- 1971LI1H H.J. Lipkin, Phys. Rev. Lett. 27 (1971) 432
- 1971LI30 W.K. Lin, Phys. Lett. B37 (1971) 480

- 1971MA13 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A165 (1971) 145
- 1971MA1C Macek et al, Bull. Amer. Phys. Soc. 16 (1971) 512
- 1971MA44 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A172 (1971) 166
- 1971MI12 D. Miljanic and V. Valkovic, Nucl. Phys. A176 (1971) 110
- 1971MO01 J.M. Mosher, R.W. Kavanagh and T.A. Tombrello, Phys. Rev. C3 (1971) 438
- 1971MO1R Mollendorff, Thesis, Univ. of Berlin (1971)
- 1971NO02 J.L. Norton and P. Goldhammer, Nucl. Phys. A165 (1971) 33
- 1971RA36 R.B. Raphael and M. Rosen, Part. Nucl. 2 (1971) 29
- 1971RE19 D. Rendic, N.D. Gabitzsch, V. Valkovic, W. von Witsch and G.C. Phillips, Nucl. Phys. A178 (1971) 49
- 1971RE1H Rej, Lett. Nuovo Cim. 1 (1971) 882
- 1971SA31 W. Salathe, E. Baumgartner and P. Huber, Helv. Phys. Acta 44 (1971) 815
- 1971SC19 M. Schaeffer, M. Suffert and D. Magnac-Valette, Nucl. Phys. A175 (1971) 217; Erratum Nucl. Phys. A196 (1972) 639
- 1971SC1N Schnabel, FRNC TH 191 (1971)
- 1971SE02 R. Seki, Phys. Rev. C3 (1971) 454
- 1971SH26 V.S. Shirley, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1255
- 1971SI1J Simons, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 597
- 1971SQ03 G.T.A. Squier, A.M. Hanna, J.B.A. England, D.L. Clough and L.H. Watson, Nucl. Phys. A170 (1971) 571
- 1971ST35 D.M. Stanojevic, M.R. Aleksic, B.Z. Stepancic and R.V. Popic, Fizika 3 (1971) 99
- 1971TR03 G.F. Trentelman, B.M. Preedom and E. Kashy, Phys. Rev. C3 (1971) 2205
- 1971TU04 V.G. Tuskiya and G.A. Chilashvili, Yad. Fiz. 14 (1971) 1158; Sov. J. Nucl. Phys. 14 (1972) 645
- 1971VA34 S.S. Vasilev, B.S. Galakhmatova, A.S. Demyanova, T.N. Mikhaleva, E.A. Romanovskii and D.L. Chuprunov, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 2364; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 2144
- 1971WA1K Watkins, Ludwig, Clegg and Dzubay, Bull. Amer. Phys. Soc. 16 (1971) 621
- 1971WA37 A.H. Wapstra and N.B. Gove, Nucl. Data Tables 9 (1971) 267
- 1971WE1L Werby, Thesis, Florida State Univ. (1971)
- 1971WI18 D.H. Wilkinson, Phys. Rev. Lett. 27 (1971) 1018
- 1971YO04 P.G. Young and R.H. Stokes, Phys. Rev. C4 (1971) 1597

- 1971ZA07 A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 273
- 1972AB19 Y. Abe and N. Takigawa, Suppl. Prog. Theor. Phys. 52 (1972) 228
- 1972AB1H Abramian et al, Yad. Fiz. 16 (1972) 739
- 1972AD04 J.C. Adloff, W.K. Lin, K.H. Souw and P. Chevallier, Phys. Rev. C5 (1972) 664
- 1972AH1B Ahrens et al., Proc. Int. Conf. on Nucl. Struct. Studies Using Electron Scattering and Photoreaction, Sendai, Japan (1972) 213
- 1972AN05 R.K. Anderson, M.R. Wilson and P. Goldhammer, Phys. Rev. C6 (1972) 136
- 1972AN09 Y.P. Antufev, V.L. Agranovich, V.B. Ganenko, V.S. Kuzmenko, I.I. Miroshnichenko, P.V. Sorokin and S.V. Shalatskii, Yad. Fiz. 15 (1972) 643; Sov. J. Nucl. Phys. 15 (1972) 357
- 1972AR22 U.R. Arifkhanov, M. Gulyamov, B.I. Islamov and E. Ergashov, Yad. Fiz. 15 (1972) 1102; Sov. J. Nucl. Phys. 15 (1972) 610
- 1972AV04 G.V. Avakov, E.I. Dolinsky and V.V. Turovtsev, Nucl. Phys. A196 (1972) 529
- 1972BE1E Bertsch and Mekjian, Ann. Rev. Nucl. Sci. (1972) 25
- 1972BE34 H.W. Bertini, Phys. Rev. C6 (1972) 631
- 1972BL12 R.E. Bluvshstein, V.M. Dubovik and A.A. Cheshkov, Yad. Fiz. 15 (1972) 100; Sov. J. Nucl. Phys. 15 (1972) 59
- 1972BO01 A. Bottino and G. Ciocchetti, Nucl. Phys. A178 (1972) 593
- 1972BR1R Bray, Cameron, Neilson and Sharma, Bull. Amer. Phys. Soc. 17 (1972) 112
- 1972BU1P Budyashov et al., Sov. Phys. JETP 35 (1972) 13
- 1972BU1R Buchnea et al, Bull. Amer. Phys. Soc. 17 (1972) 549
- 1972BU29 A.P. Bukhvostov, A.M. Chatrchyan, G.E. Dogotar, R.A. Eramzhyan, N.P. Popov and V.A. Vartanjan, Acta Phys. Pol. B3 (1972) 375
- 1972BU30 A.J. Buffa, Jr. and M.K. Brussel, Nucl. Phys. A195 (1972) 545
- 1972CA1Q Carlson, Lind and Zafiratos, Bull. Amer. Phys. Soc. 17 (1972) 71
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CE1A Cerny, At. Masses Fund. Constants, Teddington, 1971 (1972) 26
- 1972CH1N Chand and Knight, Bull. Amer. Phys. Soc. 17 (1972) 199
- 1972CH1P Chu, Han and Lin, Bull. Amer. Phys. Soc. 17 (1972) 609
- 1972CI05 D. Ciric, B. Stepanic, R. Popic, D. Stanojevic and M. Aleksic, Fizika 4 (1972) 40, 193
- 1972CL1A Clayton, Encyclopedia of the Twentieth Century (1972)
- 1972CR1E Crawford, 4th Ainslie Nucl. Phys. Conf., Sydney, 1972 (1972) 30

- 1972DE01 R.M. Devries, J.W. Sunier, J.-L. Perrenoud, M. Singh, G. Paic and I. Slaus, Nucl. Phys. A178 (1972) 417
- 1972DE02 R.M. Devries, J.-L. Perrenoud, I. Slaus and J.W. Sunier, Nucl. Phys. A178 (1972) 424
- 1972DE44 J. Decharge, G. Surget, G. Bruno and M.Y. Decharge, J. Phys. (France) 33 (1972) 485
- 1972DM01 T.A. Dmitrieva and I.B. Teplov, Yad. Fiz. 15 (1972) 478; Sov. J. Nucl. Phys. 15 (1972) 266
- 1972ES05 J.E. Esterl, D. Allred, J.C. Hardy, R.G. Sextro and J. Cerny, Phys. Rev. C6 (1972) 373
- 1972FR09 V. Franco, Phys. Rev. C6 (1972) 748
- 1972GI07 R.W. Givens, M.K. Brussel and A.I. Yavin, Nucl. Phys. A187 (1972) 490
- 1972HA57 H.H. Hackenbroich and T.H. Seligman, Phys. Lett. B41 (1972) 102
- 1972HI16 J. Hiura and R. Tamagaki, Suppl. Prog. Theor. Phys. 52 (1972) 25
- 1972HU1A W.T. Huang, C.A. Levinson and M.K. Banerjee, Phys. Rev. C5 (1972) 651
- 1972IK1A Ikeda, Marumori, Tamagaki and Tanaka, Suppl. Prog. Theor. Phys. 52 (1972) 1
- 1972JA10 J.A. Jansen, R.T. Peerdeman and C. de Vries, Nucl. Phys. A188 (1972) 337
- 1972KO1E Koike, Prog. Theor. Phys. 48 (1972) 66
- 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353
- 1972LI31 Y.C. Liu, Chin. J. Phys. (Taiwan) 10 (1972) 76
- 1972MA1H Macek, Bull. Amer. Phys. Soc. 17 (1972) 33
- 1972MA21 C. Maples and J. Cerny, Phys. Lett. B38 (1972) 504
- 1972MA47 V.A. Matusевич, V.A. Solov'ev and I.P. Chernov, Yad. Fiz. 15 (1972) 670; Sov. J. Nucl. Phys. 15 (1972) 375
- 1972MA62 A.M. MacLeod and G.R. Milne, J. Phys. A5 (1972) 1252
- 1972MC1E McDonald et al, Bull. Amer. Phys. Soc. 17 (1972) 464
- 1972ME07 F. Merchez, J. Arvieux, T. Honda and S. Morita, Nucl. Phys. A183 (1972) 417
- 1972NA05 P.T. Nang, Nucl. Phys. A185 (1972) 413
- 1972OB01 A.W. Obst, T.B. Grandy and J.L. Weil, Phys. Rev. C5 (1972) 738
- 1972OH01 I.K. Oh, C.S. Zaidins, C.D. Zafiratos and S.I. Hayakawa, Nucl. Phys. A178 (1972) 497
- 1972PA1C Parker, Astrophys. J. 175 (1972) 261
- 1972PN1A Pniewski, Few Particle Problems, UCLA, 1972 (1972) 145
- 1972QU01 J.R. Quinn, M.B. Epstein, S.N. Bunker, J.W. Verba and J.R. Richardson, Nucl. Phys. A181 (1972) 440
- 1972RA1J Rayet, Nucl. Phys. B38 (1972) 387

- 1972RA34 G.H. Rawitscher, Phys. Rev. C6 (1972) 1212
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430
- 1972SA10 W.W. Sapp, Jr., M. Eckhause, G.H. Miller and R.E. Welsh, Phys. Rev. C5 (1972) 690
- 1972SC1U Schilling et al, Bull. Amer. Phys. Soc. 17 (1972) 930
- 1972SE09 F. Seiler, Nucl. Phys. A187 (1972) 379
- 1972SE1F Seki, Bull. Amer. Phys. Soc. 17 (1972) 917
- 1972SO03 P.M. Sood and S.K. Tuli, Nucl. Phys. A187 (1972) 153
- 1972SU1E Surget, CEA R 4270 (1972)
- 1972TA31 Y. Takeichi, T. Une, K. Nakamura, T. Kohmura and T. Miyazima, Prog. Theor. Phys. 48 (1972) 858
- 1972TH12 B.W. Thomas, D.M. Crawford and H.H. Thies, Nucl. Phys. A196 (1972) 89
- 1972TH13 C. Thibault and R. Klapisch, Phys. Rev. C6 (1972) 1509
- 1972THZF H. Theissen, Springer Tracts in Mod. Phys.; Ed., Hohler, Vol. 65 (1972) 1
- 1972VO06 A.A. Vorobyov, D.M. Seliverstov, V.T. Grachev, I.A. Kondurov, A.M. Nikitin, N.N. Smirnov and Y.K. Zalite, Phys. Lett. B40 (1972) 102
- 1972WA07 G.E. Walker and R.H. Stokes, Part. Nucl. 3 (1972) 1
- 1972WA1M Watkins, Ludwig, Clegg and Dzubay, Bull. Amer. Phys. Soc. 17 (1972) 200
- 1972WE08 K.A. Weber, K. Meier-Ewert, H. Schmidt-Bocking and K. Bethge, Nucl. Phys. A186 (1972) 145
- 1972WI1C Wilkinson, Few Particle Problems, UCLA, 1972 (1972) 191
- 1972WI28 D.H. Wilkinson, Proc. Roy. Soc. Edinburgh A70 (1972) 307
- 1973AB10 S.N. Abramovich, B.Y. Guzhovskii, A.G. Zvenigorodskii and S.V. Trusillo, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1967; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 144
- 1973AH1A Ahrens et al, in Asilomar (1973) 23
- 1973AU1H Audouze and Truran, OAP-310 (1973)
- 1973BA1Y Bamberger et al, Nucl. Phys. B60 (1973) 1
- 1973BA2R Backenstoss et al, Contrib., Uppsala (1973) 146
- 1973BA2V Backenstoss et al, in Munich, 1 (1973) 317
- 1973BE19 J.C. Bergstrom, I.P. Auer, M. Ahmad, F.J. Kline, J.H. Hough, H.S. Caplan and J.L. Groh, Phys. Rev. C7 (1973) 2228
- 1973BE1N Belyaev, Dmitriev and Rumyantsev, in Munich, 1 (1973) 556
- 1973CL09 C.F. Clement, Nucl. Phys. A213 (1973) 469

- 1973CO1N Costa, in Asilomar (1973) 1319
- 1973DA1R D'Amico, Jannelli, Mazzaneres and Potenza, Nuovo Cim. A15 (1973) 723
- 1973DI1H Dillig and Huber, Contrib., Uppsala (1973) 107
- 1973DO13 P. Dougan and W. Stiefler, Z. Phys. 265 (1973) 1
- 1973GA20 N.D. Gabitzsch, G.S. Mutchler, C.R. Fletcher, E.V. Hungerford, L. Coulson, D. Mann, T. Witten, M. Furic, G.C. Phillips, B. Mayes et al, Phys. Lett. B47 (1973) 234
- 1973GE1J Geiger, van der Zwan and Werle, Phys. in Canada 29 (1973) 4
- 1973GU08 V.B. Gubin, S.S. Vasilev, B.S. Galakhmatova, T.N. Mikhaleva, E.A. Romanovskii, D.L. Chuprunov and L.I. Nikitina, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 101; Bull. Acad. Sci. USSR Phys. Ser. 37 (1973) 89
- 1973HA49 P.S. Hauge and S. Maripuu, Phys. Rev. C8 (1973) 1609
- 1973HA77 J.C. Hardy, Nucl. Data Tables A11 (1973) 327
- 1973HE06 J.C.P. Heggie and P.W. Martin, Phys. Lett. B43 (1973) 289
- 1973HE26 J.C.P. Heggie and P.W. Martin, Nucl. Phys. A212 (1973) 78
- 1973HI03 S. Hiramatsu, T. Kamae, H. Muramatsu, K. Nakamura, N. Izutsu and Y. Watase, Phys. Lett. B44 (1973) 50
- 1973HO30 H.J. Hoffman and M.B. Epstein, Nucl. Phys. A210 (1973) 126
- 1973HS1A Hsieh, Kane, Spence and Carroll, Bull. Amer. Phys. Soc. 10 (1973) 691
- 1973HS1B Hsieh et al, Contrib., Uppsala (1973) 145
- 1973HU05 A.H. Hussein and H.S. Sherif, Phys. Rev. C8 (1973) 518
- 1973HU1G Hughes, Sambell and Spicer, Asilomar (1973) Paper 2B4
- 1973JA1K James et al, Bull. Amer. Phys. Soc. 18 (1973) 1428
- 1973JU1E Julien et al, Contrib., Uppsala (1973) 171
- 1973JU2A M. Juric, G. Bohm, J. Klabuhn, U. Kreckler, F. Wysotzki, G. Coremans-Bertrand, J. Sacton, G. Wilquet, T. Cantwell, F. Esmael et al, Nucl. Phys. B52 (1973) 1
- 1973KO1D Kovar, Symp. on Heavy Ion Transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 59
- 1973KO1J Kochetov and Khrylin, JETP Lett. 18 (1973) 79
- 1973KO1M Komarov et al, Contrib., Uppsala (1973) 179
- 1973KU03 D. Kurath, Phys. Rev. C7 (1973) 1390
- 1973KU13 N. Kumar, Lett. Nuovo Cim. 6 (1973) 224
- 1973LA19 H. Laumer, S.M. Austin, L.M. Panggabean and C.N. Davids, Phys. Rev. C8 (1973) 483
- 1973LA1T Lapikas, Box and De Vries, in Munich, 1 (1973) 619

- 1973MA1K Maripuu, 5th Symp. Struct. Low-Medium Mass Nuclei (1973) 63
- 1973MA24 R.E. Marrs, D. Bodansky and E.G. Adelberger, Phys. Rev. C8 (1973) 427
- 1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
- 1973MCZW C.R. McClenahan, R.E. Segel and R.P. Redwine, Bull. Amer. Phys. Soc. 18 (1973) 651, GJ7
- 1973MI1J Miller et al, Bull. Amer. Phys. Soc. 18 (1973) 1382
- 1973MO01 D.G. Montague, R.K. Cole, P.S. Lewis, C.N. Waddell and D.L. Hendrie, Nucl. Phys. A199 (1973) 433
- 1973MU11 N.C. Mukhopadhyay, Phys. Lett. B45 (1973) 309
- 1973MU12 T. Mukoyama, Y. Isozumi, T. Kitahara and S. Shimizu, Phys. Rev. C8 (1973) 1308
- 1973NY04 E.M. Nyman, Nucl. Phys. A215 (1973) 397
- 1973OK1B Okabe, Abe and Tanaka, in Munich, 1 (1973) 116
- 1973OS01 P. Osland, Nucl. Phys. B56 (1973) 308
- 1973PE1E Pedroni et al, Contrib., Uppsala (1973) 75
- 1973RA1E Rayet, Nucl. Phys. B57 (1973) 269
- 1973RA37 G.M. Raisbeck, J. Lestringuez and F. Yiou, Nature 244 (1973) 28
- 1973RE1G Reeves, Audouze, Fowler and Schramm, Astrophys. J. 179 (1973) 909
- 1973RO2F Roeckl et al, in Munich, 1 (1973) 326
- 1973SL02 A.G. Slight, T.E. Drake and G.R. Bishop, Nucl. Phys. A208 (1973) 157
- 1973SQ01 G.T.A. Squier, M.E. Cage, G.J. Pyle, A.S. Clough, G.K. Turner, B.W. Allardyce, C.J. Batty, D.J. Baugh, W.J. McDonald, R.A.J. Riddle et al, Phys. Rev. Lett. 31 (1973) 389
- 1973TO14 I.S. Towner, Nucl. Phys. A216 (1973) 589
- 1973UL1D Ullrich, Boschitz, Engelhardt and Lewis, Contrib., Uppsala (1973) 156
- 1973VA08 S.S. Vasilev, B.S. Galakhmatova, A.S. Demyanova, T.N. Mikhaleva, E.A. Romanovskii, D.L. Chuprunov and L.I. Nikitina, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 105; Bull. Acad. Sci. USSR Phys. Ser. 37 (1973) 92
- 1973VO02 H.J. Votava, T.B. Clegg, E.J. Ludwig and W.J. Thompson, Nucl. Phys. A204 (1973) 529; Erratum Nucl. Phys. A217 (1973) 628
- 1973VO07 U. von Mollendorff, A. Janett, F. Seiler and H.R. Striebel, Nucl. Phys. A209 (1973) 323
- 1973WA28 F.M. Waterman and R. Madey, Phys. Rev. C8 (1973) 2419
- 1973WE03 H. Werle, L. Van der Zwan and K.W. Geiger, Z. Phys. 259 (1973) 275
- 1973WE19 K.A. Weaver, J.D. Anderson, H.H. Barschall and J.C. Davis, Nucl. Sci. Eng. 52 (1973) 35

1973WI07 N. Willis, I. Brissaud, Y. Le Bornec, B. Tatischeff and G. Duhamel, Nucl. Phys. A204 (1973) 454

1973WI11 D.H. Wilkinson, Nucl. Phys. A209 (1973) 470

1973WI15 J. Wilczynski, Phys. Lett. B47 (1973) 124

1973WO1E Woody et al, Bull. Amer. Phys. Soc. 18 (1973) 79

1973ZW1A Zwiaglinski, Piotrowski, Saganek and Sledzinska, Nucl. Phys. A209 (1973) 348

1974AU1A Audouze and Tinsley, Astrophys. J. 192 (1974) 487

1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1

1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1

KO67Q Unknown Source

