

Energy Levels of Light Nuclei $A = 10$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A227* (1974), p. 1. This version of $A = 10$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 31, 1973)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 10$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{10}\text{He}\$](#) , [\$^{10}\text{Li}\$](#) , [\$^{10}\text{Be}\$](#) , [\$^{10}\text{B}\$](#) , [\$^{10}\text{C}\$](#)

B. Tables of Recommended Level Energies:

[Table 10.1](#): Energy levels of ^{10}Be

[Table 10.5](#): Energy levels of ^{10}B

[Table 10.24](#): Energy levels of ^{10}C

C. [References](#)

D. Figures: [\$^{10}\text{Be}\$](#) , [\$^{10}\text{B}\$](#) , [\$^{10}\text{C}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

¹⁰He

(Not illustrated)

¹⁰He has not been observed when ²³²Th has been bombarded with 145 MeV ¹⁵N ions (1971AR21), nor in the spontaneous fission of ²⁵²Cf (1967CO36). See also (1966PO09). It is particle unstable: the calculated $(M - A) = 52.75$ MeV. The breakup energies into ⁹He + n and ⁸He + 2n are calculated to be -1.1 and -4.9 MeV, respectively (1972TH13). See also (1966GA25, 1972GA1F).

See also (1968CE01, 1973KO1D) and (1969BA55, 1970RY03, 1970RY04, 1971LO13; theor.).

¹⁰Li

(Not illustrated)

¹⁰Li is not observed in the 5.3 GeV bombardment of uranium (1966PO09). Based on the tentative locations of the $T = 2$ analogs, ¹⁰Be*(21.3) [reaction 2 in ¹⁰Be: see (1970CH1Q, 1973AB10)] and ¹⁰B*(23.0) [reaction 9 in ¹⁰B: see (1968HA1K)], the mass excess of ¹⁰Li, $(M - A) = 33.10 \pm 0.06$ MeV (1973AB10). The breakup energy into ⁹Li + n is then -0.06 ± 0.06 MeV. Using the calculated values suggested by (1969GA1G, 1972TH13), $E_b = -1.1 \pm 0.3$ MeV. See also (1966GA25) and (1968CE01).

¹⁰Be

(Figs. 19 and 22)

GENERAL: (See also (1966LA04).)

Shell model: (1961KO1A, 1968FA1B, 1969SA1A, 1970BO35, 1970KA32, 1973KU03, 1973SA30).

Cluster and alpha-particle model: (1973KU03).

Special levels: (1970BO35, 1970FR1C, 1970PE18, 1973SA30).

Electromagnetic transitions: (1965ST22, 1973SA30).

Astrophysical questions: (1967KO1F, 1967SH1B, 1971LE33, 1971RA1K, 1972SI1G, 1973LA19, 1973RA37, 1974SI1H).

Special reactions: (1967KO1F, 1967SH1B, 1968YI01, 1969AR13, 1969GA18, 1969GO1H, 1969YI1A, 1971AR02, 1971LE33, 1971RA1K, 1972AM04, 1972PN1A, 1972SI1G, 1972VO06, 1973KO1D, 1973KU03, 1973LA19, 1973VO1G, 1973WI15).

Pion capture and reactions (See also reaction 12.): (1971DA10, 1971KA62, 1972FA14, 1972FU1C, 1973KA1D).

Other topics: (1968FA02, 1968FA1B, 1968JO1E, 1968NE1C, 1969SA1A, 1970FO1B, 1970PE18, 1971BA2Y, 1971ST40, 1972AN05, 1972CA37, 1972PN1A, 1973BU20, 1973JU2A, 1973KU03, 1973MA48, 1973MU11).

Table 10.1: Energy levels of ^{10}Be

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
g.s.	$0^+; 1$	$\tau_{1/2} = (1.6 \pm 0.2) \times 10^6 \text{ y}$	β^-	1, 3, 4, 5, 12, 13, 14, 17, 18, 19, 21, 22, 24, 25, 26, 27, 28, 30
3.3680 ± 0.2	$2^+; 1$	$\tau_m = 180 \pm 17 \text{ fsec}$	γ	3, 4, 5, 12, 13, 18, 19, 21, 24, 26, 28
5.9583 ± 0.3	$2^+; 1$	$\tau_m < 80 \text{ fsec}$	γ	4, 5, 12, 13, 28
5.9599 ± 0.6	$1^-; 1$		γ	4, 12, 13, (28)
6.1793 ± 0.7	$0^+; 1$	$\tau_m = 1.1^{+0.4}_{-0.3} \text{ psec}$	π, γ	4, 12, 13
6.2633 ± 5	$2^-; 1$		γ	4, 12, 13
7.371 ± 1	$3^-; 1$	$\Gamma = 15.7 \pm 0.5 \text{ keV}$	n	4, 6, 12, 13
7.542 ± 1	$2^+; 1$	$6.3 \pm 0.8 \text{ keV}$	n	6, 12, 13
9.27	$(4^-); 1$	$\approx 100 \text{ keV}$	n	4, 6, 12, 28
(9.4)	$(2^+); 1$	$\approx 400 \text{ keV}$	n	6, 12
10.7	$\geq 1; 1$		n	4, 6
(11.75 \pm 110)				4, 12
(17.4)			n, t	2
17.79		$110 \pm 35 \text{ keV}$	n, t	2, 4
18.55		$\approx 350 \text{ keV}$	γ, n, t	2, 4
(18.6)		$\approx 1.1 \text{ MeV}$	γ, t	2
(21.3)	$(T = 2)$	400 keV	p, t	2
(24.0)		broad		24

Ground-state properties: (1968FA02, 1968FA1B, 1971ST40, 1973SA30).

1. $^{10}\text{Be}(\beta^-)^{10}\text{B}$

$$Q_m = 0.5559$$

The half-life is $(1.6 \pm 0.2) \times 10^6 \text{ y}$ (1972EM01), $(1.7 \pm 0.4) \times 10^6 \text{ y}$ (1972MC26) and $(1.5 \pm 0.3) \times 10^6 \text{ y}$ (1972YI01): we adopt $\tau_{1/2} = (1.6 \pm 0.2) \times 10^6 \text{ y}$: $\log ft = 13.42$ (B. Zimmerman and G. Fox, private communication). The spectrum is of the D₂ type (1950WU1A). For a discussion of astrophysical and nuclear chronology considerations see (1972YI01). See also (1969VA1C, 1970WA11; theor.) and (1966BR1N).

2. (a) ${}^7\text{Li}(t, \gamma){}^{10}\text{Be}$	$Q_m = 17.251$	
(b) ${}^7\text{Li}(t, n){}^9\text{Be}$	$Q_m = 10.4389$	$E_b = 17.251$
(c) ${}^7\text{Li}(t, p){}^9\text{Li}$	$Q_m = -2.397$	
(d) ${}^7\text{Li}(t, d){}^8\text{Li}$	$Q_m = -4.2249$	
(e) ${}^7\text{Li}(t, \alpha){}^6\text{He}$	$Q_m = 9.837$	

The yield of γ_0 and γ_1 (to ${}^{10}\text{Be}^*(0, 3.4)$) has been studied at 90° for $E_t = 0.5$ to 3.0 MeV. The γ_1 yield shows a peak at $E_t = 1.85$ MeV, $\Gamma_{\text{lab}} \approx 0.5$ MeV. The γ_0 yield peaks near the same energy but with a width $\Gamma_{\text{lab}} \approx 1.5$ MeV (1973BL1B).

The neutron yield exhibits a weak structure at $E_t = 0.24$ MeV and broad resonances at $E_t \approx 0.77$ MeV [$\Gamma = 160 \pm 50$ keV] and 1.74 MeV (1960SE12, 1961VA43, 1962SE1A).

The yields of ${}^9\text{Li}$ and ${}^8\text{Li}$ (reactions (c) and (d)) have been studied from threshold to $E({}^7\text{Li}) = 21$ MeV (1970CH1Q) and $E_t = 6.5$ MeV (1973AB10). A sharp minimum is observed in the ${}^9\text{Li}$ yield at $E({}^7\text{Li}) \approx 13.6$ MeV (1970CH1Q) and $E_t = 5.62$ MeV (1973AB10). It may correspond to a $T = 2$ state in ${}^{10}\text{Be}$ at $E_x = 21.2$ MeV, $\Gamma_{\text{c.m.}} = 0.4$ MeV (1970CH1Q, 1973AB10).

For reaction (e) see (1966LA04). See also ${}^6\text{He}$, ${}^8\text{Li}$, ${}^9\text{Li}$ and ${}^9\text{Be}$.

3. ${}^7\text{Li}(\alpha, p){}^{10}\text{Be}$	$Q_m = -2.5638$
---	-----------------

Angular distributions of the p_0 and p_1 groups have been measured at $E_\alpha = 30$ MeV (1960KL03, 1972ME07). See also (1966LA04).

4. ${}^7\text{Li}({}^7\text{Li}, \alpha){}^{10}\text{Be}$	$Q_m = 14.784$
---	----------------

Angular distributions are reported for $E({}^7\text{Li}) = 2.1$ to 5.75 MeV (1969CA1A; $\alpha_1, \alpha_{2+3+4+5}$), 3.5 to 5.75 MeV (1969CA1A; α_0) and 30.3 MeV (1971GL07; α -groups to ${}^{10}\text{Be}^*(0, 3.4, 6.0, 7.4, 9.4, 10.7, 11.9, 17.9)$). (1971GL07) also report an α -group corresponding to ${}^{10}\text{Be}^*(18.8)$. See also (1970OG1A), (1966RO1E; theor.) and ${}^{14}\text{C}$ in (1976AJ04).

5. ${}^9\text{Be}(n, \gamma){}^{10}\text{Be}$	$Q_m = 6.8120$
	$Q_0 = 6811.9 \pm 0.4$ keV (1966GR18);
	$Q_0 = 6812.5 \pm 1.0$ keV (1967RA24).

The thermal capture cross section is 9.2 ± 1.0 mb (1973MU14). Reported γ transitions are displayed in Table 10.2 (1960BA01, 1961JA19, 1963DR02, 1966GR18, 1969WE10). See (1968FOZY) for astrophysical considerations. See also (1968RO1H).

Table 10.2: Neutron capture γ -rays in ^{10}Be

E_γ ^b (keV)	Transition	Intensities ^a			E_x ^b (keV)
		A	B	C	
6809.4 \pm 0.4	capt. \rightarrow g.s.	62	70	65	5958.3 \pm 0.3
5955.9 \pm 0.5	5.96 ^d \rightarrow g.s.	1.4	2	\lesssim 2	
3443.3 \pm 0.3	capt. \rightarrow 3.37	11	15	11	3368.0 \pm 0.2
3367.4 \pm 0.2 ^c	3.37 \rightarrow g.s.	37	28	28	
2589.9 \pm 0.25	5.96 ^d \rightarrow 3.37	28	17	21	
853.5 \pm 0.3	capt. \rightarrow 5.96 ^d	29.4	16	24	

A: (1960BA01: see (1963DR02)).

B: (1961JA19).

C: (1963DR02).

^a Gamma rays per 100 captures.

^b (1966GR18).

^c Not Doppler broadened (1969WE10).

^d This is the 2^+ member of the doublet at $E_x = 5.96$ MeV: see reaction 13.

6. $^9\text{Be}(n, n)^9\text{Be}$

$$E_b = 6.8120$$

The older cross section data is summarized in (1964ST25). More recent measurements of the total cross section are reported for $E_n = 0.5$ to 20 MeV ((1971SC1P), and R.B. Schwartz, private communication), 1.9 to 4.6 MeV (1968JO1F), 2.5 to 15 MeV (1971FO1P, 1971FO24), 13.70 to 14.60 MeV (1968HU17, 1969AN1E, 1970AN1F), 23 to 45 MeV (1972BU1T), 36.3 to 58.9 MeV (1972AU01: at five energies in this range) and at a neutron momentum of 10 GeV/ c (1968EN1A). Angular distributions are summarized in (1970GA1A). See also ^9Be . The coherent scattering length (thermal, bound) is 7.74 ± 0.10 fm (1973MU14).

Five resonances are reported in the total cross section, at $E_n = 0.63, 0.82, 2.73, (2.85)$ and 4.3 MeV; see Table 10.3. Polarization and differential cross sections are reported for $E_n = 0.2$ to 2 MeV by (1961LA1A, 1962EL01, 1964LA04). Analysis of these data leads to the $3^-, 2^+$ assignments for $^{10}\text{Be}^*(7.37, 7.55)$ (1964LA04). Below $E_n = 0.5$ MeV the scattering cross section reflects the effect of bound 1^- and 2^- states, presumably $^{10}\text{Be}^*(5.960, 6.26)$. There is also indication of interference with s-wave background and with a broad $l = 1, J^\pi = 3^+$ state (1964LA04). The structure at $E_n = 2.73$ MeV is ascribed to two levels: a broad state at about 2.85 MeV with $J^\pi = 2^+$, and a narrow one, $\Gamma \approx 100$ keV, at $E_n = 2.73$ MeV with a probable assignment of $J^\pi = 4^-$ (1951BO45, 1966SC16). The 4^- assignment results from a study of the polarization of the n_0 group at $E_n = 2.60$ to 2.77 MeV. A rapid variation of the polarization over this interval is observed, and the data are consistent with $4^- (l = 2)$ for $^{10}\text{Be}^*(9.27)$ (1966SC16). The polariza-

Table 10.3: Resonances in ${}^9\text{Be}(n, n){}^9\text{Be}$

E_{res}^a (MeV \pm keV)	${}^{10}\text{Be}^*$ (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π	l	R (fm)	θ^2 (%)	Refs.
0.6220 ± 0.8^b	7.371	15.7 ± 0.5^b	3^-	2	5.6	7.5	(1951BO45, 1964LA04)
0.8118 ± 0.7^b	7.542	6.3 ± 0.8^b	2^+	1	5.6	0.28	(1951BO45, 1955WI25, 1964LA04)
2.73	9.27	≈ 100	(4^-)	(2)			(1951BO45, 1966SC16) ^b
(2.85)	9.4	≈ 400	(2^+)	(1)			(1951BO45)
4.3	10.7		≥ 1				(1961FO07)

^a (1962PE10) report an additional anomaly in the cross section at $E_n = 207$ keV.

^b (1971SC1P) and R.B. Schwartz, private communication.

tion of the n_0 group has also been studied at $E_n = 4$ MeV (1967GO36). See also (1966DA1B). A weak dip at $E_n \approx 4.3$ MeV is ascribed to a level with $J \geq 1$ (1961FO07).

See also (1966LA04), (1969MA39), (1966CH1C) and (1966AG1A, 1966JA08, 1966SE1E, 1967BE1F, 1967HO1F, 1972FR09; theor.).

7. (a) ${}^9\text{Be}(n, n'){}^9\text{Be}^*$

$$E_b = 6.8120$$

(b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$

$$Q_m = -1.6651$$

Data on non-elastic cross sections are summarized by (1964ST25). The processes involved include $(n, 2n)$ and $(n, n'){}^9\text{Be}^* \rightarrow n + {}^8\text{Be}$; the (n, α) process is relatively weak. The total non-elastic cross section rises rapidly from threshold to 730 ± 100 mb at $E_n = 5.2$ MeV (1968EA1B). The cross section is ≈ 600 mb at $E_n = 6$ MeV and then falls slowly to 500 mb at 14 MeV. In the range 3.5 to 6.0 MeV, ${}^9\text{Be}(n, n'){}^9\text{Be}^*(2.4)$ accounts for about half of the non-elastic cross section (1959MA34: see (1964ST25)). The $(n, 2n)$ cross section increases slowly from $E_n = 2.00$ to 2.70 (the threshold for the process ${}^9\text{Be}(n, n'){}^9\text{Be}^*(2.4) \rightarrow n + {}^8\text{Be}$) and then sharply, reaching a plateau with a value ≈ 580 mb at 6.40 MeV (1969HO45, 1973BL03). See also (1968EA1B).

See also (1966LA04), (1966JE1B) and (1966FE04, 1966SA20, 1967RO1E, 1968GR1F, 1969MA39, 1969PR17, 1971AL1E).

8. ${}^9\text{Be}(n, p){}^8\text{Li}$

$$Q_m = -12.836$$

$$E_b = 6.8120$$

See (1966LA04, 1972ED01) and (1970BA1L).

9. ${}^9\text{Be}(n, d){}^8\text{Li}$

$$Q_m = -14.664$$

$$E_b = 6.8120$$

Table 10.4: Radiative transitions in ${}^9\text{Be}(d, p){}^{10}\text{Be}$ ^a

E_x (keV)	Transition	ΔJ^π	Mult.	Branch (%)	τ_m (psec)	Γ_γ (meV)
3368.0 ± 0.2 ^b	$3.37 \rightarrow \text{g.s.}$	$2^+ \rightarrow 0^+$	E2	100	0.189 ± 0.020 ^g	3.48 ± 0.37
					0.160 ± 0.030 ^h	4.11 ± 0.78
5958.3 ± 0.3 ^{b,e}	$5.96 \rightarrow 3.37$	$2^+ \rightarrow 2^+$	M1	> 90	< 0.08 ^h	
	$5.96 \rightarrow \text{g.s.}$	$2^+ \rightarrow 0^+$	E2	< 10		
5959.9 ± 0.6 ^{c,e}	$5.96 \rightarrow \text{g.s.}$	$1^- \rightarrow 0^+$	E1	83_{-6}^{+10}	$1.1_{-0.3}^{+0.4}$ ^d	0.028 ± 0.012
	$5.96 \rightarrow 3.37$	$1^- \rightarrow 2^+$	E1	17_{-10}^{+6}		
6179.3 ± 0.7 ^d	$6.18 \rightarrow 5.96$	$0^+ \rightarrow 1^-$	E1	4.6 ± 1.5 ^d	$1.1_{-0.3}^{+0.4}$ ^d	0.57 ± 0.17
	$6.18 \rightarrow 3.37$	$0^+ \rightarrow 2^+$	E2	95 ± 2 ^d		
	$6.18 \rightarrow \text{g.s.}$	$0^+ \rightarrow 0^+$	E0	0.24 ± 0.08 ^d		
6263.3 ± 5 ^d	$6.26 \rightarrow 5.96$	$2^- \rightarrow 1^-$	M1	≤ 1 ^f	99_{-2}^{+1} ^f	$(1.4 \pm 0.6) \times 10^{-3}$
		$2^- \rightarrow 2^+$	E1			
	$6.26 \rightarrow 3.37$	$2^- \rightarrow 2^+$	E1			
	$6.26 \rightarrow \text{g.s.}$	$2^- \rightarrow 0^+$	M2	1 ± 1 ^f		

^a See also Tables 10.2 and 10.5 in (1966LA04).

^b From (1966GR18): ${}^9\text{Be}(n, \gamma){}^{10}\text{Be}$.

^c From (1966GR18, 1969RO12): see (1969AL17).

^d See (1969AL17).

^e (1966WA1C) and F.C. Young, private communication.

^f (1969RO12).

^g (1968FI09).

^h (1966WA10).

The cross section for the (n, d) reaction has been measured for $E_n = 16.3$ to 18.7 MeV (1969SC05).

$$10. \text{}^9\text{Be}(n, t)^7\text{Li} \qquad Q_m = -10.4389 \qquad E_b = 6.8120$$

See (1966JE1B, 1966LA04) and (1972BI1E).

$$11. \text{}^9\text{Be}(n, \alpha)^6\text{He} \qquad Q_m = -0.602 \qquad E_b = 6.8120$$

The cross section for production of ${}^6\text{He}$ shows a smooth rise to a broad maximum of 104 ± 7 mb at 3.0 MeV, followed by a gradual decrease to 70 mb at 4.4 MeV. No indication of resonance is found at $E_n = 2.7$ MeV (1957ST95). From $E_n = 3.9$ to 8.6 MeV, the cross section decreases smoothly from 100 mb to 32 mb (1961BA53). At $E_n = 14.4$ MeV, the total cross section for the α_0 transition is 11.7 ± 1.8 mb (1967PA03). See also (1966JE1B, 1966LA04) and (1970BA1L).

$$12. \text{}^9\text{Be}(p, \pi^+)^{10}\text{Be} \qquad Q_m = -133.539$$

At $E_p = 185$ MeV the π^+ spectrum shows groups corresponding to ${}^{10}\text{Be}^*(0, 3.37 \pm 0.12, 6.07 \pm 0.13, 7.39 \pm 0.13, 9.31 \pm 0.24, 11.75 \pm 0.11)$. Angular distributions have been obtained for the π^+ corresponding to these six states (1973DA09). See also reaction 3 in ${}^{10}\text{C}$ and (1971DA10).

$$13. \text{}^9\text{Be}(d, p)^{10}\text{Be} \qquad Q_m = 4.5873$$

$$Q_0 = 4.590 \pm 0.008 \text{ (1967SP09).}$$

Angular distributions of proton groups have been studied at many energies. Except at the lowest energies the stripping process appears to dominate: see (1966LA04) for the earlier references and (1968BE1E: 0.30 to 1.0 MeV; p_0), (1973SA1Q: 0.9 to 3.1 MeV; p_0, p_1), (1969RO12: 2.8 MeV; p_0, p_1 and p to ${}^{10}\text{Be}^*(5.958, 5.960)$), (1970PO03: 4.50 to 6.0 MeV; p_0, p_1), (1964SC12, 1967SC29: 11.8 MeV; p_0, p_1) and (1969AR1B: 15 MeV; p_0). The angular distributions show $l_n = 1$ transfer for ${}^{10}\text{Be}^*(0, 3.37, 5.958)$, $l_n = 0$ transfer for ${}^{10}\text{Be}^*(5.960, 6.26)$ and $l_n = 2$ transfer for ${}^{10}\text{Be}^*(7.37)$. ${}^{10}\text{Be}^*(6.18, 7.54)$ are also populated. See (1954JU23, 1969RO12) and Table 10.4 in (1966LA04).

Attempts to understand the γ -decay of ${}^{10}\text{Be}^*(5.96)$ and its population in ${}^9\text{Be}(n, \gamma)^{10}\text{Be}$ led to the discovery that it consisted of two states separated by 1.6 ± 0.5 keV. The lower of the two has $J^\pi = 2^+$ and decays primarily by a cascade transition via ${}^{10}\text{Be}^*(3.37)$ [it is the state fed in the

$^9\text{Be}(n, \gamma)$ decay]; the higher state has $J^\pi = 1^-$ and goes mainly by a crossover to $^{10}\text{Be}_{g.s.}$. Angular correlations measured with the γ -ray detector located normal to the reaction plane (\equiv angular distributions) lead to l_n values consistent with the assignments of 2^+ and 1^- for $^{10}\text{Be}^*(5.9658, 5.9660)$ obtained from the character of the γ -decay (1969RO12). $^{10}\text{Be}^*(6.18)$ decays primarily to $^{10}\text{Be}^*(5.9660)$: $E_\gamma = 219.4 \pm 0.3$ keV (1969AL17). See Table 10.4 for a listing of the information on radiative transitions obtained in this reaction (1969AL17, 1969RO12), and lifetime measurements (1966WA10, 1968FI09, 1969AL17). For (p, γ) correlations through $^{10}\text{Be}^*(3.37)$ see (1966LA04) and (1967FA03, 1969CA1F).

The probabilities $p_{3/2}$ for transfer of a neutron with angular momentum $\frac{3}{2}$ have been determined for the p_0 and p_1 groups using vector polarized deuterons with $E_d = 10$ and 12 MeV, and compared with the shell model calculations of (1965CO25, 1967CO32) [(1970FI07); and see also (1967SC29)]. For polarization measurements see ^{11}B in (1975AJ02) and (1966LA04).

See also (1968LA1F), (1966WA1C, 1967LE1F, 1967WA1C, 1967YO1C, 1968SC1E, 1969CO1H) and (1971JO22, 1972DZ06, 1973DO02; theor.).

$$14. \ ^9\text{Be}(t, d)^{10}\text{Be} \qquad Q_m = 0.5544$$

The d_0 group has been observed at $E_t = 1.6$ to 3.5 MeV (1967BI1E). See also (1969NA04) and ^{12}B in (1975AJ02).

$$15. \ ^9\text{Be}(\alpha, \ ^3\text{He})^{10}\text{Be} \qquad Q_m = -13.7665$$

Not reported.

$$16. \ ^9\text{Be}(^{14}\text{N}, \ ^{13}\text{N})^{10}\text{Be} \qquad Q_m = -3.742$$

See (1973TO05; theor.).

$$17. \ ^9\text{Be}(^{18}\text{O}, \ ^{17}\text{O})^{10}\text{Be} \qquad Q_m = -0.679$$

The angular distribution has been measured at $E(^{18}\text{O}) = 16$ and 20 MeV (1971KN05).

$$18. \ (a) \ ^{10}\text{Be}(p, p)^{10}\text{Be}$$

$$(b) \ ^{10}\text{Be}(d, d)^{10}\text{Be}$$

Angular distributions of the p_0 and p_1 groups have been measured at $E_p = 12.0$ to 16.0 MeV. The elastically scattered deuterons have been studied at $E_d = 12.0$ and 15.0 MeV. $B(E2)$ for the $^{10}\text{Be}^*(3.4) \xrightarrow{\gamma} ^{10}\text{Be}(0)$ transition is $23.8 e^2 \cdot \text{fm}^4$ (from a comparison of the angular distributions with DWBA predictions) (1970AU02).

19. $^{10}\text{B}(n, p)^{10}\text{Be}$ $Q_m = 0.2266$

See (1966GO1D, 1970NE03).

20. $^{10}\text{B}(t, ^3\text{He})^{10}\text{Be}$ $Q_m = -0.5372$

Not reported.

21. $^{11}\text{B}(\gamma, p)^{10}\text{Be}$ $Q_m = -11.2294$

Transitions to $^{10}\text{Be}^*(3.4)$ from the upper region of the giant resonance in ^{11}B are reported to be about twice as intense as those from the lower region (1971PA10): see ^{11}B in (1975AJ02). Angular distributions of the p_0 and p_1 groups have been measured with $E_{bs} = 18.5$ MeV (1970SO03). See also (1969MU10).

22. $^{11}\text{B}(n, d)^{10}\text{Be}$ $Q_m = -9.0047$

The angular distribution of the d_0 group has been measured at $E_n = 14.4$ MeV (1968MI10, 1971MI12).

23. $^{11}\text{B}(e, ep)^{10}\text{Be}$ $Q_m = -11.2294$

See (1969BA1F; theor.).

24. $^{11}\text{B}(p, 2p)^{10}\text{Be}$ $Q_m = -11.2294$

Structure is observed in the summed proton spectrum corresponding to $Q = -10.9 \pm 0.35$, -14.7 ± 0.4 , -21.1 ± 0.4 , -35 ± 1 MeV: see (1966LA04, 1966TY01). See also (1971RA26), (1966JA09, 1967KO1B, 1968JA1D; theor.) and ^{11}B in (1975AJ02).

25. $^{11}\text{B}(\text{d}, ^3\text{He})^{10}\text{Be}$ $Q_m = -5.7356$

Angular distributions have been obtained for the ^3He group to $^{10}\text{Be}(0)$ at $E_d = 11.8$ MeV (1967FI07) and at 22 MeV (1971ST05: also ^3He to $^{10}\text{Be}^*(3.4)$).

26. $^{11}\text{B}(\text{t}, \alpha)^{10}\text{Be}$ $Q_m = 8.585$

Angular distributions of α_0 and α_1 have been measured for $E_t = 1.0$ to 2.1 MeV (1969SI12).

27. $^{11}\text{B}(^{14}\text{N}, ^{15}\text{O})^{10}\text{Be}$ $Q_m = -3.937$

Angular distributions of the ground-state transitions have been measured at $E(^{14}\text{N}) = 41, 77$ and 113 MeV (1971LI11). See also (1973SC1J) and (1973DE35; theor.).

28. $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}$ $Q_m = -21.443$

At $E(^6\text{Li}) = 80$ MeV, $^{10}\text{Be}^*(0, 3.37, 5.95 \pm 0.05)$ are strongly populated. There is also some evidence for the excitation of $^{10}\text{Be}^*(9.27 \pm 0.1)$. Angular distributions have been measured to the first three of these states (1973CE1B).

29. $^{12}\text{C}(^{11}\text{B}, ^{13}\text{N})^{10}\text{Be}$ $Q_m = -9.286$

See (1969BR1D).

30. $^{13}\text{C}(\text{n}, \alpha)^{10}\text{Be}$ $Q_m = -3.836$

See ^{14}C in (1976AJ04).

31. (a) $^{13}\text{C}(^{11}\text{B}, ^{14}\text{N})^{10}\text{Be}$ $Q_m = -3.679$
 (b) $^{15}\text{N}(^{11}\text{B}, ^{16}\text{O})^{10}\text{Be}$ $Q_m = 0.898$
 (c) $^{16}\text{O}(^{11}\text{B}, ^{17}\text{F})^{10}\text{Be}$ $Q_m = -10.629$

See (1969BR1D).

¹⁰B
(Figs. 20 and 22)

GENERAL: (See also (1966LA04).)

Shell model: (1961KO1A, 1965CO25, 1966HA18, 1966MA1P, 1966WI1E, 1967CO32, 1967EV1C, 1967HS1A, 1967PI1B, 1968GO01, 1969VA1C, 1970CO1H, 1971NO02, 1972LE1L, 1973HA49, 1973JO1K, 1973KU03, 1973SA30).

Cluster and α -particle model: (1965NE1B, 1967TA1C, 1969BA1J, 1969HU1F, 1969NAZY, 1970NA06, 1971NO02, 1972LE1L, 1973KU03).

Special levels: (1967CO32, 1967HS1A, 1968GO01, 1968HE1G, 1969CO1H, 1969HA1F, 1971NO02, 1972VA36, 1973SA30).

Electromagnetic transitions: (1962MO1A, 1965CO25, 1967HS1A, 1967KU1E, 1968BI1C, 1968HE1G, 1968KU1D, 1969HA1F, 1969VA1C, 1972BE1E, 1972EV03, 1972LO1D, 1972TA21, 1972NA05, 1973HA49, 1973SA30).

Astrophysical questions: (1968HA1C, 1970BA1M, 1972CL1A, 1972KO1E, 1973AU1H, 1973LA19, 1973RE1G, 1974AU1A).

Special reactions: (1968YI01, 1969DA1D, 1969GA18, 1969YI1A, 1971AR02, 1971BA16, 1972OB1B, 1972PN1A, 1973KU03, 1973LA19, 1973VO1G).

Muon capture: (1967BA78, 1967BU1D, 1968BA2G, 1969WU1A, 1970FA15, 1971DE2D, 1972BU29, 1973BU20, 1973MU11, 1974MU1J).

Pion capture and reactions: (1965CH12, 1968BA2G, 1968BO32, 1968NO1A, 1968RI1H, 1968WI1B, 1968TA1C, 1969BU1C, 1969CH1C, 1969CH19, 1969GO1C, 1969KAZZ, 1969MO1E, 1970CH1F, 1970BA1E, 1971CA01, 1971CA1J, 1971KA62, 1972FU1C[†], 1972HU1A, 1972SW1A, 1973KA1D, 1973NY04, 1973UL1D).

Kaon capture and reactions: (1972BA09, 1973CH1M).

Other topics: (1965CO25, 1966HA18, 1966WI1E, 1967CA17, 1967EV1C, 1967MI1A, 1968GO01, 1968JO1C, 1968RI1H, 1969HO1M, 1970CO1H, 1970FO1B, 1972AN05, 1972CA37, 1972LE1L, 1972PN1A, 1973CL09, 1973CO16, 1973JO1K, 1973JU2A, 1973KU03, 1973MA48, 1973RO1R).

Ground-state properties: (1965CO25, 1966MA1P, 1966WI1E, 1967BA78, 1967SH05, 1967SH14, 1968PE16, 1969LE1B, 1969VA1C, 1969WU1A, 1970HI08, 1971TA1A, 1972LE1L, 1972VA36, 1973CO1P, 1973MA1K, 1973SA30, 1973SU1B).

$$J = 3 \text{ (1967BE1T);}$$

[†] (1972FU1C) observe the γ -de-excitation of ¹⁰B*(0.7, 1.7, 2.2) formed in ratio 2:1:1 when π^- are stopped in a carbon target.

Table 10.5: Energy Levels of ^{10}B ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
g.s.	$3^+; 0$	stable		1, 4, 5, 6, 7, 12, 13, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57, 58, 59
0.71832 ± 0.09	$1^+; 0$	$\tau_m = 1.013 \pm 0.015$ nsec	γ	1, 4, 5, 7, 12, 13, 18, 19, 20, 26, 27, 29, 31, 32, 39, 41, 42, 43, 47, 48, 49, 56
1.74016 ± 0.17	$0^+; 1$	< 30 fsec	γ	1, 4, 12, 13, 18, 19, 20, 26, 27, 29, 31, 39, 41, 42, 43, 47, 48, 52
2.1550 ± 1.0	$1^+; 0$	2.13 ± 0.18 psec	γ	1, 4, 12, 13, 18, 19, 26, 27, 29, 31, 32, 41, 42, 43, 47, 48, 49, 56
3.5897 ± 2.2	$2^+; 0$	147 ± 14 fsec	γ	1, 4, 5, 7, 13, 18, 19, 26, 27, 29, 31, 32, 40, 41, 42, 43, 47, 48, 49, 56
4.773 ± 3	$3^+; 0$	$\Gamma = 0.014$ keV	γ, α	1, 4, 7, 12, 18, 19, 26, 27, 29, 31, 32, 41, 42, 43, 47, 48
5.112 ± 4	$2^-; 0$	1.2	γ, α	1, 18, 19, 27, 29, 31, 32, 41, 42, 43, 48
5.166 ± 4	$2^+; 1$	0.003	γ, α	1, 18, 19, 27, 31, 40, 41, 42, 43
5.180 ± 10	$1^+; 0$	200 ± 30	γ, α	1, 3, 13, 29, 31, 32, 41, 42
5.924 ± 4	$2^+; 0$	6 ± 1	γ, α	1, 3, 18, 19, 27, 32, 41, 43, 47, 48, 56
6.025 ± 2	4^+	0.05 ± 0.03	γ, α	1, 3, 18, 19, 25, 27, 29, 31, 32, 41, 43, 47, 48, 52, 56
6.133 ± 2	3^-	2.36 ± 0.03	α	1, 3, 18, 19, 27, 29, 32, 41, 43, 47, 48, 56

Table 10.5: Energy Levels of ^{10}B ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
6.561 \pm 2	(2 ⁺)	25.1 \pm 1.1	d, α	3, 17, 18, 19, 27, 29, 31, 32, 41, 43, 48
6.881 \pm 5	1 ⁻ ; 0 + 1	120 \pm 10	γ , p, d, α	1, 13, 15, 17, 18
7.002 \pm 6	(1, 2) ⁺ ; (0)	100 \pm 10	p, d, α	3, 17, 18, 27, 29, 43, 48
7.431 \pm 7	(1) ⁻ ; 0 + 1	100 \pm 10	γ , p, d, α	1, 13, 17, 47
7.467 \pm 10	1 ⁺	65 \pm 10	(γ), p	13, 15
7.477 \pm 2	2 ⁻ ; 1	74 \pm 4	γ , p	13, 15, 18, 25, 27, 43
7.5595 \pm 1.0	0 ⁺ ; 1	2.65 \pm 0.18	γ , p	13, 15, 18, 43
(7.67 \pm 30)	(1 ⁺ ; 0)	250 \pm 20	p, d	15, 17
7.84 \pm 30	1 ⁻	260 \pm 30	γ , p, α	3, 13, 15, 18, 43
8.07 \pm 50	(2 ⁻ ; 0)	800 \pm 200	p, d, α	17, 18
(8.65)	(1 ⁺ , 2 ⁺)	\approx 300	p	15
8.889 \pm 6	3 ⁽⁻⁾ ; (1)	84 \pm 7	n, p	14, 15
8.894 \pm 2	2 ⁺ ; 1	36 \pm 2	γ , p, α	13, 15, 17, 23
9.7	$T = 1$	\approx 600	n, p, α	14, 17
10.84 ^b		\approx 500	γ , n, p	13, 14, 15, 23
11.53 \pm 40		270 \pm 50		41, 43
12.57 \pm 30	(0 ⁺ , 1 ⁺ , 2 ⁺)	90 \pm 30	γ , p	13, 15, 43
(13.3)	(0 ⁺ , 1 ⁺ , 2 ⁺)	\approx 300	γ , p	13
14.1 \pm 0.2		1100 \pm 200	α	3, 41
(14.1)	(0 ⁺ , 1 ⁺ , 2 ⁺)	\approx 250	γ , p	13
(14.6)	(0 ⁺ , 1 ⁺ , 2 ⁺)	\approx 150	γ , p	13
(15.6)	(2 ⁺ , 3 ⁺ , 4 ⁺)	\approx 400	γ , p	13
18.43	2 ⁻ ; 1	340	γ , ^3He	7
18.8	2 ⁺ ; (1)	< 600	γ , ^3He , α	7, 11
19.3	2 ⁻ ; 1	190 \pm 20	γ , n, p, ^3He , α	7, 8, 9, 11, 13
20.1 \pm 100	1 ⁻ ; 1	350 \pm 70	γ , n, p, t, ^3He , α	7, 8, 9, 10, 11, 23
(21.1)			γ , ^3He	7
(23.0)	$T = 2$	\approx 500	p, ^3He	9
23.1 \pm 100			γ , n	23

^a See also Tables 10.6 and 10.9.^b For J^π for these and many higher states see Table 10.11.

Table 10.6: Electromagnetic transitions in ^{10}B ^a

Initial state	$J^\pi; T$	Γ_γ (total) (eV)	Branching ratios (%) to final states at:					Γ_γ/Γ	Refs.
			g.s. 3 ⁺ ; 0	0.72 1 ⁺ ; 0	1.74 0 ⁺ ; 1	2.16 1 ⁺ ; 0	3.59 2 ⁺ ; 0		
0.72	1 ⁺ ; 0	6.5×10^{-7}	100						Table 10.9
1.74	0 ⁺ ; 1	> 0.02	< 0.2	100					Table 10.9, (1966SE03, 1969YO01)
2.16	1 ⁺ ; 0	3.1×10^{-4}	21.1 ± 1.6	27.3 ± 0.9	51.6 ± 1.6				Table 10.9, (1969YO01)
3.59	2 ⁺ ; 0	4.4×10^{-3}	19 ± 3	67 ± 3	< 0.3	14 ± 2			Table 10.9, (1966SE03, 1968WA15, 1969GA06, 1969YO01)
4.77	3 ⁺ ; 0	0.033 ± 0.006	0.5 ± 0.1	> 99				$(2.3 \pm 0.3) \times 10^{-3}$ ^b	Table 10.7, (1966AL06)
5.11	2 ⁻ ; 0	0.1	64 ± 7	31 ± 7	5 ± 5				Table 10.7, (1966FO05)
5.17	2 ⁺ ; 1	2.9 ± 1.1	5 ± 1	24 ± 3	< 0.5	69 ± 5	7 ± 3	0.88 ± 0.04 ^c	Table 10.7, (1966AL06, 1966FO05, 1967PA01)
5.18	1 ⁺ ; 0	0.06 ± 0.03			≈ 100				Table 10.7
5.92	2 ⁺ ; 0	≤ 54	82 ± 5	18 ± 5				≤ 0.009	Table 10.7, (1966FO05, 1969YO01)
6.03	4 ⁺	≤ 0.5	≈ 100					≤ 0.009	Table 10.7, (1969YO01)
6.13	3 ⁻	≤ 21						≤ 0.009	(1969YO01)
^e									

^a See also Table 10.7 in (1966LA04).^b $\Gamma_\alpha = 14 \pm 3$ eV.^c $\Gamma_\alpha = 0.44 \pm 0.09$ eV.^d Other branches $< 3\%$.^e For γ -decay of higher ^{10}B states [see Table 10.5], refer to Tables 10.10, 10.11 and 10.12.

$\mu = +1.8007$ nm (1969FU11, 1971SH26);
 $Q = +0.08$ b (1969FU11); [$+0.0847 \pm 0.0006$ b (1970NE05); theor.]; see also (1968SC18).

1. ${}^6\text{Li}(\alpha, \gamma){}^{10}\text{B}$ $Q_m = 4.460$

Six resonances are observed in the range $E_\alpha = 0.5$ to 3.8 MeV, corresponding to ${}^{10}\text{B}^*(4.77, 5.11, 5.17, 5.18, 5.92, 6.03)$: see Table 10.7 (1957ME27, 1961SP02, 1966AL06, 1966FO05, 1966SE02, 1971AU1H). Some weak effects corresponding to ${}^{10}\text{B}^*(6.13)$ may also be present (1966FO05). Angular distribution and branching ratio measurements are consistent with 3^+ for ${}^{10}\text{B}^*(4.77)$ and $2^-, 2^+, 1^+, 2^+$ and 4^+ for the five higher states: see (1966LA04) and (1966AL06, 1966FO05, 1966SE02). The results are in good agreement with the I.P.M. calculations of (1965CO25). See also (1967WA1C).

2. (a) ${}^6\text{Li}(\alpha, n){}^9\text{B}$ $Q_m = -3.975$ $E_b = 4.460$
 (b) ${}^6\text{Li}(\alpha, p){}^9\text{Be}$ $Q_m = -2.1251$
 (c) ${}^6\text{Li}(\alpha, d){}^8\text{Be}$ $Q_m = -1.5656$

The excitation functions for neutrons (1963ME08: from threshold to $E_\alpha = 15.5$ MeV) and for d_0 particles (1963BL20: $E_\alpha = 9.5$ to 11.4 MeV) do not show resonance structure. See also ${}^8\text{Be}$, ${}^9\text{Be}$ and ${}^9\text{B}$.

3. ${}^6\text{Li}(\alpha, \alpha){}^6\text{Li}$ $E_b = 4.460$

Reported anomalies in the elastic scattering are listed in Table 10.8 (1962DE10, 1967ME08, 1971BA41). See also (1965SI1B). The 1^+ assignment for ${}^{10}\text{B}^*(5.18)$ supports the proposal by (1961TR1B) that it is a member of the doublet formed by two-nucleon excitation into the 2s shell, whose other member is the $J^\pi = 0^+; T = 1$ state at 7.56 MeV.

Excitation functions of α_0 have been reported since (1966LA04) at $E_\alpha = 2.0$ to 3.7 MeV (1967ME08), 2.5 to 4.5 MeV (1972BO07), 3.3 to 5.0 MeV (1971BA41) and 13.3 to 18.0 MeV (1970BI1B, 1971BI12). In the high-energy work, the yield curves are generally monotonically decreasing functions of energy except in the region $E_\alpha = 15.7$ to 17 MeV, where a broad resonance structure is observed which may correspond to weak excitation of ${}^{10}\text{B}^*(14.0)$ (1970BI1B, 1971BI12). Excitation functions for α_0 and α_1 [to ${}^6\text{Li}^*(2.19)$] at $E_\alpha = 9.5$ to 12.5 MeV do not show resonance structure (1963BL20). See also (1969TR1B, 1972RI10; theor.).

4. ${}^6\text{Li}({}^6\text{Li}, d){}^{10}\text{B}$ $Q_m = 2.987$

Table 10.7: Levels of ^{10}B from $^6\text{Li}(\alpha, \gamma)^{10}\text{B}$ ^a

E_{res} (keV)	E_x (MeV)	$J^\pi; T$	Γ_{lab} (keV)	Decay to E_f	Branch (%)	$\omega\gamma$ (eV)	Γ_γ (eV)	$ M ^2$ ^g
500 ± 25	4.760	$3^+; 0$		0	0.5 ± 0.1			
				0.72	> 99		0.033 ± 0.006 ^c	
1085	5.112	$2^-; 0$	2 ^b	0	64 ± 7	0.059 ± 0.012 ^d		1.5 ^e 9×10^{-4} ^k
				0.72	31 ± 7	0.028 ± 0.008		1.5 ^e 6×10^{-4} ^k
				1.74	5 ± 5	0.005 ± 0.005		100 ± 100 ^e
1175 ⁱ	5.166	$2^+; 1$	< 0.5 ^b	0	5 ± 1	0.035 ± 0.007	0.15 ^g	0.8 ^f 0.06 ^h
				0.72	24 ± 3	0.17 ± 0.04	0.56 ^g	4 ^f 0.4 ^h
				1.74	2 ± 1	0.017 ± 0.008		160 ± 80 ^f
				2.15	69 ± 5	0.47 ± 0.09	1.20 ^g	80 ^f 3.8 ^h
				3.58	9 ± 2 ^g		0.19 ^g	
1210 ± 35	5.187	$1^+; 0$	340 ± 50	1.74	≈ 100		0.06 ± 0.03	
2435	5.922	2^+	10 ± 1	0	82 ± 5	0.31 ± 0.06		$0.27 \rightarrow 25$ ^{f,j}
				0.72	18 ± 5	0.07 ± 0.02		$0.1 \rightarrow 9$ ^{f,j}
				1.74		< 0.02		
2605	6.024	4^+	< 1.5 ^b	0	≈ 100	0.57 ± 0.08		20 ^f 4×10^{-3} ^h
				0.72		< 0.02		
4032 ± 5	6.881 ^{l,m}	$T = 0 + 1$		0	6 ± 2			
				0.72	19 ± 3			

Table 10.7: Levels of ^{10}B from $^6\text{Li}(\alpha, \gamma)^{10}\text{B}$ ^a (continued)

E_{res} (keV)	E_x (MeV)	$J^\pi; T$	Γ_{lab} (keV)	Decay to E_f	Branch (%)	$\omega\gamma$ (eV)	Γ_γ (eV)	$ M ^2$ ^g
4950 ± 10	7.432 ^l			1.74 2.15 n	52 ± 3 13 ± 4			

^a (1957ME27, 1961SP02, 1966AL06, 1966FO05, 1966SE02, 1971AU1H). See also (1966LA04).

^b See (1958ME81) and $^9\text{Be}(\text{d}, \text{n})^{10}\text{B}$.

^c $\Gamma_\gamma/\Gamma = (2.3 \pm 0.3) \times 10^{-3}$, $\Gamma_\alpha = 14 \pm 3$ eV (1966AL06).

^d Absolute errors only.

^e M2.

^f E2.

^g (1966FO05).

^h M1.

ⁱ $\Gamma_\gamma = 2.9 \pm 1.1$ eV, $\Gamma_\alpha = 0.44 \pm 0.09$ eV, $\omega\gamma = 0.63 \pm 0.13$ eV; α branch is $13 \pm 4\%$ (1966AL06).

^j E2/M1 = 0.01 or 10 (1957ME27).

^k E1.

^l (1971AU1H) and V. Meyer, private communication.

^m $\Gamma_{\text{c.m.}} = 110 \pm 10$ keV.

ⁿ $\Gamma_{\text{c.m.}} = 90 \pm 10$ keV. At 0° the branches to $^{10}\text{B}^*(0, 0.72)$ are equally strong. $J^\pi = 2^-$; $T = 0 + 1$ is suggested (V. Meyer, private communication).

Table 10.8: ^{10}B levels from $^6\text{Li}(\alpha, \alpha)^6\text{Li}$

E_α (keV)	E_x (MeV)	Γ_{lab} (keV)	$J^\pi; T$	Refs.
1210 ± 30	5.19	175	$1^+; 0$	(1962DE10)
2440	5.93	≈ 30	$2^+; 0$	(1962DE10)
2606.0 ± 1.5	6.025	0.09 ± 0.04	4^+	(1967ME08)
2785.5 ± 1.5	6.133	3.93 ± 0.05	3^-	(1967ME08)
3498.5 ± 1.6 ^a	6.561	41.8 ± 1.9	$4^-, 2^-$ ^d	(1967ME08) ^b
4250 ± 15 ^a	7.012	183 ± 25	$(2)^+; (0)$ ^d	(1971BA41)
(5500)	(7.76)			(1965SI1B) ^c
16000	14.0	broad		(1970BI1B, 1971BI12)

^a There is evidence of broad structure near these states (1971BA41).

^b See also (1971BA41).

^c This result was published in an abstract.

^d See, however, (1973SI27) in reaction 17.

Angular distributions of deuteron groups have been determined at $E(^6\text{Li}) = 2.4$ to 9.0 MeV (d_0, d_1, d_3) and 7.35 and 9.0 MeV (d_4, d_5) (1966KI09). The d_2 group is also observed but its intensity is weak: see (1960MO17, 1961MO02, 1966KI09). For τ_m measurements see Table 10.9 (1969TH01). See also (1971PO1D), (1966BR1G), (1966RO1E; theor.) and ^{12}C in (1975AJ02).

$$5. \ ^6\text{Li}(^7\text{Li}, t)^{10}\text{B} \quad Q_m = 1.993$$

Angular distributions of the t_0 and t_1 groups have been measured at $E(^6\text{Li}) = 3.3$ MeV (1967GA06) and $E(^7\text{Li}) = 3.78$ to 5.95 MeV (1967KI03). For τ_m measurements see Table 10.9 (1969TH01). See also (1968DA20), (1972GA1E), (1966LA04) and ^{13}C in (1976AJ04).

$$6. \ ^6\text{Li}(^9\text{Be}, ^5\text{He})^{10}\text{B} \quad Q_m = 1.99$$

See (1962MC12, 1963NO02).

$$7. \ ^7\text{Li}(^3\text{He}, \gamma)^{10}\text{B} \quad Q_m = 17.788$$

Table 10.9: Lifetimes of ^{10}B states ^a

^{10}B state	τ_m	Reactions	Refs.
0.72	1.013 ± 0.015 nsec	mean of values quoted in Table 10.20	(1966LA04)
1.74	< 40 fsec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1968DO01)
	< 30 fsec	$^9\text{Be}(^3\text{He}, \text{d})$	(1966FI01, 1968FI09)
	< 22 fsec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1969JA1N)
	< 30 fsec	“best” value	
2.16 ^b	$2.1_{-0.5}^{+0.8}$ psec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1968DO01)
	$2.7_{-0.4}^{+0.5}$ psec	$^9\text{Be}(\text{d}, \text{n}), ^9\text{Be}(^3\text{He}, \text{d}), ^{10}\text{B}(\text{p}, \text{p}')$	(1968FI09)
	2.0 ± 0.6 psec	$^9\text{Be}(\text{d}, \text{n})$	(1969AL17)
	2.0 ± 0.25 psec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1969JA1N)
	2.15 ± 0.45 psec	$^7\text{Li}(\alpha, \text{n})$	(1970GA01)
	2.13 ± 0.18 psec	mean of above values	
3.59	152 ± 21 fsec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1969JA1N)
	100 ± 30 fsec	$^6\text{Li}(^6\text{Li}, \text{d}), ^6\text{Li}(^7\text{Li}, \text{t})$	(1969TH01)
	115 ± 40 fsec ^d	$^9\text{Be}(\text{d}, \text{n})$	(1966WA10)
	150 ± 15 fsec	$^9\text{Be}(\text{d}, \text{n}), ^{10}\text{B}(\text{p}, \text{p}')$	(1968FI09)
	170 ± 70 fsec	$^{11}\text{B}(^3\text{He}, \alpha)$	(1968DO01)
	147 ± 14 fsec	mean of last three values	
5.17 ^c	< 80 fsec	$^9\text{Be}(\text{d}, \text{n})$	(1966WA10)

^a See also (1966LA04).

^b See also (1966FI01, 1969TH01).

^c See also (1967PA01).

^d Based on τ_m of $^{10}\text{Be}^*(3.37) = 180$ fsec.

Capture γ -rays have been observed for $E(^3\text{He}) = 0.8$ to 3.0 MeV (1965PA02, 1971LI20) and 3.0 to 6.0 MeV (1968LI03). The γ_0 and γ_5 yields [to $^{10}\text{B}^*(0, 4.77)$] show resonances at $E(^3\text{He}) = 1.1$ and 2.2 MeV [$E_{\text{res}} = 0.92$ and 2.1 MeV: see (1971LI20)], the γ_1 and γ_4 yields [to $^{10}\text{B}^*(0.72, 3.59)$] at 1.4 MeV and the γ_4 yield at 3.4 MeV: see Table 10.10 (1965PA02, 1968LI03, 1971LI20). Both the 1.1 and 2.2 MeV resonances [$^{10}\text{B}^*(18.4, 19.3)$] appear to result from s-wave capture; the subsequent decay is to two 3^+ states [$^{10}\text{B}^*(0, 4.77)$]. Therefore the most likely assignment is $J^\pi = 2^-$; $T = 1$ for both [there appears to be no decay of these states via α_2 to $^6\text{Li}^*(3.56)$ which has $J^\pi = 0^+$; $T = 1$: see reaction 11] (1965PA02, 1971LI20). The assignment for $^{10}\text{B}^*(18.8)$ [1.4 MeV resonance] is 1^+ or 2^+ but there appears to be α_2 decay and therefore $J^\pi = 2^+$. $^{10}\text{B}^*(20.2)$ [3.4 MeV resonance] has an isotropic angular distribution of γ_4 and therefore $J^\pi = 0^+, 1^-, 2^-$. The γ_2 group resonates at this energy which eliminates 2^- , and 0^+ is eliminated on the basis of the strength of the transition which is too large for E2 (1971LI20).

8. $^7\text{Li}(^3\text{He}, n)^9\text{B}$

$$Q_m = 9.353$$

$$E_b = 17.788$$

The excitation curve is smooth up to $E(^3\text{He}) = 1.8$ MeV (1962SE1A, 1963DU12, 1966DI04) and the n_0 yield shows resonance behavior at $E(^3\text{He}) = 2.2$ and 3.25 MeV, $\Gamma_{\text{lab}} = 270 \pm 30$ and 500 ± 100 keV. No other resonances are observed up to $E(^3\text{He}) = 5.5$ MeV (1966DI04). See also Table 10.10.

9. $^7\text{Li}(^3\text{He}, p)^9\text{Be}$

$$Q_m = 11.2027$$

$$E_b = 17.788$$

The yield of p_0 and p_2 has been measured for $E(^3\text{He}) = 0.60$ to 1.25 MeV (1971ST35): it is suggested that $^{10}\text{B}^*(18.4)$ is formed. The yield of protons is relatively flat for $E(^3\text{He}) = 1.8$ to 4.8 MeV with some indication of weak maxima at ≈ 2.3 and 3.3 MeV (1961WO05, 1972LI31). See also (1969SA04, 1970DI1F). The yield of γ -rays from the decay of the $T = \frac{3}{2}$ state of ^9Be at 14.39 MeV to $^9\text{Be}^*(0, 2.4)$ has been studied at $E(^3\text{He}) = 5.0$ to 10.0 MeV: some suggestion is reported of a $T = 2$ state at $E_x \approx 23$ MeV, $\Gamma \approx 0.5$ MeV (1968HA1K: unpublished results; W.D. Harrison, private communication).

10. (a) $^7\text{Li}(^3\text{He}, d)^8\text{Be}$

$$Q_m = 11.762$$

$$E_b = 17.788$$

(b) $^7\text{Li}(^3\text{He}, t)^7\text{Be}$

$$Q_m = -0.8804$$

Yields of tritons have been measured for $E(^3\text{He}) = 0.65$ to 1.25 MeV (1971ST35; d_0), 2.0 to 4.2 MeV (1969OR01; t_0), 8.0 to 10.0 MeV (1968MA1W; t_0, t_1) and 10 to 16 MeV (1969NU1A; t_0, t_1). A broad peak is reported at $E(^3\text{He}) \approx 3.5$ MeV (1969OR01). For reaction (a) see ^8Be .

11. $^7\text{Li}(^3\text{He}, \alpha)^6\text{Li}$

$$Q_m = 13.3279$$

$$E_b = 17.788$$

Table 10.10: Resonances in ${}^7\text{Li} + {}^3\text{He}$

E_{res} (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	E_x (MeV)	$J^\pi; T$	Γ_γ (eV) for transition to				Γ_α (keV)	Γ_n	Γ_p	Γ_t	Refs.
				g.s.	0.72	3.59	4.77					
0.92	340	18.43	$2^-; 1$	≥ 3			≥ 17					(1965PA02, 1971LI20)
1.4	< 600	18.8	$2^+, 1^+$		≥ 20	$\geq 20^b$		res. < 80				(1965PA02, 1965PA03, 1971LI20)
2.1	280 ^a	19.3	$2^-; 1$	≥ 12			≥ 49	res. < 20	res. n_0	(p)		(1963DU12, 1965PA03, 1966DI04, 1968LI03, 1971LI20, 1972LI31)
3.4 (4.7)	910 ^a	20.2 (21.1)	$1^-; 1$			≥ 350		res. α_2	res. n_0	(p)	res. t_0	(1961WO05, 1966DI04, 1968LI03, 1971LI20) (1968LI03)

^a (1966DI04) report $\Gamma_{\text{c.m.}} = 190 \pm 20$ and 350 ± 70 keV, respectively, from the n_0 yield.

^b Assumes isotropy of angular distribution (1971LI20).

Excitation functions have been measured at $E(^3\text{He}) = 1.3$ to 5.4 MeV (1965FO07; $\alpha_0, \alpha_1, \alpha_2$), 2.0 to 4.2 MeV (1969OR01; α_0), 2.5 to 4.2 MeV (1969OR01; α_2), 5.0 to 8.0 MeV (1970OR03; $\alpha_0, \alpha_1, \alpha_2$) and 16.0 to 18.0 MeV (1971ZA07; α_0). The α_0 group (at 8°) shows a broad maximum at ≈ 2 MeV, a minimum at 3 MeV, followed by a steep rise which flattens off between $E(^3\text{He}) = 4.5$ and 5.5 MeV. Integrated α_0 and α_1 yields rise monotonically to 4 MeV and then tend to decrease (1965FO07). Angular distributions give evidence of the resonances at $E(^3\text{He}) = 1.4$ and 2.1 MeV seen in $^7\text{Li}(^3\text{He}, \gamma)^{10}\text{B}$: $J^\pi = 2^+$ or 1^- ; $T = (1)$ for both [see, however, reaction 7]: Γ_α is small (1965PA03). The α_2 yield [to $^6\text{Li}^*(3.56)$, $J^\pi = 0^+$; $T = 1$] shows some structure at $E(^3\text{He}) = 1.4$ MeV and a broad maximum at ≈ 3.3 MeV (1965FO07, 1969OR01): see Table 10.10.



Angular distributions have been measured for the n_0 group at $E_\alpha = 4.78$ to 7.85 MeV (1972VA02) and 13.5 and 13.9 MeV (1962KJ05), and for the n_1 group [to $^{10}\text{B}^*(0.7)$] at $E_\alpha = 6.71$ to 7.85 MeV (1972VA02). The n_2 group has also been seen in this reaction (1972VA02). Slow-neutron threshold measurements have been reported corresponding to the formation of $^{10}\text{B}^*(0, 0.72, (4.77), (6.42))$: see (1957BI84, 1963ME08) and (1966LA04). The lifetime of $^{10}\text{B}^*(2.16)$, $\tau_m = 2.16 \pm 0.43$ psec: the “best” value for τ_m based on this and on other measurements [see Table 10.9] is 2.4 ± 0.3 psec corresponding to $\Gamma_\gamma = 275 \pm 35$ μeV . The γ -decay of $^{10}\text{B}^*(2.15)$ involves $E_\gamma = 415.1 \pm 0.5$ keV ($2.15 \rightarrow 1.74$), 1435.6 ± 1.0 keV ($2.15 \rightarrow 0.72$) and 719.1 ± 0.6 keV ($0.72 \rightarrow 0$). The excitation energies for the first three excited states are then 719.1 ± 0.6 , 1739.7 ± 1.5 and 2154.8 ± 1.2 keV (1970GA01). See also (1972DA32).



Parameters of observed resonances are listed in Tables 10.11 and 10.12. Table 10.6 summarizes the γ -transitions from this and other reactions.

The $E_p = 0.33$ MeV resonance ($^{10}\text{B}^* = 6.88$ MeV) is ascribed to s-wave protons because of its comparatively large proton width [see $^9\text{Be}(p, p)$] and because of the isotropy of the γ -radiation. The strong transition to $^{10}\text{B}^*(1.74)$ requires E1 and hence $J^\pi = 1^-$; $T = 0$. $T = 0$ is also indicated by the large deuteron width. On the other hand, the strength of E1 transitions to $^{10}\text{B}^*(0.7, 2.1)$ indicates $T = 1$. The amplitudes for the $T = 0$ and $T = 1$ parts of the wave function for $^{10}\text{B}^*(6.88)$ are 0.92 and 0.39 , respectively (1972RE07). See also (1969CO1G, 1969RO12, 1971AU1H).

The proton capture data near $E_p = 1$ MeV appears to require at least five resonant states, at $E_p = 938, (980), 992, 1083$ and 1290 keV. The narrow $E_p = 1083$ keV level ($^{10}\text{B}^* = 7.56$ MeV) is formed by p-wave protons, $J^\pi = 0^+$ [see $^9\text{Be}(p, p)$, $^9\text{Be}(p, \alpha)$]. The isotropy of the gamma rays supports this assignment (1961TA02). The strong M1 transitions to $J^\pi = 1^+$; $T = 0$ levels at $0.7, 2.16$ and 5.18 MeV (Table 10.12) indicate $T = 1$ (1959WA16). See also (1970BO1Y).

The excitation function for ground-state radiation shows resonance at $E_p = 992$ ($\Gamma = 80$ keV) and 1290 keV ($\Gamma = 230$ keV) (1962EL06, 1964HO02). Elastic scattering studies indicate

Table 10.11: Resonances in ${}^9\text{Be}(p, \gamma){}^{10}\text{B}$ ^a

E_p (MeV \pm keV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	Γ_p/Γ	Γ_γ (eV)	Refs.
0.330	6.88	130 ± 5 ^f	$1^-; 0 + 1$ ^b	0.30	4.8	(1956CL69, 1969CO1G)
0.938 ± 10 (0.98)	7.429 (7.47)	140 ± 30	$(1)^-; 0 + 1$ ^b (2^+)	0.7	2.4	(1962EL06, 1964HO02, 1969CO1G) (1962EL06)
0.992 ± 2	7.477	72 ± 4	$2^-; 1$	≈ 0.65	25.8	(1964HO02)
1.0832 ± 0.4	7.5595	2.65 ± 0.18	$0^+; 1$	1.0	8.5	(1964BO13, 1964HO02, 1972HA63)
1.29	7.75	210 ± 60	$2^-; (1)$ ^c	≈ 0.65	8.5	(1964HO02)
2.567 ± 2	8.894	36 ± 2	$2^+; 1$			(1953MA1A, 1956MA55)
4.72	10.83	≈ 500	$2^+, 3^+, 4^+$			(1952HA10, 1970FI1B) ^{d,e}
6.7 (7.0)	12.6 (12.9)	< 200 (≈ 100)	$0^+, 1^+, 2^+$ ($\pi = +$)			(1970FI1B) ^e
7.5	13.3	≈ 300	$0^+, 1^+, 2^+$			(1970FI1B) ^e
8.4	14.1	≈ 250	$0^+, 1^+, 2^+$			(1970FI1B) ^e
8.9	14.6	≈ 150	$2^+, 3^+, 4^+$			(1970FI1B) ^e
10.0	15.6	≈ 400	$2^+, 3^+, 4^+$			(1970FI1B) ^e
14.6	19.7	≈ 500	$2^-, 3^-, 4^-$			(1970FI1B) ^e

^a See Table 10.12 for decay schemes.

^b See also (1969RO12).

^c See, however, (1973RO24) in reaction 15.

^d See also (1969FI1C).

^e G.A. Fisher, private communication.

^f V. Meyer, private communication.

s-wave formation and $J^\pi = 2^-$ for both (1956MO90). For the lower level ($E_x = 7.48$ MeV) the intensity of the g.s. capture radiation, $\Gamma_\gamma = 25$ eV (1964HO02) indicates E1 and $T = 1$. The angular distribution of γ -rays, $1 + 0.1 \sin^2 \theta$, is consistent with s-wave formation with some d-wave admixture (1953PA22) or with some contribution from a nearby p-wave resonance (1956MO90); possibly a $J^\pi = 2^+$ level at $E_p = 980$ keV (1956MO90, 1962EL06: see, however, (1964HO02)).

The angular distribution of ground-state radiation at $E_p = 1330$ keV is isotropic and $\Gamma_\gamma = 8.5$ eV (1964HO02), supporting E1, $T = 1$ for this level ($E_x = 7.75$ MeV). See, however, (1973RO24): reaction 15.

Transitions to ${}^{10}\text{B}^*(0.7) [\gamma_1]$ show resonance at $E_p = 992, 1290$ and 938 keV, $\Gamma = 155$ keV (1962EL06, 1964HO02). The latter is presumably also a resonance for (p, d) and (p, α). An assignment of $J^\pi = 2^-; T = 0$ is consistent with the data, although the E1 radiation then seems somewhat too strong for a $\Delta T = 0$ transition (1964HO02). See also (1973RO24) in reaction 15.

A resonance for capture radiation at $E_p = 2.567 \pm 0.003$ ($E_x = 8.894$ MeV) has a width of 40 ± 2 keV and decays mainly via ${}^{10}\text{B}^*(0.7)$ (1953MA1A). It appears from the width that this resonance corresponds to that observed in ${}^9\text{Be}(p, \alpha)$, $J^\pi = 2^+; T = 1$ and not to the ${}^9\text{Be}(p, n)$

Table 10.12: Radiative transitions in ${}^9\text{Be}(p, \gamma){}^{10}\text{B}$

Initial state (MeV)	Γ_γ (tot) (eV)	Relative intensities to final states									Refs.
		ground $3^+; 0$	0.72 $1^+; 0$	1.74 $0^+; 1$	2.16 $1^+; 0$	3.59 $2^+; 0$	5.11 $2^-; 0$	5.17 $2^+; 1$	5.18 $1^+; 0$	5.92 $2^+; 0$	
6.88 $E_p = 0.33$ $1^-; 0$	4.8	0.05 < 0.7	0.35 ± 0.04 0.20 ± 0.02 1.0 6.4	0.48 ± 0.04 0.53 ± 0.02 2.6 15	0.11 ± 0.04 0.13 ± 0.01 0.5 4.5	< 0.01 ≈ 1	0.04 ± 0.01	0.02 ± 0.01	< 0.01 0.65 < 2.9	0.035 ± 0.005 W.u.	(1972RE07) A (1956CL69) (1959ME85) (1972RE07)
7.43 $E_p = 0.94$ $(1)^-; 0$	[2.4]	< 2	1.3 0.013	[< 0.14]	0.62 0.013	0.5 0.03			[< 1]		(1964HO02) (1964HO02)
7.48 $E_p = 0.99$ $2^-; 1$	[25.8]	400 25 0.19	< 10 0.3 0.003	19 [< 0.14]	22 0.49 0.10	< 7 0			< 13 [< 1]		(1959ME85) (1964HO02) (1964HO02)
7.56 $E_p = 1.08$ $0^+; 1$	6.6 [8.5]	< 3 < 0.2	100 100 6.7 1.0	< 8 < 2 < 0.3	< 8 10 0.8 0.24	< 4 < 0.2			23 40 1.0 3.5		(1959ME85) (1962ME1A) (1964HO02) (1964HO02)
7.75 ^a $E_p = 1.29$ $2^-; 1$	14.6 [8.5]	83% 6.6 0.044	 0.9 0.008	 < 0.08	 0.3 0.006	 0.3 0.013			 0.4 0.08		(1963FU11) (1964HO02) (1964HO02)

A: V. Meyer, private communication.

^a See, however, Table 10.14.

resonance at the same energy (1956MA55). A further resonance is reported at $E_p = 4.72 \pm 0.01$ MeV, $\Gamma \approx 0.5$ MeV (1952HA10).

In the range $E_p = 4$ to 18 MeV, the γ_0 yield at 90° shows the resonance at $E_p = 4.7$ MeV ($E_x = 10.7$ MeV) and shows fluctuations suggesting states at $E_x \approx 14.6, 15.6$ and 19.7 MeV. It is suggested that $^{10}\text{B}^*(19.7)$ decays via E1 and therefore $J^\pi = 2^-, 3^-, 4^-$. The other three states presumably decay by M1 and therefore $J^\pi = 2^+, 3^+, 4^+$. These fluctuations appear on a nearly constant γ_0 yield with a 90° differential cross section $\approx 1.5 \mu\text{b/sr}$. The average yield of γ_1 is $\approx \frac{2}{3}$ of the γ_0 yield. The broad giant resonance peak is centered at $E_x \approx 14.5$ MeV. Fluctuations in the γ_1 yield are reported at $E_x \approx 12.6, 13.3$ and 14.1 MeV. These states presumably decay by M1 to $^{10}\text{B}^*(0.7)$ [$J_f^\pi = 1^+$] and therefore $J_i^\pi = 0^+, 1^+, 2^+$. The weak γ_2 yield (to $^{10}\text{B}^*(1.74)$ [$J^\pi = 0^+; T = 1$]) seems to exhibit a broad peak centered near $E_x = 15$ MeV (maximum 90° differential cross section $\approx 0.5 \mu\text{b/sr}$) and possibly some structure near $E_x = 20$ MeV. The γ_3 yield (to $^{10}\text{B}^*(2.16)$ [$J^\pi = 1^+$]) increases to $\approx 0.4 \mu\text{b/sr}$ at $E_x \approx 16$ MeV and seems to remain constant beyond that energy, with some suggestion of a fluctuation corresponding to $E_x \approx 12.9$ MeV. $^{10}\text{B}^*(12.9)$ appears to have positive parity. Angular distributions of $\gamma_0, \gamma_1, \gamma_2$ and γ_3 are also reported (1970FI1B; G.A. Fisher, private communication). See also (1969FI1C).

The magnetic moment of $^{10}\text{B}^*(0.72)$ has been studied via γ - γ correlations from $^{10}\text{B}^*(7.56)$: $g = +0.63 \pm 0.12$ (1972AV01). For measurements of the mean life of $^{10}\text{B}^*(0.72)$, see Table 10.20. See also (1965ZO1A, 1966ED1A) and (1966LA04, 1973SU1E).

14. $^9\text{Be}(p, n)^9\text{B}$

$$Q_m = -1.8498$$

$$E_b = 6.5853$$

Resonances in neutron yield occur at $E_p = 2.56$ and 4.7 MeV: see Table 10.13. There is some indication of a broad maximum near $E_p = 3.5$ MeV; a peak reported at $E_p = 4.9$ MeV for n_1 neutrons may reflect the effect of this level (1959MA20). A sharp break at $E_p = 6.55 \pm 0.03$ MeV is ascribed to a level in ^9B at 4.04 MeV (1964BA16). See also (1972VO17).

The $E_p = 2.56$ MeV resonance is considerably broader than that observed at the same energy in $^9\text{Be}(p, \alpha)$ and $^9\text{Be}(p, \gamma)$ and the two resonances are believed to be distinct (1956MA55). The shape of the resonance and the magnitude of the cross section can be accounted for with $J^\pi = 3^-$ or 3^+ : the former assignment is in better accord with $^{10}\text{Be}^*(7.37)$. For $J^\pi = 3^-$, $\theta_n^2 = 0.135$, $\theta_p^2 = 0.115$ ($R = 4.47$ fm). The $J^\pi = 2^+$ level should contribute about 10% to the cross section at $E_p = 2.56$ MeV (1962AL1A).

There are no resonances at threshold as had been reported by (1969SI20, 1970SI12): see (1970CO06). See also reaction 15.

For angular distributions see ^9B . For polarization measurements see (1966LA04) and (1972SH1K). See also (1969VE02, 1970DA26, 1971JU05, 1973NE1G). For astrophysical considerations see (1969BA1N).

15. (a) $^9\text{Be}(p, p)^9\text{Be}$

$$E_b = 6.5853$$

(b) $^9\text{Be}(p, pn)^8\text{Be}$

$$Q_m = -1.6651$$

Table 10.13: Resonances in ${}^9\text{Be}(p, n){}^9\text{B}$

E_{res} (MeV \pm keV)	Γ_{lab} (keV)	E_x (MeV)	$J^\pi; T$	σ_{max} (mb)	Refs.
2.562 ± 6	85 ± 10	8.889	$3^{(-)}; (1)$		(1956MA55, 1962AL1A)
	100 ± 10			160	(1959GI47)
3.5	≈ 700	9.7			(1956MA55, 1959MA20)
				240	(1959GI47)
4.72 ± 10	500	10.84			(1952HA10)
4.68 ± 30				470	(1955MA84, 1959GI47)
4.62 ± 30					(1959MA20)
4.94 ± 30	≈ 100			510	(1955MA84, 1959GI47)
4.82 ± 60				a	(1959MA20)

^a Resonance for n_1 only; possibly due to 3.5 MeV resonance.

Elastic scattering has been studied for $E_p = 0.2$ to 2.7 MeV by (1956DE33, 1956MO90, 1969MO29, 1973RO24). Below $E_p = 0.7$ MeV only s-waves are present exhibiting resonance at $E_p = 330$ keV [${}^{10}\text{B}^*(6.88)$], $J^\pi = 1^-$. Between $E_p = 0.8$ to 1.6 MeV polarization and cross-section measurements are well fitted by a phase-shift analysis, using only the ${}^3\text{S}_1$, ${}^5\text{S}_2$, ${}^5\text{P}_1$ and ${}^5\text{P}_2$ phases. Four levels satisfy the data, 1^+ and 2^- states at $E_x = 7.48$ MeV, a sharp 0^+ state at $E_x = 7.56$ MeV, and a 1^- state at 7.82 MeV: see Table 10.14 (1956MO90, 1973RO24).

Pronounced minima at $E_p = 2.48$ and 2.55 are observed in the polarization (p_0): these are ascribed to $T = 1$ analogs of the 3^- and 2^+ states ${}^{10}\text{Be}^*(7.37, 7.52)$ (1969AN27). A resonance corresponding to ${}^{10}\text{B}^*(8.89)$ is also reported by (1956DE33). The yield of p_0 , p_1 and p_2 has been measured for $E_p = 4.2$ to 6.0 MeV by (1972YA06): structures are reported at $E_p \approx 4.7$ MeV ($\Gamma \approx 0.3$ MeV) in the p_2 yield and at $E_p \approx 5.1$ MeV in the p_1 yield (1972YA06). Excitation curves for the p_0 group have been measured for $E_p = 6.0$ to 15.0 MeV (1972VO17, 1973VO02): a strong anomaly is observed at $E_p = 6.7$ MeV: see Table 10.14. (1973VO02) find that the p_0 differential cross sections and polarization analyzing power are adequately described by a spherical optical model potential for $E_p = 13$ to 30 MeV: only the volume real potential depth V_R and the surface imaginary potential depth W_S need vary with energy. When coupled-channels analyses were made (1973VO02) found that a quadrupole-deformed optical model potential with a deformation $\beta = 1.1$ gives an improved description of the (p , p_0) data and good fits to data obtained for (p , p_2). See also (1971WE07).

The reports of anomalies in the yield of protons near the threshold for the (p , n) reaction, by (1968SI07, 1969SI20, 1970SI12), are in error: see (1970CO06, 1971GO33). See also (1965HU10, 1967CA1K, 1967MC1E, 1970BO1Y).

Cross section measurements have also been reported at $E_p = 6.36$ to 6.48 MeV (1971VA34; yield of p_1), 16.2 to 29.1 MeV (1973MO02; σ_R , and also σ for p_2 and p_3), 24.4 to 45.9 MeV

Table 10.14: Resonances in ${}^9\text{Be}(p, p){}^9\text{Be}$

E_{res} (keV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π	Γ_p/Γ	Refs.
330	6.88	145	1^-	0.30	(1956MO90, 1969MO29)
980 ± 10	7.467	65 ± 10	$1^+{}^a$	1.0	(1956MO90, 1969MO29, 1973RO24)
$980 \pm 10{}^a$	7.467	80 ± 8	2^-	0.90 ± 0.05	(1956DE33, 1956MO90, 1969MO29, 1973RO24)
1084 ± 2	7.560	2.7	0^+	1.0	(1956MO90, 1969MO29)
(1200 \pm 30)	(7.67)	250 ± 20	(1^+)	0.30 ± 0.10	(1969MO29)
1370 ± 20	7.817	265 ± 30	$1^-{}^a$	0.90 ± 0.05	(1956MO90, 1969MO29, 1973RO24)
(2300)	(8.65)	≈ 300	($1^+, 2^+$)		(1973RO24)
(2480)	(8.82)		($3^-; 1$)		(1969AN27)
2560	8.89		$\geq 2; (1)$	large	(1956DE33, 1969AN27)
(4700)	(10.8)	≈ 300			(1972YA06)
(5100)	(11.2)				(1972YA06)
6700	12.6	broad			(1972VO17, 1973VO02)

^a See, however, (1969MO29). (1969MO29) also suggest states at $E_x = 7.44$ and 8.1 MeV with $J^\pi = 1^-$ and 2^+ , respectively and $\Gamma_{\text{c.m.}} = 130 \pm 10$ and ≈ 80 keV, respectively: see discussion in (1973RO24).

(1969MC1A; σ_R), and 232 to 553 MeV (1972RE06; σ_R). See also (1966LA04) for the earlier work.

Polarization measurements involving the elastic group have been carried out at $E_p = 0.9$ to 2.7 MeV (1973RO24), 0.9 to 3.0 MeV (1969AN27), 2.2 to 3.8 MeV (1968BL1F), 3.0 to 12.0 MeV (1970LO03), 7.0 , 8.5 and 10.0 MeV (1967KO1E; also p_2 at $E_p = 11.0$ MeV), 14.5 MeV (1965RO22), 17.8 MeV (1966BA59), 25 MeV (1973BI1H, 1973BI1N), 30.3 MeV (1967WA31), 42.0 MeV (1966CA01; p), 49.3 MeV (1970CL10; also p_2), 49.8 MeV (1971MA13, 1971MA44; also p_2), 138.2 and 145 MeV (1966JA08, 1966JA1F) and 990 MeV (1972VO20). Polarization measurements on the proton group to ${}^9\text{Be}^*(2.4)$ [p_2] have also been carried out at $E_p = 8$ to 10 MeV (1969BE1Q). See also (1966LA04).

For reaction (b) see (1967WA1H, 1972WA30). For spallation see (1970KO25). For astrophysical considerations see (1971EL1B). See also (1966RO1B, 1969CO1J, 1970SL1B) and (1967SA1C, 1969GU1K, 1969WA11, 1971BA87, 1971IN05, 1973GU08, 1973HU05; theor.).

16. ${}^9\text{Be}(p, t){}^7\text{Be}$

$$Q_m = -12.0831$$

$$E_b = 6.5853$$

See ${}^7\text{Be}$.

17. (a) ${}^9\text{Be}(p, d){}^8\text{Be}$

$$Q_m = 0.5595$$

$$E_b = 6.5853$$

(b) ${}^9\text{Be}(p, \alpha){}^6\text{Li}$

$$Q_m = 2.1251$$

Knowledge of the cross sections of these two reactions at low energies is of importance for power generation and astrophysical considerations: see (1973SI27). Absolute cross sections for the d_0 and α_0 groups have been measured for $E_p = 28$ to 697 keV with $\pm 5 - 6\%$ uncertainty. The value of $S_{c.m.}(E = 0)$ for the combined cross sections is estimated to be (35^{+45}_{-15}) MeV \cdot b. At the 0.33 MeV resonance ($J^\pi = 1^-$), $\sigma_{\alpha_0} = 360 \pm 20$ mb and $\sigma_{d_0} = 470 \pm 30$ mb. The data (including angular distributions), analyzed by an R -matrix compound nucleus model, were fitted by assuming three states at $E_p(c.m.) = -20$ keV ($J^\pi = 2^+$; 3^+ possible) [$E_x = 6.57$ MeV], 310 keV (1^-) and 410 keV (1^+ ; 2^+ or 3^+ possible) (1973SI27). Measurements of the yields of d_0 , α_0 and α_1 for $E_p = 0.2$ to 2.0 MeV have been reported by (1969TU1A): the total cross section is reported to show definite peaks at $E_p = 0.330, 0.450, 0.935$ and 1.80 MeV. The yields of α_0 and d_0 have also been measured for $E_p = 0.3$ to 0.9 MeV by (1968BE1N), and the yield of d_0 has been obtained at several angles for $E_p = 6$ to 15 MeV (1972VO1H). The total cross sections for d_0 and d_1 have been measured for $E_p = 17.0, 21.0, 25.0$ and 29.1 MeV (1973MO02). See also (1970BO1Y, 1971GU23). Polarization measurements have been made at $E_p = 0.30$ to 1.25 MeV (1971KE1D; d_0, α_0), 0.33 MeV (1967RO07; d_0), 1.6 to 3.8 MeV (1965DA05, 1968FR10; d_0), 3, 4 and 5 MeV (1966VE03; d_0), 4.91, 6.90, 8.27, 9.80 MeV (1967IV01; d_0), 5 to 12 MeV (1969LO1E; d_0), 13.0 and 15.0 MeV (1972VO1H; d_0) and 185 MeV (1968IN02, 1969IN1C; d to ${}^8\text{Be}^*(16.9, 19.2)$). Previous studies are described in (1966LA04). Observed resonances are exhibited in Table 10.15. Deuteron production has been studied for $E_p = 1$ to 3 GeV by (1969ED02). See also (1966MI1E, 1969CO1J), and (1969KO1P, 1971MC1H, 1971SA1J).

18. ${}^9\text{Be}(d, n){}^{10}\text{B}$

$$Q_m = 4.3607$$

Neutron groups are observed corresponding to the ${}^{10}\text{B}$ states listed in Table 10.16. Thresholds for slow-neutron production corresponding to ${}^{10}\text{B}$ states from 4.77 to 6.56 MeV are reported in Table 10.17. Angular distributions have been measured at many energies: see (1959AJ76, 1966LA04) for earlier references and (1965BU10: $E_d = 2.6$ to 3.2, and 7 MeV; see Table 10.16), (1967FI01: $E_d = 3.0, 3.5$ and 5.5 MeV; see Table 10.16), (1967FU04: $E_d = 5.5$ MeV; n_0, n_2, n_3) and (1973PA14: $E_d = 7.0, 12.0, 15.0, 16.0$ MeV; $E_x < 6.6$ MeV). See also (1969RO14, 1971DE2C). The values of l_p and spectroscopic factors obtained in the analyses by (1965BU10, 1967FI01, 1973PA14) are shown in Table 10.16. The spectroscopic factors measured for ${}^{10}\text{B}^*(1.74)$ ($J^\pi = 0^+$; $T = 1$) are brought into accord with shell model predictions when the $\bar{t} \cdot \bar{T}$ interaction is considered (1967TA1A): see also (1966SI02, 1967FU04, 1973PA14).

There have been many reports of additional neutron groups [see (1959AJ76, 1966LA04)], in particular a state at 2.86 MeV. (1967GL03: $E_d = 1.7$ and 1.8 MeV) have shown that such a state does not exist: the intensity of a neutron group to such a state $\leq 0.4\%$ relative to ${}^{10}\text{B}^*(3.59)$ and $\leq 0.3\%$ relative to ${}^{10}\text{B}^*(2.16)$. See also (1965MA50).

Observed γ -transitions are listed in Tables 10.6 and 10.18 (1949RA02, 1963WA17, 1964WA05, 1969GA06). Reported values of τ_m for a number of ${}^{10}\text{B}$ states are displayed in Table 10.9 (1966WA10, 1968FI09, 1969AL17).

Table 10.15: Resonances in ${}^9\text{Be}(p, d){}^8\text{Be}$ and ${}^9\text{Be}(p, \alpha){}^6\text{Li}$

E_p (MeV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	Γ_p/Γ	θ_p^2	θ_d^2	θ_α^2	Refs.
0.34	6.89		$1^-; 0$	0.30	0.34	0.15	0.055	(1956MO90, 1969TU1A, 1973SI27)
0.46	7.00		$1^+(2^+, 3^+)^b$		0.3	0.3	0.1	(1949TH05, 1951NE03, 1969TU1A, 1973SI27)
(0.68)	(7.20)							(1949TH05, 1951NE03)
0.94	7.43	140	$(2^-; 0)$	0.7	0.04	0.02		(1956WE37, 1964HO02, 1969TU1A)
1.15	7.62	225 ± 50	$(1^+; 0)$	≈ 0.4	≈ 0.1			(1956WE37, 1964HO02)
1.65	8.07	800 ± 200	$(2^-; 0)$	≈ 0.07	0.18	0.21		(1956WE37, 1964HO02)
(1.80)	(8.20)							(1969TU1A)
(2.3)	(8.7)	(≈ 220)						(1956WE37, 1965MO27)
2.56	8.89	36 ± 2	$2^+; 1$				^a	(1956WE37)
3.5	9.7		$; 1$				^a	(1959MA20)
4.49	10.62		$; 1$				^a	(1959MA20)

^a Resonance for α_2 to ${}^6\text{Li}^*(3.56)$, $J^\pi = 0^+; T = 1$.

^b See, however, (1971BA41) in reaction 3.

From all the various experiments the following picture emerges: the first five states of ^{10}B have even parity [from l_p]. The ground state is known to have $J = 3$, by direct measurement, and $^{10}\text{B}^*(1.74)$ has $J^\pi = 0^+$ and is the $T = 1$ analogue of the $^{10}\text{C}_{\text{g.s.}}$ [from the β^+ decay of ^{10}C]. Then looking at the branching ratios and lifetimes of the other states, the sequence for $^{10}\text{B}^*(0, 0.72, 1.74, 2.16, 3.59)$ is $J^\pi = 3^+, 1^+, 0^+, 1^+, 2^+$, respectively [see discussion in (1966LA04, 1966WA10)].

See also ^{11}B and (1965FO1C, 1966FI01, 1970SR01, 1970WI1C, 1972SR04, 1972SR02, 1973WE1T, 1973WE19), (1966WA1C, 1967LE1F) and (1970FO1B, 1970MI1G, 1973CO16; theor.).

$$19. \ ^9\text{Be}(^3\text{He}, d)^{10}\text{B} \quad Q_m = 1.0916$$

Deuteron groups have been seen corresponding to a number of states of ^{10}B : see Table 10.19 (1960HI08, 1966FO08, 1966SI02, 1967CR04). See also (1970CA28). No evidence is seen for previously reported states at $E_x = 5.58$ and 6.40 MeV (1966FO08). Angular distributions obtained at $E_d = 10$ MeV (1967CR04), 10.2 MeV (1960HI08), 17 MeV (1966SI02) [see also (1973PA14)], 22 MeV (1969MA1R) and 25 MeV (1960WE04, 1967SI1A) have been analyzed by DWBA and lead to the l_p and the spectroscopic factors shown in Table 10.19. See also (1971BI1D), (1967LE1F) and (1968TA1A, 1971DZ07, 1973CO16; theor.). For τ_m measurements see Table 10.9 (1966FI01, 1968FI09).

$$20. \ ^9\text{Be}(\alpha, t)^{10}\text{B} \quad Q_m = -13.229$$

Angular distributions have been obtained at $E_\alpha = 27$ MeV (1973KE1E; t_0, t_1, t_3), 28.3 MeV (1965KA14; t_0, t_1) and $E_\alpha = 43$ MeV (1967DE1K, 1967SI1A; t_0, t_2). At $E_\alpha = 27$ MeV, $^{10}\text{B}^*(1.7)$ is not observed (1973KE1E). See also (1966LA04).

$$21. \ ^9\text{Be}(^{14}\text{N}, ^{13}\text{C})^{10}\text{B} \quad Q_m = -0.9654$$

See ^{13}C in (1970AJ04, 1976AJ04)

$$22. \ ^{10}\text{Be}(\beta^-)^{10}\text{B} \quad Q_m = 0.5559$$

See ^{10}Be .

$$\begin{aligned} 23. \ (a) \ ^{10}\text{B}(\gamma, n)^9\text{B} & \quad Q_m = -8.435 \\ (b) \ ^{10}\text{B}(\gamma, p)^9\text{Be} & \quad Q_m = -6.5853 \\ (c) \ ^{10}\text{B}(\gamma, d)^8\text{Be} & \quad Q_m = -6.0258 \\ (d) \ ^{10}\text{B}(\gamma, \alpha)^6\text{Li} & \quad Q_m = -4.460 \end{aligned}$$

Table 10.16: Levels of ^{10}B from $^9\text{Be}(d, n)^{10}\text{B}$ ^a

(1965BU10)			(1973PA14)		(1967FI01)	
$^{10}\text{B}^*$ (MeV \pm keV)	l_p ^d	S_{rel} ^e	S_{rel} ^k	$^{10}\text{B}^*$ (MeV \pm keV)	l_p ^g	S_{rel} ^h
0	1	1.0	1.0	0	1	1.0
0.72 ^b	1	2.3	1.97	0.72 \pm 10	1	2.8
1.74 ^b	1	1.0 ^f	1.42	1.74 \pm 10	1	1.4 ^f
2.16 ^b	1	0.41	0.41	2.15 \pm 10	1	0.5
3.59 ^b	1	0.45	0.10	3.58 \pm 10	(1)	0.3
4.77 ^b	(≥ 2)	0.43		4.77 \pm 10	undetermined	0.07 if $l = 1$ 0.4 if $l = 2$
5.11 \pm 20	0	1.3	0.14	5.11 \pm 12	0	1.7 ⁱ
5.16 \pm 20	1		0.43	5.17 \pm 14	1	0.6
5.90 \pm 80	(1)	1.3	0.49	5.93 \pm 10	1	0.7
6.10 \pm 80	(2)			{ 6.03 \pm 12 6.14 \pm 10	(2)	0.3 ^{i,j}
6.35 \pm 50	(0)	(0.86), 0.51				
6.50 \pm 50	1			6.57 \pm 10	(3)	
6.95 \pm 30 ^c	0	1.8		{ 6.89 \pm 15 7.00 \pm 12	(1) (1)	
7.50 \pm 50				7.48 \pm 15		
7.60 \pm 50				7.56 \pm 25		
(7.85 \pm 50)						
(8.07 \pm 50)						
(8.12 \pm 50)						

^a See also Table 10.17 in (1966LA04).

^b E_x not determined for the first five excited states.

^c $\Gamma \approx 200$ keV.

^d $E_d = 2.6 - 3.2$ MeV and 7 MeV.

^e $E_d = 7$ MeV.

^f See, however, discussion in text and (1967TA1A).

^g $E_d = 3.0, 3.5, 5.5$ MeV.

^h Average of values shown by (1967FI01).

ⁱ PWBA.

^j If $J = 3$.

^k Average of values shown by (1973PA14) at $E_d = 12.0, 15.0$ and 16.0 MeV.

Table 10.17: Slow-neutron thresholds
in ${}^9\text{Be}(d, n){}^{10}\text{B}$ (1954BO79)

E_d (MeV)	E_x (MeV)	Γ (keV)
0.52	4.79	< 10
0.92 ^a	5.11	< 10
0.99 ^a	5.17	< 10
1.92	5.93	< 10
2.08	6.06	< 10
2.20	6.16	< 20
2.53	6.43	
2.70	6.57	≈ 30

^a 0.921 ± 0.009 and 0.989 ± 0.009 MeV
(1962WA21): $E_x = 5.113$ and 5.169 MeV.

Absolute measurements have been made of the ${}^{10}\text{B}(\gamma, \text{all } n)$ cross section from threshold to 28 MeV: the giant resonance is broad with the major structure contained in two peaks at $E_x = 20.1 \pm 0.1$ and 23.1 ± 0.1 MeV ($\sigma \approx 6.5$ mb) (1973HU09, 1973HU1D). The integrated cross section to 29 MeV is 66.7 MeV \cdot mb (1965HA19). See also (1964GR40) and (1973AR1L). Using 12.5 MeV bremsstrahlung (1968SH21) report peaks in the (γ, p) cross section corresponding to ${}^{10}\text{B}$ states at 8.8 ± 0.1 [$J^\pi = 2^+$], 9.2 ± 0.1 [3^+] and (≈ 10.5) [$\pi = +$] MeV, with $T = 1$. See (1959AJ76, 1966LA04) for reactions (c) and (d).

24. ${}^{10}\text{B}(\gamma, \gamma){}^{10}\text{B}$

See (1967LO1B).

25. ${}^{10}\text{B}(e, e){}^{10}\text{B}$

The quadrupole contribution to the elastic form factor is best accounted for by the undeformed shell model, $Q = 7.45$ ($\pm 20\%$) fm^2 , $\langle r^2 \rangle^{1/2} = 2.45$ fm (1966ST12). The magnetic form factors for $E_e = 70$ to 200 MeV have been determined for the elastic group (+ a small contribution due to ${}^{10}\text{B}^*(0.7)$): there is no evidence of an M3 contribution – the magnetic octupole moment $\lesssim 1.6$ fm^2 , $a_0 = 1.4 \pm 0.2$ fm (1966RA29). At $E_e = 29.2$ to 57.8 MeV (1966SP02) has studied the inelastic groups corresponding to $E_x = 6.014 \pm 0.020$ ($J^\pi = 4^+$, E2) and 7.477 ± 0.020 MeV

Table 10.18: Gamma rays observed in ${}^9\text{Be}(d, n){}^{10}\text{B}$

Transition	E_γ (keV)	Branching ratio ^d (%)
0.7 → g.s.	716.6 ± 1 ^a	100
1.7 → g.s.		< 2
1.7 → 0.7	1022 ± 2 ^a	100
2.16 → g.s.	2152 ± 15 ^a	22
2.16 → 0.7	1433 ± 5 ^a	27
2.16 → 1.7	413.5 ± 1 ^a	51
3.59 → g.s.	3583 ± 13 ^{b,c}	19 ± 4 ^e
3.59 → 0.7	2872 ± 15 ^{b,c}	70 ± 7 ^e
3.59 → 1.74		< 1 ^e
3.59 → 2.16		11 ± 2 ^e
5.17 → g.s.	5159 ± 16 ^b	5.5 ± 0.7 ^b
5.17 → 0.7	4461 ± 13 ^{b,c}	29.5 ± 2 ^b
5.17 → 2.16	3028 ± 15 ^b	65 ± 2 ^b

^a (1949RA02).

^b (1963WA17): Doppler corrected.

^c M1 + E2 (1964WA05).

^d See Table 10.6.

^e (1969GA06).

($J^\pi = 2^+$, M1) [‡]: $\Gamma_{\gamma_0} = 0.122 \pm 0.020$ eV and 12.0 ± 2.2 eV [§], and $\Gamma_{\gamma_0}/\Gamma_w = 14$ and 1.4 , respectively. $\Gamma_{\text{tot}} \approx 40$ keV for ${}^{10}\text{B}^*(7.48)$ (1966SP02). A number of other inelastically scattered electron groups are reported by (1966KO08) at $E_e = 50$ MeV: Γ_{γ_0} values are given ($\pm 50\%$). See also (1965FR07, 1965VA1G).

See also (1966PE1E), (1966IS1A, 1967LE1E, 1968GO1J) and (1967KA1A, 1969UB1B, 1969VI02, 1970LA13; theor.).

26. ${}^{10}\text{B}(n, n){}^{10}\text{B}$

Elastic angular distributions have been measured for $E_n = 1.5$ to 5.0 MeV by (1970PO1E, 1973NE1H). At higher energies in addition to the elastic group (not resolved from the group

[‡] See, however, Table 10.11.

[§] Reanalysis of the data, including Coulomb distortion effects, leads to $\Gamma_{\gamma_0} = 11.0 \pm 2.2$ eV (1969CH1A) for ${}^{10}\text{B}^*(7.48)$. See also (1972THZF).

Table 10.19: Levels of ${}^9\text{Be}({}^3\text{He}, \text{d}){}^{10}\text{B}$

E_x (MeV \pm keV)	J^π	l_p ^e	S_{rel} ^f	S_{rel} ⁱ
0	3^+	1	1.0	1.0
0.717 ± 10 ^a	1^+	1	1.8	1.8
1.744 ± 10 ^a	0^+	1	2.6 ^g	2.6 ^g
2.156 ± 10 ^a	1^+	1	0.71	0.55
3.59 ^b	2^+	1	0.30	
4.77 ^c			0.15	
5.11 ^d				
			0.66 ^h	
5.17 ^{c,d}				
5.92 ^{c,d}	2^+	1	1.2	
6.03 ^d				
6.13 ^{c,d}			0.90 ^h	
6.56 ^{c,d}			0.86 ^h	

^a (1960HI08). See also (1970CA28).

^b The E_x of this state and of the states listed below have not been determined.

^c (1967CR04).

^d (1966FO08).

^e See text.

^f $E_d = 10$ MeV (1967CR04).

^g The $\bar{t} \cdot \bar{T}$ term significantly changes S_{rel} (1968TA1A).

^h $(2J + 1)S_{\text{rel}}$.

ⁱ $E_d = 17$ MeV (1966SI02); see also (1973PA14).

to $^{10}\text{B}^*(0.7)$), angular distributions have been obtained for the neutrons to $^{10}\text{B}^*(1.74 \pm 2.16, 3.59, 4.77, \text{unresolved states near } 5.0, 6.0, 6.6)$: see (1969HO1G: $E_n = 7.02$ and 7.55 MeV), (1970CO12: $E_n = 9.72$ MeV) and (1968AL1E, 1970VA19: $E_n = 14.1$ MeV). The gamma decay of ^{10}B states with $E_n < 6.1$ MeV has been studied by (1970NE03). See also (1969HA1R), (1966LA04) and ^{11}B in (1975AJ02).

27. $^{10}\text{B}(p, p')^{10}\text{B}$

Angular distributions of elastically scattered protons have been measured at $E_p = 3.0$ to 10.5 MeV (1970BO17), 5.1 to 16.5 MeV (1968WA1H, 1969WA23), 33.6 MeV (1970KU1D) and 49.5 MeV (1970SQ01). Observed inelastic groups are displayed in Table 10.20 (1953BO70, 1962AR02, 1964AR04). Angular distributions for the proton groups to the first five excited states and to $^{10}\text{B}^*(6.04)$ have been measured in the range $E_p = 5.1$ to 16.5 MeV (1968WA1H, 1969WA23). See also (1966LA04). The 4^+ state at 6.03 MeV is the most strongly excited of the ^{10}B states: see (1962SC12, 1965HA17, 1969DE1N, 1969WA23): $B(E2\uparrow) = 26 \pm 5 \text{ fm}^4$ and $\Gamma(E2\downarrow) = 0.14 \text{ eV}$ (1965JA1A). For optical model parameters see (1969WA11, 1970SQ01).

Gamma rays have been observed with $E_\gamma = 718.5 \pm 0.2$ keV and 1021.5 ± 0.5 keV (1966FR09: from decay of $E_x = 718.5 \pm 0.2$ and 1740.0 ± 0.6 keV) and $E_\gamma = 720.1 \pm 2.0, 1022.0 \pm 2.0, 1435.1, 2155.6 \pm 2.0$ and 2868.5 ± 2.0 keV (1969PA09: from decay of $E_x = 720.4 \pm 1.9, 1742.3 \pm 2.3, 2155.4 \pm 1.9$ and 3589.7 ± 2.2 keV). See also reaction 39. (1968MA18) report $\delta(E2/M1) = -(0.23_{-0.05}^{+0.06})$ or $-(4.1_{-0.7}^{+1.0})$ for the $2.15 \rightarrow 0.72$ transition. The $1.74 \rightarrow \text{g.s.}$ and $3.59 \rightarrow 1.74$ transitions have not been observed: $< 0.2\%$ and $< 0.3\%$, respectively (1966SE03). The branching ratio for $5.17 \rightarrow 1.74$ is $< 0.5\%$ and $5.17 \rightarrow 3.59$ is $(4.5 \pm 1)\%$ [see Table 10.6] (1967PA01). See Table 10.9 for other τ_m measurements (1966FI01, 1968FI09). See also (1971SC1N) and (1969TI02, 1973HU05, 1973KA04, 1973ZW1A; theor.). See also ^{11}C in (1975AJ02).

- | | |
|--|-----------------|
| 28. (a) $^{10}\text{B}(p, 2p)^9\text{Be}$ | $Q_m = -6.5853$ |
| (b) $^{10}\text{B}(p, pn)^9\text{B}$ | $Q_m = -8.435$ |
| (c) $^{10}\text{B}(p, p\alpha)^6\text{Li}$ | $Q_m = -4.460$ |

The summed proton spectrum (reaction (a)), observed at $E_p = 460$ MeV, shows peaks corresponding to the removal of an $l \neq 0$ proton at $Q = -6.7 \pm 0.5, -11.9 \pm 0.5$ and -17.1 ± 0.6 MeV; for removal of an $l = 0$ proton, $Q = -30.5 \pm 0.6$ MeV (1966TY01). See also (1966JA1A, 1967JA1E, 1968JA1G, 1969KO1J; theor.) and (1966LA04). For reaction (b) see (1970TH1F) and for reaction (c) see (1971GA1J).

29. $^{10}\text{B}(d, d')^{10}\text{B}^*$

Table 10.20: ^{10}B levels from $^{10}\text{B}(\text{p}, \text{p}')$, $^{10}\text{B}(\text{d}, \text{d}')$ and $^{10}\text{B}({}^3\text{He}, {}^3\text{He}')$

E_x (MeV \pm keV)			$\Gamma_{\text{c.m.}}$ (keV)	L	β_L
(1953BO70) ^a	(1962AR02) ^a	(1964AR04) ^a	(1964AR04)	(1968SQ01) ^h	(1968SQ01) ^h
0	0				
0.717 ± 5 ^b	0.72 ± 10			2	0.37 ± 0.04
1.739 ± 5 ^c	1.74 ± 10 ^c			(3)	(0.69 ± 0.09)
2.152 ± 5	2.15 ± 10			2	0.36 ± 0.04
3.583 ± 5	3.58 ± 10			2	0.36 ± 0.04
4.771 ± 5	4.77 ± 10				
	5.11 ± 10				
	5.16 ± 10 ^d				
	5.18 ± 10 ^e		110 ± 10 ^g		
		5.92 ± 10	< 5		
		6.03 ± 10	< 5		
		6.13 ± 10	< 5	2	0.62 ± 0.03
		6.55 ± 10	25 ± 5		
		7.00 ± 10	95 ± 10		
		7.48 ± 10 ^f	90 ± 15		

^a (p, p') and (d, d').

^b 719 ± 1.6 keV (1952CR30), 718 ± 5 keV (1954DA20).

^c This state not observed at $E_d = 7.6$ MeV (1953BO70) nor at $E_d = 10.0$ MeV (1962AR02). Intensity of this d_2 group $\lesssim 1\%$ of d_3 group for $E_d = 6$ to 9 MeV (1972ST1M).

^d Not observed in (d, d') at $E_d = 10$ MeV (1962AR02).

^e Not reported in (p, p') at $E_p = 10$ MeV (1962AR02).

^f Relative intensity in (d, d') at $E_d = 11.4$ MeV $< 2\%$.

^g (1962AR01).

^h (${}^3\text{He}, {}^3\text{He}'$): $E({}^3\text{He}) = 32.5$ MeV.

Deuteron groups have been observed corresponding to twelve states of ^{10}B : see Table 10.20 (1953BO70, 1962AR02, 1964AR04, 1972ST1M). The very low intensity of the group to $^{10}\text{B}^*(1.74)$ (1972ST1M) and the absence of the group to $^{10}\text{B}^*(5.17)$ (1962AR02) is good evidence of their $T = 1$ character. Angular distributions have been reported at $E_d = 4, 6$ and 8 MeV (1965LE1B, 1967DI01; d_0), 7.0 and 9.0 MeV (1972ST1M; d_2 and d_3), 11.8 MeV (1967FI07; $d_0, d_1, d_3, d_4, d_5, d_6$ and d to unresolved 6 MeV states) and 28 MeV (1968GA13, 1968ME1E; d_0 ; see for optical parameters). See also (1966BR1G), (1968NE1A, 1969NE08; theor.) and ^{12}C in (1975AJ02).

30. $^{10}\text{B}(t, t)^{10}\text{B}$

Angular distributions of elastically scattered tritons have been measured at $E_t = 1.5$ MeV (1963HO19, 1969HE08) and $2.5, 2.7, 3.1$ and 3.3 MeV (1971GE09).

31. $^{10}\text{B}(^3\text{He}, ^3\text{He})^{10}\text{B}$

Angular distributions of elastically scattered ^3He have been measured at $E(^3\text{He}) = 4, 8, 10, 12, 15$ and 18 MeV (1970DU07), 9.8 MeV (1967PA1H), $13.2, 17.2$ and 24.3 MeV (1972BU30), 14 MeV (1970NU02) and 32.5 MeV (1968SQ01). See also (1970BA1P). For derived optical model parameters see (1968SQ01, 1970DU07, 1970NU02, 1972BU30). Angular distributions have also been measured at $E(^3\text{He}) = 32.5$ MeV to $^{10}\text{B}^*(0.72, 1.74, 2.16, 3.59, 4.77, 5.11 + 5.17 + 5.18, 6.03, 6.56)$. $L = 2$ gives a good fit to the distributions of ^3He ions to $^{10}\text{B}^*(0.72, 2.16, 3.59, 6.03)$: derived β_L are shown in Table 10.20 (1968SQ01). See also (1967CO1J, 1970BA1P) and (1968HO1C).

32. (a) $^{10}\text{B}(\alpha, \alpha')^{10}\text{B}^*$

$$(b) \ ^{10}\text{B}(\alpha, 2\alpha)^6\text{Li} \quad Q_m = -4.460$$

Angular distributions of elastically scattered α -particles have been measured at $E_\alpha = 5$ to 30 MeV (1972DA04) and at 56 MeV (1969GA11). For optical model parameters see (1970ZE03, 1972DA04). Inelastic α -groups are also reported to $^{10}\text{B}^*(0.72, 2.16, 3.59, 4.77, 5.11 + 5.17 + 5.18, 5.92 + 6.03, 6.13, 6.56)$ (1972DA04). $^{10}\text{B}^*(1.74)$ is not seen (1967CO1P, 1972DA04).

At $E_\alpha = 24$ MeV, analysis of the angular distributions of the α -particles emitted in the decay of $^{10}\text{B}^*(4.77, 5.92)$ suggests 4^+ for the former (see, however, Table 10.5) and 2^+ for the latter (1973EI03). See also (1966GE12, 1971GA1J, 1974DO1G) and (1969GA01) in ^{14}N in (1970AJ04).

33. (a) $^{10}\text{B}(^6\text{Li}, ^6\text{Li})^{10}\text{B}$
(b) $^{10}\text{B}(^7\text{Li}, ^7\text{Li})^{10}\text{B}$

For reaction (a) see (1970BO1W). The elastic scattering in reaction (b) has been studied at $E(^7\text{Li}) = 24$ MeV (1972WE08). See also (1969RO1G; theor.).

34. $^{10}\text{B}(^{10}\text{B}, ^{10}\text{B})^{10}\text{B}$

See (1967BE1T).

35. $^{10}\text{B}(^{12}\text{C}, ^{12}\text{C})^{10}\text{B}$

The elastic scattering angular distribution has been measured at $E(^{10}\text{B}) = 18$ MeV (1968VO1A, 1969VO10).

36. $^{10}\text{B}(^{14}\text{N}, ^{14}\text{N})^{10}\text{B}$

See (1970IS1A).

37. (a) $^{10}\text{B}(^{16}\text{O}, ^{16}\text{O})^{10}\text{B}$
(b) $^{10}\text{B}(^{18}\text{O}, ^{18}\text{O})^{10}\text{B}$

The elastic scattering in reaction (a) has been studied with $E(^{16}\text{O}) = 15.0$ to 32.5 MeV: see (1968OK06, 1968OK1B, 1969KR03); that in reaction (b) has been studied for $E(^{18}\text{O}) = 20, 24$ and 30.5 MeV (1971KN05). See also (1970BL1E).

38. $^{10}\text{B}(^{19}\text{F}, ^{19}\text{F})^{10}\text{B}$

The elastic scattering has been studied for $E(^{19}\text{F}) = 20$ and 24 MeV (1971KN05).

39. $^{10}\text{C}(\beta^+)^{10}\text{B}$ $Q_m = 3.650$

Table 10.21: The beta decay of ^{10}C

Decay to $^{10}\text{B}^*$ (MeV)	Branching ratio (%)	$\log ft$	Refs.
0.72	98.53 ± 0.02	3.047^{d}	(1972RO03) ^a
1.74	1.465 ± 0.014	$3.499^{+0.005}_{-0.006}$ ^b	(1972RO03) ^a
	1.52 ± 0.08		(1969BR13)
2.15	$\leq 8 \times 10^{-4}$ ^c	≥ 5.67	(1972GO1A)

^a See also (1966FR09, 1969FR02, 1969FR09).

^b Includes radiative corrections (1972RO03). (1973TO04) using the same input data find $3.497^{+0.005}_{-0.006}$, including both radiative and charge dependent corrections.

^c See also (1969BR13).

^d B. Zimmerman and G. Fox, private communication.

The half-life is 19.48 ± 0.05 sec (1962EA02), 19.27 ± 0.08 sec (1963BA52): the weighted mean is 19.42 ± 0.06 sec [¶]. The decay is to $^{10}\text{B}^*(0.7, 1.7)$: see Table 10.21 for branching ratios and $\log ft$ (1969BR13, 1972GO1A, 1972RO03). The vector coupling constant determined from the decay to the analog state $^{10}\text{B}^*(1.7)$ is $G_{\beta}^{\text{V}} = (1.396 \pm 0.009) \times 10^{-49}$ erg \cdot cm³ (1972RO03). See also (1972FR1L). The excitation energies of $^{10}\text{B}^*(0.7, 1.7)$ are $E_x = 718.32 \pm 0.09$ and 1740.16 ± 0.17 keV [$E_{\gamma} = 718.29 \pm 0.09$ and 1021.78 ± 0.14 keV] (1969FR02). See also (1966FR09). See also (1967MU1A, 1969FR09) and (1965CO25, 1966BA1A, 1968BO1U, 1968MO1F, 1969BL1E, 1969BL1D, 1969KA1B, 1970DA21, 1972BE04, 1972WI1C, 1973HA49, 1973TO04, 1973WI11; theor.).

$$40. \ ^{11}\text{B}(\gamma, n)^{10}\text{B} \quad Q_{\text{m}} = -11.4560$$

The intensities of the transitions to $^{10}\text{B}^*(3.59, 5.17)$ [$T = 0$ and 1 , respectively] depend on the region of the giant dipole resonance in ^{11}B from which the decay takes place: it is suggested that the lower energy region consists mainly of $T = \frac{1}{2}$ states and the higher energy region of $T = \frac{3}{2}$ states (1971PA10). See also reaction 22 in ^{10}Be , and ^{11}B in (1975AJ02). See also (1969MU10).

$$41. \ ^{11}\text{B}(\text{p}, \text{d})^{10}\text{B} \quad Q_{\text{m}} = -9.2314$$

Angular distributions of deuteron groups have been measured at $E_{\text{p}} = 19$ MeV (1963LE03; $\text{d}_0, \text{d}_1, \text{d}_2, \text{d}_3, \text{d}_4$), 33.6 MeV (1968KU04, 1970KU1D; $\text{d}_0, \text{d}_1, \text{d}_2, \text{d}_3, \text{d}_4, \text{d}_5$ and deuterons to states

[¶] See also Table 10.24.

at 5.18 (unres.), 6.04 (unres.) and 154.8 MeV (1969BA05, 1969TO1A; d_0, d_1, d_2, d_3, d_4 and d to $5.1 \pm 0.1, 11.4 \pm 0.2$ and 14.1 ± 0.2 MeV). The weak excitation of states at 6.56 and 7.5 MeV is also reported (1968KU04), and $\Gamma_{c.m.}$ of $^{10}\text{B}^*(11.4) = 1.1 \pm 0.2$ MeV (1969BA05). Spectroscopic factors have been extracted by [(1963LE03): PWBA; (1968KU04): DWBA; (1969BA05) and (1969TO1A): DWBA].

42. $^{11}\text{B}(d, t)^{10}\text{B}$ $Q_m = -5.1984$

Angular distributions have been measured at $E_d = 11.8$ MeV (1967FI07; $t_0, t_1, t_2, t_3; l = 1; S = 1.88, 0.94, 1.35, 1.35$, respectively) and at $E_d = 21.6$ MeV (1967DE1M; $t_0 \rightarrow t_5$ and t to $^{10}\text{B}^*(5.1); S_{rel}$ extracted). A dependence of the angular distribution on the isospin of the final state is discussed by (1967FU04).

43. (a) $^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B}$ $Q_m = 9.1225$
 (b) $^{11}\text{B}(^3\text{He}, 2\alpha)^6\text{Li}$ $Q_m = 4.662$

Reported levels are listed in Table 10.22 (1965GO05, 1967PU04). See also (1966LA04). Angular distributions have been measured at $E(^3\text{He}) = 1.0, 1.8, 2.15$ MeV (1966LO15; $\alpha_0 \rightarrow \alpha_3$), 8, 10, 12 MeV (1973CO19; α_0) and 33 MeV (1969DE10: see Table 10.22). See also the earlier work of (1960TA12, 1965FO06). The decays of many of the states have been studied by (1966AL06, 1968WA15, 1969YO01): see Table 10.6. Lifetime measurements are summarized in Table 10.9 (1968DO01, 1969JA1N).

Alpha- α angular correlations (reaction (b)) have been measured for the transitions via $^{10}\text{B}^*(5.92, 6.03, 6.13, 6.56, 7.00)$. The results are consistent with $J^\pi = 2^+$ and 4^+ for $^{10}\text{B}^*(5.92, 6.03)$ and require $J^\pi = 3^-$ for $^{10}\text{B}^*(6.13)$. There is substantial interference between levels of opposite parity for the α -particles due to $^{10}\text{B}^*(6.56, 7.00)$: the data are fitted by $J^\pi = 3^+$ for $^{10}\text{B}^*(7.00)$ and $(3, 4)^-$ for $^{10}\text{B}^*(6.56)$ [the $^6\text{Li}(\alpha, \alpha)$ results then require $J^\pi = 4^-$] (1971YO05). See, however, (1973SI27) in reaction 17. See also (1970LI1K).

44. $^{11}\text{B}(^{16}\text{O}, ^{17}\text{O})^{10}\text{B}$ $Q_m = -7.314$

See (1968OK06) and (1969BR1D).

45. $^{12}\text{C}(\gamma, d)^{10}\text{B}$ $Q_m = -25.1885$

See (1972SK08).

Table 10.22: ^{10}B levels from $^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B}$

E_x^a (MeV \pm keV)	E_x^b (MeV \pm keV)	$\Gamma_{\text{c.m.}}^b$ (keV)	l^c	S_{rel}^c
0			1	1.0
0.718 \pm 7			1	0.22
1.744 \pm 7			1	0.73
2.157 \pm 6			1	0.44
3.587 \pm 6			1	0.09
4.777 \pm 5			1	0.09
5.114 \pm 5				
5.166 \pm 5			1	1.81
5.923 \pm 5				
6.028 \pm 5				
6.131 \pm 5				
6.573 \pm 8	6.566 \pm 10	30 \pm 10		
	7.002 \pm 10	95 \pm 10		
7.475 \pm 10				
7.567 \pm 10				
	7.87 \pm 40	240 \pm 50		
	11.53 \pm 40	270 \pm 50		
	12.57 \pm 30	90 \pm 30		

^a (1965GO05).

^b (1967PU04).

^c (1969DE10): 33 MeV; JULIE code.

46. $^{12}\text{C}(\text{n}, \text{t})^{10}\text{B}$ $Q_{\text{m}} = -18.9309$

Not reported.

47. (a) $^{12}\text{C}(\text{p}, ^3\text{He})^{10}\text{B}$ $Q_{\text{m}} = -19.6948$

(b) $^{12}\text{C}(\text{p}, \text{pd})^{10}\text{B}$ $Q_{\text{m}} = -25.1885$

Angular distributions of ^3He ions have been measured at $E_{\text{p}} = 39.8$ MeV (1973HO10: to $^{10}\text{B}^*(0, 0.72, 1.74, 2.16, 3.59)$) and 185 MeV (1968TI1A, 1972DA26: to $^{10}\text{B}^*(0, 1.0, 1.7, 2.5, 3.5, 4.8, 6.0, 7.3)$). See also (1966KI07) and (1968SA1H, 1971BA61; theor.). For reaction (b) see (1972BO17), (1968PA1J) and (1966LH1A, 1971BA16; theor.).

48. $^{12}\text{C}(\text{d}, \alpha)^{10}\text{B}$ $Q_{\text{m}} = -1.3409$

Alpha groups have been observed to the known states of ^{10}B below $E_{\text{x}} = 7.1$ MeV: see Table 10.23 (1962AR02, 1965PE17, 1970AN1J). Angular distributions have been determined for $\alpha_0, \alpha_1, \alpha_3$ and α_4 at 9.2 to 13.9 MeV (1965BA06, 1966BA32, 1968KL06), 11.4 and 12.4 MeV (1965DO08), 12 MeV (1970AN1J), 13 MeV (1965ME13), 14.7 to 19.7 MeV (1961YA08, 1963YA1B) and 24 MeV (1965PE17). Angular distributions have also been measured at $E_{\text{d}} = 5.00$ to 8.40 MeV (1969CO02; α_0, α_1), 6.2 to 6.8 MeV (1965NE10; α_0, α_1), 12 MeV (1970AN1J; α to $^{10}\text{B}^*(4.77, 5.11, 5.17, 5.92, 6.03, 6.13, 6.56)$), 24 MeV (1965PE17; α to $^{10}\text{B}^*(4.77, 5.17, 6.03)$). See also (1969CU08).

The population of the isospin-forbidden group to $^{10}\text{B}^*(1.74) [\alpha_2]$ has been studied with E_{d} up to 30 MeV. For $E_{\text{d}} \lesssim 15$ MeV the reaction appears to proceed via compound nucleus formation [see ^{14}N]; at the higher energies direct interaction mechanism may be involved: see (1966LA04) and (1966ME09, 1968JA09, 1969SM03, 1971JA04, 1971RI15, 1971VO04). See also (1966HA09). Angular distributions of the α_2 group have been reported at $E_{\text{d}} = 7.0$ to 14.0 MeV (1969SM03, 1971RI15), 9.0 to 12.5 MeV (1966ME09), 14.0 to 17.0 MeV (1971VO04), 14.8, 18.0, 21.0, 29.1 MeV (1968JA09, 1971JA04).

See also (1966PA1J, 1967BR1B, 1969DE29), (1966BR1G, 1969SC1F) and (1966DR1C, 1966GR1G, 1966HO1D, 1966ME1E, 1966WA1G, 1967NO1A, 1968NO1C, 1968ZE1B, 1969NO1B, 1969NO1C, 1971BA61; theor.).

49. $^{12}\text{C}(\alpha, ^6\text{Li})^{10}\text{B}$ $Q_{\text{m}} = -23.715$

At $E_{\alpha} = 42$ MeV angular distributions have been measured for the transitions to $^6\text{Li}_{\text{g.s.}} + ^{10}\text{B}_{\text{g.s.}}$, $^6\text{Li}_{\text{g.s.}} + ^{10}\text{B}_{0.72}^*$, $(^6\text{Li}_{2.19}^* + ^{10}\text{B}_{\text{g.s.}}) + (^6\text{Li}_{\text{g.s.}} + ^{10}\text{B}_{2.15}^*)$, $^6\text{Li}_{\text{g.s.}} + ^{10}\text{B}_{2.15}^*$, $^6\text{Li}_{2.19}^* + ^{10}\text{B}_{0.72}^*$ (1972RU03). See also ^{16}O in (1977AJ02).

Table 10.23: ^{10}B levels from $^{12}\text{C}(d, \alpha)^{10}\text{B}$

E_x^a (MeV \pm keV)	E_x^c (MeV \pm keV)
	0
0.72 ± 10 b	0.72 ± 20 d
2.15 ± 10	2.15 ± 20
3.58 ± 10	3.59 ± 20
4.77 ± 10	4.77 ± 30
5.11 ± 10 b	5.17 ± 50 e
	6.04 ± 50 e
	6.67 ± 110
	7.05 ± 100

^a (1962AR02).

^b Absent: intensity of $E_x = 5.17$ MeV group $< 1\%$ relative to average of $T = 0$ levels.

^c (1965PE17).

^d See discussion in text.

^e The excitation of $^{10}\text{B}^*(5.92, 6.13, 6.56)$ is also reported by (1970AN1J).

$$50. \ ^{12}\text{C}(^{10}\text{B}, 3\alpha)^{10}\text{B} \quad Q_m = -7.2748$$

See (1965SH1A, 1967BI1F).

$$51. \text{ (a) } ^{12}\text{C}(^{11}\text{B}, ^{13}\text{C})^{10}\text{B} \quad Q_m = -6.5096$$

$$\text{ (b) } ^{12}\text{C}(^{14}\text{N}, ^{16}\text{O})^{10}\text{B} \quad Q_m = -4.4518$$

See (1969BR1D, 1972MO1E, 1973SC1J).

$$52. {}^{13}\text{C}(\text{p}, \alpha){}^{10}\text{B} \quad Q_{\text{m}} = -4.0627$$

Angular distributions have been measured at $E_{\text{p}} = 43.7$ and 50.5 MeV for the α -particle groups to ${}^{10}\text{B}^*(1.74)$ [$L = 1$] and ${}^{10}\text{B}^*(6.03)$ [$L = 3$] ([1972MA21](#)).

$$53. {}^{13}\text{C}({}^{11}\text{B}, {}^{14}\text{C}){}^{10}\text{B} \quad Q_{\text{m}} = -3.2790$$

See ([1969BR1D](#), [1972MO1E](#), [1973SC1J](#)).

$$54. \text{(a) } {}^{14}\text{N}(\gamma, \alpha){}^{10}\text{B} \quad Q_{\text{m}} = -11.6134$$

$$\text{(b) } {}^{14}\text{N}(\text{p}, \alpha\text{p}){}^{10}\text{B} \quad Q_{\text{m}} = -11.6134$$

For reaction (a) see ([1959AJ76](#)); for reaction (b) see ([1961CL09](#)).

$$55. {}^{14}\text{N}(\text{d}, {}^6\text{Li}){}^{10}\text{B} \quad Q_{\text{m}} = -10.140$$

See ([1970MC1G](#)).

$$56. {}^{14}\text{N}({}^3\text{He}, {}^7\text{Be}){}^{10}\text{B} \quad Q_{\text{m}} = -10.027$$

At $E({}^3\text{He}) = 41$ MeV, the population of ${}^{10}\text{B}^*(0, 0.72, 2.16, 3.59, 6.1)$ has been observed. The transition to ${}^{10}\text{B}^*(1.74)$ [$T = 1$] is very weak ([1971DE37](#)).

$$57. {}^{15}\text{N}({}^{11}\text{B}, {}^{16}\text{N}){}^{10}\text{B} \quad Q_{\text{m}} = -8.966$$

See ([1969BR1D](#)).

$$58. {}^{16}\text{O}(\text{p}, {}^7\text{Be}){}^{10}\text{B} \quad Q_{\text{m}} = -20.533$$

See ([1969HO1H](#)).

$$59. {}^{20}\text{Ne}({}^{11}\text{B}, {}^{21}\text{Ne}){}^{10}\text{B} \quad Q_{\text{m}} = -4.695$$

See ([1969BR1D](#)).

^{10}C
(Figs. 21 and 22)

GENERAL: (See also (1966LA04).)

Model calculations: (1968FA1B, 1969SA1A, 1969SO08, 1973SA30).

Special reactions: (1967AU1B, 1969YI1A, 1971AR02).

Pion reactions (See also reaction 3.): (1973AL1E, 1973CH20).

Other topics: (1968FA02, 1968FA1B, 1969SO08, 1969SO1E, 1970FO1B, 1972AN05, 1972CA37).

Ground-state properties: (1966KE16, 1968FA02, 1968FA1B, 1969GA1G, 1973SA30).

1. $^{10}\text{C}(\beta^+)^{10}\text{B}$ $Q_m = 3.650$

^{10}C decays with a half-life of 19.42 ± 0.06 sec to $^{10}\text{B}^*(0.7, 1.7)$: the branching ratios are $98.53 \pm 0.02\%$ and $1.465 \pm 0.014\%$, respectively (1972RO03): see reaction 39 in ^{10}B and Tables 10.21 and 10.24.

2. $^7\text{Be}(^3\text{He}, 2p)^4\text{He}^4\text{He}$ $Q_m = 11.274$ $E_b = 14.999$

See (1972PA1C; astrophys.).

3. $^9\text{Be}(p, \pi^-)^{10}\text{C}$ $Q_m = -136.634$

At $E_p = 185$ MeV the π^- spectrum shows groups corresponding to $^{10}\text{C}^*(0, 3.36 \pm 0.07, 5.28 \pm 0.06, 6.63 \pm 0.15)$. Angular distributions have been obtained for the π^- corresponding to these four states. The π^+/π^- intensity ratio is usually $\gg 1$ for analog states in ^{10}Be and ^{10}C (1973DA09). See also (1971DA10, 1972RO22, 1973DI1J).

4. $^9\text{Be}(^3\text{He}, 2n)^{10}\text{C}$ $Q_m = -5.566$

See (1973MO1N).

5. $^{10}\text{B}(p, n)^{10}\text{C}$ $Q_m = -4.433$

Table 10.24: Energy levels of ^{10}C

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$0^+; 1$	$\tau_{1/2} = 19.42 \pm 0.06$ sec ^b	β^+	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
3.3527 ± 1.5	$2^+; 1$	$\tau_m = 155 \pm 25$ fsec	γ	3, 5, 6, 7, 9, 11
5.28 ± 40	$(2^+); 1$	$\Gamma = 300 \pm 50$		3, 5, 6, 9
6.60 ± 40		$\Gamma = 300 \pm 50$		3, 5, 6, 9
(10.2 ± 200)		$\Gamma \approx 1500$		9
10.72 ^a				11

^a Three states with $E_x \approx 10$ MeV are reported in reaction 11 (preliminary results).

^b (1974AZ01) report $\tau_{1/2} = 19.28 \pm 0.02$ sec.

$$E_{\text{thresh.}} = 4880.0 \pm 2.0 \text{ keV};$$

$$Q_0 = -4432.8 \pm 1.8 \text{ keV (1966FR09)}.$$

The first excited state of ^{10}C is located at $E_x = 3.3527 \pm 0.0015$ MeV (1969PA09: from γ -decay), 3.38 ± 0.03 MeV (1963EA01: neutron threshold), 3.35 ± 0.01 MeV (1966SE03: from γ -decay). τ_m for $^{10}\text{C}^*(3.35) = 155 \pm 25$ fsec; $\Gamma_\gamma = 4.25 \pm 0.69$ meV (1968FI09). Angular distributions have been measured for the n_0 and n_1 groups and for the neutrons corresponding to $^{10}\text{C}^*(5.2 \pm 0.3)$ at $E_p = 30$ and 50 MeV. The excitation of $^{10}\text{C}^*(6.5 \pm 0.3, 8.3 \pm 0.3, 8.9 \pm 0.3)$ is also reported (1970CL01). See also (1969BR13, 1973GO1V), (1966LA04) and ^{11}C in (1975AJ02).

6. $^{10}\text{B}(^3\text{He}, t)^{10}\text{C}$ $Q_m = -3.669$

At $E(^3\text{He}) = 14$ MeV the angular distribution of the t_0 group has been measured (1970NU02). At $E(^3\text{He}) = 30.6$ and 40 MeV triton groups have been observed to $^{10}\text{C}^*(0, 3.37 \pm 0.03, (5.03 \pm 0.06), 5.28 \pm 0.06, (5.60 \pm 0.06), 6.58 \pm 0.06)$ (1966MA36). At $E(^3\text{He}) = 30.3$ MeV, the energy of the first excited state is determined to be 3.344 ± 0.008 MeV, and the triton groups corresponding to the two states at $E_x = 5.03$ and 5.60 MeV suggested by (1966MA36) are assigned to ^{11}C (1968BR23). See also ^{13}N in (1976AJ04).

7. $^{10}\text{B}(^6\text{Li}, ^6\text{He})^{10}\text{C}$ $Q_m = -7.160$

Both $^{10}\text{C}^*(0, 3.35)$ are strongly populated at $E(^6\text{Li}) = 30$ MeV, $\theta = 16^\circ$ (1972LE1P).

8. $^{12}\text{C}(\gamma, 2n)^{10}\text{C}$ $Q_m = -31.846$

See (1959OC07).

9. $^{12}\text{C}(\text{p}, \text{t})^{10}\text{C}$ $Q_m = -23.364$

Angular distributions have been reported at $E_p = 30.0$ to 54.1 MeV (1967CO23; t_0), 43 MeV (1967BE27; t_0, t_1 , and to $^{10}\text{C}^*(5.3)$), 43.7 MeV (1966CE05; t_0, t_1) and 49.5 MeV (1970NE17; t_0, t_1 ; polarized protons). $L = 0, 2$ and 2 , respectively for $^{10}\text{C}^*(0, 3.35, 5.3)$, and therefore $J^\pi = 0^+, 2^+$ and 2^+ : see (1967BE27). [$^{10}\text{C}^*(5.3)$ may correspond to several unresolved states: see ^{10}Be .] At $E_p = 43$ and 52 MeV (1967BE27) report the excitation of $^{10}\text{C}^*(5.29 \pm 0.06, 6.61 \pm 0.06)$ [$\Gamma_{\text{c.m.}} = 300 \pm 50$ keV for both these states]. The states at $E_x = 7.2 \pm 0.2$ and 10.2 ± 0.2 MeV reported by (1965BA17) are not observed. See also (1965CA1C, 1968TA1P) and (1971KA04; theor.).

10. $^{12}\text{C}(^{12}\text{C}, ^{10}\text{C})^{14}\text{C}$ $Q_m = -18.723$

See (1972MO1E, 1973SC1J).

11. $^{13}\text{C}(^3\text{He}, ^6\text{He})^{10}\text{C}$ $Q_m = -15.243$

At $E(^3\text{He}) = 70.3$ MeV the angular distributions of the ^6He ions corresponding to the population of $^{10}\text{C}^*(0, 3.35)$ have been measured. The group to $^{10}\text{C}^*(3.35)$ is much more intense than the ground state group: multi-step processes may be important (1973KA16). Three states of ^{10}C are observed with $E_x \approx 10$ MeV. One of these [$E_x = 10.72$ MeV] is very strongly populated. A search for a $T = 2$ state with $E_x \approx 22$ MeV was unsuccessful (1973BE2J).

12. $^{14}\text{N}(^3\text{He}, ^7\text{Li})^{10}\text{C}$ $Q_m = -12.816$

See (1971DE37).

References

(Closed December 31, 1973)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1949RA02 V. K. Rasmussen, W.F. Hornyak and T. Lauritsen, Phys. Rev. 76 (1949) 581
- 1949TH05 R.G. Thomas, S. Rubin, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 75 (1949) 1612
- 1950WU1A Wu, Rev. Mod. Phys. 22 (1950) 386
- 1951BO45 C.K. Bockelman, D.W. Miller, R.K. Adair and H.H. Barschall, Phys. Rev. 84 (1951) 69
- 1951NE03 J.A. Neuendorffer, D.R. Inglis and S.S. Hanna, Phys. Rev. 82 (1951) 75
- 1952CR30 D.S. Craig, D.J. Donahue and K.W. Jones, Phys. Rev. 88 (1952) 808
- 1952HA10 T.M. Hahn, C.W. Snyder, H.B. Willard, J.K. Bair, E.D. Klema, J.D. Kington and F.P. Green, Phys. Rev. 85 (1952) 934
- 1953BO70 C.K. Bockelman, C.P. Browne, W.W. Buechner and A. Sperduto, Phys. Rev. 92 (1953) 665
- 1953MA1A Mackin, Thesis, CalTech (1953)
- 1953PA22 E.B. Paul and H.E. Gove, Proc. Roy. Soc. Canada 47 (1953) 145A
- 1954BO79 T.W. Bonner and C.F. Cook, Phys. Rev. 96 (1954) 122
- 1954DA20 R.B. Day and T. Huus, Phys. Rev. 95 (1954) 1003
- 1954JU23 J.J. Jung and C.K. Bockelman, Phys. Rev. 96 (1954) 1353
- 1955MA84 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 91
- 1955WI25 H.B. Willard, J.K. Bair and J.D. Kington, Phys. Rev. 98 (1955) 669
- 1956CL69 A.B. Clegg, Phil. Mag. 1 (1956) 1116
- 1956DE33 G. Dearnaley, Phil. Mag. 1 (1956) 821
- 1956MA55 J.B. Marion, Phys. Rev. 103 (1956) 713
- 1956MO90 F.S. Mozer, Phys. Rev. 104 (1956) 1386
- 1956WE37 G. Weber, L.W. Davis and J.B. Marion, Phys. Rev. 104 (1956) 1307
- 1957BI84 H. Bichsel and T.W. Bonner, Phys. Rev. 108 (1957) 1025
- 1957ME27 L. Meyer-Schutzmeister and S.S. Hanna, Phys. Rev. 108 (1957) 1506
- 1957ST95 P.H. Stelson and E.C. Campbell, Phys. Rev. 106 (1957) 1252

1958ME81 W.E. Meyerhof and L.F. Chase, Jr., Phys. Rev. 111 (1958) 1348
 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
 1959MA20 J.B. Marion and J.S. Levin, Phys. Rev. 115 (1959) 144
 1959MA34 J.B. Marion, J.S. Levin and L. Cranberg, Phys. Rev. 114 (1959) 1584
 1959ME85 W.E. Meyerhof, N.W. Tanner and C.M. Hudson, Phys. Rev. 115 (1959) 1227
 1959OC07 J. O'Connell, P. Dyal and J. Goldemberg, Phys. Rev. 116 (1959) 173
 1959WA16 E.K. Warburton, Phys. Rev. 113 (1959) 595
 1960BA01 W.D. Barfield, H.T. Motz and R.E. Carter, Bull. Amer. Phys. Soc. 5 (1960) 44, MA4
 1960HI08 S. Hinds and R. Middleton, Proc. Phys. Soc. 75 (1960) 754
 1960KL03 P.R. Klein, N. Cindro, L.W. Swenson and N.S. Wall, Nucl. Phys. 16 (1960) 374
 1960MO17 G.C. Morrison, Phys. Rev. Lett. 5 (1960) 565
 1960SE12 R. Seltz and D. Magnac-Valette, Compt. Rend. 251 (1960) 2006
 1960TA12 I.J. Taylor, F.de S. Barros, P.D. Forsyth, A.A. Jaffe and S. Ramavataram, Proc. Phys. Soc. A75 (1960) 772
 1960WE04 H.E. Wegner and W.S. Hall, Phys. Rev. 119 (1960) 1654
 1961BA53 R. Bass, T.W. Bonner and H.P. Haenni, Nucl. Phys. 23 (1961) 122
 1961CL09 A.B. Clegg, K.J. Foley, G.L. Salmon and R.E. Segel, Proc. Phys. Soc. 78 (1961) 681
 1961FO07 D.B. Fossan, R.L. Walter, W.E. Wilson and H.H. Barschall, Phys. Rev. 123 (1961) 209
 1961JA19 von L. Jarczyk, J. Lang, R. Muller and W. Wolffi, Helv. Phys. Acta 34 (1961) 483
 1961KO1A Koltun, Phys. Rev. 124 (1961) 1162
 1961LA1A Lane, Langsdorf, Monahan and Elwyn, Ann. Phys. 12 (1961) 135
 1961MO02 G.C. Morrison, Phys. Rev. 121 (1961) 182
 1961SP02 E.L. Sprenkel, J.W. Olness and R.E. Segel, Phys. Rev. Lett. 7 (1961) 174
 1961TA02 N.W. Tanner and S.S. Hanna, Nucl. Phys. 23 (1961) 319
 1961TR1B True and Warburton, Nucl. Phys. 22 (1961) 426
 1961VA43 A.K. Val-ter, P.I. Vatsset, L.Y. Kolesnikov, S.G. Tonapetyan, K.K. Chernyavskii and A.I. Shpetnyi, Atomn. Energ. (USSR) 10 (1961) 577; Sov. J. At. Energy 10 (1962) 574; Correction At. Energy 123 (2018) 432
 1961WO05 E.A. Wolicki and A.R. Knudson, Bull. Amer. Phys. Soc. 6 (1961) 415, B2
 1961YA08 T. Yanabu, J. Phy. Soc. Jpn. 16 (1961) 2118
 1962AL1A Altman, MacDonald and Marion, Nucl. Phys. 35 (1962) 85

1962AR01 B.H. Armitage and R.E. Meads, Phys. Lett. 1 (1962) 284; Erratum Phys. Lett. 2 (1962) 164

1962AR02 B.H. Armitage and R.E. Meads, Nucl. Phys. 33 (1962) 494

1962DE10 G. Dearnaley, D.S. Gemmell and S.S. Hanna, Nucl. Phys. 36 (1962) 71

1962EA02 L.G. Earwaker, J.G. Jenkin and E.W. Titterton, Nature 195 (1962) 271

1962EL01 A.J. Elwyn and R.O. Lane, Nucl. Phys. 31 (1962) 78

1962EL06 R.J. Ellison and B. Dickinson, Nucl. Phys. 35 (1962) 606

1962KJ05 J. Kjellman, A. Nilsson and S. Schwarz, Ark. Fys. 21 (1962) 169

1962MC12 R.L. McGrath, Phys. Rev. 127 (1962) 2138

1962ME1A Mehta and McIldowie, Nucl. Phys. 33 (1962) 502

1962MO1A Morpurgo, Nucl. Spectroscopy; Ed., Racah (1962)

1962PE10 A. Perrin, G. Surget, C. Thibault and F. Verriere, Compt. Rend. 255 (1962) 277

1962SC12 G. Schrank, E.K. Warburton and W.W. Daehnick, Phys. Rev. 127 (1962) 2159

1962SE1A Serov and Guzhovskii, Atomn. Energ. (USSR) 12 (1962) 5

1962WA21 W.K. Warburton and L.F. Chase, Jr., Nucl. Phys. 34 (1962) 517

1963BA52 F.J. Bartis, Phys. Rev. 132 (1963) 1763

1963BL20 H.R. Blieden, G.M. Temmer and K.L. Warsh, Nucl. Phys. 49 (1963) 209

1963DR02 J.E. Draper and C.O. Bostrom, Nucl. Phys. 47 (1963) 108

1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336

1963EA01 L.G. Earwaker, J.G. Jenkin and E.W. Titterton, Nucl. Phys. 42 (1963) 521

1963FU11 B. Furubayashi, E. Teranishi and M. Kageyama, J. Phys. Soc. Jpn. 18 (1963) 1235

1963HO19 H.D. Holmgren, L.M. Cameron and R.L. Johnston, Nucl. Phys. 48 (1963) 1

1963LE03 J.C. Legg, Phys. Rev. 129 (1963) 272

1963ME08 M.K. Mehta, W.E. Hunt, H.S. Plendl and R.H. Davis, Nucl. Phys. 48 (1963) 90

1963NO02 E. Norbeck, S.A. Coon, R.R. Carlson and E. Berkowitz, Phys. Rev. 130 (1963) 1971

1963WA17 E.K. Warburton, D.E. Alburger and D.H. Wilkinson, Phys. Rev. 132 (1963) 776

1963YA1B Yanabu et al., J. Phys. Soc. Jpn. 18 (1963) 747

1964AR04 B.H. Armitage and R.E. Meads, Phys. Lett. 8 (1964) 346

1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209

1964BO13 R.O. Bondelid and J.W. Butler, Nucl. Phys. 53 (1964) 618

1964GR40 L. Green and D.J. Donahue, Phys. Rev. 135 (1964) B701

1964HO02 W.F. Hornyak, C.A. Ludemann and M.L. Roush, Nucl. Phys. 50 (1964) 424

- 1964LA04 R.O. Lane, A.J. Elwyn and A. Langsdorf, Jr., Phys. Rev. 133 (1964) B409
- 1964SC12 U. Schmidt-Rohr, R. Stock and P. Turek, Nucl. Phys. 53 (1964) 77
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Ed., Suppl. Vol. 1 (1964)
- 1964WA05 E.K. Warburton, D.E. Alburger, A. Gallmann, P. Wagner and L.F. Chase, Jr., Phys. Rev. 133 (1964) B42
- 1965BA06 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klages, B. Kuhn and G. Stiller, Nucl. Phys. 64 (1965) 55
- 1965BA17 C.D. Bachelier, M. Bernas, I. Brissaud, P. Radvanyi and M. Roy, Phys. Lett. 16 (1965) 304
- 1965BU10 S.G. Buccino and A.B. Smith, Phys. Lett. 19 (1965) 234
- 1965CA1C Carlson et al., Bull. Amer. Phys. Soc. 10 (1965) 1194
- 1965CH12 G. Charpak, G. Gregoire, L. Massonnet, J. Saudinos, J. Favier, M. Gusakow and M. Jean, Phys. Lett. 16 (1965) 54
- 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707
- 1965DA05 S.E. Darden and A.J. Froelich, Phys. Rev. 140 (1965) B69
- 1965DO08 V.K. Dolinov, Y.V. Melikov and A.F. Tulinov, Pisma Zh. Eksp. Teor. Fiz. 2 (1965) 120; JETP Lett. 2 (1965) 74
- 1965FO06 P.D. Forsyth, I.J. Taylor and R.R. Perry, Nucl. Phys. 66 (1965) 376
- 1965FO07 P.D. Forsyth and R.R. Perry, Nucl. Phys. 67 (1965) 517
- 1965FO1C Foster and Glasgow, Nucl. Instrum. Meth. 36 (1965) 1
- 1965FR07 G. Fricke, G.R. Bishop and D.B. Isabelle, Nucl. Phys. 67 (1965) 187
- 1965GO05 S. Gorodetzky, A. Gallmann and R. Rebmeister, Phys. Rev. 137 (1965) B1466
- 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
- 1965HA19 E. Hayward and T. Stovall, Nucl. Phys. 69 (1965) 241
- 1965HU10 H. Hulubei, M. Ivascu, A. Berinde, I. Neamu, N. Scintei, I. Francz, N. Martalogu and E. Marincu, Rev. Roum. Phys. 10 (1965) 403
- 1965JA1A Jacmart, Cahiers Phys. (France) 19 (1965) 1
- 1965KA14 S. Kakigi, J. Phys. Soc. Jpn. 20 (1965) 1967
- 1965LE1B Lee and Siemssen, Bull. Amer. Phys. Soc. 10 (1965) 510
- 1965MA50 D. Maydan and D.G. Vass, Proc. Phys. Soc. 86 (1965) 817
- 1965ME13 Y.V. Melikov and A.F. Tulinov, Yad. Fiz. 2 (1965) 64; Sov. J. Nucl. Phys. 2 (1966) 44
- 1965MO27 S. Morita, T. Tohei, T. Nakagawa, T. Hasegawa, H. Ueno and C.-C. Hsu, Nucl. Phys. 66 (1965) 17

- 1965NE10 Y.A. Nemiřov, V.S. Sadkovskii, E.D. Teterin, A.E. Denisov and R.P. Kolalis, *Izv. Akad. Nauk SSSR Ser. Fiz.* 29 (1965) 1197; *Bull. Acad. Sci. USSR Phys.Ser.* 29 (1966) 1202
- 1965NE1B Neudachin and Smirnov, *At. Energy Rev.* 3 (1965) 157
- 1965PA02 P. Paul, S.L. Blatt and D. Kohler, *Phys. Rev.* 137 (1965) B493
- 1965PA03 P. Paul, S.L. Blatt and D. Kohler, *Phys. Rev.* 137 (1965) B499
- 1965PE17 R.H. Pehl, J. Cerny, E. Rivet and B.G. Harvey, *Phys. Rev.* 140 (1965) B605
- 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, *Ann. Phys.* 34 (1965) 96
- 1965SH1A Shumilov, Klyucharev and Rutkevich, *Yad. Fiz.* 2 (1965) 1030
- 1965SI1B Singh and Gemmell, *Bull. Amer. Phys. Soc.* 10 (1965) 538
- 1965ST22 P.H. Stelson and L. Grodzins, *Nucl. Data A1* (1965) 21
- 1965VA1G Van Praet and Kossanyi-Demay, *Nuovo Cim.* 39 (1965) 388
- 1965ZO1A Zobel, Maienschein and Scroggs, ORNL 3506 (1965)
- 1966AG1A Agee and Rosen, LA-3538-MS (1966)
- 1966AL06 D.E. Alburger, P.D. Parker, D.J. Bredin, D.H. Wilkinson, P.F. Donovan, A. Gallmann, R.E. Pixley, L.F. Chase, Jr. and R.E. McDonald, *Phys. Rev.* 143 (1966) 692
- 1966BA1A Bahcall, *Nucl. Phys.* 75 (1966) 10
- 1966BA32 F. Baldeweg, V. Bredel, H. Guratzsch, R. Klables, B. Kuhn and G. Stiller, *Nucl. Phys.* 85 (1966) 171
- 1966BA59 D.J. Baugh, J.A.R. Griffith and S. Roman, *Nucl. Phys.* 83 (1966) 481
- 1966BR1G Browne, F. S. U. Isobaric Spin Conf. (1966) 136
- 1966BR1N Brahmavar, *Nucl. Phys.* 75 (1966) 79
- 1966CA01 T.A. Cahill, J.R. Richardson and R.P. Haddock, *Phys. Rev.* 144 (1966) 932
- 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, *Phys. Rev.* 152 (1966) 950
- 1966CH1C Chien and Smith, *Nucl. Sci. Eng.* 26 (1966) 500
- 1966DA1B Darden, *Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965* (1966) 433
- 1966DI04 G.U. Din and J.L. Weil, *Nucl. Phys.* 86 (1966) 509
- 1966DR1C Drachman, *Phys. Rev. Lett.* 17 (1966) 1017
- 1966ED1A J.A. Edgington and B. Rose, *Nucl. Phys.* 89 (1966) 523
- 1966FE04 F.F. Felber, Jr., D.R. Farmelo and V.C. Van Sickle, *Nucl. Sci. Eng.* 25 (1966) 1
- 1966FI01 T.R. Fisher, S.S. Hanna and P. Paul, *Phys. Rev. Lett.* 16 (1966) 850
- 1966FO05 P.D. Forsyth, H.T. Tu and W.F. Hornyak, *Nucl. Phys.* 82 (1966) 33

- 1966FO08 P.D. Forsyth, A.R. Knudson and F.C. Young, Nucl. Phys. 85 (1966) 153
- 1966FR09 J.M. Freeman, J.G. Jenkin and G. Murray, Phys. Lett. 22 (1966) 177
- 1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197
- 1966GE12 L.B. Geesaman, H.B. Knowles and J.B. Mead, Nucl. Phys. 88 (1966) 177
- 1966GO1D Gotz, Mittenbacher, Prager and Muhle, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122 (1966) 201
- 1966GR18 R.C. Greenwood, Phys. Lett. 23 (1966) 482
- 1966GR1G T.A. Griffy, Phys. Lett. 21 (1966) 693
- 1966HA09 E.T. Hazzard, F. Ajzenberg-Selove and P.V. Hewka, Nucl. Phys. 75 (1966) 592; Erratum Nucl. Phys. 89 (1966) 706
- 1966HA18 E.C. Halbert, Y.E. Kim and T.T.S. Kuo, Phys. Lett. 20 (1966) 657
- 1966HO1D Hodgson, Proc. Conf. Nucl. Reactions, Rossendorf, 1966; Ed., J. Schintlmeister; ZFK-122 (1966) 71
- 1966IS1A Isabelle, J. Phys. (Paris) 27 (1966) C1-17
- 1966JA08 O.N. Jarvis, Nucl. Phys. 79 (1966) 305
- 1966JA09 D.F. Jackson, Nuovo Cim. B41 (1966) 86
- 1966JA1A Jacob and Maris, Rev. Mod. Phys. 38 (1966) 121
- 1966JA1F O.N. Jarvis, B. Rose and A.E. Taylor, Nucl. Phys. 79 (1966) 275
- 1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KI07 F.G. Kingston, R.J. Griffiths, A.R. Johnston, W.R. Gibson and E.A. McClatchie, Phys. Lett. 22 (1966) 458
- 1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932
- 1966KO08 P. Kossanyi-Demay and G.J. Vanpraet, Nucl. Phys. 81 (1966) 529
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LH1A Lhuillier and Benoist-Gueutal, J. Phys. (Paris) 27 (1966) C1-78
- 1966LO15 J.A. Lonergan and D.J. Donahue, Nucl. Phys. 88 (1966) 465
- 1966MA1P MacFarlane, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 411
- 1966MA36 N. Mangelson, F. Ajzenberg-Selove, M. Reed and C.C. Lu, Nucl. Phys. 88 (1966) 137
- 1966ME09 L. Meyer-Schutzmeister, D. Von Ehrenstein and R.G. Allas, Phys. Rev. 147 (1966) 743
- 1966ME1E Melikov, Vestnik Mosk. Univ. Fiz. Astron. 6 (1966) 102

- 1966MI1E Miller, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 410
- 1966PA1J Parish, Rawlins and Shin, Bull. Amer. Phys. Soc. 11 (1966) 27
- 1966PE1E Peterson, Ziegler and Cole, Bull. Amer. Phys. Soc. 11 (1966) 120
- 1966PO09 A.M. Poskanzer, S.W. Cospers, E.K. Hyde and J. Cerny, Phys. Rev. Lett. 17 (1966) 1271
- 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246
- 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 253
- 1966RO1E Rotter and Zhusupov, Ann. Phys. 17 (1966) 57
- 1966SA20 O.A. Salnikov, N.I. Fetisov, G.N. Lovchikova, G.V. Kotelnikova, V.B. Anufrienko and B.V. Devkin, Yad. Fiz. 4 (1966) 1154; Sov. J. Nucl. Phys. 4 (1967) 831
- 1966SC16 G. Schirmer, H. Pose and H. Hansgen, Nucl. Phys. 84 (1966) 201
- 1966SE02 R.E. Segel and R.H. Siemssen, Phys. Lett. 20 (1966) 295
- 1966SE03 R.E. Segel, P.P. Singh, S.S. Hanna and M.A. Grace, Phys. Rev. 145 (1966) 736
- 1966SE1E Seth and Tabony, Ann. Phys. 40 (1966) 183
- 1966SI02 R.H. Siemssen, G.C. Morrison, B. Zeidman and H. Fuchs, Phys. Rev. Lett. 16 (1966) 1050
- 1966SP02 E. Spamer, Z. Phys. 191 (1966) 24
- 1966ST12 T. Stovall, J. Goldemberg and D.B. Isabelle, Nucl. Phys. 86 (1966) 225
- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692
- 1966VE03 V.V. Verbinski and M. Sibtain Bokhari, Phys. Rev. 143 (1966) 688
- 1966WA10 E.K. Warburton, J.W. Olness, K.W. Jones, C. Chasman, R.A. Ristinen and D.H. Wilkinson, Phys. Rev. 148 (1966) 1072
- 1966WA1C Warburton and Alburger, Nucl. Spin-Parity Assignments; Ed., Gove (1966) 114
- 1966WA1G Wang, Ching and Tsiao, Acta Phys. Sinica (China) 22 (1966) 611
- 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967BA78 G. Backenstoss, S. Charalambus, H. Daniel, H. Koch, G. Poelz, H. Schmitt and L. Tauscher, Phys. Lett. B25 (1967) 547
- 1967BE1F Beery, Harper, Stovall and Rosen, LA-3788 (1967)
- 1967BE1T Bethge, Gunther, Schubert and Speth, Z. Phys. 202 (1967) 70

- 1967BE27 W. Benenson, G.M. Crawley, J.D. Dreisbach and W.P. Johnson, Nucl. Phys. A97 (1967) 510
- 1967BI1E Bing, Seltz, Gerardin and Magnac-Valette, Private Communication (1967)
- 1967BI1F Birnbaum and Bromley, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 157
- 1967BR1B Browne and O'Donnell, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 508
- 1967BU1D Bukhvostov and Popov, Yad. Fiz. 6 (1967) 1241
- 1967CA17 P. Camiz, Nuovo Cim. B51 (1967) 190
- 1967CA1K Carlson et al., Bull. Amer. Phys. Soc. 12 (1967) 483
- 1967CO1J Coop, Poate and Titterton, Aust. J. Phys. 20 (1967) 609
- 1967CO1P Coop, Poate and Titterton, Nature 213 (1967) 587
- 1967CO23 S.W. Cospers, H. Brunnader, J. Cerny and R.L. McGrath, Phys. Lett. B25 (1967) 324
- 1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1
- 1967CO36 S.W. Cospers, J. Cerny and R.C. Gatti, Phys. Rev. 154 (1967) 1193
- 1967CR04 M.A. Crosby and J.C. Legg, Nucl. Phys. A95 (1967) 639
- 1967DE1K Dehnhard and Siemssen, Bull. Amer. Phys. Soc. 12 (1967) 17
- 1967DE1M Dehnhard, Morrison and Vager, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 112
- 1967DI01 G.U. Din, M.A. Nagarajan and R. Pollard, Nucl. Phys. A93 (1967) 190
- 1967EV1C J.A. Evans and R.P.J. Perazzo, Nucl. Phys. A103 (1967) 225
- 1967FA03 M.A. Farouk, M.N.H. Comsan, A.Z. El-Behay, A.A. El-Kamhawy, A.N. Lvov and A.I. Popov, Z. Phys. 201 (1967) 52
- 1967FI01 A.A. Fife, G.C. Neilson and W.K. Dawson, Nucl. Phys. A91 (1967) 164
- 1967FI07 W. Fitz, R. Jahr and R. Santo, Nucl. Phys. A101 (1967) 449
- 1967FU04 H. Fuchs and R. Santo, Phys. Lett. B24 (1967) 234
- 1967GA06 D.S. Gale and J.M. Blair, Phys. Rev. 161 (1967) 1061
- 1967GL03 D.W. Glasgow and D.G. Foster, Jr., Nucl. Phys. A99 (1967) 170
- 1967GO36 G.V. Gorlov, N.S. Lebedeva and V.M. Morozov, Yad. Fiz. 6 (1967) 910; Sov. J. Nucl. Phys. 6 (1968) 663
- 1967HO1F P.E. Hodgson, Nucl. Phys. A103 (1967) 127
- 1967HS1A L.S. Hsu, Phys. Lett. B25 (1967) 588
- 1967IV01 M. Ivanovich, H. Cords and G.U. Din, Nucl. Phys. A97 (1967) 177
- 1967JA1E B.K. Jain and D.F. Jackson, Nucl. Phys. A99 (1967) 113
- 1967KA1A Kabachnik and Grishanova, Sov. J. Nucl. Phys. 4 (1967) 583

- 1967KI03 K.G. Kibler, Phys. Rev. 155 (1967) 1110
- 1967KO1B Kolybasov and Smorodinskaya, Yad. Fiz. 5 (1967) 777
- 1967KO1E Kostoff, Jahns, Terrell and Bernstein, Bull. Amer. Phys. Soc. 12 (1967) 893
- 1967KO1F Kohman and Bender, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Chen (1967) 169
- 1967KU1E D. Kurath and R.D. Lawson, Phys. Rev. 161 (1967) 915
- 1967LE1E Lehmann, Proc. Int. Conf. Electromag. Inter. Dubna, 1967, Vol. 3 (1967) 66
- 1967LE1F Lee, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 31
- 1967LO1B Loiseaux, Maison and Langevin, J. Phys. (France) 28 (1967) 11
- 1967MC1E McGill et al., Bull. Amer. Phys. Soc. 12 (1967) 1144
- 1967ME08 V. Meyer, R.E. Pixley and P. Truol, Nucl. Phys. A101 (1967) 321
- 1967MI1A Mitler, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 59
- 1967MU1A G. Murray, Phys. Lett. B24 (1967) 268
- 1967NO1A J.V. Noble, Phys. Rev. 162 (1967) 934
- 1967PA01 P. Paul, T.R. Fisher and S.S. Hanna, Phys. Lett. B24 (1967) 51
- 1967PA03 G. Paic, D. Rendic and P. Tomas, Nucl. Phys. A96 (1967) 476
- 1967PA1H Patterson et al., Proc. Phys. Soc. 90 (1967) 577
- 1967PI1B Pinkston, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 419
- 1967PU04 J.D. Purvis, F. Ajzenberg-Selove and L.M. Polsky, Phys. Rev. 162 (1967) 1005
- 1967RA24 N.C. Rasmussen, V.J. Orphan and Y. Hukai, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 278
- 1967RO07 D.C. Robinson, Nucl. Phys. A95 (1967) 663
- 1967RO1E Robrock and Hecker, Nukleonik 10 (1967) 257
- 1967SA1C Sannes and Stairs, Can. J. Phys. 45 (1967) 1497
- 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
- 1967SH05 Y.Y. Sharon, Nucl. Phys. A99 (1967) 321
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SH1B Shapiro and Silberberg, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 37
- 1967SI1A R.H. Siemssen and D. Dehnhard, Phys. Rev. Lett. 19 (1967) 377
- 1967SP09 A. Sperduto, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 657

- 1967TA1A Tamura, Phys. Rev. Lett. 19 (1967) 321
- 1967TA1C Takimoto, Mem. Coll. Sci. Univ. Kyoto A31 (1967) 267
- 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 43
- 1967WA1H J.W. Wachter, W.R. Burrus and W.A. Gibson, Phys. Rev. 161 (1967) 971
- 1967WA31 D.L. Watson, J. Lowe, J.C. Dore, R.M. Craig and D.J. Baugh, Nucl. Phys. A92 (1967) 193
- 1967YO1C Young, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 109
- 1968AL1E Alder, Vaucher and Joseph, Helv. Phys. Acta 41 (1968) 433
- 1968BA2G Backenstoss et al., Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968BE1E Bertrand, Grenier and Pernet, Comm. A L'energie Atomique Rept. CEA 3504 (1968)
- 1968BE1N Bertrand, Grenier and Pernet, Comm. A L'energie Atomique Rept. CEA 3575 (1968)
- 1968BI1C Bizzeti, Suppl. Nuovo Cim. 6 (1968) 664
- 1968BL1F Blue, Prior, Keyser and Ramirez, Bull. Amer. Phys. Soc. 13 (1968) 1449
- 1968BO1U Bohr, Damgard and Mottelson, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 178
- 1968BO32 N.E. Booth, A. Beretvas, R.E.P. Davis, C. Dolnick, R.E. Hill, M. Raymond and D. Sherden, Nucl. Phys. A119 (1968) 233
- 1968BR23 H. Brunnader, J.C. Hardy and J. Cerny, Phys. Rev. 174 (1968) 1247
- 1968CE01 J. Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968DA20 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Yad. Fiz. 7 (1968) 758; Sov. J. Nucl. Phys. 7 (1968) 463
- 1968DO01 D.J. Donahue, M.J. Wozniak, R.L. Hershberger, J.E. Cummings and J.A. Lonergan, Phys. Rev. 165 (1968) 1071
- 1968EA1B Eaton and Walker, Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 169
- 1968EN1A J. Engler, K. Horn, J. Konig, F. Monnig, P. Schludecker, H. Schopper, P. Sievers, H. Ullrich and K. Runge, Phys. Lett. B28 (1968) 64
- 1968FA02 A. Faessler, P.U. Sauer and M.M. Stingl, Phys. Rev. 167 (1968) 1013
- 1968FA1B Faessler, Sauer and Stingl, Z. Phys. 212 (1968) 1
- 1968FI09 T.R. Fisher, S.S. Hanna, D.C. Healey and P. Paul, Phys. Rev. 176 (1968) 1130
- 1968FOZY D.T. Fowler, Neutron Cross Sections Tech., NBS Special Pub. 299, Vol. I (1968) 1
- 1968FR10 A.J. Froelich and S.E. Darden, Nucl. Phys. A119 (1968) 97

- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GO1J Goldemberg, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 379
- 1968GR1F Green, Nucl. Sci. Eng. 32 (1968) 30
- 1968HA1C Hayakawa, Suppl. Prog. Theor. Phys. (1968) 156
- 1968HA1K Harrison, COO-1265-57 (1968)
- 1968HE1G E.M. Henley, Phys. Lett. B28 (1968) 1
- 1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, 1967 (1968) 41
- 1968HU17 I. Hunyadi and I. Angeli, Nucl. Phys. A119 (1968) 525
- 1968IN02 A. Ingemarsson, A. Johansson and G. Tibell, Phys. Lett. B27 (1968) 439
- 1968JA09 J. Janecke, T.F. Yang, R.M. Polichar and W.S. Gray, Phys. Rev. 175 (1968) 1301
- 1968JA1D Jackson, Adv. Phys. 17 (1968) 481
- 1968JA1G Jackson, in Tokyo (1968) 105
- 1968JO1C Joshi, Nucl. Phys. Solid State Phys. Symp. (1968)
- 1968JO1E K.P. Joshi, G.K. Mehta and Y.R. Waghmare, Phys. Rev. 176 (1968) 1154
- 1968JO1F Johnson et al., Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 851
- 1968KL06 R. Klabes, F. Baldeweg and G. Stiller, Nucl. Phys. A121 (1968) 113; Erratum Nucl. Phys. A127 (1969) 693
- 1968KU04 L.A. Kull and E. Kashy, Phys. Rev. 167 (1968) 963
- 1968KU1D Kurath, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 393
- 1968LA1F Lancman et al., Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968LI03 S.C. Ling, A.M. Young and S.L. Blatt, Nucl. Phys. A108 (1968) 221
- 1968MA18 J.R. MacDonald, D.F.H. Start and D.H. Wilkinson, Nucl. Phys. A113 (1968) 129
- 1968MA1W Mancuso, Knudson and Wolicki, Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968ME1E Meyer, Audebert, Elbaz and Lafoucriere, Compt. Rend. B266 (1968) 969
- 1968MI10 D. Miljanic, M. Furic and V. Valkovic, Nucl. Phys. A119 (1968) 379
- 1968MO1F Morita, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 419
- 1968NE1A NeuV.G. Neudatchin, N.S. Zelenskaya, E.Zh. Magzumov and E.A. Romanovsky, Phys. Lett. B27 (1968) 490

- 1968NE1C Nemirovskii, Sov. J. Nucl. Phys. 6 (1968) 29
- 1968NO1A M.E. Nordberg, K.F. Kinsey and R.L. Burman, Phys. Rev. 165 (1968) 1096
- 1968NO1C J.V. Noble, Phys. Rev. 173 (1968) 1034
- 1968OK06 Y. Okuma, J. Phys. Soc. Jpn. 25 (1968) 1
- 1968OK1B Okuma, J. Phys. Soc. Jpn. 24 (1968) 677
- 1968PA1J Palevsky, Proc. of Symp. on Use of Nimrod, 1968, RHEL-R166 (1968) 19
- 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
- 1968RI1H A. Richter and R.A. Chatwin, Phys. Lett. B27 (1968) 181
- 1968RO1H Roturier, NP 17794 (1968)
- 1968SA1H Sakamoto, Nuovo Cim. B56 (1968) 359
- 1968SC18 H.F. Schaefer III, R.A. Klemm and F.E. Harris, Phys. Rev. 176 (1968) 49
- 1968SC1E Schwarzschild and Warburton, Ann. Rev. Nucl. Sci. 18 (1968) 265
- 1968SH21 A.K. Shardanov and B.A. Yurev, Yad. Fiz. 8 (1968) 424; Sov. J. Nucl. Phys. 8 (1969) 245
- 1968SI07 V.S. Siksinev, A.I. Baz, V.N. Domoratskii, A.N. Voronin, V.S. Marchenko, G.E. Kosinskii, A.N. Kudryavtsev and G.D. Batyshchev, Yad. Fiz. 7 (1968) 950; Sov. J. Nucl. Phys. 7 (1968) 571
- 1968SQ01 G.T.A. Squier, E.A. McClatchie, A.R. Johnston, R.J. Batten, J.B.A. England and F.G. Kingston, Nucl. Phys. A119 (1968) 369
- 1968TA1A T. Tamura, Phys. Rev. 165 (1968) 1123
- 1968TA1C Tanner, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 91
- 1968TA1P Taketani, Yamaguchi, Kokame and Muto, in Tokyo (1968) 270
- 1968TI1A Tibell, Proc. Symp. on Use of Nimrod for Nucl. Struct. Phys., Rutherford High Energy Lab. (1968) 127
- 1968VO1A W. Von Oertzen, H.H. Gutbrod, M. Muller, U. Voos and R. Bock, Phys. Lett. B26 (1968) 291
- 1968WA15 E.K. Warburton, J.W. Olness, S.D. Bloom and A.R. Poletti, Phys. Rev. 171 (1968) 1178
- 1968WA1H Watson, Thesis, Indiana Univ. (1968)
- 1968WI1B Wilkinson, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 469
- 1968YI01 F. Yiou, Ann. Phys. (Paris) 3 (1968) 169
- 1968ZE1B Zelenskaya et al., Sov. J. Nucl. Phys. 6 (1968) 47

- 1969AL17 D.E. Alburger, E.K. Warburton, A. Gallmann and D.H. Wilkinson, Phys. Rev. 185 (1969) 1242
- 1969AN1E I. Angeli, J. Csikai and I. Hunyadi, Phys. Lett. B29 (1969) 36
- 1969AN27 G.B. Andreev, I.D. Lopatko, V.I. Borovlev, V.A. Lutsik, R.P. Slabospitskii and A.Y. Taranov, Yad. Fiz. 10 (1969) 902; Sov. J. Nucl. Phys. 10 (1970) 521
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969AR1B Armstrong, Beery, Keaton and Veaser, LA-4177 (1969)
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BA1F V.V. Balashov, N.M. Kabachnik and V.I. Markov, Nucl. Phys. A129 (1969) 369
- 1969BA1J Balashov, Bochum Conf. STI/PUB/232 IAEA (1969) 59
- 1969BA1N Bahcall and Fowler, Astrophys. J. 157 (1969) 659
- 1969BA55 A.I. Baz, V.F. Demin and M.V. Zhukov, Yad. Fiz. 9 (1969) 1184; Sov. J. Nucl. Phys. 9 (1969) 693
- 1969BE1Q Bernstein, Bonner and Michelman, Bull. Amer. Phys. Soc. 14 (1969) 488
- 1969BL1D Blin-Stoyle, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 115
- 1969BL1E R.J. Blin-Stoyle, Phys. Lett. B29 (1969) 12
- 1969BR13 D.R. Brown, S.M. Ferguson and D.H. Wilkinson, Nucl. Phys. A135 (1969) 159
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 242
- 1969BU1C Burman and Nordberg, Bull. Amer. Phys. Soc. 14 (1969) 537
- 1969CA1A R.R. Carlson and H.W. Wyborny, Phys. Rev. 178 (1969) 1529
- 1969CA1F Calvert and Conde, Contrib., Montreal (1969) 282
- 1969CH19 D.L. Cheshire and S.E. Sobottka, Phys. Lett. B30 (1969) 244
- 1969CH1A B.T. Chertok, Phys. Rev. 187 (1969) 1340
- 1969CH1C D.T. Chivers, E.M. Rimmer, B.W. Allardyce, R.C. Witcomb, J.J. Domingo and N.W. Tanner, Nucl. Phys. A126 (1969) 129
- 1969CO02 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A127 (1969) 95
- 1969CO1G Cooper, Roush, Connors and Hornyak, Bull. Amer. Phys. Soc. 14 (1969) 489
- 1969CO1H Connors et al., Contrib., Montreal (1969) 327
- 1969CO1J P. Connors et al., Contrib., Montreal (1969) 328
- 1969CU08 J.R. Curry, W.R. Coker and P.J. Riley, Phys. Rev. 185 (1969) 1416
- 1969DA1D C.N. Davids, H. Laumer and S.M. Austin, Phys. Rev. Lett. 22 (1969) 1388

- 1969DE10 D. Dehnhard, N. Williams and J.L. Yntema, Phys. Rev. 180 (1969) 967; Erratum Phys. Rev. C1 (1970) 366
- 1969DE1N de Swiniarski et al., Bull. Amer. Phys. Soc. 14 (1969) 851
- 1969DE29 A.E. Denisov, R.P. Kolalis, Y.A. Nemilov, V.S. Sadkovskii and E.U. Teterin, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 1376; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1271
- 1969ED02 R.D. Edge and H.H. Knox, Phys. Rev. 184 (1969) 1034
- 1969FI1C Fisher, Hasinoff, Riess and Hanna, Bull. Amer. Phys. Soc. 14 (1969) 1167
- 1969FR02 J.M. Freeman, J.G. Jenkin and G. Murray, Nucl. Phys. A124 (1969) 393
- 1969FR09 J.M. Freeman, D.C. Robinson and G.L. Wick, Phys. Lett. B30 (1969) 240
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA01 A. Gallmann, F. Hibou and P. Fintz, Nucl. Phys. A123 (1969) 27
- 1969GA06 R.B. Galloway and D.G. Vass, Nucl. Phys. A127 (1969) 538
- 1969GA11 P. Gaillard, R. Bouche, L. Feuveys, M. Gaillard, A. Guichard, M. Gusakov, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
- 1969GA18 J. Galin, B. Gatty, M. Lefort, J. Peter, X. Tarrago and R. Basile, Phys. Rev. 182 (1969) 1267
- 1969GA1G Garvey, Ann. Rev. Nucl. Sci. 19 (1969) 433
- 1969GO1C Golovanova and Zelenskaya, Sov. J. Nucl. Phys. 8 (1969) 158
- 1969GO1H Goel, Nature 223 (1969) 1263
- 1969GU1K I.A. Gubkin, Nucl. Phys. A126 (1969) 355
- 1969HA1F Hanna, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 591
- 1969HA1R Haduat, Lachkar, Patin and Sigaud, Neutron Capture Gamma-Ray Spectroscopy, Studsvik Symp., STI/PUB/235 IAEA (1969) 119
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
- 1969HO1G Hopkins and Drake, Nucl. Sci. Eng. 36 (1969) 275
- 1969HO1H Holman et al., Bull. Amer. Phys. Soc. 14 (1969) 1168
- 1969HO1M T.H. Ho and A.B. Volkov, Phys. Lett. B30 (1969) 303
- 1969HO45 M. Holmberg and J. Hansen, Nucl. Phys. A129 (1969) 305
- 1969HU1F Hutzelmeyer and Hackenbroich, Bochum Conf. STI/PUB/232 IAEA (1969) 286
- 1969IN1C Ingemarsson and Sundqvist, Ark. Fys. 39 (1969) 429
- 1969JA1N Jackson, Thesis, Univ. of Toronto (1969)
- 1969KA1B R.W. Kavanagh, Nucl. Phys. A129 (1969) 172

1969KAZZ R.S. Kaushal, Nucl. Phys. Solid State Phys. Symp., Roorkee, India (1969)
 1969KO1J V.M. Kolybasov and N.Ya. Smorodinskaya, Nucl. Phys. A136 (1969) 165
 1969KO1P K. Kolltveit and K. Nagatani, Nucl. Phys. A124 (1969) 287
 1969KR03 E. Krubasik, H. Voit, E. Blatt, H.-D. Helb and G. Ischenko, Z. Phys. 219 (1969) 185
 1969LE1B Leonardi, Cargese Lect. in Phys.; Ed., M. Jean, Vol. 3 (1969) 97
 1969LO1E Loyd and Haeberli, Bull. Amer. Phys. Soc. 14 (1969) 37
 1969MA1R Mairle et al., Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 291
 1969MA39 S.C. Mathur, P.S. Buchanan and I.L. Morgan, Phys. Rev. 186 (1969) 1038
 1969MC1A McGill, UCLA P-83 (1969)
 1969MO1E L. Moyer and D.S. Koltun, Phys. Rev. 182 (1969) 999
 1969MO29 T. Mo and W.F. Hornyak, Phys. Rev. 187 (1969) 1220
 1969MU10 K.M. Murray, Phys. Rev. Lett. 23 (1969) 1461
 1969NA04 K.S. Nam and G.M. Osetinskii, Yad. Fiz. 9 (1969) 487; Sov. J. Nucl. Phys. 9 (1969) 279
 1969NAZY K.L. Narayana, Nucl. Phys. Solid State Phys. Symp., Roorkee, India (1969) 356
 1969NE08 V.G. Neudachin, E.Z. Magzumov and E.A. Romanovskii, Yad. Fiz. 9 (1969) 94; Sov. J. Nucl. Phys. 9 (1969) 56
 1969NO1B J.V. Noble, Phys. Rev. Lett. 22 (1969) 473, Erratum Phys. Rev. Lett. 22 (1969) 1028
 1969NO1C Noble, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 233
 1969NU1A Nusslin, Werner and Zimmerer, Bochum Conf. STI/PUB/232 IAEA (1969) 229
 1969OR01 H. Orihara, T. Nakagawa, H. Ueno, T. Tohei, T. Yamaya, Y. Nakagome, M. Baba and S. Morita, Nucl. Phys. A139 (1969) 226
 1969PA09 R.A. Paddock, S.M. Austin, W. Benenson, I.D. Proctor and F.S. Amant, Phys. Rev. 182 (1969) 1104
 1969PR17 G.O. Prokopets and V.I. Strizhak, Ukr. Fiz. Zh. USSR 14 (1969) 687
 1969RO12 M.L. Roush, F.C. Young, P.D. Forsyth and W.F. Hornyak, Nucl. Phys. A128 (1969) 401
 1969RO14 L.-H. Rosier, Y. Deschamps and R. Tamisier, Compt. Rend. B268 (1969) 1149
 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
 1969SA04 J. Sanada, Y.C. Liu, Y. Sugiyama and O. Mikoshiba, J. Phys. Soc. Jpn. 26 (1969) 853
 1969SA1A Sauer, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 717
 1969SC05 W. Scobel, Z. Naturforsch. A24 (1969) 289
 1969SC1F Schiffer, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 665

1969SI12 J.D. Silverstein and G.H. Herling, Phys. Rev. 181 (1969) 1512
 1969SI20 V.S. Siskin, V.N. Domoratskii, L.V. Donetskov, A.N. Voronin, M.M. Novikov, A.N. Kudryavtsev, G.D. Batyshev and V.F. Tseshkovskaya, Yad. Fiz. 10 (1969) 1130; Sov. J. Nucl. Phys. 10 (1970) 643
 1969SM03 H.V. Smith, Jr. and H.T. Richards, Phys. Rev. Lett. 23 (1969) 1409
 1969SO08 B. Sorensen, Nucl. Phys. A134 (1969) 1
 1969SO1E Sorensen, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 239
 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
 1969TI02 G. Tibell, Phys. Lett. B28 (1969) 638
 1969TO1A I.S. Towner, Nucl. Phys. A126 (1969) 97
 1969TR1B P. Truoel and W. Bierter, Phys. Lett. B29 (1969) 21
 1969TU1A Tu and Hornyak, Bull. Amer. Phys. Soc. 14 (1969) 489
 1969UB1B H. Uberall and P. Uginčius, Phys. Rev. 178 (1969) 1565
 1969VA1C S. Varma and P. Goldhammer, Nucl. Phys. A125 (1969) 193
 1969VE02 V.V. Verbinski and W.R. Burrus, Phys. Rev. 177 (1969) 1671
 1969VI02 D. Vinciguerra and T. Stovall, Nucl. Phys. A132 (1969) 410
 1969VO10 U.C. Voos, W. von Oertzen and R. Bock, Nucl. Phys. A135 (1969) 207
 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
 1969WA23 B.A. Watson, R.E. Segel, J.J. Kroepfl and P.P. Singh, Phys. Rev. 187 (1969) 1351
 1969WE10 K.J. Wetzel, Phys. Rev. 186 (1969) 1292
 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
 1969YI1A Yiou, Seide and Bernas, J. Geophys. Res. 74 (1969) 2447
 1969YO01 F.C. Young and W.F. Hornyak, Nucl. Phys. A124 (1969) 469
 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
 1970AN1F Angeli, Csikai and Hunyadi, Acta Phys. Acad. Sci. Hung. 28 (1970) 629
 1970AN1J Andrade, Rendic, Otte and Phillips, Bull. Amer. Phys. Soc. 15 (1970) 1653
 1970AU02 D.L. Auton, Nucl. Phys. A157 (1970) 305
 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
 1970BA1L Baumann, Huck and Walter, Bull. Amer. Phys. Soc. 15 (1970) 567
 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119
 1970BA1P Barenholtz, Thesis, Univ. of Illinois (1970)
 1970BI1B Bingham, Thesis, Florida State Univ. (1970)

- 1970BL1E Blair, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 1
- 1970BO17 F. Boreli, R. Little and A. Rego, Fizika 2 (1970) 19
- 1970BO1W Boyd et al., Bull. Amer. Phys. Soc. 15 (1970) 1653
- 1970BO1Y Botemps, NP 18438 (1970)
- 1970BO35 M. Bouten, M.-C. Bouten, H. Depuydt and L. Schotsmans, Nucl. Phys. A158 (1970) 561
- 1970CA28 J. Catala, A. Garcia, V. Martinez and F. Senent, An. Fis. 66 (1970) 297
- 1970CH1F Cheshire and Sobottka, Nucl. Phys. A146 (1970) 129
- 1970CH1Q Chen, Harrison and Tombrello, Bull. Amer. Phys. Soc. 15 (1970) 1598
- 1970CL01 A.S. Clough, C.J. Batty, B.E. Bonner and L.E. Williams, Nucl. Phys. A143 (1970) 385
- 1970CL10 N.M. Clarke, E.J. Burge, D.A. Smith and J.C. Dore, Nucl. Phys. A157 (1970) 145
- 1970CO06 M.D. Cooper, A.S. Figuera, G. Pfeufer and G. Randers-Pehrson, Nucl. Phys. A142 (1970) 445
- 1970CO12 J.A. Cookson and J.G. Locke, Nucl. Phys. A146 (1970) 417
- 1970CO1H Cohen and Kurath, Nucl. Phys. A141 (1970) 145
- 1970DA21 A. D'Andrea and M. Scalia, Nuovo Cim. A69 (1970) 702
- 1970DA26 V.K. Daruga and E.S. Matusевич, Atomn. Energ. (USSR) 29 (1970) 456; Sov. J. At. Energ. 29 (1970) 1233
- 1970DI1F Dixon, Thesis, Univ. of South Carolina (1970)
- 1970DU07 J.L. Duggan, J.Y. Park, S.D. Danielopoulos, P.D. Miller, J. Lin, M.M. Duncan and R.L. Dangle, Nucl. Phys. A151 (1970) 107
- 1970FA15 D. Favart, F. Brouillard, L. Grenacs, P. Igo-Kemenes, P. Lipnik and P.C. Macq, Phys. Rev. Lett. 25 (1970) 1348
- 1970FI07 D. Fick, R. Kankowsky, K. Kilian and E. Salzbom, Phys. Rev. Lett. 24 (1970) 1503
- 1970FI1B Fisher, Thesis, Stanford Univ. (1970)
- 1970FO1B Fonda, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 283
- 1970FR1C Frahn, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 297
- 1970GA01 A. Gallmann, G. Frick, B. Heusch, F. Haas and E.K. Warburton, Nucl. Phys. A143 (1970) 326
- 1970GA1A Garber et al., BNL 400, 3rd Edition, Vol. 1 (1970)
- 1970HI08 K.D. Hildenbrand, H.H. Gutbrod, W. von Oertzen and R. Bock, Nucl. Phys. A157 (1970) 297

- 1970IS1A Ischenko, Reichardt, Voit and Held, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 105
- 1970KA32 I. Kakkar, G.K. Mehta and Y.R. Waghmare, Nucl. Phys. A156 (1970) 199
- 1970KO25 V.I. Komarov, G.E. Kosarev and O.V. Savchenko, Yad. Fiz. 11 (1970) 711; Sov. J. Nucl. Phys. 11 (1970) 399
- 1970KU1D Kull, Thesis, Michigan State Univ. (1970)
- 1970LA13 J. Langworthy and H. Uberall, Phys. Rev. C2 (1970) 911
- 1970LI1K Lim, Nucl. Phys. A148 (1970) 299
- 1970LO03 D.H. Loyd and W. Haeberli, Nucl. Phys. A148 (1970) 236
- 1970MC1G McGrath et al., Bull. Amer. Phys. Soc. 15 (1970) 629
- 1970MI1G Miller, Morrison and Westley, Bull. Amer. Phys. Soc. 15 (1970) 1315
- 1970NA06 J. Nalda, T. Barriuso and J. Casanova, Nucl. Phys. A149 (1970) 350
- 1970NE03 D.O. Nellis, W.E. Tucker and I.L. Morgan, Phys. Rev. C1 (1970) 847
- 1970NE05 R.K. Nesbet, Phys. Rev. Lett. 24 (1970) 1155
- 1970NE17 J.M. Nelson, N.S. Chant and P.S. Fisher, Nucl. Phys. A156 (1970) 406
- 1970NU02 F. Nusslin and P. Braun-Munzinger, Z. Phys. 240 (1970) 217
- 1970OG1A Ogloblin, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 231
- 1970OR03 H. Orihara, M. Baba, M. Akiyama, S. Iwasaki, T. Nakagawa, H. Ueno and M. Watanabe, J. Phys. Soc. Jpn. 29 (1970) 533
- 1970PE18 A.K. Petrauskas and L.Y. Sabalyauskas, Yad. Fiz. 12 (1970) 492; Sov. J. Nucl. Phys. 12 (1970) 269
- 1970PO03 D.L. Powell, G.M. Crawley, B.V.N. Rao and B.A. Robson, Nucl. Phys. A147 (1970) 65
- 1970PO1E Porter, Coles and Wyld, Rept. AWRE 045/70, Aldermaston (1970)
- 1970RY03 V.F. Rybachenko and A.A. Sadovoi, Yad. Fiz. 12 (1970) 710; Sov. J. Nucl. Phys. 12 (1971) 384
- 1970RY04 V.F. Rybachenko and A.A. Sadovoi, Izv. Akad. Nauk SSSR Ser. Fiz. 34 (1970) 1784; Bull. Acad. Sci. USSR Phys. Ser. 34 (1971) 1586
- 1970SI12 V.S. Siksin, V.N. Domoratskii, L.V. Donetskov, A.N. Voronin, M.M. Novikov, A.N. Kudryavtsev, G.D. Batyshchev, S.K. Barvin and V.F. Tseshkovskaya, Yad. Fiz. 12 (1970) 15; Sov. J. Nucl. Phys. 12 (1971) 9
- 1970SL1B Slaus, Theory of Nucl. Struct., Trieste, 1969, IAEA STI/PUB/249 (1970) 17
- 1970SO03 Y.I. Sorokin, A.K. Shardanov, V.G. Shevchenko and B.A. Yurev, Yad. Fiz. 11 (1970) 8; Sov. J. Nucl. Phys. 11 (1970) 4

- 1970SQ01 G.T.A. Squier, A.R. Johnston, E.W. Spiers, S.A. Harbison and N.M. Stewart, Nucl. Phys. A141 (1970) 158
- 1970SR01 M.L. Srivastava and J.P. Sah, Current Sci. (India) 39 (1970) 273
- 1970TH1F Theus et al., Bull. Amer. Phys. Soc. 15 (1970) 1695
- 1970VA19 B. Vaucher, J.C. Alder and C. Joseph, Helv. Phys. Acta 43 (1970) 237
- 1970WA11 E.K. Warburton, G.T. Garvey and I.S. Towner, Ann. Phys. (New York) 57 (1970) 174
- 1970WI1C K. Wiesner, K. Trutzschler, D. Schubert and R. Neumann, Nucl. Instrum. Meth. 78 (1970) 131
- 1970ZE03 B. Zeidman, H.T. Fortune and A. Richter, Phys. Rev. C2 (1970) 1612
- 1971AL1E Alexander, Fieno and Ford, Nucl. Sci. Eng. 44 (1971) 95
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AR21 A.G. Artukh, V.V. Avdeichikov, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A168 (1971) 321
- 1971AU1H Auwarter, Meyer and Zamboni, Helv. Phys. Acta 44 (1971) 597
- 1971BA16 V.V. Balashov and V.I. Markov, Nucl. Phys. A163 (1971) 465
- 1971BA2Y Baus-Baghdikian, Univ. libre Bruxelles, Bull. No. 45 (1971)
- 1971BA41 M. Balakrishnan, M.K. Mehta and A.S. Divatia, Nuovo Cim. A1 (1971) 205
- 1971BA61 A.G. Baryshnickov and L.D. Blokhintsev, Phys. Lett. B36 (1971) 205
- 1971BA87 C.J. Batty, Nucl. Phys. A178 (1971) 17
- 1971BI12 H.G. Bingham, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A175 (1971) 374
- 1971BI1D Bingham, Holden and Fortune, Bull. Amer. Phys. Soc. 16 (1971) 621
- 1971CA01 P.J. Castleberry, L. Coulson, R.C. Minehart and K.O.H. Ziock, Phys. Lett. B34 (1971) 57
- 1971CA1J Castleberry, Thesis, Univ. of Virginia (1971)
- 1971DA10 S. Dahlgren, B. Hoistad and P. Grafstrom, Phys. Lett. B35 (1971) 219
- 1971DE2C Deschamps, FRNC TH 182 (1971)
- 1971DE2D Devons, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 619
- 1971DE37 C. Detraz, C.D. Zafiratos, C.E. Moss and C.S. Zaidins, Nucl. Phys. A177 (1971) 258
- 1971DZ07 P.O. Dzhamalov and E.I. Dolinskii, Yad. Fiz. 14 (1971) 753; Sov. J. Nucl. Phys. 14 (1972) 423
- 1971EL1B Elwyn, Monahan and Schiffer, Nucl. Fusion 11 (1971) 551
- 1971FO1P D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 604

- 1971FO24 D.G. Foster, Jr. and D.W. Glasgow, Phys. Rev. C3 (1971) 576
- 1971GA1J Gaillard, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (1971)
- 1971GE09 C. Gerardin, R. Seltz and D. Magnac-Valette, Nucl. Phys. A169 (1971) 521
- 1971GL07 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, Yad. Fiz. 13 (1971) 277; Sov. J. Nucl. Phys. 13 (1971) 154
- 1971GO33 P.M. Gopych, V.I. Salatskii and I.V. Sizov, Yad. Fiz. 14 (1971) 247; Sov. J. Nucl. Phys. 14 (1972) 138
- 1971GU23 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Semenescu, Phys. Rev. C4 (1971) 1092
- 1971IN05 A. Ingemarsson and G. Tibell, Phys. Scr. 4 (1971) 235
- 1971JA04 J. Janecke, T.F. Yang, W.S. Gray and R.M. Polichar, Phys. Rev. C3 (1971) 79
- 1971JO22 R.C. Johnson and F.D. Santos, Part. Nucl. 2 (1971) 285
- 1971JU05 J.A. Jungerman, F.P. Brady, W.J. Knox, T. Montgomery, M.R. McGie, J.L. Romero and Y. Ishizaki, Nucl. Instrum. Meth. 94 (1971) 421
- 1971KA04 S. Kahana and D. Kurath, Phys. Rev. C3 (1971) 543
- 1971KA62 V.V. Karapetyan and G.Y. Korenman, Yad. Fiz. 14 (1971) 1199; Sov. J. Nucl. Phys. 14 (1972) 668
- 1971KE1D Keck, Schober and Jochim, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haeberli (1971) 601
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971LE33 J. Lestringuez, G.M. Raisbeck, F. Yiou and R. Bernas, Phys. Lett. B36 (1971) 331
- 1971LI11 M. Liu, W. von Oertzen, J.C. Jacmart, F. Pougheon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A165 (1971) 118
- 1971LI20 S.C. Ling and S.L. Blatt, Nucl. Phys. A174 (1971) 375
- 1971LO13 R.J. Lombard, Phys. Lett. B35 (1971) 493
- 1971MA13 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A165 (1971) 145
- 1971MA44 G.S. Mani, D. Jacques and A.D.B. Dix, Nucl. Phys. A172 (1971) 166
- 1971MC1H McNally, Nucl. Fusion 11 (1971) 554
- 1971MI12 D. Miljanic and V. Valkovic, Nucl. Phys. A176 (1971) 110
- 1971NO02 J.L. Norton and P. Goldhammer, Nucl. Phys. A165 (1971) 33
- 1971PA10 B.H. Patrick, H.A. Medicus, G.K. Mehta, E.M. Bowey and D.B. Gayther, Phys. Lett. B34 (1971) 488

- 1971PO1D Pornet and Ulpat, CEA R 4252 (1971)
- 1971RA1K Raisbeck and Yiou, Nature 233 (1971) 73
- 1971RA26 G.M. Raisbeck and F. Yiou, Phys. Rev. Lett. 27 (1971) 875
- 1971RI15 H.T. Richards and H.V. Smith, Jr., Phys. Rev. Lett. 27 (1971) 1735
- 1971SA1J Santos, Polarization Phenom. in Nucl. Reactions, Madison, 1970; Eds., H.H. Barschall and W. Haerberli (1971) 729
- 1971SC1N Schnabel, FRNC TH 191 (1971)
- 1971SC1P Schwartz, Schrack and Heaton, Bull. Amer. Phys. Soc. 16 (1971) 495
- 1971SH26 V.S. Shirley, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1255
- 1971ST05 R.H. Stokes and P.G. Young, Phys. Rev. C3 (1971) 984
- 1971ST35 D.M. Stanojevic, M.R. Aleksic, B.Z. Stepancic and R.V. Popic, Fizika 3 (1971) 99
- 1971ST40 A.I. Steshenko and G.F. Filippov, Yad. Fiz. 14 (1971) 715; Sov. J. Nucl. Phys. 14 (1972) 403
- 1971TA1A Talmi, Proc. Int. Conf. Hyperfins Interactions Detected by Nucl. Radiation, Israel, 1970 (1971) 1133
- 1971VA34 S.S. Vasilev, B.S. Galakhmatova, A.S. Demyanova, T.N. Mikhaleva, E.A. Romanovskii and D.L. Chuprunov, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 2364; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 2144
- 1971VO04 D. von Ehrenstein, L. Meyer-Schutzmeister, J.E. Monahan, A. Richter and J.C. Stoltzfus, Phys. Rev. Lett. 27 (1971) 107
- 1971WE07 M.F. Werby, S. Edwards and W.J. Thompson, Nucl. Phys. A169 (1971) 81
- 1971YO05 F.C. Young, R.A. Lindgren and W. Reichart, Nucl. Phys. A176 (1971) 289
- 1971ZA07 A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 273
- 1972AM04 B.S. Amin, S. Biswas, D. Lal and B.L.K. Somayajulu, Nucl. Phys. A195 (1972) 311
- 1972AN05 R.K. Anderson, M.R. Wilson and P. Goldhammer, Phys. Rev. C6 (1972) 136
- 1972AU01 M. Auman, F.P. Brady, J.A. Jungerman, W.J. Knox, M.R. McGie and T.C. Montgomery, Phys. Rev. C5 (1972) 1
- 1972AV01 R. Avida, I. Ben-Zvi, G. Goldring, S.S. Hanna, P.N. Tandon and Y. Wolfson, Nucl. Phys. A182 (1972) 359
- 1972BA09 W.A. Bardeen and E.W. Torigoe, Phys. Lett. B38 (1972) 135
- 1972BE04 H. Behrens and W. Buhring, Nucl. Phys. A179 (1972) 297
- 1972BE1E Bertsch and Mekjian, Ann. Rev. Nucl. Sci. (1972) 25

- 1972BI1E Biro, Sudar and Csikai, Conf. Nucl. Struct. Study with Neutrons, Budapest, 1972 (1972) 130
- 1972BO07 H. Bohlen, N. Marquardt, W. von Oertzen and P. Gorodetzky, Nucl. Phys. A179 (1972) 504
- 1972BO17 V.V. Boikov, B.V. Velberg, I.I. Loshchakov and V.I. Ostroumov, Yad. Fiz. 15 (1972) 425; Sov. J. Nucl. Phys. 15 (1972) 239
- 1972BU1T Bubb, 4th Ains Nucl. Phys. Conf., Sydney, 1972 (1972) 32
- 1972BU29 A.P. Bukhvostov, A.M. Chatrchyan, G.E. Dogotar, R.A. Eramzhyan, N.P. Popov and V.A. Vartanjan, Acta Phys. Pol. B3 (1972) 375
- 1972BU30 A.J. Buffa, Jr. and M.K. Brussel, Nucl. Phys. A195 (1972) 545
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CL1A Clayton, Encyclopedia of the Twentieth Century (1972)
- 1972DA04 P. David, J. Debrus, H. Mommsen and A. Riccato, Nucl. Phys. A182 (1972) 234
- 1972DA26 S. Dahlgren, A. Johansson, P.-U. Renberg and G. Tibell, Phys. Scr. 6 (1972) 5
- 1972DA32 V.K. Daruga and E.S. Matusевич, Atomn. Energ. (USSR) 33 (1972) 757; Sov. At. Energy 33 (1973) 868
- 1972DZ06 P.O. Dzhamalov, E.I. Dolinskii and A.M. Mukhamedzhanov, Yad. Fiz. 15 (1972) 258; Sov. J. Nucl. Phys. 15 (1972) 147
- 1972ED01 G. Eder, G. Winkler and P. Hille, Z. Phys. 253 (1972) 335
- 1972EM01 J.F. Emery, S.A. Reynolds, E.I. Wyatt and G.I. Gleason, Nucl. Sci. Eng. 48 (1972) 319
- 1972EV03 N.N. Evstafev, I.N. Rozantsev and I.V. Estulin, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 2567; Bull. Acad. Sci. USSR Phys. Ser. 36 (1973) 2230
- 1972FA14 G. Faldt and H. Pilkuhn, Phys. Lett. B40 (1972) 613
- 1972FR09 V. Franco, Phys. Rev. C6 (1972) 748
- 1972FR1L Freeman et al., At. Masses Fund. Constants, Teddington, 1971 (1972) 105
- 1972FU1C Funsten, Kossler and Stronach, Bull. Amer. Phys. Soc. 17 (1972) 464
- 1972GA1E Garrett, Symp. on Two-Nucleon Transfer and Pairing Excitations, Argonne (1972) 232; CONF-720309
- 1972GA1F Garvey, Comments Nucl. Part. Phys. 5 (1972) 85
- 1972GO1A D.R. Goosman and D.E. Alburger, Phys. Rev. C5 (1972) 1893
- 1972HA63 S.S. Hanna, M. Hass, Z. Shkedi and Y.S. Horowitz, Nucl. Phys. A195 (1972) 91
- 1972HU1A W.T. Huang, C.A. Levinson and M.K. Banerjee, Phys. Rev. C5 (1972) 651
- 1972KO1E Koike, Prog. Theor. Phys. 48 (1972) 66

1972LE1L Lee and Cusson, *Ann. Phys.* 72 (1972) 353
 1972LE1P Lee et al., *Communications, Proc. Aix-en-Provence Conf.* 2 (1972) 72
 1972LI31 Y.C. Liu, *Chin. J. Phys. (Taiwan)* 10 (1972) 76
 1972LO1D Lobov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 36 (1972) 881
 1972MA21 C. Maples and J. Cerny, *Phys. Lett.* B38 (1972) 504
 1972MC26 E.M. McMillan, *Phys. Rev.* C6 (1972) 2296
 1972ME07 F. Merchez, J. Arvieux, T. Honda and S. Morita, *Nucl. Phys.* A183 (1972) 417
 1972MO1E Morrison, *J. Phys. (Paris)* 33 (1972) C5-111
 1972NA05 P.T. Nang, *Nucl. Phys.* A185 (1972) 413
 1972OB1B Oberg, Bodansky, Chamberlin and Jacobs, *Bull. Amer. Phys. Soc.* 17 (1972) 893
 1972PA1C Parker, *Astrophys. J.* 175 (1972) 261
 1972PN1A Pniewski, *Few Particle Problems, UCLA, 1972* (1972) 145
 1972RE06 P.U. Renberg, D.F. Measday, M. Pepin, P. Schwaller, B. Favier and C. Richard-Serre, *Nucl. Phys.* A183 (1972) 81
 1972RE07 M.J. Renan, J.P.F. Sellschop, R.J. Keddy and D.W. Mingay, *Phys. Rev.* C6 (1972) 12
 1972RI10 A.S. Rinat, L.P. Kok and M. Stingl, *Nucl. Phys.* A190 (1972) 328
 1972RO03 D.C. Robinson, J.M. Freeman and T.T. Thwaites, *Nucl. Phys.* A181 (1972) 645
 1972RO22 J. Rohlin, K. Gabathuler, N.W. Tanner, C.R. Cox and J.J. Domingo, *Phys. Lett.* B40 (1972) 539
 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, *Nucl. Phys.* A188 (1972) 430
 1972SH1K Sherwood, McDaniel and McEllistrem, *Bull. Amer. Phys. Soc.* 17 (1972) 923
 1972SI1G Silberberg, *Bull. Amer. Phys. Soc.* 17 (1972) 96
 1972SK08 D.M. Skopik, et al., *Nucl. Phys.* A197 (1972) 185
 1972SR02 M.L. Srivastava and J.P. Sah, *Current Sci. (India)* 41 (1972) 136
 1972SR04 M.L. Srivastava and J.P. Sah, *Indian J. Pure Appl. Phys.* 10 (1972) 200
 1972ST1M Stocker, Jolivette and Browne, *Bull. Amer. Phys. Soc.* 17 (1972) 464
 1972SW1A Swannack, *Thesis, Carnegie Mellon Univ.* (1972)
 1972TA21 V.K. Tartakovskii and A.V. Fursaev, *Yad. Fiz.* 15 (1972) 51; *Sov. J. Nucl. Phys.* 15 (1972) 31
 1972TH13 C. Thibault and R. Klapisch, *Phys. Rev.* C6 (1972) 1509
 1972THZF H. Theissen, *Springer Tracts in Mod. Phys.*; Ed., Hohler, Vol. 65 (1972) 1
 1972VA02 L. van der Zwan and K.W. Geiger, *Nucl. Phys.* A180 (1972) 615

- 1972VA36 J.F.A. Van Hienen and P.W.M. Glaudemans, Phys. Lett. B42 (1972) 301
- 1972VO06 A.A. Vorobyov, D.M. Seliverstov, V.T. Grachev, I.A. Kondurov, A.M. Nikitin, N.N. Smirnov and Y.K. Zalite, Phys. Lett. B40 (1972) 102
- 1972VO17 H.J. Votava and W.J. Thompson, Phys. Lett. B41 (1972) 405
- 1972VO1H Votava, Thesis, Univ. of North Carolina (1972)
- 1972VO20 V.G. Vovchenko, A.A. Zhdanov, V.M. Zheleznyakov, O.Y. Fedorov and A.V. Shvedchikov, Yad. Fiz. 16 (1972) 1145; Sov. J. Nucl. Phys. 16 (1973) 628
- 1972WA30 J.W. Wachter, W.A. Gibson and W.R. Burrus, Phys. Rev. C6 (1972) 1496
- 1972WE08 K.A. Weber, K. Meier-Ewert, H. Schmidt-Bocking and K. Bethge, Nucl. Phys. A186 (1972) 145
- 1972WI1C Wilkinson, Few Particle Problems, UCLA, 1972 (1972) 191
- 1972YA06 M. Yasue, T. Ohsawa, N. Fujiwara, S. Kakigi, D.C. Nguyen and S. Yamashita, J. Phys. Soc. Jpn. 33 (1972) 265
- 1972YI01 F. Yiou and G.M. Raisbeck, Phys. Rev. Lett. 29 (1972) 372
- 1973AB10 S.N. Abramovich, B.Y. Guzhovskii, A.G. Zvenigorodskii and S.V. Trusillo, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1967; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 144
- 1973AL1E Alster et al., Contrib., Uppsala (1973) 172
- 1973AR1L Arad and Ben-David, Rev. Mod. Phys. 45 (1973) 230
- 1973AU1H Audouze and Truran, OAP-310 (1973)
- 1973BE2J Benenson, Kashy, Moalem and Nann, Bull. Amer. Phys. Soc. 18 (1973) 1382
- 1973BI1H Birchall et al., in Munich, 1 (1973) 371
- 1973BI1N Birchall et al., Bull. Amer. Phys. Soc. 18 (1973) 1394
- 1973BL03 M. Bloser, Atomkernenergie 20 (1972) 309
- 1973BL1B Blatt, Suffert, Degre and Schaeffer, Asilomar (1973) Paper 884
- 1973BU20 A.P. Bukhvostov, S. Ciechanowicz, G.E. Dogotar, R.A. Eramzhyan, N.P. Popov and V.A. Vartanjan, Acta Phys. Pol. B4 (1973) 495
- 1973CE1B Cerny et al., in Munich, 1 (1973) 465
- 1973CH1M D. Chattarji and P. Ghosh, Phys. Rev. C8 (1973) 2115
- 1973CH20 Il-Tong Cheon, Phys. Rev. C8 (1973) 534
- 1973CL09 C.F. Clement, Nucl. Phys. A213 (1973) 469
- 1973CO16 S. Cotanch and D. Robson, Nucl. Phys. A209 (1973) 301
- 1973CO19 W.R. Coker, J. Lin, J.L. Duggan and P.D. Miller, Phys. Lett. B45 (1973) 321
- 1973CO1P Cohen, J. Phys. Soc. Jpn. Suppl. 34 (1973) 63

- 1973DA09 S. Dahlgren, P. Grafstrom, B. Hoistad and A. Asberg, Nucl. Phys. A204 (1973) 53
- 1973DE35 R.M. DeVries, Phys. Rev. C8 (1973) 951
- 1973DI1J Dillig and Huber, in Munich, 1 (1973) 673
- 1973DO02 E.I. Dolinsky, P.O. Dzhamalov and A.M. Mukhamedzhanov, Nucl. Phys. A202 (1973) 97
- 1973EI03 H. Eichner, H. Stehle and P. Heiss, Nucl. Phys. A205 (1973) 249
- 1973GO1V Goosman, Fielding and Lind, Bull. Amer. Phys. Soc. 18 (1973) 1419
- 1973GU08 V.B. Gubin, S.S. Vasilev, B.S. Galakhmatova, T.N. Mikhaleva, E.A. Romanovskii, D.L. Chuprunov and L.I. Nikitina, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 101; Bull. Acad. Sci. USSR Phys. Ser. 37 (1973) 89
- 1973HA49 P.S. Hauge and S. Maripuu, Phys. Rev. C8 (1973) 1609
- 1973HO10 C.G. Hoot, D.K. Olsen, R.E. Brown, J.R. Maxwell and A. Scott, Nucl. Phys. A203 (1973) 339
- 1973HU05 A.H. Hussein and H.S. Sherif, Phys. Rev. C8 (1973) 518
- 1973HU09 R.J. Hughes and E.G. Muirhead, Nucl. Phys. A215 (1973) 147
- 1973HU1D Hughes and Spicer, Asilomar (1973) Paper 2B4
- 1973JO1K John and Kramer, Nucl. Phys. A204 (1973) 203
- 1973JU2A M. Juric, G. Bohm, J. Klabuhn, U. Krecker, F. Wysotzki, G. Coremans-Bertrand, J. Sacton, G. Wilquet, T. Cantwell, F. Esmael et al., Nucl. Phys. B52 (1973) 1
- 1973KA04 V.V. Karapetyan, V.N. Mileev and N.N. Titarenko, Nucl. Phys. A203 (1973) 561
- 1973KA16 E. Kashy, W. Benenson, I.D. Proctor, P. Hauge and G. Bertsch, Phys. Rev. C7 (1973) 2251
- 1973KA1D Karapetyan, Korenman and Popov, Contrib., Uppsala (1973) 154
- 1973KE1E Kemper et al., Bull. Amer. Phys. Soc. 18 (1973) 1427
- 1973KO1D Kovar, Symp. on Heavy Ion Transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 59
- 1973KU03 D. Kurath, Phys. Rev. C7 (1973) 1390
- 1973LA19 H. Laumer, S.M. Austin, L.M. Panggabean and C.N. Davids, Phys. Rev. C8 (1973) 483
- 1973MA1K Maripuu, 5th Symp. Struct. Low-Medium Mass Nuclei (1973) 63
- 1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
- 1973MO02 D.G. Montague, R.K. Cole, M. Makino and C.N. Waddell, Nucl. Phys. A199 (1973) 457
- 1973MO1N Moss and Zaidins, Bull. Amer. Phys. Soc. 18 (1973) 603
- 1973MU11 N.C. Mukhopadhyay, Phys. Lett. B45 (1973) 309

1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
 1973NE1G Nelson and Lane, Bull. Amer. Phys. Soc. 18 (1973) 1395
 1973NE1H Nelson, Hausladen and Lane, Bull. Amer. Phys. Soc. 18 (1973) 244
 1973NY04 E.M. Nyman, Nucl. Phys. A215 (1973) 397
 1973PA14 Y.S. Park, A. Niiler and R.A. Lindgren, Phys. Rev. C8 (1973) 1557; Erratum Phys. Rev. C9 (1974) 1673
 1973RA37 G.M. Raisbeck, J. Lestringuez and F. Yiou, Nature 244 (1973) 28
 1973RE1G Reeves, Audouze, Fowler and Schramm, Astrophys. J. 179 (1973) 909
 1973RO1R Robson, Nucl. Phys. A204 (1973) 523
 1973RO24 U. Rohrer and L. Brown, Nucl. Phys. A210 (1973) 465
 1973SA1Q Saganek, Sledzinska, Wilhelmi and Zwieglinski, in Munich, 1 (1973) 436
 1973SA30 R. Saayman, P.R. de Kock and J.H. van der Merwe, Z. Phys. 265 (1973) 69
 1973SC1J Scott, Symp. on Heavy Ion transfer Reactions, ANL Phy-1973B, Vol. 1 (1973) 97
 1973SI27 A.J. Sierk and T.A. Tombrello, Nucl. Phys. A210 (1973) 341
 1973SU1B Sugimoto, J. Phys. Soc. Jpn. Suppl. 34 (1973) 197
 1973SU1E Suffert, in Asilomar (1973) 741
 1973TO04 I.S. Towner and J.C. Hardy, Nucl. Phys. A205 (1973) 33
 1973TO05 W. Tobocman, R. Ryan, A.J. Baltz and S.H. Kahana, Nucl. Phys. A205 (1973) 193
 1973UL1D Ullrich, Boschitz, Engelhardt and Lewis, Contrib., Uppsala (1973) 156
 1973VO02 H.J. Votava, T.B. Clegg, E.J. Ludwig and W.J. Thompson, Nucl. Phys. A204 (1973) 529; Erratum Nucl. Phys. A217 (1973) 628
 1973VO1G Vorobyov et al., in Munich, 1 (1973) 716
 1973WE19 K.A. Weaver, J.D. Anderson, H.H. Barschall and J.C. Davis, Nucl. Sci. Eng. 52 (1973) 35
 1973WE1T Weaver, Anderson, Barschall and Davis, Phys. Med. Bio. 18 (1973) 64
 1973WI11 D.H. Wilkinson, Nucl. Phys. A209 (1973) 470
 1973WI15 J. Wilczynski, Phys. Lett. B47 (1973) 124
 1973ZW1A Zwieglinski, Piotrowski, Saganek and Sledzinska, Nucl. Phys. A209 (1973) 348
 1974AU1A Audouze and Tinsley, Astrophys. J. 192 (1974) 487
 1974AZ01 G. Azuelos, J.E. Crawford and J.E. Kitching, Phys. Rev. C9 (1974) 1213
 1974DO1G Dollhopf, Perdrisat, Kitching and Olsen, Bull. Amer. Phys. Soc. 19 (1974) 43
 1974MU1J Mukhopadhyay and Cannata, Bull. Amer. Phys. Soc. 19 (1974) 44
 1974SI1H Silberberg, Bull. Amer. Phys. Soc. 19 (1974) 41

- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1
1977AJ02 F. Ajzenberg-Selove, Nucl. Phys. A281 (1977) 1

