

Energy Levels of Light Nuclei $A = 13$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 13$ –15 was published in *Nuclear Physics A360* (1981), p. 1. This version of $A = 13$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. Also, [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed August 1, 1980)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 13$

Below is a list of links for items found within the PDF document. Figures from this evaluation are not included in the PDF document but have been scanned in and are available on this website or via the link below.

A. Nuclides: [¹³Li](#), [¹³Be](#), [¹³B](#), [¹³C](#), [¹³N](#), [¹³O](#), [¹³F](#), [¹³Ne](#)

B. Tables of Recommended Level Energies:

[Table 13.1](#): Energy levels of ¹³B

[Table 13.4](#): Energy levels of ¹³C

[Table 13.18](#): Energy levels of ¹³N

[Table 13.23](#): Energy levels of ¹³O

C. [References](#)

D. Figures: [¹³B](#), [¹³C](#), [¹³N](#), [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

¹³Li
(Not illustrated)

¹³Li has not been observed. ¹³Li is predicted to have an atomic mass excess of 61.56 MeV: it is then unstable for breakup into ¹²Li + n and ¹¹Li + 2n by 0.6 and 4.5 MeV, respectively (1974TH01). The modified mass equation leads to a calculated atomic mass excess of 60.34 MeV: ¹³Li would then be stable with respect to breakup into ¹²Li + n by 0.7 MeV, assuming the a.m.u. of ¹²Li to be 52.94 MeV, but would be unstable with respect to ¹¹Li + 2n by 3.3 MeV (1975JE02). See also (1976AJ04).

¹³Be
(Not illustrated)

¹³Be has not been observed. ¹³Be is predicted to have an atomic mass excess of 35.35 MeV (1974TH01), 34.60 MeV (1975JE02). It is then unstable with respect to decay into ¹²Be + n by 2.20 MeV or by 1.45 MeV, respectively, based on the atomic mass excess of ¹²Be, 25.078 MeV (1978AL29). See also (1976AJ04) and (1977SE1D; theor.).

¹³B
(Figs. 1 and 4)

GENERAL: (See also (1976AJ04).)

Experimental work on complex reactions in which ¹³B is observed: (1976BU16, 1977AR06, 1978GE1C, 1978KO01, 1979LEZR).

Reviews and theoretical papers: (1976AB04, 1976VA29, 1977DO06, 1978AB08, 1978DE15, 1979AL22, 1979BE1H, 1979BO22, 1980MA1F, 1980MU1B).

$$Q = 0.0478 \pm 0.0046 \text{ b (1973HAVZ, 1978LEZA).}$$

$$\mu = +3.17778 \pm 0.00051 \text{ nm (1978LEZA).}$$

1. ¹³B(β^-)¹³C $Q_m = 13.437$

The half-life of ¹³B is 17.33 ± 0.17 msec (1971WI07). The mean of this and a previously reported measurement is 17.36 ± 0.16 msec (1971WI07). The branching ratios of various ¹³C states are shown in Table 13.2: they indicate $J^\pi = \frac{1}{2}^-$ and $\frac{3}{2}^-$ for ¹³B_{g.s.}. See also (1976AJ04) and (1977AZ02, 1977MA16, 1977RI08, 1979DE15; theor.).

Table 13.1: Energy levels of ^{13}B

| E_x (MeV \pm keV) | $J^\pi; T$ | τ or Γ_{cm} (keV) | Decay | Reactions |
|-----------------------|------------------------------|--------------------------------------|------------|---------------------------|
| g.s. | $\frac{3}{2}^-; \frac{3}{2}$ | $\tau_{1/2} = 17.36 \pm 0.16$ msec | β^- | 1, 2, 3, 4, 5, 6, 7, 8, 9 |
| 3.483 ± 5 | ^a | | γ | 2, 4 |
| 3.5346 ± 3.1 | ^a | $\tau_m > 0.3$ psec | γ | 2, 4 |
| 3.681 ± 5 | ^a | | (γ) | 2, 4, 7 |
| 3.7126 ± 4.5 | ^a | $\tau_m < 0.38$ psec | (γ) | 2, 4, 7 |
| 4.131 ± 6 | | $\tau_m = 0.062 \pm 0.050$ psec | γ | 2, 4 |
| 4.834 ± 7 | | | (γ) | 2, 4 |
| 5.029 ± 7 | | | | 2, 4 |
| 5.106 ± 10 | | $\Gamma = 60 \pm 10$ keV | | 4 |
| 5.392 ± 8 | | 10 ± 10 | | 2, 4 |
| (5.557 ± 7) | | | | 2 |
| 6.167 ± 7 | | | | 2, 4 |
| 6.425 ± 7 | | 36 ± 5 | | 2, 4 |
| 6.934 ± 9 | ^a | 55 ± 15 | | 2, 4 |
| (7.516 ± 8) | | | | 2 |
| (7.859 ± 20) | | | | 2 |
| 8.134 ± 7 | | 100 ± 15 | | 2, 4 |
| 8.683 ± 7 | | 89 ± 20 | | 2, 4 |
| 9.44 ± 30 | | 81 ± 25 | | 4 |
| 10.22 ± 20 | | 210 ± 20 | | 4 |
| 10.89 ± 20 | | | | 4 |
| (11.80) | | | | 4 |

^a See Table 13.3.

Table 13.2: Beta decay of ^{13}B ^a

| Decay to $^{13}\text{C}^*$ (MeV) | J^π | Branch (%) | $\log ft$ ^b |
|----------------------------------|-------------------|-------------------|------------------------|
| 0 | $\frac{1}{2}^-$ | 92.1 ± 0.8 | 4.01 ± 0.01 |
| 3.09 | $\frac{1}{2}^+$ | ≤ 0.7 | ≥ 5.7 |
| 3.68 | $\frac{3}{2}^-$ | 7.6 ± 0.8 | 4.45 ± 0.04 |
| 3.85 | $\frac{5}{2}^+$ | ≤ 0.7 | ≥ 5.5 |
| 7.55 ^c | $\frac{5}{2}^-$ | 0.094 ± 0.020 | 5.33 ± 0.08 |
| 8.86 | $\frac{1}{2}^-$ | 0.16 ± 0.03 | 4.60 ± 0.08 |
| 9.50 | $(\frac{3}{2}^-)$ | < 0.01 | > 5.2 |
| 9.90 | $\frac{3}{2}^-$ | 0.022 ± 0.007 | 4.95 ± 0.12 |

^a (1969JO21, 1974AL12). See also Table 13.24.

^b Log ft shown here are based on $\tau_{1/2} = 17.33 \pm 0.17$ msec (1971WI07). See also (1977RI08; theor.).

^c $E_x = 7.577 \pm 0.030$ MeV (1974AL12).

2. $^7\text{Li}(^7\text{Li}, \text{p})^{13}\text{B}$ $Q_m = 5.965$

Observed proton groups are shown in Table 13.3 (1972WY01). For angular distribution measurements see (1969CA1A). The lifetimes of $^{13}\text{B}^*$ (3.53, 3.71, 4.13) are, respectively, > 0.3 , < 0.38 and 0.062 ± 0.050 psec: $E_\gamma = 3536.3 \pm 4.2$ and 4133.4 ± 7.8 keV (1969TH01). See also ^{14}C .

3. $^{10}\text{Be}(\alpha, \text{p})^{13}\text{B}$ $Q_m = -8.818$

See ^{14}C .

4. $^{11}\text{B}(\text{t}, \text{p})^{13}\text{B}$ $Q_m = -0.233$

Observed proton groups are displayed in Table 13.3. Angular distributions have been measured at $E_t = 11$ MeV (1964MI04) and 23 MeV (1978AJ02).

5. $^{12}\text{C}(^9\text{Be}, ^8\text{B})^{13}\text{B}$ $Q_m = -28.136$

Table 13.3: Proton groups from ${}^7\text{Li}({}^7\text{Li}, p){}^{13}\text{B}$ and ${}^{11}\text{B}(t, p){}^{13}\text{B}$

| ${}^7\text{Li}({}^7\text{Li}, p){}^{13}\text{B}$ (1972WY01) | ${}^{11}\text{B}(t, p){}^{13}\text{B}$ | | | | |
|----------------------------------------------------------------|----------------------------------------|-----------------------|-----------------------------------------------|------------------|-----------------------|
| | (1964MI04) | | | (1978AJ02) | |
| | E_x (MeV \pm keV) | E_x (MeV \pm keV) | L | J^π | E_x (MeV \pm keV) |
| 0 | 0 | 0 | $\frac{3}{2}^-$ | 0 | |
| | 3.483 ± 5 | 1 | $(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+ g$ | 3.482 ± 10 | |
| a | 3.533 ± 5 | 2 | $(\frac{1}{2}, \frac{5}{2}, \frac{7}{2})^- g$ | 3.531 ± 10 | |
| | 3.681 ± 5 | 1 | $(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+ g$ | 3.681 ± 10 | |
| a,b | 3.712 ± 5 | 2 | $(\frac{1}{2}, \frac{5}{2}, \frac{7}{2})^- g$ | 3.715 ± 10 | |
| c,d | 4.13 ± 10 | 2 | $(\frac{1}{2}, \frac{5}{2}, \frac{7}{2})^- g$ | 4.128 ± 10 | |
| 4.833 ± 10 | 4.82 ± 10 | | | 4.834 ± 10 | |
| 5.033 ± 8 | 5.01 ± 10 | 1 | $(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+ g$ | 5.023 ± 10 | |
| | | | | 5.106 ± 10 | 60 ± 10 |
| 5.391 ± 8 | 5.38 ± 10^e | | | 5.393 ± 10 | 10 ± 10 |
| 5.557 ± 8 | | | | f | |
| 6.169 ± 8 | 6.17 ± 20 | | | 6.164 ± 10 | |
| 6.419 ± 8 | | | | 6.434 ± 10 | 36 ± 5 |
| 6.939 ± 15 | | | | 6.932 ± 10^h | 55 ± 15 |
| 7.516 ± 8 | | | | f | |
| 7.859 ± 20 | | | | f | |
| 8.129 ± 10 | | | | 8.138 ± 10 | 100 ± 15 |
| 8.682 ± 9 | | | | 8.684 ± 10 | 89 ± 20 |
| | | | | 9.44 ± 30 | 81 ± 25 |
| | | | | 10.22 ± 20 | 210 ± 20 |
| | | | | 10.89 ± 20 | |
| | | | | (11.80) | |

^a See text.

^b The decay is primarily by γ_0 : the upper limit to the cascade via ${}^{13}\text{B}^*(3.5)$ is 10%.

^c The decay is $75 \pm 10\%$, $25 \pm 10\%$ and $< 10\%$, respectively to ${}^{13}\text{B}^*(0, 3.5, 3.7)$.

^d All values in this column are based on $E_x = 4131$ keV for ${}^{13}\text{B}^*(4.13)$.

^e $\Gamma = 15 \pm 5$ keV.

^f Not observed.

^g See, however, (1978AJ02), p. 1289.

^h $L \geq 4$.

See (1975WI26).

$$6. {}^{13}\text{C}(\pi^-, \gamma){}^{13}\text{B} \quad Q_m = 126.130$$

See (1979MAZW) and (1979TR1B).

$$7. {}^{14}\text{C}(\text{d}, {}^3\text{He}){}^{13}\text{B} \quad Q_m = -15.338$$

At $E_d = 52$ MeV angular distributions for ${}^{13}\text{B}^*(0, 3.7)$ have been reported (1975MA41).

$$8. {}^{14}\text{C}(\text{t}, \alpha){}^{13}\text{B} \quad Q_m = -1.017$$

At $E_t = 23$ MeV the angular distribution for α_0 has been analyzed by (1979SE07): the DWBA fit is poor. The extracted $S(\frac{3}{2}) = 5.7$, near that predicted for a filled $p_{3/2}$ shell.

$$9. {}^{14}\text{C}({}^{14}\text{C}, {}^{13}\text{B}){}^{15}\text{N} \quad Q_m = -10.624$$

See (1980NA14).

¹³C
(Figs. 2 and 4)

GENERAL: (See also (1976AJ04).)

Shell model: (1977TE01, 1978BO31, 1979HO17).

Collective, rotational and deformed models: (1976BR26, 1977ME1E).

Cluster model: (1977SA19).

Special levels: (1977ME1E, 1977TE01, 1977RI08, 1978BO31, 1978MI04, 1979RO1E, 1980BA54).

Electromagnetic transitions: (1977ME1E, 1977YO1D, 1978KI08, 1978KR19, 1978MI04, 1980BA54).

Giant resonances: (1977AL18, 1977MA06, 1979DO17, 1979HO17).

Applied work: (1975GA1E, 1977AN1F, 1977SH1E, 1978EM1A, 1978KE1E, 1978LE1P, 1979AL1P, 1979EN1D, 1979JA1F, 1979KU20, 1979SM1B).

Special reactions: (1976AB04, 1976BE1K, 1976BU16, 1976HI05, 1976LE12, 1976LE1F, 1977AR06, 1977RE08, 1977SH1D, 1977UD1A, 1977YA1B, 1978AB08, 1978BI08, 1978GE1C, 1978HE1C, 1978KO01, 1978LE15, 1979AL22, 1979BO22, 1979HE1D, 1979PO16, 1979SA27, 1979ST1D, 1980MI01).

Astrophysical questions: (1975LO1E, 1976BE1C, 1976BO1M, 1976BR1K, 1976DE1F, 1976DE1G, 1976DE1H, 1976EP1A, 1976FI1E, 1976FU1C, 1976GA1F, 1976HI1D, 1976MA1N, 1976NO1C, 1976QU1A, 1976RE1B, 1976SP1B, 1976TO1D, 1976VA1D, 1976VI1B, 1976WA1G, 1977AU1E, 1977AU1F, 1977CA1H, 1977CA1J, 1977CA1K, 1977CH1J, 1977CL1B, 1977CL1C, 1977CO1J, 1977DA1E, 1977DE1N, 1977DI1C, 1977FU1L, 1977HA1L, 1977JO1D, 1977LA1F, 1977LA1G, 1977LI1H, 1977PA1D, 1977PR1E, 1977SC1H, 1977SH09, 1977ST1H, 1977TI1A, 1977TR1D, 1977WA1N, 1977WA1P, 1977WI1D, 1978BU1B, 1978DE1R, 1978DO1B, 1978IB1A, 1978LU1C, 1978SN1A, 1978ST1C, 1978ST1D, 1978TR1C, 1978VA1B, 1978WA1E, 1979BE1V, 1979BJ1A, 1979GE1D, 1979GU1D, 1979LA1H, 1979MC1B, 1979ME1D, 1979OL1B, 1979PE1E, 1979RA1C, 1979SA1M, 1979SW1B, 1979TU1A, 1979WI1H, 1980HE1D, 1980LA1G, 1980MC1G, 1980PE1F).

Muon and neutrino capture and reactions: (1977BA1P, 1977DO06, 1977MU1A, 1978DE15, 1978LE04, 1979BE1N, 1979DE01, 1980MU1B).

Pion capture and reactions (See also reactions 35 and 53.): (1976AS1B, 1976AU1E, 1976BO2C, 1976DI11, 1976EN02, 1976GA06, 1976GA21, 1976JA1D, 1976LI04, 1976NI05, 1976OS06, 1977AU01, 1977DI1B, 1977FU10, 1977GI06, 1977HO1B, 1977LO19, 1977WA02, 1978DY01, 1978KI08, 1978KI13, 1978MI02, 1978RO1H, 1978WA02, 1979AL1J, 1979ALZW, 1979AN1J, 1979ANZM, 1979BA2D, 1979BA2E, 1979BE1N, 1979BO1U, 1979CO1H, 1979CO1C, 1979DE34, 1979GI1D, 1979GR1K, 1979GY1A, 1979GY1B, 1979HI1A, 1979HO1F, 1979IN1A, 1979JO1C,

1979JO08, 1979LA22, 1979LEZR, 1979LI1H, 1979MAZW, 1979MO1N, 1979MO1P, 1979MO1Q, 1979PI1C, 1979RE1A, 1979SA21, 1979ST1J, 1979TR1B, 1979TRZR, 1980BO03, 1980CH08, 1980ER01, 1980GA12, 1980JO06, 1980LA1C, 1980LE02, 1980MA1F, 1980NA1B, 1980PE1C, 1980SA04, 1980TH01).

Table 13.4: Energy levels of ^{13}C ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m or Γ_{cm} (keV) | Decay | Reactions |
|-----------------------|------------------------------|---------------------------------|-------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| g.s. | $\frac{1}{2}^-; \frac{1}{2}$ | | stable | 7, 8, 9, 13, 15, 16, 17, 24, 25, 26, 27, 29, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 48, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 75, 77, 78, 79, 80, 81, 83, 85, 86, 87, 88, 89, 91, 92, 93, 94 |
| 3.089443 ± 0.020 | $\frac{1}{2}^+$ | $\tau_m = 1.55 \pm 0.15$ fsec | γ | 8, 9, 13, 15, 24, 26, 35, 36, 37, 40, 43, 44, 52, 53, 54, 55, 56, 57, 58, 61, 69, 70, 71, 78, 80, 81, 83, 87, 88, 91 |
| 3.684482 ± 0.023 | $\frac{3}{2}^-$ | 1.59 ± 0.13 fsec | γ | 7, 8, 9, 13, 15, 16, 24, 26, 29, 35, 36, 37, 39, 40, 47, 52, 53, 54, 55, 56, 57, 58, 61, 69, 70, 71, 77, 78, 79, 80, 81, 87, 88, 91 |
| 3.853783 ± 0.022 | $\frac{5}{2}^+$ | 12.5 ± 0.3 psec | γ | 8, 9, 13, 15, 24, 26, 35, 36, 37, 38, 39, 40, 42, 43, 44, 52, 53, 54, 55, 56, 57, 58, 61, 69, 70, 71, 80, 81, 87, 91 |
| 6.864 ± 3 | $\frac{5}{2}^+$ | $\Gamma = 6$ | γ, n | 7, 8, 9, 14, 15, 24, 26, 30, 35, 36, 52, 55, 57, 58, 81 |
| 7.492 ± 10 | $(\frac{7}{2}^+)$ | < 5 | | 7, 9, 14, 24, 26, 36, 43, 53, 58, 80, 81 |
| 7.547 ± 3 | $\frac{5}{2}^-$ | 1.2 ± 0.3 | γ, n | 7, 9, 14, 16, 24, 26, 30, 35, 36, 47, 49, 52, 53, 55, 57, 58, 79, 80, 81, 87 |

Table 13.4: Energy levels of ^{13}C ^a (continued)

| E_x (MeV \pm keV) | J^π, T | τ_m or Γ_{cm} (keV) | Decay | Reactions |
|--------------------------------|--------------------------------|----------------------------------------|---------------------|--------------------------------------------------|
| 7.686 \pm 6 | $\frac{3}{2}^+$ | 70 \pm 5 | γ, n | 14, 24, 26, 30, 36, 49, 58, 81 |
| 8.2 \pm 100 | $\frac{3}{2}^+$ | 1000 \pm 300 | γ, n | 9, 30, 36, 49 |
| 8.860 \pm 20 | $\frac{1}{2}^-$ | 150 \pm 30 | γ, n | 24, 30, 36, 47, 49, 52, 57, 77, 80, 81, 87, 88 |
| 9.4998 \pm 0.1 | $\frac{9}{2}^+$ | 5 | n | 7, 9, 14, 24, 30, 35, 36, 52, 53, 57, 80, 81, 87 |
| 9.897 \pm 5 | $\frac{3}{2}^-$ | 26 \pm 3 | γ, n | 7, 14, 24, 30, 36, 47, 49, 51, 52 |
| 10.46 | | 200 | n | 30 |
| 10.753 \pm 4 | $\frac{7}{2}^-$ | 55 \pm 2 | n | 14, 24, 30, 36, 81 |
| 10.818 \pm 5 | $(\frac{5}{2}^-)$ | 24 \pm 3 | n | 7, 9, 14, 24, 36, 81 |
| 10.996 \pm 6 | $\frac{1}{2}^+$ | 37 \pm 4 | γ, n, α | 4, 24, 30, 36, 49, 80, 81 |
| 11.080 \pm 5 | $(\frac{1}{2}^-)$ | < 4 | γ, n, α | 4, 24, 30, 36, 52, 81, 87 |
| 11.748 \pm 10 | | 107 \pm 14 | n | 24, 36, 53, 81 |
| 11.851 \pm 5 | $(\frac{3}{2}^-)$ | 68 \pm 4 | n | 30, 36, 52, 57, 77, 80, 87 |
| 11.95 \pm 40 | $(\frac{5}{2}, \frac{7}{2})^+$ | \approx 200 | n, α | 4, 6, 36, 55, 80 |
| 12.106 \pm 5 | $> \frac{7}{2}$ | 81 \pm 8 | n, α | 4, 30, 36, 81 |
| (12.187 \pm 10) | | 110 \pm 50 | | 52 |
| 12.438 \pm 12 | $\frac{7}{2}^-$ | 160 \pm 40 | n, α | 4, 30, 52, 87 |
| 13.0 \pm 1000 | | broad | γ, n | 49 |
| (13.28) | $(\frac{3}{2}^-)$ | 340 | α | 6 |
| 13.41 | $(\frac{9}{2}^-)$ | 35 \pm 3 | n, α | 4, 6 |
| 13.56 | $(\frac{3}{2}, \frac{5}{2})^+$ | \approx 600 | n, α | 4, 6, 30 |
| 13.76 | $(\frac{5}{2}, \frac{3}{2})^+$ | \approx 300 | n, α | 4, 6 |
| 14.12 | $(\frac{5}{2}^-)$ | \approx 150 | n, α | 4, 6 |
| 14.390 \pm 15 | $(\frac{1}{2}, \frac{5}{2})^-$ | 280 \pm 70 | n, α | 4, 52 |
| 14.582 \pm 10 | | 230 \pm 50 | n, α | 4, 52 |
| 14.983 \pm 10 | $(\frac{3}{2}^+)$ | 380 \pm 60 | n, α | 4, 52 |
| 15.1082 \pm 1.2 ^b | $\frac{3}{2}^-; \frac{3}{2}$ | 5.49 \pm 0.25 | γ, n, α | 4, 6, 7, 24, 49, 52, 53, 57, 87 |
| 15.526 \pm 11 | | 150 \pm 30 | n, α | 4, 52 |

Table 13.4: Energy levels of ^{13}C ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m or Γ_{cm} (keV) | Decay | Reactions |
|-----------------------|------------|----------------------------------------|-----------------------------------|--------------------|
| 16.080 \pm 7 | | 150 \pm 15 | n, α | 4, 52 |
| 16.15 \pm 50 | | 230 | n, α | 4, 52, 53 |
| 16.95 \pm 50 | | 330 | n, α | 4 |
| 17.36 \pm 100 | | 190 | n, α | 4, 53 |
| 17.699 \pm 5 | | 170 | n, α | 4, 53 |
| 18.30 \pm 50 | | 300 | n, α | 4 |
| 18.699 \pm 5 | | 100 \pm 15 | γ , n, p, α | 4, 50, 52 |
| 19.5 | | \approx 450 | n | 30 |
| 19.9 | | \approx 600 | n, p, d | 18, 19 |
| 20.021 \pm 13 | | 230 \pm 30 | (γ), n, (p), d, α | 17, 18, 22, 50, 52 |
| 20.429 \pm 8 | | 116 \pm 10 | (γ), n, p, d | 17, 18, 19, 32, 52 |
| 21.28 \pm 15 | | 159 \pm 15 | n, p, d | 18, 19, 53, 55 |
| 21.466 \pm 8 | | 270 \pm 20 | | 52 |
| 21.81 \pm 20 | | 114 \pm 21 | n, d | 18, 55 |
| 22 | | \approx 1000 | n, d | 18, 52, 53 |
| 23 | | \approx 1000 | n | 30 |
| 24 | | \approx 4000 | γ , n, p | 49, 50, 52 |
| 25.5 | | broad | γ , p | 50, 52 |
| 26.8 | | | n, d | 18 |
| 27.5 | | | n, d | 18, 52 |
| 30 | | | γ , n | 49, 52 |

^a See also Tables 13.5 and 13.6.

^b See Table 13.7.

Other topics: (1976BI1A, 1976BR26, 1976FE1B, 1976VA1C, 1978DA1A, 1978LI1D, 1978MC04, 1978PO1A, 1978SH1B, 1978SO1A, 1979BE1H, 1979HE1F).

Ground state of ^{13}C : (1976BR26, 1976DU04, 1976FU06, 1976JO1B, 1977AN21, 1977NO07, 1977YO1D, 1978AN07, 1978HE1D, 1978MI04, 1978TA09, 1978ZA1D, 1979SA27).

$$\mu = +0.702411 \pm 0.000001 \text{ nm (1978LEZA)}.$$

Radius of ^{13}C : The rms nuclear charge radius is 2.51 ± 0.04 fm (1979BE1N; prelim.: see also reaction 52). The neutron rms radius is 2.35 ± 0.03 fm (1979JO08). See also (1979SA27; theor.).

Table 13.5: Lifetimes of low-lying states of ^{13}C ^a

| $^{13}\text{C}^*$ (MeV) | τ_m | Refs. |
|-------------------------|----------------------|---------------|
| 3.09 | 1.50 ± 0.17 fsec | (1968RO02) |
| | 1.7 ± 0.3 fsec | (1975RA22) |
| 3.68 | 1.55 ± 0.15 fsec | weighted mean |
| | 1.50 ± 0.15 fsec | (1969RA20) |
| | 1.83 ± 0.25 fsec | (1970WI04) |
| 3.85 | 1.59 ± 0.13 fsec | weighted mean |
| | 10.7 ± 1 psec | (1969HE22) |
| | 9.9 ± 0.9 psec | (1970GA01) |
| | 12.4 ± 0.8 psec | (1974BE48) |
| | 13.0 ± 0.4 psec | (1975RA29) |
| | 12.6 ± 0.3 psec | (1977HE12) |
| | 12.5 ± 0.3 psec | weighted mean |

^a See also Table 13.5 in (1976AJ04).

$^{13}\text{C}^*(3.85) [J^\pi = \frac{5}{2}^+]$: g is negative (1976DY05).

| | | |
|---------------------------------------------------|----------------|----------------|
| 1. (a) $^6\text{Li}(^7\text{Li}, n)^{12}\text{C}$ | $Q_m = 20.924$ | $E_b = 25.870$ |
| (b) $^6\text{Li}(^7\text{Li}, p)^{12}\text{B}$ | $Q_m = 8.337$ | |
| (c) $^6\text{Li}(^7\text{Li}, 2n)^{11}\text{C}$ | $Q_m = 2.203$ | |
| (d) $^6\text{Li}(^7\text{Li}, d)^{11}\text{B}$ | $Q_m = 7.192$ | |
| (e) $^6\text{Li}(^7\text{Li}, t)^{10}\text{B}$ | $Q_m = 1.994$ | |
| (f) $^6\text{Li}(^7\text{Li}, \alpha)^9\text{Be}$ | $Q_m = 15.223$ | |

The yield curves for $d_0(E(^6\text{Li}) = 4 \text{ to } 14 \text{ MeV})$, $t_0(E(^7\text{Li}) = 5 \text{ to } 14 \text{ MeV})$ and $\alpha_0(E(^6\text{Li}) = 4 \text{ to } 14 \text{ MeV})$ show broad, uncorrelated structures. Energy-averaged differential cross sections are also reported for a number of ^{12}B , ^{11}B and ^{10}B states (1971WY01). Total cross section measurements have been measured for $E(^7\text{Li}) = 3.8 \text{ to } 6.0 \text{ MeV}$ for $p_0 \rightarrow p_2, p_{3+4}, p_5$; $d_0 \rightarrow d_3, d_{4+5}, d_6$; $t_0 \rightarrow t_2$; and α_0 : total cross sections generally increase smoothly with energy without showing any structure (1967KI03). The ^{11}C yield has been measured for $E(^6\text{Li}) = 1.2 \text{ to } 3.6 \text{ MeV}$ by (1961NO05). See also ^9Be and ^{10}B in (1979AJ01) and ^{11}B and ^{12}B in (1980AJ01).

Table 13.6: Summary of results on the total radiation widths of the low-lying levels of $^{13}\text{C}-^{13}\text{N}$ ^a

| $J_i^\pi \rightarrow J_f^\pi$ | $^{13}\text{C}^*$ (MeV) | Γ_γ (eV) | Refs. | $^{13}\text{N}^*$ (MeV) | Γ_γ (eV) | Refs. |
|-------------------------------------------|-------------------------|---------------------------------------------|------------|-------------------------|----------------------|------------|
| $\frac{1}{2}^+ \rightarrow \frac{1}{2}^-$ | 3.09 ^b | 0.43 ± 0.04 | Table 13.5 | 2.37 | 0.45 ± 0.05 | (1968RI16) |
| $\frac{3}{2}^- \rightarrow \frac{1}{2}^-$ | 3.68 ^c | 0.41 ± 0.04 | Table 13.5 | 3.51 ^f | 0.70 | (1963YO06) |
| $\frac{5}{2}^+ \rightarrow \frac{1}{2}^-$ | 3.85 ^d | $(5.3 \pm 0.2) \times 10^{-5}$ ^e | Table 13.5 | 3.55 | $< 2 \times 10^{-3}$ | (1963YO06) |

^a See also Tables 13.15 and 13.21.

^b $E_x = 3089.443 \pm 0.020$ keV, $E_\gamma = 3089.049 \pm 0.020$ keV* (1980WA24: here, and in footnote ^d measured values are starred (*); the others are derived).

^c Branching ratio for cascade via $^{13}\text{C}^*(3.09)$ is $0.75 \pm 0.04\%$ (1980WA24). See also (1975TR07). $E_x = 3684.482 \pm 0.023$ keV, $E_\gamma = 3683.921 \pm 0.023$ keV. $\delta(E2/M1) = -0.094 \pm 0.009$. E_γ for transition to $^{13}\text{C}^*(3.09)$ is 595.013 ± 0.011 keV (1980WA24).

^d Branching ratios for cascades via $^{13}\text{C}^*(3.68, 3.09)$ are $36.3 \pm 0.6\%$ and $1.20 \pm 0.04\%$, respectively (1980WA24). See also (1975TR07). $E_x = 3853.783 \pm 0.022$ keV, $E_\gamma = 3853.170 \pm 0.022$ keV; E_γ for the transitions to $^{13}\text{C}^*(3.09, 3.68)$ are 764.316 ± 0.010 keV* and 169.300 ± 0.004 keV* (1980WA24).

^e The ground state branching ratio is $62.5 \pm 0.6\%$ (1980WA24) and $\delta(E3/M2) = +0.12 \pm 0.03$ (1966PO11).

^f Branching ratio for cascade via $^{13}\text{N}^*(2.37)$ is $8 \pm 1\%$ (1974RO29). See also footnote ^j in Table 13.21.

$$2. \ ^7\text{Li}(^7\text{Li}, n)^{13}\text{C} \quad Q_m = 18.620$$

See (1978KA1H) and (1970AJ04).

$$3. \ ^9\text{Be}(\alpha, \gamma)^{13}\text{C} \quad Q_m = 10.6479$$

See (1976AJ04).

$$4. \ (a) \ ^9\text{Be}(\alpha, n)^{12}\text{C} \quad Q_m = 5.7015 \quad E_b = 10.6479$$

$$(b) \ ^9\text{Be}(\alpha, 2n)^{11}\text{C} \quad Q_m = -13.020$$

Resonances for n_0 and n_1 , for γ -rays from $^{12}\text{C}^*(4.4, 12.7, 15.1)$ and resonances in the total neutron cross section are given in Table 13.8. The yield of neutrons to $^{12}\text{C}^*(7.65, 9.64)$ has been measured in the range $E_\alpha = 2.9-6.4$ MeV (1972OB01). The n_0 and n_1 excitation functions exhibit weak resonance anomalies at $E_\alpha = 6.44$ MeV corresponding to the $J^\pi = \frac{3}{2}^-$, $T = \frac{3}{2}$ state at $E_x = 15.11$ MeV: see Tables 13.7 and 13.8 (1978HI06). Based on all available information the charge asymmetry of the reduced transition strengths, $\delta \equiv B(^{13}\text{C})/B(^{13}\text{N}) - 1$, $\delta(M1) = -0.15 \pm 0.07$ and $\delta(E2) = 1.0 \pm 0.6$ (1978HI06).

Table 13.7: Parameters of the first $T = \frac{3}{2}$ states in ^{13}C and ^{13}N ^a

| | $^{13}\text{C}^*(15.11)$ | $^{13}\text{N}^*(15.06)$ |
|-----------------------------------------------------|------------------------------------------------------|-----------------------------------------------------------------------|
| E_x (MeV) | 15.1082 ± 0.0012 | 15.0646 ± 0.0009 |
| J^π | $\frac{3}{2}^-$ | $\frac{3}{2}^-$ |
| Γ_{cm} (keV) | 5.49 ± 0.25 | 0.932 ± 0.028 |
| Γ_{γ_0} (eV) | 22.4 ± 1.5 (M1), 0.6 ± 0.1 (E2) ^k | 24.2 ± 1.5 (M1) ^e 0.32 ± 0.12 (E2) ^f |
| Γ_{γ_1} (eV) | 4.12 ± 0.74 | $\leq 2.82 \pm 0.30$ ^g |
| $\Gamma_{\gamma_{2+3}}$ (eV) | 18.2 ± 2.4 | 19.6 ± 1.4 ^h |
| Γ_{γ_0}/Γ (%) | 0.396 ± 0.030 b | |
| $\Gamma_{p_0}\Gamma_{\gamma_0}/\Gamma$ (eV) | | 5.79 ± 0.20 |
| $\Gamma_{\gamma_0}/\Gamma_{p_0}$ (%) | | 12.1 ± 1.1 |
| Γ_{n_0} or Γ_{p_0} (keV) ^c | 0.38 ± 0.10 | 0.228 ± 0.016 ⁱ |
| Γ_{n_1} or Γ_{p_1} (keV) ^c | 1.43 ± 0.18 | 0.140 ± 0.014 ⁱ |
| Γ_{n_2} or Γ_{p_2} (keV) ^c | 0.14 ± 0.10 | 0.049 ± 0.015 ⁱ |
| Γ_{p_3} (keV) ^c | | 0.089 ± 0.014 ⁱ |
| Γ_{p_5} (keV) ^c | | 0.15 ± 0.04 ⁱ j |
| Γ_{α_0} (keV) ^d | 0.104 ± 0.028 | 0.046 ± 0.026 ⁱ |
| Γ_{α_1} (keV) ^d | | 0.036 ± 0.036 ⁱ |
| Γ_{α_2} (keV) ^d | | 0.067 ± 0.042 ⁱ |

^a (1973AD02, 1973HU07, 1975HI07, 1975MA21, 1977MA16, 1978HI06, 1979AD01, 1980TH05).

^b The decay width to $^{13}\text{C}^*(7.55)$ is < 0.9 eV (1977MA16).

^c Widths for $^{13}\text{C}^*(15.11) \rightarrow ^{12}\text{C}_{\text{g.s.}} + n_0$ or $^{13}\text{N}^*(15.06) \rightarrow ^{12}\text{C}_{\text{g.s.}} + p_0$ [$n_1, p_1; n_2, p_2$; and p_3 and p_5 refer to the decays to $^{12}\text{C}^*(4.4, 7.7, 9.6, 10.8)$ respectively].

^d Widths for $^{13}\text{C}^*(15.11) \rightarrow ^9\text{Be}_{\text{g.s.}} + \alpha_0$ or $^{13}\text{N}^*(15.06) \rightarrow ^9\text{B}_{\text{g.s.}} + \alpha_0$ [α_1 and α_2 refer to the decays to $^9\text{B}^*((1.6), 2.4)$].

^e $\delta = -0.07 \pm 0.13$ (1975MA21). Here $\delta \equiv B(^{13}\text{C})/B(^{13}\text{N}) - 1$.

^f $\delta = 0.82^{+1.2}_{-0.6}$ (1975MA21).

^g $\delta \geq 0.83 \pm 0.29$ (1975MA21).

^h $\delta = -0.04 \pm 0.14$ (1975MA21).

ⁱ Based on measured branching ratios and on $\Gamma_{\text{cm}} = 0.932 \pm 0.028$ keV (1980TH05).

^j The decay width to $^{12}\text{C}^*(12.71)$ is < 0.13 keV. It is expected to be ≈ 0.03 keV. The sum of the branching ratios for all measured decays of $^{13}\text{N}^*(15.06)$ is $92 \pm 8\%$. It is apparent from the character of the decay modes of this state that 2s-1d shell isospin admixtures are important (1979AD01).

^k See also (1979JU01).

Table 13.8: Resonances in ${}^9\text{Be}(\alpha, n){}^{12}\text{C}$

| E_α ^a (MeV) | E_α ^b (MeV) | E_α ^c (MeV) | Γ_{cm} (keV) | J^π | ${}^{13}\text{C}^*$ ^d (MeV) | Refs. |
|-------------------------------|-------------------------------|-------------------------------|----------------------------|----------------------------------|----------------------------------------|---------------|
| 0.52 | 0.52 | | ≈ 55 ^e | $(\frac{1}{2}^+)$ | 11.01 | A |
| 0.60 | 0.60 | | < 4 ^c | | 11.06 | A |
| 1.9 | 1.905 | 1.92 | 130 | $(\frac{7}{2}^-)$ | 11.97 | A |
| 2.24 | | 2.25 | 280 | | 12.20 | A |
| 2.58 | 2.6 | 2.58 | ≈ 200 | $(\frac{1}{2}^-)$ | 12.43 | A |
| 4.00 | 3.98 | 4.00 | 35 ± 3 ^g | | 13.41 | A |
| 4.18 | | | 570 | $(\frac{3}{2}^+)$ | 13.54 | A, (1973DE14) |
| 4.50 | 4.47 | 4.50 | ≈ 350 | $(\frac{5}{2}^+)$ | 13.76 | A, (1973DE14) |
| 5.0 | 5.02 | 5.0 | ≈ 200 | | 14.12 | A |
| 5.40 ± 0.10 | 5.3 ^f | | 260 | $(\frac{1}{2}^-, \frac{5}{2}^-)$ | 14.39 ± 0.1 | A, (1973DE14) |
| | 5.75 | 5.75 | 210 | | 14.63 | A |
| 6.20 ± 0.05 | | | 380 | $(\frac{3}{2}^+)$ | 14.94 ± 0.05 | A, (1973DE14) |
| | 6.44 ^h | | | $\frac{3}{2}^-; T = \frac{3}{2}$ | 15.109 | (1978HI06) |
| 7.10 ± 0.05 | 7.00 | | 220 | | 15.56 ± 0.05 | A |
| | 7.75 | 7.8 | 210 | | 16.01 | A |
| 7.95 ± 0.05 | | | 230 | | 16.15 ± 0.05 | A |
| 9.10 ± 0.05 | | 9.1 | 330 | | 16.95 ± 0.05 | A |
| 9.7 ± 0.10 | 9.70 | | 190 | | 17.36 ± 0.1 | A |
| 10.2 ± 0.05 | | | 170 | | 17.71 ± 0.05 | A |
| 11.05 ± 0.05 | | | 300 | | 18.30 ± 0.05 | A |
| 11.70 ± 0.03 | 11.60 | | 70 | | 18.75 ± 0.03 | A |

A: References for this resonance are displayed in Table 13.8 (1976AJ04).

^a Resonances in total neutron yield.

^b Resonances in n_1 group and for 4.4 MeV γ -rays.

^c Resonances in total cross section.

^d Not corrected for effects of Coulomb barrier penetration.

^e $\omega\gamma = 3.79$ and 0.88 eV, respectively (1968DA05).

^f (1973DE14) suggest the possibility that ${}^{13}\text{C}^*(14.4)$ is composed of two states with the same J^π .

^g J.L. Weil, private communication.

^h Anomalies in n_0 and n_1 yields at $E_\alpha = 6443.5 \pm 2.0$ keV: see Table 13.7 for parameters of 15.11 MeV state (1978HI06).

Polarization measurements have been reported for E_α to 22.9 MeV (see (1976AJ04)) and also at $E_\alpha = 2.4$ and 2.8 MeV (1976NI01; n_0, n_1) and 100 MeV (1978LE10). Reaction (b) has been studied at a number of energies in the range $E_\alpha = 17$ to 44 MeV (1970KR09). See also (1975GE19, 1976GE1A, 1977LI19).

| | | |
|--------------------------------------------------|-----------------|-----------------|
| 5. (a) ${}^9\text{Be}(\alpha, p){}^{12}\text{B}$ | $Q_m = -6.886$ | $E_b = 10.6479$ |
| (b) ${}^9\text{Be}(\alpha, d){}^{11}\text{B}$ | $Q_m = -8.0308$ | |
| (c) ${}^9\text{Be}(\alpha, t){}^{10}\text{B}$ | $Q_m = -13.229$ | |

Excitation curves have been measured for $E_\alpha = 15$ to 25 MeV (1975VA19; d_0, d_1) and 26.5 to 27.5 MeV (1975PU01; $d_0 \rightarrow d_3, d_{4+5}, d_6$): no sharp structures are observed. See also ${}^{11}\text{B}$ in (1980AJ01). At $E_\alpha = 26.0$ to 27.5 MeV yield curves have been obtained for the t_0, t_1 and t_3 groups: no structure is observed (1974KE06). See also (1976LE1K) and (1968AJ02) for reaction (a).

| | |
|-------------------------------------------------|-----------------|
| 6. ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$ | $E_b = 10.6479$ |
|-------------------------------------------------|-----------------|

A number of excitation functions have been measured for elastically scattered α -particles (reaction (a)) for $E_\alpha = 1.4$ to 20 MeV: these show considerable resonance structure with the variations being most prominent below 10 MeV but persisting up to 20 MeV. The parameters resulting from a best-fit of the excitation functions are displayed in Table 13.9: see the footnotes to that table for a summary of the most important caveats (1973GO15). A weak resonance is observed in the α_0 yield at $E_\alpha = 6.44$ MeV corresponding to the excitation of the first $T = \frac{3}{2}$ state at $E_x = 15.11$: see Table 13.7 for the parameters of that state (1978HI06). See also (1974LO1B, 1979AJ01).

| | |
|-----------------------------------------------------|---------------|
| 7. ${}^9\text{Be}({}^6\text{Li}, d){}^{13}\text{C}$ | $Q_m = 9.174$ |
|-----------------------------------------------------|---------------|

At $E({}^6\text{Li}) = 23.8$ MeV angular distributions are reported to ${}^{13}\text{C}^*(0, 3.1, 3.6, 6.9, 7.5, 9.9, 10.8, 12.0, 13.3)$. There is some indication of the population of ${}^{13}\text{C}^*(9.5, 13.9, 14.5, 15.2, 16.7, 18.5)$ (1971GO24).

| | |
|-----------------------------------------------------|---------------|
| 8. ${}^9\text{Be}({}^7\text{Li}, t){}^{13}\text{C}$ | $Q_m = 8.181$ |
|-----------------------------------------------------|---------------|

At $E({}^7\text{Li}) = 5.6$ to 6.2 MeV angular distributions are reported for t_0, t_1, t_{2+3}, t_4 (1969SN02). See also (1976AJ04).

Table 13.9: Resonances in ${}^9\text{Be}(\alpha, \alpha_0)$ ^a

| E_α (MeV) | Γ_{cm} (keV) | l_α | J^π | ${}^{13}\text{C}^*$ (MeV) |
|-------------------|----------------------------|------------|----------------------------------|---------------------------|
| 1.93 ^b | 180 ^b | 1, 0 | $\frac{5}{2}^+$ | 11.98 |
| 3.80 | 343 | 0, 2 | $\frac{3}{2}^-$ ^c | 13.28 |
| 4.00 | 58 | (4, 6) | $(\frac{9}{2}^-)$ | 13.42 |
| 4.20 | 685 | 1, 3 | $\frac{5}{2}^+$ ^d | 13.56 |
| 4.50 | 247 | 1, 3 | $\frac{3}{2}^+$ ^d | 13.76 |
| 5.00 | 75 | 2, 4 | $\frac{5}{2}^-$ ^e | 14.11 |
| 5.075 | 73 | 3, 5 | $\frac{7}{2}^+$ ^e | 14.162 |
| (5.50) | 400 | (1, 3) | $(\frac{5}{2}^+)$ | (14.46) |
| 6.44 | f | | $\frac{3}{2}^-; T = \frac{3}{2}$ | 15.11 |

^a (1973GO15): from analysis in the single-level approximation. This assumes the J^π ordering suggested by (1965LI09).

^b This resonance is reported by (1974SA16). It is not clear whether the Γ is in the cm or lab systems. Γ_α is given as 72 keV.

^c Favored by the analysis but the assignment is not certain and more than one state may be involved.

^d (1973DE14) suggest the opposite ordering [$\frac{3}{2}^+, \frac{5}{2}^+$]: see Table 13.8.

^e An equally good fit to the data is obtained with a $\frac{7}{2}^-$ state at 5.0 MeV and a $(\frac{3}{2}, \frac{5}{2}, \frac{7}{2})^+$ state at 5.08 MeV.

^f Weak anomaly at $E_\alpha = 6443.5 \pm 2.0$ keV: see Table 13.7 for parameters of 15.11 MeV state, and reaction 4 (1978HI06).

$$9. {}^9\text{Be}({}^{12}\text{C}, {}^8\text{Be}){}^{13}\text{C} \quad Q_m = 3.2812$$

Angular distributions have been measured at $E({}^{12}\text{C}) = 12$ and 15 MeV (1970BA49, 1971BA68) and $E({}^9\text{Be}) = 20$ MeV (1979JA1G, 1979BO1K) for the transitions to ${}^{13}\text{C}^*(0, 3.09, 3.85)$ and in addition to ${}^{13}\text{C}^*(3.68)$ in the latter experiment. The neutron spectroscopic factors are 1.15, 0.95, 0.20, 1.02 for the first four states of ${}^{13}\text{C}$ (1979JA1G). See also (1976AJ04). At $E({}^9\text{Be}) = 50$ MeV ${}^{13}\text{C}^*(6.86, 7.5, 8.2, 9.5, 10.8)$ are also populated (1977ST20). For yield measurements [total cross sections, and transitions involving ${}^{13}\text{C}^*(0, 3.09, 3.68, 3.85, 6.86, 7.5)$], see (1977SW05, 1978CH02, 1978MA44, 1979BO06). See also (1977UH1A).

$$10. {}^9\text{Be}({}^{14}\text{N}, {}^{10}\text{B}){}^{13}\text{C} \quad Q_m = -0.9653$$

See (1970AJ04).

$$11. \text{}^{10}\text{Be}(\alpha, n)\text{}^{13}\text{C} \quad Q_m = 3.8361$$

See (1976EP1A; astrophys.).

$$12. \begin{array}{lll} \text{(a) } \text{}^{10}\text{B}(\text{t}, \text{p})\text{}^{12}\text{B} & Q_m = 6.343 & E_b = 23.8766 \\ \text{(b) } \text{}^{10}\text{B}(\text{t}, \text{d})\text{}^{11}\text{B} & Q_m = 5.1979 & \\ \text{(c) } \text{}^{10}\text{B}(\text{t}, \alpha)\text{}^9\text{Be} & Q_m = 13.2287 & \end{array}$$

The p_0 and p_1 yields from reaction (a), the d_0 yield from reaction (b) and the α_0 yield from reaction (c) have been determined for $E_t = 0.8$ to 2.0 MeV. There is no evidence of resonance behavior (1963HO19). See also (1977ST1N; p_0, p_1, p_5, p_6) and (1977CI1A; $\alpha_0 \rightarrow \alpha_3$), both in the interval $E_t = 0.5$ to 1.25 MeV.

$$13. \text{}^{10}\text{B}(\alpha, \text{p})\text{}^{13}\text{C} \quad Q_m = 4.0626$$

Proton groups have been observed to the first four states of ^{13}C : see (1976AJ04). Angular distributions of p_0 have been measured at many energies up to $E_\alpha = 30.4$ MeV: see (1970AJ04) and ^{14}N . For $\tau_{1/2}$ measurements see Table 13.5 in (1976AJ04). For the decay of $^{13}\text{C}^*(3.85)$ see Table 13.6 (1980WA24). See also (1976EP1A; astrophys.).

$$14. \text{}^{10}\text{B}(\text{}^6\text{Li}, \text{}^3\text{He})\text{}^{13}\text{C} \quad Q_m = 8.083$$

Comparisons of the relative intensities of the ^3He groups in this reaction and of the triton groups in the mirror reaction (see reaction 9 in ^{13}N) at $E(^6\text{Li}) = 18$ MeV suggest that the following states are analogs: $6.86 - 6.36, 7.49 - 7.16, 9.50 - 9.00, 9.90 - 9.48, (10.82 + 10.75) - (10.36 + 10.36)$ [the first (set of) E_x is in ^{13}C , the second in ^{13}N]. An angular distribution is also reported for $^{13}\text{C}^*(9.50)$ and the widths for $^{13}\text{C}^*(7.55, 7.69, 9.50)$ are determined to be $< 30, 60 \pm 30$ and < 30 keV, respectively (1974HO06).

$$15. \text{}^{10}\text{B}(\text{}^7\text{Li}, \alpha)\text{}^{13}\text{C} \quad Q_m = 21.4099$$

Angular distributions have been measured at $E(^7\text{Li}) = 5.20$ MeV for the $\alpha_0, \alpha_1, \alpha_{2+3}$ and α_4 groups (1966MC05).

16. $^{10}\text{B}(^{14}\text{N}, ^{11}\text{C})^{13}\text{C}$ $Q_m = 1.140$

At $E(^{10}\text{B}) = 100$ MeV angular distributions are reported for the transitions to $^{13}\text{C}^*(0, 3.68, 7.5, 11.8)$ (1975NA15, 1976NA09). See also (1979MO14).

17. $^{11}\text{B}(\text{d}, \gamma)^{13}\text{C}$ $Q_m = 18.6787$

The $90^\circ \gamma_0$ excitation curve measured for $E_d = 1.0$ to 4.2 MeV shows a resonance at $E_d = 2.0 \pm 0.1$ MeV ($^{13}\text{C}^*(20.4)$) with $\Gamma_{\text{lab}} \approx 0.6$ MeV. The angular distributions of γ_0 are isotropic to within 10% on and off resonance, consistent with E1 radiation (1973WE12).

18. $^{11}\text{B}(\text{d}, \text{n})^{12}\text{C}$ $Q_m = 13.7323$ $E_b = 18.6787$

The yield of neutrons and γ -rays has been measured for $E_d = 0.2$ to 11 MeV: observed resonances are displayed in Table 13.10. For a listing of polarization measurements see Table 13.11 in (1976AJ04) and (1975SI22; $E_d = 0.9$ and 1.2 MeV; n_0, n_1). See also (1974LO1B, 1977SE09) and ^{12}C in (1980AJ01).

19. $^{11}\text{B}(\text{d}, \text{p})^{12}\text{B}$ $Q_m = 1.145$ $E_b = 18.6787$

Some measurements show that the thin-target yield rises smoothly from $E_d = 0.3$ to 3.1 MeV with no evidence of resonances: see (1970AJ04). However, (1964BR1A) and (1968CH05) report a resonance in the yields of p_0, p_1 and p_2 , and of γ_1 and γ_2 : see Table 13.10. For polarization measurements see (1976AJ04) and (1976TA07; 1.0 to 4.0 MeV; p_0). See also (1974FI1D, 1976HA1J), (1977SA25; theor.) and ^{12}B in (1980AJ01).

20. $^{11}\text{B}(\text{d}, \text{d})^{11}\text{B}$ $E_b = 18.6787$

A polarization measurement has been reported at $E_d = 12.6$ MeV (1971ZA04). See also ^{11}B in (1980AJ01).

21. (a) $^{11}\text{B}(\text{d}, \text{t})^{10}\text{B}$ $Q_m = -5.1979$ $E_b = 18.6787$
 (b) $^{11}\text{B}(\text{d}, ^3\text{He})^{10}\text{Be}$ $Q_m = -5.7352$

Table 13.10: Resonant structure in $^{11}\text{B} + \text{d}$

| Resonant structure in yield of | | | | | | | | | Γ_{cm} (keV) | E_x (MeV) |
|--------------------------------|--------------------|--------------------|--------------------|--------------------|------------------------------|----------------|-------------------------------------------|-----------------------|----------------------------|-------------------|
| γ_0 ^a | n_0 ^b | n_1 ^b | n_2 ^b | n_3 ^b | $\gamma_{15.1}$ ^c | p | $\gamma_{0.9}, \gamma_{1.7}$ ^d | α ^e | | |
| (MeV \pm keV) | | | | | | | | | | |
| 2.0 ± 100 | | 1.2 | | | | | | | | 19.7 ⁱ |
| | | 1.45 | | | | | 1.5 ^g | | ≈ 600 | 19.90 |
| | | 1.6 | 1.8 ^f | | | | | | ≈ 200 | 20.24 |
| | | | 2.2 ^f | | | 2.180 ± 10 | 2.2 ^{g,h} | ≈ 2.1 | 116 ± 10 | 20.52 |
| | | | | | | 3.080 ± 15 | 3.0 ^g | | 159 ± 15 | 21.28 |
| | | 3.6 | | | | 3.71 ± 20 | | | 114 ± 21 | 21.81 |
| | | 4.23 | 4.0 | 4.1 | | 4.4 | | | broad | 22.23 |
| | | | (5.2) | | | | | | | (23.1) |
| | | 9.6 | 9.6 | 9.6 | 9.6 | | | | | 26.8 |
| | 10.4 | | 10.4 | 10.4 | | | | | 27.5 | |

^a (1973WE12): $\Gamma_{\text{lab}} \approx 600$ keV.

^b (1965AL17, 1967DI01, 1972TH14).

^c (1958KA31, 1964KU09).

^d Broad resonance in yields of $\gamma_{0.95}$ and $\gamma_{1.67}$ (1968CH05).

^e Yield of $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ (1969FR03).

^f (1965AL17) report a resonance at 1.8 MeV while (1967DI01) report one at 2.2 MeV, in addition to a sharper structure at 1.2 MeV.

^g Resonance in polarization of ^{12}B recoils (1967PF02).

^h Yield of p_0, p_1 and p_2 (1964BR1A).

ⁱ (1971RI19, 1972SE09) suggest $J^\pi = \frac{5}{2}^-$.

See (1976AJ04) and ^{10}B and ^{10}Be in (1979AJ01).

22. $^{11}\text{B}(\text{d}, \alpha)^9\text{Be}$

$$Q_m = 8.0308$$

$$E_b = 18.6787$$

At low energies the excitation functions for α_0 and α_1 increase monotonically: see (1970AJ04). Then at $E_d = 1.85$ MeV a pronounced resonance is observed in the $\alpha_0, \alpha_1, \alpha_2$ and α_3 yields (1969FR03). Some gross structure is also observed in these yields for $E_d = 1.0$ to 3.2 MeV (1964BR1A, 1969FR03). See also (1978GR07) and ^9Be in (1979AJ01).

23. $^{11}\text{B}(\text{t}, \text{n})^{13}\text{C}$

$$Q_m = 12.4214$$

See ^{14}C .

24. $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$

$$Q_m = 13.1851$$

Levels derived from proton groups are displayed in Table 13.11. The decay parameters for the first $T = \frac{3}{2}$ state at $E_x = 15.11$ MeV are displayed in Table 13.7 (1973AD02, 1975MA21, 1977MA16). See also reactions 11 and 13 in ^{13}N . The reduced asymmetries $B(^{13}\text{C})/B(^{13}\text{N}) - 1$ are -0.07 ± 0.13 , $0.82_{-0.6}^{+1.2} \leq 0.83 \pm 0.29$ and -0.04 ± 0.14 for the $\gamma_0(\text{M1})$, $\gamma_0(\text{E2})$, $\gamma_1(\text{E1})$ and $\gamma_2(\text{M1})$ transitions. Changes in the wave functions due to binding energy differences in ^{13}C and ^{13}N do not account for the observed asymmetry of the E1 decays of the first excited states in ^{13}C and ^{13}N (1977MA16): see Table 13.6.

Angular distributions of proton groups have been measured at many energies to $E(^3\text{He}) = 12$ MeV: see (1970AJ04, 1976AJ04). See also (1979HAYZ) and (1976EP1A; astrophys.).

25. $^{11}\text{B}(\alpha, \text{d})^{13}\text{C}$

$$Q_m = -5.1680$$

Angular distributions of the d_0 group have been measured at $E_\alpha = 15.1$ to 25.2 MeV (1975VA19) and 23 and 25 MeV (1967AL16). See also (1976LE1K).

26. $^{11}\text{B}(^6\text{Li}, \alpha)^{13}\text{C}$

$$Q_m = 17.205$$

Angular distributions are reported at $E(^6\text{Li}) = 4.72$ MeV for $\alpha_0, \alpha_1, \alpha_{2+3}, \alpha_4, \alpha_{5+6+7}$ (1966MC05).

27. $^{11}\text{B}(^{14}\text{N}, ^{12}\text{C})^{13}\text{C}$

$$Q_m = 8.4063$$

Angular distributions involving the ground state transitions have been measured at $E(^{14}\text{N}) = 41, 77$ and 113 MeV (1971LI11). See also (1978DZ1A; theor.).

28. $^{11}\text{B}(^{16}\text{O}, ^{14}\text{N})^{13}\text{C}$

$$Q_m = -2.0576$$

See (1976AJ04).

29. $^{12}\text{C}(\text{n}, \gamma)^{13}\text{C}$

$$Q_m = 4.94639$$

$$Q_0 = 4946.32 \pm 0.05 \text{ keV (1978HA14).}$$

Table 13.11: Levels of ^{13}C from $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$ ^a

| E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) |
|-----------------------------|------------------------------|
| 0 | |
| 3.09 | |
| 3.68 | < 5 |
| 3.85 | < 5 |
| 6.871 ± 12 ^b | < 10 |
| 7.500 ± 12 | < 5 |
| 7.554 ± 12 | < 5 |
| 7.694 ± 14 | 70 ± 10 |
| 8.869 ± 36 | 150 ± 30 |
| 9.509 ± 12 | < 10 |
| 9.896 ± 12 ^c | < 10 |
| 10.76 ± 10 | |
| 10.82 ± 10 | |
| 11.01 ± 10 | |
| 11.09 ± 10 | |
| (11.72) ^d | |
| 15.11 ^e | |

^a A number of higher states were reported in Table 13.7 of (1970AJ04): however, the references for these levels have not been published. References for the data shown here are displayed in Table 13.12 (1976AJ04).

^b Decay is by n-emission to $^{12}\text{C}_{\text{g.s.}}$: branching ratio = 0.99 ± 0.09 (1973AD02).

^c Branching ratios for neutron decay to $^{12}\text{C}^*(0, 4.4) = 1.0 \pm 0.2$ and < 0.15 , respectively (1973AD02).

^d For this state these branching ratios are 0.67 ± 0.16 and 0.33 ± 0.08 , respectively (1973AD02).

^e See Table 13.7 for the parameters of this state.

See also previous values listed in (1976AJ04) and (1974SP04, 1979GR1J [preliminary value 4946.329 ± 0.024 keV]). The thermal capture cross section is 3.4 ± 0.3 mb (1973MU14). The capture is $32 \pm 1\%$ via $^{13}\text{C}^*(3.68)$ and $68 \pm 1\%$ to $^{13}\text{C}_{\text{g.s.}}$ (1968SP01). $^{13}\text{C}^*(3.68)$ decays with $E_\gamma = 3684.28 \pm 0.14$ keV (1968SP01), 3683.94 ± 0.17 keV (1967PR10): see Table 13.13 in (1976AJ04) for other data. See also (1978AR1N), (1980DO1C; astrophys.) and (1977MA06; theor.).

30. (a) $^{12}\text{C}(n, n)^{12}\text{C}$ $E_b = 4.94639$
 (b) $^{12}\text{C}(n, n')^{12}\text{C}^*$
 (c) $^{12}\text{C}(n, n')^4\text{He}^4\text{He}^4\text{He}$ $Q_m = -7.2748$

The coherent scattering length (thermal, bound) $a_{\text{coh}} = 6.6535 \pm 0.0014$ fm; $\sigma_{\text{scatt}} = 4.7456 \pm 0.0020$ b (1979KO26). See (1978FU1G) for an R -matrix analysis of all the data below the inelastic channel and for best values of the elastic scattering cross sections.

Total cross section measurements in the range 1 keV to 273 GeV/ c are listed in Table 13.14 of (1976AJ04) and displayed in (1976GAYV). Recent σ_t measurements are reported by (1979SM08; 0.1 to 4.5 MeV), (1979AU07; 1.2 to 13.9 MeV) and (1980CI03; 4.5 to 8 MeV). Cross sections for populating $^{12}\text{C}^*(4.4)$ [by observing n_1 or 4.4 MeV γ -rays] have recently been reported by (1975RO30; 4.8 \rightarrow 20 MeV), (1977DI11; 5 \rightarrow 19 MeV), (1978HA1P; 8.0 \rightarrow 14.5 MeV; also n_0), (1978NO04; 8.1 \rightarrow 9.0 MeV) and (1976GL11; 9 \rightarrow 15 MeV; also n_0). For the earlier measurements see (1959AJ76, 1970AJ04, 1976AJ04, 1976GAYV).

Observed resonances are displayed in Tables 13.12 and 13.13. The parameters of the resonances with $E_n > 6.3$ MeV are poorly known.

Polarization measurements have recently been reported at 4.60, 4.82, 5.17 MeV (1979WE1E, 1979WEZP; n_0), 14.2 MeV (1976BR1J, 1976CA13; n_0, n_1) and 15.85 MeV (1976LE1M, 1976TH1B; n_0, n_1). For the earlier work see Table 13.15 in (1976AJ04). For reaction (c) see (1976CO1N) and ^{12}C in (1980AJ01). See also (1975BL07, 1976HA1N, 1976TO1C, 1978ME12, 1978SU1E, 1979BE50, 1979FI1B, 1979LAZP, 1979LAZQ, 1979URZZ, 1980LA1D), (1975HO1F, 1976KI1F, 1976LI1H, 1976WA1B, 1976WA1C, 1977LA1H), (1977SM1D; applied) and (1976TH10, 1977PU02, 1978AD1A, 1978CA03, 1978MI1E, 1978MI04, 1979RO1E, 1980SH01; theor.).

31. $^{12}\text{C}(n, 2n)^{11}\text{C}$ $Q_m = -18.721$ $E_b = 4.94639$

See (1975AC1A) and (1976AJ04, 1976GAYV).

32. $^{12}\text{C}(n, p)^{12}\text{B}$ $Q_m = -12.587$ $E_b = 4.94639$

Table 13.12: Resonances in $^{12}\text{C}(n, n)^{12}\text{C}$ ^a

| E_{res} (MeV \pm keV) | Γ_{cm} (keV) | $^{13}\text{C}^*$ (MeV) | l_n | J^π | |
|----------------------------------|-----------------------------|-------------------------|-------|----------------------------------|-------------------------------------------------|
| | | 3.09 | 0 | $\frac{1}{2}^+$ | $\theta^2 = 0.185$ ^b |
| 2.079 ± 3 | 6 | 6.864 | 2 | $\frac{5}{2}^+$ | ^c |
| 2.819 ± 3 | 1.2 ± 0.3 | 7.547 | | | |
| 2.94 ± 10 | 124 ± 7 ^d | 7.66 | 2 | $\frac{3}{2}^+$ | |
| 3.472 ± 15 | 1000 ± 50 ^d | 8.149 | 2 | $\frac{3}{2}^+$ | $\theta^2 = 0.35$ ^c |
| 4.259 ± 15 | 210 ± 15 ^d | 8.875 | 1 | $\frac{1}{2}^-$ | $\Gamma_{\text{el}}/\Gamma = 1.00$ ^c |
| 4.93707 ± 0.07 ^e | 1.9 ± 0.15 ^e | 9.4998 | 1 | $(\frac{1}{2}^-, \frac{3}{2}^-)$ | 1.00 ^c |
| 5.368 ± 5 | 26 ± 3 | 9.897 | 1 | $\frac{3}{2}^-$ | 0.70 ± 0.10 ^c |
| 6.294 ± 5 | 53 ± 4 | 10.751 | 3 | $\frac{7}{2}^-$ | 0.70 ± 0.10 |
| 6.5 | | 10.9 | | | |
| 6.558 ± 8 | 37 ± 4 | 10.994 | (0) | $(\frac{1}{2}^+)$ | 0.40 ± 0.10 |
| 6.7 | | 11.1 | | | |
| (7.4) | (250) | (11.8) | | $(\geq \frac{5}{2})$ | |
| 7.759 ± 8 | (200) | 12.102 | | $(> \frac{7}{2})$ | |
| (8.1) | (150) | (12.4) | | | |
| 9.3 | 370 | 13.5 | | | |
| 11.1 | 450 | 15.2 | | $(\geq \frac{3}{2})$ | |
| 12.1 | 230 | 16.1 | | | |
| 15.8 | ≈ 460 | 19.5 | 1 | $(\frac{1}{2}, \frac{3}{2})^-$ | |
| 19.6 ± 200 | ≈ 1000 | 23.0 | | | |

^a See Tables 13.10 in (1970AJ04) and 13.16 in (1976AJ04) for references. See also (1973MU14).

^b $\gamma_n^2 = 540$ keV, radius = 4.80 fm (1970ME1C: single bound state + hard sphere scattering).

^c See also Table 1 in (1973DA17).

^d Γ_n for $^{13}\text{C}^*(7.69, 8.2, 8.86)$ are reported to be 170, 1110 and 170 keV, respectively (1980HO11; $^{13}\text{C}(\gamma, n_0)$).

^e Derived from a lorentzian probability plot (1980CI03).

Table 13.13: Resonances in $^{12}\text{C}(n, n'\gamma_{4.4})^{12}\text{C}$ ^a

| E_n (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | E_x in ^{13}C (MeV) |
|--------------|------------------------------|--------------------------------|
| 4.96 | < 80 | 9.52 |
| 5.42 | < 80 | 9.95 |
| 5.98 | 200 | 10.46 |
| 6.35 | 120 | 10.80 |
| 6.57 | < 80 | 11.01 |
| 6.65 | < 80 | 11.08 |
| 7.50 | 260 | 11.86 |
| 7.81 | 180 | 12.15 |
| 8.14 | 220 | 12.45 |
| 9.31 | 500 | 13.53 |
| 15.8 | ≈ 450 | 19.5 |
| ≈ 19 | | 22.5 ^b |

^a The first ten resonances are from (1959HA13), the two highest are reported by (1968BO34).

^b See, however, (1970DE14).

The cross section exhibits a weak resonance corresponding to $E_x \approx 20.5$ MeV and a stronger structure at $E_x \approx 21.5$ MeV: see (1976AJ04). Polarization measurements are reported at $E_n = 50$ MeV (1977FI1B). See also ^{12}B in (1980AJ01) and (1974BO1E, 1975NI1C, 1976GAYV, 1976SL2A, 1979ROZV).

$$33. \text{ (a) } ^{12}\text{C}(n, d)^{11}\text{B} \quad Q_m = -13.7323 \quad E_b = 4.94639$$

$$\text{ (b) } ^{12}\text{C}(n, t)^{10}\text{B} \quad Q_m = -18.9302$$

For reaction (a) see (1976AJ04). For reaction (b) see (1978QA01). For both see (1976GAYV, 1976SL2A).

$$34. ^{12}\text{C}(n, \alpha)^9\text{Be} \quad Q_m = -5.7015 \quad E_b = 4.94639$$

The cross section for the α_0 group shows a broad structure at $E_n \approx 8$ MeV (1963DA12). See also (1978SU1D) and (1976AJ04, 1976GAYV, 1976SL2A).

35. $^{12}\text{C}(\text{p}, \pi^+)^{13}\text{C}$

$$Q_m = -135.403$$

At $E_p = 185$ MeV, angular distributions have been observed for the π^+ groups to the ground state of ^{13}C and to $E_x = 3.08 \pm 0.09, 3.80 \pm 0.12, 6.87 \pm 0.18, 7.57 \pm 0.16$ and 9.51 ± 0.22 MeV. The shapes of the distributions and the differential cross sections to the various states are quite different: several kinds of reaction processes appear to be involved (1973DA24). Polarization measurements are reported at $E_p = 200$ MeV to $^{13}\text{C}^*(0, 3.09 + 3.68 + 3.85)$ (1978AU07)[†]. See also (1979PIZZ, 1979SOZY, 1979TSZZ), (1975HU1D, 1976AJ04), (1976MI14, 1976NO1D, 1977KU1G, 1977LE1H, 1978WE1G, 1979CE1A; theor.) and the “Pion capture and reactions” section here.

36. (a) $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$

$$Q_m = 2.7218$$

(b) $^{12}\text{C}(\text{d}, \text{np})^{12}\text{C}$

$$Q_m = -2.2246$$

Measurements of the proton groups are summarized in Table 13.14. Angular distributions have been studied at many energies to $E_d = 80.2$ MeV: see Table 13.19 in (1976AJ04) for a listing of earlier work and (1979OS11; $E_d = 1.58 \rightarrow 2.20$ MeV; p_1). Observed γ -rays are listed in Table 13.16 of (1970AJ04) and in Table 13.6 here: ground state decays have been observed for $^{13}\text{C}^*(3.09, 3.68, 3.85)$. $^{13}\text{C}^*(3.68)$ decays via $^{13}\text{C}^*(3.09)$ with a $0.75 \pm 0.04\%$ branch (1980WA24). $^{13}\text{C}^*(3.85)$ also decays via $^{13}\text{C}^*(3.68)$ with a $36.0 \pm 0.7\%$ branch (1975TR07), $36.3 \pm 0.6\%$ (1980WA24) with $E_\gamma = 169.300 \pm 0.004$ keV (1980WA24) and via $^{13}\text{C}^*(3.09)$ with a $1.20 \pm 0.04\%$ branch (1980WA24). Mixing ratios are given in (1970AJ04). Lifetimes of the first three excited states are displayed in Table 13.5: see (1976AJ04) and (1977HE1D, 1977HE12). The nuclear g factor of $^{13}\text{C}^*(3.85)$ is 0.59 ± 0.05 (1974BE48).

Reaction (b) has been studied in a kinematically complete experiment at $E_d = 5.00$ to 9.85 MeV: there is some evidence for p-n final state interaction (1973SA03). See also (1966YO1A, 1976DY05, 1979SI07), (1976EP1A; astrophys.), (1977FI1A, 1978BO1L; applied), (1976HE17, 1976OS07, 1976SE1F, 1976SH13, 1976TA1D, 1977AN1G, 1977TA08, 1978HA1Q, 1979NE1B, 1979ZE1B; theor.) and ^{14}N .

37. $^{12}\text{C}(\text{t}, \text{d})^{13}\text{C}$

$$Q_m = -1.3109$$

At $E_t = 12$ MeV angular distributions have been studied for the $\text{d}_0 \rightarrow \text{d}_3$ groups (1966GL01).

38. $^{12}\text{C}(^3\text{He}, 2\text{p})^{13}\text{C}$

$$Q_m = -2.7718$$

[†] For cross section measurements near threshold see (1979MA38, 1979MA39).

Table 13.14: Levels of ^{13}C from $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$

| ^{13}C (MeV \pm keV) | | Γ_{cm} (keV) | l_n^b | J^π^b | S |
|---------------------------------|----------------------------------------|-------------------------------|---------|-------------------|---------------------------------------|
| (1955MC75) | (1973GO03, 1980WA24) | | | | |
| 0 | | | 1 | $\frac{1}{2}^-$ | 0.58 ± 0.04^c 1.1 ^d |
| 3.09 | 3.089443 ± 0.020^g | | 0 | $\frac{1}{2}^+$ | 0.36 ± 0.02^c 1.1 ^d |
| 3.68 | 3.684482 ± 0.023^g | | 1 | $\frac{3}{2}^-$ | 0.10 ^d |
| 3.85 | 3.853783 ± 0.022^g | | 2 | $\frac{5}{2}^+$ | 1.1 ^d |
| 6.86 | | | 2 | $\frac{5}{2}^+$ | 0.04 ^d |
| 7.470 ± 20 | | | | | |
| 7.533 ± 20 | | | | | |
| 7.641 ± 20 | | 70 ± 15 | | | |
| 8.4 ± 300 | 8.25 ^f 8.86 ^f | 1100 ± 300 | 2 | $\frac{3}{2}^+$ | 1.0 ^{d,e} |
| 9.500 ± 20 | | | 1 | $\frac{1}{2}^-$ | 0.5 ^{d,e} |
| 9.897 ± 20 | | | (1) | $(\frac{3}{2}^-)$ | |
| 10.759 ± 20 | | | 1 | $\frac{3}{2}^-$ | 0.1 ^{d,e} |
| | 10.755 ± 5 | 56 ± 2 | | | |
| | 10.818 ± 5 | 24 ± 3 | | | |
| | 10.997 ± 8 | 82 ± 15 | | | |
| | 11.080 ± 5 | 8 | | | |
| | 11.748 ± 10 | 107 ± 14 | | | |
| | 11.851 ± 5 | 68 ± 4 | | | |
| | 11.97 ± 40^a | ≈ 260 | | | |
| | 12.108 ± 5 | 81 ± 8 | | | |

^a May correspond to unresolved states.

^b See (1970AJ04, 1973DA17).

^c (1972PE11).

^d (1973DA17) [HD parameters]. See also Table 13.14 in (1970AJ04).

^e $\Gamma_{\text{d,p}}/\Gamma_n = 0.68, 0.91$ and 2.2 for $^{13}\text{C}^*(8.2, 8.86, 9.90)$, respectively (1973DA17).

^f Nominal energies (1973DA17).

^g From E_γ measurements (1980WA24). See also (1975TR07).

The angular distributions for the transitions to $^{13}\text{C}^*(0, 3.85)$ have been studied at $E(^3\text{He}) = 40$ MeV (1971ST21). The proton decay of various ^{14}N states to $^{13}\text{C}_{\text{g.s.}}$ has been studied by (1974NO01): see ^{14}N .

$$39. \ ^{12}\text{C}(\alpha, ^3\text{He})^{13}\text{C} \quad Q_{\text{m}} = -15.6314$$

Angular distributions of the ^3He particles have been measured at $E_{\alpha} = 56$ MeV (1969GA11; to $^{13}\text{C}^*(0, 3.68 + 3.85)$), 104 MeV (1972HA08; to $^{13}\text{C}(0)$) and 139 MeV (1973SM03; to $^{13}\text{C}^*(0, 3.85)$). A detailed comparison has been made between the angular distributions of the ground state ^3He particles and of the tritons from the mirror reaction $^{12}\text{C}(\alpha, \text{t})^{13}\text{N}$ (1969GA11, 1972HA08).

$$40. \ ^{12}\text{C}(^7\text{Li}, ^6\text{Li})^{13}\text{C} \quad Q_{\text{m}} = -2.304$$

Angular distributions at $E(^7\text{Li}) = 48$ MeV to $^{13}\text{C}^*(0, 3.09, 3.68, 3.85)$ have been analyzed and yield spectroscopic factors of 0.80, 0.44, 0.17 and 0.74 for the first four states of ^{13}C (1979ZE01). See (1973SC26) for earlier work at $E(^7\text{Li}) = 36$ MeV.

$$41. \ ^{12}\text{C}(^{11}\text{B}, ^{10}\text{B})^{13}\text{C} \quad Q_{\text{m}} = -6.5088$$

See (1974AN36) and (1976AJ04).

$$42. \ ^{12}\text{C}(^{12}\text{C}, ^{11}\text{C})^{13}\text{C} \quad Q_{\text{m}} = -13.775$$

Angular distributions at $E(^{12}\text{C}) = 93.8$ MeV have been measured for the transitions to $^{13}\text{C}^*(0, 3.85)$ (1979FU04). See also (1979HEZX) and (1976AJ04).

$$43. \ ^{12}\text{C}(^{14}\text{N}, ^{13}\text{N})^{13}\text{C} \quad Q_{\text{m}} = -5.607$$

Angular distributions have been reported at $E(^{14}\text{N}) = 78, 100$ and 154.8 MeV involving $^{13}\text{C}^*(0, 3.85)$ [and at 100 MeV $^{13}\text{C}^*(3.09, 7.3 \pm 0.3)$]: see (1976AJ04) and (1976NA09). More recently angular distributions have been studied at $E(^{14}\text{N}) = 28, 32, 34$ and 36 MeV (1976BA16). The ratio of the differential cross section to $^{13}\text{C}_{\text{g.s.}}$ and that to $^{13}\text{N}_{\text{g.s.}}$ (observed via $^{12}\text{C}(^{14}\text{N}, ^{13}\text{C})^{13}\text{N}$), as a function of θ is approximately 1 as expected if mirror symmetry holds (1969VO01, 1970VO02, 1975VO05). See also (1978MA1F) and (1977WE1H, 1978DZ1A, 1978NA15, 1978WE1F, 1979DO13; theor.).

44. $^{12}\text{C}(^{16}\text{O}, ^{15}\text{O})^{13}\text{C}$ $Q_m = -10.7175$

Angular distributions are reported at $E(^{16}\text{O}) = 128$ MeV to $^{13}\text{C}^*(0, 3.85)$ (1979PR07).

45. (a) $^{12}\text{C}(^{17}\text{O}, ^{16}\text{O})^{13}\text{C}$ $Q_m = 0.802$

(b) $^{12}\text{C}(^{18}\text{O}, ^{17}\text{O})^{13}\text{C}$ $Q_m = -3.098$

Angular distributions have been studied at $E(^{17}\text{O}) = 30.5$ and 33.8 MeV (to $^{13}\text{C}^*(0, 3.09)$) and at $E(^{18}\text{O}) = 32.3$ and 35 MeV (to $^{13}\text{C}_{\text{g.s.}}$) (1978CH16).

46. $^{12}\text{C}(^{19}\text{F}, ^{18}\text{F})^{13}\text{C}$ $Q_m = -5.4849$

See (1970AJ04).

47. $^{13}\text{B}(\beta^-)^{13}\text{C}$ $Q_m = 13.437$

See ^{13}B and Table 13.2.

48. $^{13}\text{C}(\gamma, \gamma)^{13}\text{C}$

For τ_m studied by resonance scattering see Table 13.5.

49. (a) $^{13}\text{C}(\gamma, n)^{12}\text{C}$ $Q_m = -4.94639$

(b) $^{13}\text{C}(\gamma, 2n)^{11}\text{C}$ $Q_m = -23.668$

The main features of the cross section are a sharp peak corresponding to the $T = \frac{3}{2}$ state $^{13}\text{C}^*(15.11)$ [$\Gamma_{\gamma_0} = 19.7 \pm 2.0$ eV], the broad pigmy resonance at $E_x = 13$ MeV [on which peaks are superimposed at $E_x = 11.0, 13.8, 16.5$ and 17.8 MeV] and the giant resonance at $E_x = 24$ MeV ($\sigma_{\text{max}} = 9.5$ mb) [surrounded by shoulder resonances at $E_x = 20.8$ and ≈ 30 MeV, both of which appear to decay substantially to highly excited states of ^{12}C]. There is also some evidence for a weak resonance at ≈ 37 MeV superimposed on the high energy tail of the GDR (1979JU01: monoenergetic photons). A recent study of the angular distributions of n_0 suggests states at $E_x = 7.70$ ($\frac{3}{2}^+$), 7.95 ($\frac{3}{2}^+$), 8.95 ($(\frac{1}{2}^-)$), 10.0 ($(\frac{3}{2}^-)$), 11.0 ($(\frac{1}{2}^+)$) and 12.05 MeV ($(\frac{3}{2}^+)$) (1979WO06). Γ_n and Γ_{γ_0} deduced from (γ, n_0) measurements are displayed in Tables 13.12 and

Table 13.15: Electromagnetic transitions ^a in ¹³C from ¹³C(e, e')¹³C ^b

| E_x (MeV \pm keV) | J^π | Mult. | Γ_{γ_0} (eV) | $\Gamma_{\gamma_0}/\Gamma_W$ (W.u.) |
|------------------------------|-----------------|-------|----------------------------------|-------------------------------------|
| 3.08 \pm 30 ^g | $\frac{1}{2}^+$ | E1 | 0.68 \pm 0.23 | 0.062 |
| 3.69 \pm 20 ^g | $\frac{3}{2}^-$ | E2 | $(3.61 \pm 0.40) \times 10^{-3}$ | 3.52 |
| 3.85 ^g | | M1 | 0.358 \pm 0.047 | 0.339 |
| 6.85 \pm 60 ^g | $\frac{5}{2}^+$ | M2 | $(6.9 \pm 3.6) \times 10^{-5}$ | 0.055 |
| 7.54 \pm 20 | $\frac{5}{2}^-$ | M3 | (1.01×10^{-5}) | (35) |
| | | E2 | 0.115 \pm 0.007 ^e | 3.15 |
| 8.86 \pm 20 ^{d,g} | $\frac{1}{2}^-$ | M1 | 3.36 \pm 0.47 ^f | 0.230 |
| | | E0 | 2.09 ^c | |
| 9.90 \pm 30 | $\frac{3}{2}^-$ | E2 | $(6.3 \pm 1.1) \times 10^{-3}$ | 0.045 |
| | | M1 | 0.324 \pm 0.038 | 0.0159 |
| 11.07 \pm 20 | $\frac{1}{2}^-$ | M1 | 1.02 \pm 0.12 | 0.0359 |
| | | E0 | 2.62 ^c | |
| | $\frac{3}{2}^-$ | E2 | 0.256 \pm 0.047 | 1.03 |
| | | M1 | 0.172 \pm 0.020 | 0.006 |
| 15.11 \pm 20 | $\frac{3}{2}^-$ | E2 | 0.6 \pm 0.1 | 0.50 |
| | | M1 | 22.4 \pm 1.5 | 0.31 |

^a See also Tables 13.6, 13.7 and 13.16.

^b (1969WI22, 1970WI04). (1978SO1B, 1979CR1C, 1979CRZZ) also report the population of ¹³C*(3.85, 9.50).

^c Monopole matrix element in fm².

^d $\Gamma = 190 \pm 35$ keV.

^e 0.11 \pm 0.015 eV (1980HO11; ¹³C(γ , n)).

^f 5.4 \pm 0.5 eV (1980HO11); Γ_{γ_0} for ¹³C*(7.69, 8.2) are reported to be 0.6 \pm 0.1 and 7.0 \pm 0.9 eV, respectively (1980HO11).

^g Observed by (1980SO1B).

13.15 (1980HO11). For other differential cross section measurements derived from bremsstrahlung measurements see (1977WO04; n_0 , 6.0 to 37 MeV; n_1 , 10.5 to 35.5 MeV). For reports of other structures see (1976AJ04) and (1976KO22). See also (1977AL18; theor.). For reaction (b) see (1979JU01) and (1976AJ04). See also (1976BE1H, 1976CR1C), (1977DA1B, 1979BE1X) and (1977HO32, 1977MA06, 1979HO17, 1979JOZP; theor.).

$$50. {}^{13}\text{C}(\gamma, p){}^{12}\text{B} \quad Q_m = -17.533$$

(1964DE12) report structures at $E_\gamma = 18.5, 20.0, 23.5, 26.0$ and 29.0 MeV. The main part of the cross section is in the 23.5 MeV peak which has $\Gamma \approx 3$ MeV. A broad maximum near 25.5 MeV has been reported by (1956CO72, 1957CO57). The cross section to ${}^{12}\text{B}^*(0.95)$ shows a broad maximum at $E_\gamma = 25$ MeV (1975PA09). See also (1977HO32, 1977MA06, 1979HO17; theor.).

$$51. {}^{13}\text{C}(\gamma, \alpha){}^9\text{Be} \quad Q_m = -10.6479$$

See (1970AJ04).

$$52. \text{(a) } {}^{13}\text{C}(e, e'){}^{13}\text{C} \\ \text{(b) } {}^{13}\text{C}(e, ep){}^{12}\text{B} \quad Q_m = -17.533$$

The elastic scattering has been studied for $E_e = 120$ to 750 MeV (1970HE24). A number of inelastic groups are seen: see Tables 13.15 and 13.16 (1970WI04, 1971BE51, 1980SO1B). A distinct splitting of the giant resonance into two large peaks near $E_x = 20.5$ and 24.5 MeV, with widths of ≈ 3 and ≈ 4 MeV, respectively, is observed. It is suggested that these are groupings of narrower peaks [see Table 13.16]. The $E_x = 20.5$ and 24.5 MeV resonances are probably $T = \frac{1}{2}$ and $T = \frac{3}{2}$, although the 4 MeV splitting is somewhat smaller than expected (1971BE51). See also reactions 49 and 50. For radius measurements see (1976AJ04) and the “GENERAL” section here. See also (1977AL1H, 1977AL1J, 1979KI1J, 1979SE06, 1980ER01; theor.).

At $E_e = 43$ MeV the energy and angular distributions of the protons emitted in reaction (b) suggest the involvement of ${}^{13}\text{C}^*(21.8?, 24.2, 25.5, 27.5, 31.5)$ (1971SH09).

$$53. \text{(a) } {}^{13}\text{C}(\pi^-, \pi^-){}^{13}\text{C} \\ \text{(b) } {}^{13}\text{C}(\pi^+, \pi^+){}^{13}\text{C}$$

Table 13.16: States of ^{13}C from $^{13}\text{C}(e, e')^{13}\text{C}^*$ (1980SO1B)^a

| E_x (MeV \pm keV) | Γ (keV) | E_x (MeV \pm keV) | Γ (keV) |
|----------------------------|-----------------|-----------------------|----------------|
| 9.500 ± 7 ^b | | 18.699 ± 5 | 98 ± 11 |
| 11.845 ± 5 | 144 ± 5 | 20.021 ± 13 | 232 ± 27 |
| 12.187 ± 10 | 109 ± 48 | 20.429 ± 8 | 112 ± 23 |
| 12.438 ± 12 | 160 ± 37 | 21.466 ± 8 | 268 ± 14 |
| 14.390 ± 15 | 281 ± 65 | 22.2 | 1100 |
| 14.582 ± 10 | 227 ± 41 | 24.7 | 600 |
| 14.983 ± 10 | 380 ± 53 | 25.5 | 500 |
| 15.526 ± 11 | 147 ± 23 | 27.3 | 600 |
| 16.080 ± 7 | 148 ± 13 | 28.1 | 500 |
| (16.183 ± 28) | (40 ± 20) | (29.4) | 1200 |
| (18.497 ± 10) | (91 ± 23) | | |

^a $E_e = 165.8, 221.5$ and 276.8 MeV. See also Tables 13.4 and 13.15.

^b Data are consistent with $J^\pi = \frac{9}{2}^+$.

^c (1971BE34).

The elastic scattering of π^+ has been studied at $E_{\pi^+} = 48.6$ MeV by (1978DY01). At $E_\pi = 162$ MeV the inelastic transition to $^{13}\text{C}^*(9.5)$ is very strongly enhanced in π^- scattering. The enhancement factor $\sigma(\pi^-)/\sigma(\pi^+) \approx 9$ is consistent with a pure neutron transition and a $\frac{9}{2}^+$ assignment. Also enhanced in π^- scattering (but less strongly) are $^{13}\text{C}^*(0, 3.09, 3.68 + 3.85), (7.49), ^{13}\text{C}^*(7.55)$ and a group of states close to 16 MeV are enhanced in π^+ scattering. The centroid of a group of states near 22 MeV is found in $(\pi^-, \pi^{-'})$ at a higher excitation energy than in $(\pi^+, \pi^{+'})$ (1979DE34). At $E_\pi = 180$ MeV angular distributions are reported for the transitions to $^{13}\text{C}^*(0, 3.68, 7.55, 9.5, 11.7, 15.0, 17.5)$: the transitions to $^{13}\text{C}^*(9.5)$ is enhanced by a factor of $4.0_{-0.7}^{+2.0}$ (1979SC25) in π^- scattering. Preliminary work is also reported for the inelastic scattering to $^{13}\text{C}^*(16.1, 21.4)$ [large M4 strength] (1980TR1B). See also the “GENERAL” section here.

54. $^{13}\text{C}(n, n)^{13}\text{C}$

Angular distributions have been measured at $E_n = 1.6$ to 6.6 MeV for the n_0 group, and for the n_1, n_2 and n_3 groups at several energies in that range (1978LA1H, 1979KOZU; abstracts). See also (1977NO07; theor.) and ^{14}C .

55. (a) $^{13}\text{C}(p, p)^{13}\text{C}$



Angular distributions have been studied at $E_p = 1.37$ to 30.4 MeV [see (1976AJ04)] and at $E_p = 6$ MeV (1975DE26, 1977SA1B; p_0, p_1, p_3), $E_p = 10.0$ to 17.5 MeV (1978WE13; p_0), $E_p = 35.2$ MeV (1980FA07; p_0), 135 MeV (1980RI1A; to $^{13}\text{C}^*(9.50)$), 800 MeV (1978BL02; to $^{13}\text{C}^*(0, 3.09, 3.68, 6.86, 7.55, 11.92 \pm 0.06)$) and 1 GeV (1979AL26; p_0). The work of (1978BL02) suggests $J^\pi = (\frac{5}{2}, \frac{7}{2})^+$ for $^{13}\text{C}^*(11.92)$. Inelastic scattering to $^{13}\text{C}^*(21.3, 21.8)$, the former with $\Gamma \approx 0.3$ MeV [may be due to unresolved states] is observed at $E_p = 135$ MeV (1980DE1L; prelim.). Microscopic analyses of the angular distribution to $^{13}\text{C}^*(9.50)$ [$J^\pi = \frac{9}{2}^+$] show that the transition is an essentially pure $1p_{3/2}$ to $1d_{5/2}$ neutron transition with a total angular momentum transfer of 4, consistent with the results of the pion scattering (see reaction 53) (1980RI1A).

Reaction (b) has been studied at $E_p = 7.9$ to 12 MeV (1971OT02: one or more of the ^{13}C states at 7.5 MeV seem to be involved) and at $E_p = 46$ MeV (1974MI05). See also ^{14}N , (1977BA2G, 1978AL36) and (1977AM1C; theor.).



Angular distributions have been measured at $E_d = 0.71$ to 15 MeV [see (1976AJ04)] and at $E_d = 13$ MeV (1978DA17; $d_0 \rightarrow d_3$) and $E_d = 13.6$ MeV (1975ZA08; d_0). See also ^{15}N and (1979DO03; theor.).



For reaction (a) see (1980DE1M). Angular distributions of elastically scattered ^3He have been studied at $E(^3\text{He}) = 12$ to 40 MeV [see (1976AJ04)], at $18, 20$ and 24 MeV (1976MA26), at 20 MeV (1977PE23) and at 41 MeV (1980TR02). Angular distributions at $E(^3\text{He}) = 40$ MeV have also been reported to ^{13}C states at $E_x = 3.09, 3.68, 3.85, 6.86, 7.55 \pm 0.03, 8.86 \pm 0.03$ and 11.84 ± 0.03 MeV. ^3He groups to $^{13}\text{C}^*(9.50 \pm 0.03)$ and (15.11) are also reported: the latter is weak (1969BA06). At $E(^3\text{He}) = 43.6$ MeV angular distributions to $^{13}\text{C}^*(9.498)$ and to $^{13}\text{N}^*(8.918)$ [$J^\pi = \frac{1}{2}^-$] (in the $(^3\text{He}, t)$ reaction) are identical but the cross section in the latter reaction is 1.25 ± 0.15 , rather than the factor of 2 expected from isospin symmetry (1980PE1E; prelim.). See also (1980DE1M) and (1977FU1G, 1977FU1H; theor.).



Elastic angular distributions have been measured at $E_\alpha = 15$ to 40.5 MeV. In the range $E_\alpha = 24$ to 40.5 MeV angular distributions have been reported to $^{13}\text{C}^*(3.09, 3.68 + 3.85, 6.86, 7.5)$. At $E_\alpha = 22$ MeV, $E_x = 7686$ keV, $\Gamma_{\text{cm}} = 70 \pm 5$ keV (1980FU04; also line shapes). (1976WO11) report $S_\alpha = 0.67$ for $^{12}\text{C}_{\text{g.s.}}$. The excitation of $^4\text{He}^*(20.1)$ [$J^\pi = 0^+$] has been studied at $E_\alpha = 65$ MeV (1976JA17). See also (1976PA25, 1977DM1A, 1977SA19, 1978ZE03; theor.).

59. (a) $^{13}\text{C}(^6\text{Li}, ^6\text{Li})^{13}\text{C}$
 (b) $^{13}\text{C}(^7\text{Li}, ^7\text{Li})^{13}\text{C}$

Angular distributions of elastically scattered Li ions have been studied at $E(\text{Li}) = 4.5$ to 34 MeV [see (1976AJ04)], at $E(\bar{\text{Li}}) = 9$ MeV (1978DR07) and at $E(^6\text{Li}) = 40$ MeV (1979ZE01). See also (1971SC21) and (1976ST22, 1979SU1F; theor.).

60. $^{13}\text{C}(^9\text{Be}, ^9\text{Be})^{13}\text{C}$

The elastic scattering has been studied at $E(^{13}\text{C}) = 28.1$ and 36.2 MeV (1979JA04, 1979UN01).

61. (a) $^{13}\text{C}(^{12}\text{C}, ^{12}\text{C})^{13}\text{C}$
 (b) $^{13}\text{C}(^{13}\text{C}, ^{13}\text{C})^{13}\text{C}$
 (c) $^{13}\text{C}(^{14}\text{C}, ^{14}\text{C})^{13}\text{C}$

Angular distributions for reaction (a) have been studied at $E(^{12}\text{C}) = 10$ to 87 MeV [see (1976AJ04)], 15 MeV (1980VO05; elastic, and to $^{13}\text{C}^*(3.09, 3.85)$), 20 to 35.5 MeV (1978CH29; elastic, and to $^{13}\text{C}^*(3.09, 3.68 + 3.85)$ from 30 MeV), 80 MeV (1979TA1E; elastic), and for the elastic scattering at $E(^{13}\text{C}) = 12$ MeV (1976WE28, 1977GU07) and 36 MeV (1976WE21). The spectroscopic factor for $^{13}\text{C}_{\text{g.s.}} = 0.81 \pm 0.04$ (1977GU07). For fusion studies see (1976ST12, 1979KO20). See also ^{12}C in (1980AJ01), (1978LE1N), (1977WI1C), (1978RO1D; astrophys.) and (1977TR1A, 1978AV1A, 1978CH30, 1978PA1B, 1978TA1B, 1979IM02, 1979GR20, 1979ZE1B; theor.).

For reaction (b) elastic angular distributions have been measured at $E(^{13}\text{C}) = 15$ to 24 MeV (1973HE12), while for reaction (c) the elastic distribution has been measured at $E(^{13}\text{C}) = 15$ MeV (1972BO68). For a fusion study see (1980CH01). See also (1978TE02, 1980SC1D; theor.).

62. $^{13}\text{C}(^{14}\text{N}, ^{14}\text{N})^{13}\text{C}$

The elastic angular distribution has been studied at $E(^{14}\text{N}) = 19.3$ MeV (1971VO01). See also (1976AJ04) and (1980WI09) for fusion measurements.

63. (a) $^{13}\text{C}(^{16}\text{O}, ^{16}\text{O})^{13}\text{C}$
(b) $^{13}\text{C}(^{17}\text{O}, ^{17}\text{O})^{13}\text{C}$
(c) $^{13}\text{C}(^{18}\text{O}, ^{18}\text{O})^{13}\text{C}$

Elastic angular distributions have been measured for reaction (a) at $E(^{16}\text{O}) = 10$ to 30 MeV [see (1976AJ04)] and 14 MeV (1976DU04), and at $E(^{13}\text{C}) = 36$ MeV (1976WE21) and 105 MeV (1979BR03). For fusion measurements see (1979CH07). Elastic scattering distributions in reaction (b) are reported at $E(^{17}\text{O}) = 29.8$ and 32.3 MeV (1977CH22, 1978CH03; see also ^{17}O in (1982AJ01)). For fusion studies see (1980WI09). Elastic scattering in reaction (c) has been studied at $E(^{18}\text{O}) = 15, 20$ and 24 MeV (1971KN05) and 31 MeV (1978CH03). For a fusion study see (1978CO07). See also (1979GAZV) and (1978SC1E, 1979PAZY; theor.).

64. $^{13}\text{C}(^{24}\text{Mg}, ^{24}\text{Mg})^{13}\text{C}$

See (1979DU10).

65. $^{13}\text{C}(^{28}\text{Si}, ^{28}\text{Si})^{13}\text{C}$

Backward elastic angular distributions are reported at $E(^{28}\text{Si}) = 77.6$ to 94.6 MeV (1979OS01). See also (1975RA33, 1979DE1N).

66. $^{13}\text{C}(^{32}\text{S}, ^{32}\text{S})^{13}\text{C}$

See (1975RA33).

67. (a) $^{13}\text{C}(^{40}\text{Ca}, ^{40}\text{Ca})^{13}\text{C}$
(b) $^{13}\text{C}(^{48}\text{Ca}, ^{48}\text{Ca})^{13}\text{C}$

See (1978HO1C) and (1977FU1K, 1977UD1A, 1978BA26, 1978PE1E, 1979SA27; theor.).

68. $^{13}\text{N}(\beta^+)^{13}\text{C}$ $Q_m = 2.221$

See ^{13}N .

69. $^{14}\text{C}(\text{p}, \text{d})^{13}\text{C}$ $Q_m = -5.9519$

Angular distributions have been measured at $E_p = 12$ to 18.5 MeV [see (1976AJ04)] and at 27 MeV (1975CE04; $\text{d}_0 \rightarrow \text{d}_3$). The spectroscopic factors for the first four states of ^{13}C are 1.4 , 0.02 , 1.8 and 0.13 , respectively (1975CE04).

70. $^{14}\text{C}(\text{d}, \text{t})^{13}\text{C}$ $Q_m = -1.9192$

Angular distributions for $\text{t}_0 \rightarrow \text{t}_3$ have been studied at $E_d = 12$ MeV (1966GL01) and $E_d = 14$ MeV (1976WE01). The spectroscopic factors for $^{13}\text{C}^*(0, 3.09, 3.68, 3.85)$ are 1.00 , 0.06 , 1.0 and 0.08 , respectively (1976WE01). See also ^{16}N in (1982AJ01).

71. $^{14}\text{C}(^3\text{He}, \alpha)^{13}\text{C}$ $Q_m = 12.4013$

Angular distributions have been measured at $E(^3\text{He}) = 3$ to 4.5 MeV (α_0, α_1) and 3 to 6.6 MeV (α_2, α_3) (1972KE08), at $6, 8$ and 10 MeV (1971CO14; α_0) and 44.8 MeV (1966BA13; α_0, α_2 and α to $^{13}\text{C}^*(15.11)$). See also ^{17}O .

72. $^{14}\text{C}(^{11}\text{B}, ^{12}\text{B})^{13}\text{C}$ $Q_m = -4.807$

See (1976AJ04).

73. $^{14}\text{C}(^{12}\text{C}, ^{13}\text{C})^{13}\text{C}$ $Q_m = -3.2302$

See (1976AJ04).

74. $^{14}\text{C}(^{14}\text{N}, ^{13}\text{C})^{15}\text{N}$ $Q_m = 2.6568$

See (1976AJ04).

75. $^{14}\text{C}(^{16}\text{O}, ^{17}\text{O})^{13}\text{C}$ $Q_m = -4.0322$

Angular distributions have been measured at $E(^{16}\text{O}) = 20, 25$ and 30 MeV (1975SC35, 1975SC42): they are well described by DWBA (fixed range approximation). See also (1977GO1D, 1977WE1H; theor.).

76. $^{14}\text{C}(^{18}\text{O}, ^{19}\text{O})^{13}\text{C}$ $Q_m = -4.220$

See (1976AJ04).

77. (a) $^{14}\text{N}(\gamma, \text{p})^{13}\text{C}$ $Q_m = -7.5506$
 (b) $^{14}\text{N}(\gamma, \text{np})^{12}\text{C}$ $Q_m = -12.4970$

For $E_{\text{bs}} = 15.5$ to 29.5 MeV a large fraction of the neutron yield appears to be associated with sequential decay to ^{12}C via $^{13}\text{C}^*(7.75, 8.86, 11.80)$ (1972GE11, 1970SH06). Angular distributions measured in the giant resonance region of ^{14}N are consistent with the proton decay of $(\text{p}_{1/2})^{-1}(2\text{s}1\text{d})$ giant dipole states to $^{13}\text{C}_{\text{g.s.}}$ and of $(\text{p}_{3/2})^{-1}(2\text{s}1\text{d})$ states to $^{13}\text{C}^*(3.68)$ (1974BA37). The population of $^{13}\text{C}^*(3.09, 3.85)$ has also been reported (1970TH01; $E_{\text{bs}} = 29$ MeV). See also ^{14}N .

78. $^{14}\text{N}(\text{n}, \text{d})^{13}\text{C}$ $Q_m = -5.3260$

Angular distributions of d_0 have been determined at $E_n = 14.1$ to 14.7 MeV: see (1970AJ04). Differential cross sections for formation of $^{13}\text{C}^*(3.09, 3.68)$ have been measured for $E_n = 10.1$ to 11.0 MeV (1970DI1A). See also ^{15}N and (1976KI1D, 1978NE1B).

79. (a) $^{14}\text{N}(\text{e}, \text{ep})^{13}\text{C}$ $Q_m = -7.5506$
 (b) $^{14}\text{N}(\text{p}, 2\text{p})^{13}\text{C}$ $Q_m = -7.5506$

For reaction (a) see (1969BA1F). At $E_p = 46$ MeV, the summed proton spectrum shows transitions to $^{13}\text{C}^*(0, 3.68, 7.5, 11.9)$ (1970WE09). See also (1976AJ04) and (1977KO1P, 1979MA20; theor.).

80. $^{14}\text{N}(\text{d}, ^3\text{He})^{13}\text{C}$ $Q_m = -2.0571$

At $E_d = 52$ MeV, angular distributions have been measured for the ^3He particles to $^{13}\text{C}^*(0, 3.09, 3.68, 7.5, 8.86, 9.50, 11.9 \pm 0.15)$ and analyzed by DWBA: $J^\pi = \frac{5}{2}^-, \frac{1}{2}^-, \frac{3}{2}^-$ and $\frac{3}{2}^-$, respectively, are assigned to $^{13}\text{C}^*(7.5, 8.86, 9.50, 11.9)$ (1968HI01). As expected, angular distributions of ^3He and of tritons (from $^{14}\text{N}(d, t)^{13}\text{N}$) to analog states are closely the same: this has been shown for the ground state ^3He and triton groups (1966DE1C, 1968GA13: $E_d = 28.5$ MeV; (1974LU06: $E_d = 15$ MeV) and for the groups to $^{13}\text{C}^*(8.9 + 9.5)$ and $^{13}\text{N}^*(9.2)$ (1968HI01; $E_d = 52$ MeV). See also ^{16}O in (1982AJ01).

$$81. \ ^{14}\text{N}(t, \alpha)^{13}\text{C} \quad Q_m = 12.2634$$

Observed particle groups at $E_t = 2.6$ MeV are displayed in Table 13.22 of (1976AJ04). See also ^{16}O in (1977AJ02).

$$82. \ ^{14}\text{N}(\alpha, \alpha p)^{13}\text{C} \quad Q_m = -7.5506$$

See ^{14}N .

$$83. \ ^{14}\text{N}(^6\text{Li}, ^7\text{Be})^{13}\text{C} \quad Q_m = -1.944$$

An angular distribution has been obtained at $E(^6\text{Li}) = 32$ MeV for the transition to $^{13}\text{C}_{g.s.}$ and $^7\text{Be}^*(0, 0.43)$ [for the mirror reaction see reaction 40 in ^{13}N]: the relative cross sections show a deviation from isospin symmetry which is attributed to Coulomb effects. $^{13}\text{C}^*(3.09)$ was also populated (1971GR44).

$$84. \ ^{14}\text{N}(^{14}\text{N}, ^{15}\text{O})^{13}\text{C} \quad Q_m = -0.2536$$

See (1976AJ04).

$$85. \ ^{15}\text{N}(\gamma, d)^{13}\text{C} \quad Q_m = -16.1594$$

See (1979SK04) and ^{15}N .

$$86. \ ^{15}\text{N}(n, t)^{13}\text{C} \quad Q_m = -9.9020$$

Table 13.17: Energy levels of ^{13}C from $^{15}\text{N}(p, ^3\text{He})^{13}\text{C}$ (1968FL03)

| E_x in ^{13}C (MeV \pm keV) | J^π |
|------------------------------------------|-------------------|
| 0 | $\frac{1}{2}^-$ |
| 3.08 ± 20 | $\frac{1}{2}^+$ |
| 3.68^a | $\frac{3}{2}^-$ |
| 6.86^a | $\frac{5}{2}^+$ |
| 7.55 ± 20 | $\frac{5}{2}^-$ |
| 8.86 ± 60 | $\frac{1}{2}^-$ |
| 9.52 ± 30 | $(\frac{3}{2}^-)$ |
| 11.09 ± 50 | $(\frac{1}{2}^-)$ |
| 11.80 ± 30 | $(\frac{3}{2}^-)$ |
| 12.40 ± 50 | $\frac{7}{2}^-$ |
| 15.103 ± 45^b | $\frac{3}{2}^-$ |

^a Observed but E_x not determined.

^b (1966CE02): see Table 13.7.

See (1978BA1M) and ^{16}N in (1982AJ01).

$$87. \ ^{15}\text{N}(p, ^3\text{He})^{13}\text{C} \quad Q_m = -10.6658$$

At $E_p = 43.7$ MeV ^3He groups have been observed to eleven states of ^{13}C : see Table 13.17 (1966CE02, 1968FL03). For a summary of the analysis see (1976AJ04). A study of both angular distributions and analyzing powers in this reaction and in the mirror (p, t) reaction shows that DWBA generally fails to predict the analyzing powers (1974MA12; $E_{\bar{p}} = 43.8$ MeV). Angular distributions have also been measured for the ^3He groups to $^{13}\text{C}^*(0, 3.68, 7.55)$ at several energies in the range $E_p = 24.0$ to 43.5 MeV (1974PI05, 1975MI01). See also ^{16}O in (1982AJ01).

$$88. \ ^{15}\text{N}(d, \alpha)^{13}\text{C} \quad Q_m = 7.6874$$

For α_0 angular distributions see (1970AJ04). $^{13}\text{C}^*(3.09, 3.68, 8.80)$ are also populated: see (1976AJ04).

89. $^{15}\text{N}(\alpha, ^6\text{Li})^{13}\text{C}$ $Q_m = -14.6859$

At $E_\alpha = 42$ MeV, the angular distribution of the ^6Li particles to $^{13}\text{C}_{\text{g.s.}}$ has been measured (1968MI05).

90. $^{15}\text{N}(^{11}\text{B}, ^{13}\text{C})^{13}\text{C}$ $Q_m = 2.5193$

See (1976AJ04).

91. $^{16}\text{O}(\text{n}, \alpha)^{13}\text{C}$ $Q_m = -2.2156$

Angular distributions have been measured for E_n to 18.8 MeV for $\alpha_0, \alpha_1, \alpha_{2+3}$: see (1970AJ04, 1976AJ04). See also (1976NO1F, 1979SUZW).

92. $^{16}\text{O}(\alpha, ^7\text{Be})^{13}\text{C}$ $Q_m = -21.207$

At $E_\alpha = 42$ MeV the angular distribution of the reaction involving $^{13}\text{C}(0)$ and $^7\text{Be}^*(0 + 0.43)$ has been measured (1972RU03).

93. $^{16}\text{O}(^{14}\text{N}, ^{17}\text{F})^{13}\text{C}$ $Q_m = -6.9503$

This reaction has been studied at $E(^{14}\text{N}) = 76$ and 79 MeV (1976MO03, 1977MO1A) and 155 MeV (1975NA15, 1976NA09). See also (1979MO14).

94. $^{16}\text{O}(^{17}\text{O}, ^{20}\text{Ne})^{13}\text{C}$ $Q_m = -1.629$

At $E(^{17}\text{O}) = 35$ MeV angular distributions have been measured involving $^{13}\text{C}_{\text{g.s.}}$ and $^{20}\text{Ne}^*(0, 1.63)$ (1977KA26).

95. (a) $^{17}\text{O}(\text{d}, ^6\text{Li})^{13}\text{C}$ $Q_m = -4.886$

(b) $^{18}\text{O}(\text{d}, ^7\text{Li})^{13}\text{C}$ $Q_m = -5.680$

See (1976AJ04).

96. $^{20}\text{Ne}(\text{n}, 2\alpha)^{13}\text{C}$ $Q_m = -6.9465$

See (1976AJ04).

^{13}N
(Figs. 3 and 4)

GENERAL: (See also (1976AJ04).)

Shell model: (1977TE01).

Collective, rotational and deformed models: (1976BR26, 1977ME1E).

Special levels: (1977ME1E, 1977RI08, 1977TE01, 1979RO1E, 1980BA54).

Electromagnetic transitions: (1977ME1E, 1977YO1D, 1978KR19, 1980BA54).

Applied work: (1975PA1F, 1977JO1C, 1977ME1D, 1977ST1M, 1978FI1F, 1978HI1D, 1978PE1F, 1978TI1A, 1978TI1B, 1978WI1F, 1978WO1C, 1979DE1H, 1979LI1G, 1979WI1G).

Special reactions: (1976AB04, 1976BE1K, 1976BU16, 1976LE1F, 1976MA62, 1977AR06, 1977AS03, 1977SC1G, 1978GE1C, 1978HE1C, 1979KA07, 1980MI01).

Astrophysical questions: (1976DE1G, 1977BA1V, 1977SI1D, 1979PE1E).

Pion capture and reactions (See also reaction 29.): (1975KA1G, 1976AU1E, 1976BU1D, 1976ED1A, 1976GA06, 1976GA21, 1976LI04, 1976NI05, 1976OS06, 1977AU01, 1977BA2H, 1977FU10, 1977LO19, 1977MA1M, 1977WA02, 1978WA02, 1979AL1J, 1979ALZW, 1979GI1D, 1979HI1A, 1979LA22, 1979MO1N, 1979MO1P, 1979MO1Q, 1979SA21, 1980BO03, 1980CH08, 1980LE02, 1980LI1E, 1980SA04).

Other topics: (1976BR26, 1978SH1B, 1979BE1H, 1979RO1E).

Ground state of ^{13}N : (1976BR26, 1976FU06, 1976JO1B, 1977YO1D).

$$\mu = -0.32224 \pm 0.00035 \text{ nm (1964BE24, 1978LEZA).}$$

1. $^{13}\text{N}(\beta^+)^{13}\text{C}$ $Q_m = 2.2206$

The weighted mean of the three measurements of $\tau_{1/2} = 9.965 \pm 0.005$ min (1960JA12), 9.963 ± 0.009 min (1968RI15) and 9.965 ± 0.010 min (1977AZ01) is 9.965 ± 0.004 min. See also Table 13.22 of (1970AJ04). The decay is entirely to $^{13}\text{C}_{\text{g.s.}}$: $\log ft = 3.667 \pm 0.001$ from the tables of (1971GO40). The positrons are completely polarized: see (1970AJ04). See also (1978MA1P), (1976AJ04, 1977BA48, 1978RA2A), (1979DA1D; astrophys.) and (1977AZ02, 1977RI08, 1977YO1E, 1979DE15; theor.).

2. $^{10}\text{B}(^3\text{He}, \gamma)^{13}\text{N}$ $Q_m = 21.637$

Table 13.18: Energy levels of ^{13}N

| E_x (MeV \pm keV) | $J^\pi; T$ | Γ_{cm} (keV) | Decay | Reactions |
|--------------------------|------------------------------|-------------------------------------|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| g.s. | $\frac{1}{2}^-; \frac{1}{2}$ | $\tau_{1/2} = 9.965 \pm 0.004$ min | β^+ | 1, 2, 8, 10, 11, 12, 13, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 51 |
| 2.3648 ± 1.1 | $\frac{1}{2}^+$ | $\Gamma_{\text{cm}} = 33.7 \pm 0.9$ | γ, p | 8, 11, 13, 19, 20, 27, 30, 31, 37, 39, 40, 46 |
| 3.511 ± 2 | $\frac{3}{2}^-$ | 62 ± 4 | γ, p | 2, 8, 10, 11, 13, 19, 20, 22, 25, 30, 31, 32, 34, 37, 38, 39, 44, 46 |
| 3.547 ± 4 | $\frac{5}{2}^+$ | 47 ± 7 | p | 2, 8, 10, 11, 14, 19, 20, 22, 25, 27, 30, 31, 32, 37, 39, 44 |
| 6.364 ± 9 | $\frac{5}{2}^+$ | 11 | p | 9, 11, 14, 20, 31, 39, 44 |
| 6.886 ± 8 | $\frac{3}{2}^+$ | 115 ± 5 | p | 9, 11, 14, 20, 39 |
| 7.155 ± 5 | $\frac{7}{2}^+$ | 9 ± 0.5 | p | 9, 11, 14, 20, 31, 39 |
| 7.376 ± 9 | $\frac{5}{2}^-$ | 75 ± 5 | p | 9, 10, 11, 14, 20, 31, 34, 37, 38, 39, 44 |
| 7.9 | $\frac{3}{2}^+$ | ≈ 1500 | p | 14, 20, 39 |
| 8.918 ± 11 | $\frac{1}{2}^-$ | 230 | p | 11, 14, 31, 34, 37, 38, 39, 44 |
| 9.00 | $(\frac{9}{2}^+)$ | 280 ± 30 | | 9, 38 |
| 9.476 ± 8 | $\frac{3}{2}^-$ | 30 | p | 9, 11, 14, 31, 34, 38 |
| 10.25 ± 150 | $(\frac{1}{2}^+)$ | ≈ 280 | γ, p | 13 |
| 10.36 | $\frac{5}{2}^-$ | 30 | p | 9, 11, 14, 20, 34 |
| 10.36 | $\frac{7}{2}^-$ | 76 | p | 9, 14, 20 |
| 10.833 ± 9 | $(\frac{1}{2}^-)$ | | | 9, 11, 31, 44 |
| 11.530 ± 12 | $\frac{5}{2}^+$ | 430 ± 35 | p | 9, 11, 14 |
| 11.70 ± 30 | $\frac{5}{2}^-$ | 115 ± 30 | p | 14 |
| 11.74 ± 40 | $\frac{3}{2}^+$ | 240 ± 30 | γ, p | 13, 14 |
| 11.74 ± 50 | $\frac{3}{2}^-$ | 530 ± 80 | p | 14 |
| 11.878 ± 12 | $(\frac{3}{2}^-)$ | 380 ± 50 | p | 10, 11, 14, 31, 37, 38, 39, 44 |
| 12.13 ± 50 | $\frac{7}{2}^-$ | 250 ± 30 | p | 14, 20, 46 |
| 12.558 ± 23 | | > 400 | | 11 |
| 12.937 ± 24 | | > 400 | | 11 |
| 13.5 ± 200 | $\frac{3}{2}^+$ | ≈ 6500 | γ, p | 13, 14 |
| 14.05 ± 20 | $\frac{3}{2}^+; \frac{1}{2}$ | 165 ± 20 | γ, p, α | 13, 14, 17 |
| 15.0645 ± 1.0^a | $\frac{3}{2}^-; \frac{3}{2}$ | 0.86 ± 0.12 | γ, p, α | 11, 13, 14, 17, 31, 44 |

Table 13.18: Energy levels of ^{13}N (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | Γ_{cm} (keV) | Decay | Reactions |
|--------------------------|-------------------------------------------------|-------------------------------|----------------------------------------|-------------|
| 15.3 \pm 200 | $(\frac{3}{2}^+)$ | 350 \pm 150 | γ, p | 13 |
| 15.99 \pm 30 | $\frac{7}{2}^+; \frac{1}{2}$ | 135 \pm 90 | p, α | 14, 17, 31 |
| 16.0 | | \approx 500 | p | 14 |
| 17.5 | | | γ, p | 13, 14 |
| 18.15 \pm 30 | $\frac{3}{2}^+; \frac{1}{2}$ | 320 \pm 80 | p | 14 |
| 18.17 \pm 20 | $\frac{1}{2}^-; \frac{1}{2}$ | 225 \pm 50 | p, α | 14, 17 |
| 18.406 \pm 5 | $\frac{3}{2}^+; \frac{3}{2}$ | 66 \pm 8 | p, α | 11, 14, 17 |
| 18.961 \pm 10 | $\frac{3}{2}^-$ or $\frac{7}{2}^+; \frac{3}{2}$ | 23 \pm 5 | p, α | 11, 14, 17 |
| 19.83 | $\frac{5}{2}^-; \frac{1}{2}$ | 1000 | p, α | 14, 17 |
| 19.88 | $\frac{7}{2}^+; \frac{1}{2}$ | 750 | p | 14 |
| 20.2 | $\frac{5}{2}^-$ | 1000 | p | 14 |
| 20.9 \pm 300 | $\frac{1}{2}^+$ | 1200 | γ, p | 13, 14 |
| 21.4 | $\frac{3}{2}^-$ | 750 | p | 14 |
| 21.7 | $\frac{3}{2}^+$ | | p | 14 |
| 22.4 \pm 500 | $\frac{1}{2}^+$ | | p | 14 |
| 23 | | | γ, p | 13 |
| 23.3 | | 400 | $p, {}^3\text{He}$ | 4 |
| 23.83 \pm 50 | | 350 \pm 50 | $p, {}^3\text{He}$ | 4, 13 |
| 24.4 | | 700 | $p, {}^3\text{He}$ | 4, 13, 14 |
| (24.6) | | 120 | $p, {}^3\text{He}$ | 4 |
| 25.6 \pm 100 | $(\frac{3}{2})^-$ | 240 \pm 80 | $p, {}^3\text{He}$ | 4, 14 |
| 25.9 | | 1000 | $(n), p, d, {}^3\text{He}, \alpha$ | 3, 4, 7, 14 |
| 26.84 | | | p | 14 |
| 28 | | | $(\gamma), p, {}^3\text{He}, (\alpha)$ | 2, 4, 7 |
| (31) | | | p | 14 |
| 32 | | \approx 2000 | $\gamma, d, {}^3\text{He}, \alpha$ | 2, 5, 7, 13 |

^a See also Table 13.7.

The 90° cross sections for γ_0 and γ_{2+3} have been measured for $E({}^3\text{He}) = 4.8$ to 14 MeV: no pronounced structures are observed (1972BE05). See also (1976AJ04) and (1974LO1B).

3. $^{10}\text{B}({}^3\text{He}, n)^{12}\text{N}$

$$Q_m = 1.574$$

$$E_b = 21.637$$

Activation cross sections have been reported for $E({}^3\text{He}) = 1$ to 30.6 MeV: there is some evidence of broad structures. See (1976AJ04), (1974LO1B) and ^{12}N in (1980AJ01).

Table 13.19: Structure in $^{10}\text{B} + ^3\text{He}$

| E_{res} (MeV \pm keV) | Γ (keV) | Res. in | $^{13}\text{N}^*$ (MeV) | Refs. |
|-------------------------------------|-------------------|-------------------------------------------------------------------|----------------------------|--------------------------------|
| 2.1 ^a | 500 | $p_0, (p_1)$ | 23.3 | (1956SC01, 1966PA10) |
| 2.85 ± 50 | 450 ± 50 | $\gamma_{15.1}$ | 23.83 | (1964KU09) |
| 3.6 ^a | 700 | p_0, p_1 | 24.4 | (1956SC01, 1966PA10) |
| 3.9 | 120 | p_0 | 24.6 | (1956SC01, 1966PA10) |
| 4.6 ^a | 150 | $p_0, (p_1)$ | 25.2 | (1956SC01) |
| 5.2 ± 100 ^b | 240 ± 80 | $p_0, \gamma_{15.1}, p_2, p_3$ | 25.6 | (1964KU09) |
| 5.6 | 1000 ^d | $(n), p_0, p_2, p_3, \gamma_{12.7}, \gamma_{15.1}, d_0, \alpha_0$ | 25.9 | (1965PA05, 1972BE05, 1972BE56) |
| 8.5 | ^e | $(\gamma_0), p_0, \gamma_{12.7}, \gamma_{15.1}, (\alpha_0)$ | 28 | (1972BE05, 1972BE56) |
| 13.5 ^c | ≈ 2000 | $(\gamma_0), d_{4+5}, \alpha_1$ | 32 | (1972BE05, 1972BE56) |

^a See, however, (1964KU09).

^b See, however, (1966BA01, 1966PA10).

^c This may correspond to more than one state (1972BE56).

^d $J \geq \frac{3}{2}$ (1965PA10, 1966PA10).

^e $J \geq \frac{7}{2}$ (1966PA10).



$$Q_m = 19.6940$$

$$E_b = 21.637$$


Observed resonances in the yields of proton groups and of 12.7 and 15.1 MeV γ -rays are displayed in Table 13.19. For polarization measurements see (1970AJ04) and (1976LA1F). See also (1974LO1B) and ^{12}C in (1980AJ01).



$$Q_m = 3.197$$

$$E_b = 21.637$$

Activation cross sections (1976GA27) and excitation functions have been measured for $E(^3\text{He}) = 1$ to 19 MeV: for resonances see Table 13.19. See also (1974LO1B) and ^{11}C in (1980AJ01).



$$Q_m = -3.670$$

$$E_b = 21.637$$



See (1976AJ04) and ^{10}B and ^{10}C in (1979AJ01).

$$7. \text{}^{10}\text{B}(\text{}^3\text{He}, \alpha)\text{}^9\text{B} \quad Q_m = 12.142 \quad E_b = 21.637$$

Excitation functions for α_0 and α_1 have been measured for $E(^3\text{He}) = 2$ to 19 MeV. Observed resonances are displayed in Table 13.19. See also (1974LO1B) and ^9B in (1979AJ01).

$$8. \text{}^{10}\text{B}(\alpha, n)\text{}^{13}\text{N} \quad Q_m = 1.060$$

Angular distributions have been measured for $E_\alpha = 1.5$ to 4.6 MeV (n_0) and ≈ 4.6 MeV (n_1, n_{2+3}) (1973VA25), and at 18.0, 19.0 and 20.2 MeV (1976DU08; n_0, n_1). See also ^{14}N and (1976EP1A; astrophys.).

$$9. \text{}^{10}\text{B}(\text{}^6\text{Li}, t)\text{}^{13}\text{N} \quad Q_m = 5.843$$

At $E(^6\text{Li}) = 18$ MeV the known states of ^{13}N with $6.3 < E_x < 11.7$ MeV are observed, with the exception of $^{13}\text{N}^*(7.9, 8.92)$. In addition, evidence is presented for a ^{13}N state at $E_x = 9.00$ MeV with $\Gamma_{\text{cm}} = 280 \pm 30$ keV: it is very strongly excited and its angular distribution is similar to that for $^{13}\text{C}^*(9.50)$ in the mirror reaction ($^6\text{Li}, ^3\text{He}$) suggesting that these two states are analogs. Other analog assignments made on the basis of corresponding intensities in the mirror reaction are given in reaction 14 of ^{13}C . The widths of $^{13}\text{N}^*(6.89, 7.38)$ are, respectively, 120 ± 30 and 70 ± 30 keV (1974HO06). See also (1970AJ04).

$$10. \text{}^{10}\text{B}(\text{}^{14}\text{N}, \text{}^{11}\text{B})\text{}^{13}\text{N} \quad Q_m = 0.902$$

At $E(^{10}\text{B}) = 100$ MeV, angular distributions are reported for the transitions to $^{13}\text{N}^*(0, 3.5, 7.4, 11.9)$ (1975NA15). See also (1976NA09, 1977MO1A) and (1976AJ04).

$$11. \text{}^{11}\text{B}(\text{}^3\text{He}, n)\text{}^{13}\text{N} \quad Q_m = 10.182$$

Neutron groups have been observed to a number of states of ^{13}N : see Table 13.20. The parameters of the first $T = \frac{3}{2}$ state at $E_x = 15.06$ MeV are displayed in Table 13.7 where they are compared with the corresponding quantities for $^{13}\text{C}^*(15.11)$ (1973AD02, 1975MA21, 1977MA16, 1979AD01). Angular distributions for n_0 have recently been reported at $E(^3\text{He}) = 1.70$ and 1.90 MeV (1979OS10) and 5.2 to 12 MeV (1977DA18). See also ^{14}N and (1977OS08; theor.).

$$12. \text{}^{11}\text{B}(\text{}^{11}\text{B}, \text{}^9\text{Li})\text{}^{13}\text{N} \quad Q_m = -12.965$$

Table 13.20: States of ^{13}N from $^{11}\text{B}(^3\text{He}, n)^{13}\text{N}$

| E_x^a (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | L^a | $J\pi^a$ |
|----------------------------|---------------------------------|-------|----------------------------------|
| 0 | | 2 | $\frac{1}{2}^-$ |
| 2.358 ± 10 | | 1 | $\frac{1}{2}^+$ |
| 3.502 ± 10 | | 0, 2 | $\frac{3}{2}^-$ |
| 3.55 ± 18 | | | |
| 6.353 ± 9 | | 1, 3 | $\frac{5}{2}^+$ |
| 6.875 ± 10 | | 1, 3 | $\frac{3}{2}^+$ |
| 7.145 ± 9 | | 3, 5 | $\frac{7}{2}^+$ |
| 7.363 ± 8 | | 2, 4 | $\frac{5}{2}^-$ |
| 8.2 ± 22 | | | |
| 8.918 ± 11 | | | |
| 9.476 ± 8 | | 0, 2 | $\frac{3}{2}^-$ |
| 10.381 ± 8 | | 2, 4 | $\frac{5}{2}^-$ |
| 10.833 ± 9 | | | |
| 11.530 ± 12 | | | |
| 11.878 ± 12 | | 0, 2 | $\frac{3}{2}^-$ |
| 12.558 ± 23 | > 400 | | |
| 12.937 ± 24 | > 400 | | |
| $15.068 \pm 8^{b,c}$ | < 15 | | $\frac{3}{2}^-; T = \frac{3}{2}$ |
| 18.44 ± 40^b | | | $T = \frac{3}{2}$ |
| 18.98 ± 20^b | 40 ± 20 | | $T = \frac{3}{2}$ |

^a (1971HS03) except for those states labeled ^b; $E(^3\text{He}) = 4.7, 6.1$ and 6.49 MeV.

^b (1969AD02): $E(^3\text{He}) = 7.0$ to 13.5 MeV.

^c See also Table 13.7.

Table 13.21: Resonances in $^{12}\text{C}(p, \gamma)^{13}\text{N}$ ^a

| E_p (MeV \pm keV) | Γ_{cm} (keV) | Γ_{γ_0} (eV) | $^{13}\text{N}^*$ (MeV) | Res. in yield of | J^π | Refs. |
|-----------------------------|-------------------------------|------------------------------|----------------------------|--------------------------|----------------------------------|--------------------------------|
| 0.4568 \pm 0.5 | 33.7 \pm 0.9 ^b | 0.45 \pm 0.05 ^c | 2.3648 \pm 0.0011 | γ_0 | $\frac{1}{2}^+$ | ^a |
| 1.699 \pm 2 | 62 \pm 4 | 0.64 ^j | 3.511 | γ_0 | $\frac{3}{2}^-$ | ^a , (1974RO29) |
| 9.01 \pm 150 | \approx 280 | ^d | 10.25 | γ_0 | $(\frac{1}{2}^+)$ | (1976BE28) |
| 10.62 \pm 120 | 200 \pm 50 | \approx 4.2 ^e | 11.74 | γ_0 | $\frac{3}{2}^+$ | (1973ME12) |
| 12.5 \pm 200 | 6500 | \geq 1100 | 13.5 | γ_0 | $\frac{3}{2}^+$ | (1973ME12) |
| 13.12 \pm 90 | 160 \pm 20 | 3.7 \pm 1.0 ^f | 14.04 | γ_0 | $\frac{3}{2}^+$ | (1973ME12) |
| 14.2 | [see Table 13.7] | | 15.0 | γ_0, γ_{2+3} | $\frac{3}{2}^-; T = \frac{3}{2}$ | (1968DI04, 1975MA21, 1977MA16) |
| 14.5 \pm 200 ^g | 350 \pm 140 | \geq 0.5 | 15.3 | γ_1 | $(\frac{3}{2}^+)$ | (1973ME12) |
| 16.9 | | | 17.5 | γ_0 | | (1976BE28) |
| 20 ^h | | | 20 | γ_1, γ_{2+3} | | (1976BE28) |
| 20.5 ⁱ | \approx 3700 | | 20.8 | γ_0 | | (1963FI07, 1976BE28, 1976FE1C) |
| 23 | | | 23 | γ_0 | | (1976BE28) |
| 24.5 | | | 24.5 | γ_{2+3} | | (1963FI07) |
| 32.5 | broad | | 31.9 | γ_0, γ_{2+3} | | (1963FI07, 1976FE1C) |

^a See also Table 13.26 in (1976AJ04) and (1980BA54; theor.).

^b Weighted mean of values from (1968RI16, 1974BL06).

^c (1968RI16).

^d See (1973ME12).

^e A value of 0.30 ± 0.05 is assumed for Γ_{p_0}/Γ : see Table 13.22.

^f A value of 126 keV is taken for Γ_{p_0} (1969LE18).

^g This peak may be due to an unresolved doublet.

^h Giant resonance for γ_1 .

ⁱ Main dipole strength is concentrated in this peak (1976BE28).

^j Recalculated on basis of total $\Gamma_{\text{lab}} = 67 \pm 4$ keV. I am indebted to Prof. F.C. Barker for his comments [see (1980BA54)].

See (1974AN36).

13. (a) $^{12}\text{C}(p, \gamma)^{13}\text{N}$ $Q_m = 1.9434$
 (b) $^{12}\text{C}(p, \gamma p')^{12}\text{C}$
 $Q_0 = 1943.31 \pm 0.32 \text{ keV}$ (1977FR20);
 $Q_0 = 1944.01 \pm 0.22 \text{ keV}$ (1977HE26).

Resonances for capture radiation are displayed in Table 13.21. No resonance is observed at $E_p = 1.73 \text{ MeV}$ [$^{13}\text{N}^*(3.55)$]: $\omega\Gamma_\gamma < 0.006 \text{ eV}$ (1963YO06).

Excitation functions have been measured for $E_p = 150$ to 2500 keV . In addition to the first two resonances, direct radiative capture is observed. In reaction (b), studied for $E_p = 610$ to 2700 keV the capture γ -ray yield is dominated by a direct capture process to $^{13}\text{N}^*(2.36)$. The cascade decay $^{13}\text{N}^*(3.51 \rightarrow 2.36)$ has an intensity of $8 \pm 1\%$ (1974RO29). Extrapolating the cross section to $E_{\text{cm}} = 25 \text{ keV}$ yields a cross section factor $S = 1.45 \pm 0.20 \text{ keV} \cdot \text{b}$ (1974RO29). A reanalysis of the data by (1980BA54) suggests $S = 1.54_{-0.10}^{+0.15} \text{ keV} \cdot \text{b}$. For other discussions of astrophysical questions see (1976AJ04) and (1975ZIIA, 1976BR1H, 1976RO1Q).

At $E_p = 14.2 \text{ MeV}$, capture radiation from the first $T = \frac{3}{2}$ state, $^{12}\text{N}^*(15.06)$, is reported: see Table 13.7 for the parameters and the decay modes of this state (1968DI04, 1975MA21, 1977MA16). The angular distributions of the γ -rays determine $J = \frac{3}{2}$ for $^{13}\text{N}^*(15.06)$ (1968DI04).

Excitation functions for γ -rays have also been measured at $E_p = 8.7$ to 24.4 MeV (γ_0) and 19.9 to 24.4 MeV (γ_1, γ_{2+3}) (1976BE28) and at 16 to 37 MeV (γ_0) and 23 to 37 MeV (γ_{2+3}) (1976FE1C). Angular distributions of γ_0 are reported for $16.5 < E_p < 27.0 \text{ MeV}$ by (1980WO1A; prelim.). At $E_p = 40$ to 100 MeV most of the γ strength is due to transitions to $^{13}\text{N}^*(3.5)$, probably to $^{13}\text{N}^*(3.55)$ [$J^\pi = \frac{5}{2}^+$] because of its single particle character. Transitions to higher states may also be indicated (1979KO05, 1979AR1G). Differential cross sections for the transitions to the ground state have been measured for $E_p = 10$ to 17 MeV . The total E2 capture cross section are $\approx 0.2 \mu\text{b}$ and no resonance effects are observed. The E2 energy-weighted sum rule depleted over this energy range is $(8.5 \pm 3.3)\%$ (1980HE04). The interference between the M1(E2) $T = \frac{3}{2}$ resonance ($^{13}\text{N}^*(15.06)$) and the E1 GDR has been studied by (1980SN01): the E1 capture is found to be predominantly d-wave (1980SN01). At $E_p = 28.5 \text{ MeV}$, the capture is dominated by γ_{2+3} but γ_0 and a transition to $^{13}\text{N}^*(8.1)$ are also observed (1980BL1B).

See also reaction 14, (1976AJ04), (1976TSZZ, 1977TS1A, 1979BL1F), (1974LO1B, 1979SN1A), (1977YO1F, 1978GO1F, 1979RO12; applications) and (1977AL18, 1977BA29, 1979JOZP; theor.).

14. (a) $^{12}\text{C}(p, p)^{12}\text{C}$ $E_b = 1.9434$
 (b) $^{12}\text{C}(p, 2p)^{11}\text{B}$ $Q_m = -15.9569$
 (c) $^{12}\text{C}(p, p\alpha)^8\text{Be}$ $Q_m = -7.3667$
 (d) $^{12}\text{C}(p, 3p)^{10}\text{Be}$ $Q_m = -27.1857$
 (e) $^{12}\text{C}(p, \pi^+)^{13}\text{C}$ $Q_m = -135.403$

Table 13.22: ^{13}N levels from $^{12}\text{C}(p, p)$, $^{12}\text{C}(p, p')$ and $^{12}\text{C}(p, \alpha)$ ^a

| E_p (MeV \pm keV) | $^{13}\text{N}^*$ (MeV) | Γ_{cm} (keV) | l_p | J^π | | Refs. |
|--------------------------|----------------------------|-------------------------------|-------|----------------------------------|---------------------|----------------------|
| | | | | | d | |
| 0.461 ± 3 | 2.369^b | 31^b | 0 | $\frac{1}{2}^+$ | $\theta^2 = 0.54$ | A, (1979KR18) |
| 1.686 ± 6 | $3.499^{b,c}$ | 63^b | 1 | $\frac{3}{2}^-$ | 0.031 | A |
| 1.734 ± 6 | $3.543^{b,c}$ | 74^b | 2 | $\frac{5}{2}^+$ | 0.21 | A |
| 4.808 ± 10 | 6.378 | 11 | 2 | $\frac{5}{2}^+$ | 0.0031 | A |
| 5.370 ± 10 | 6.896 | 115 ± 5 | 2 | $\frac{3}{2}^+$ | 0.13 | A |
| 5.65 ± 10 | 7.155 | 9 ± 0.5 | 4 | $\frac{7}{2}^+$ | 0.016 | A |
| 5.891 | 7.38 | 75 ± 5 | 3 | $\frac{5}{2}^-$ | 0.069 | A |
| 6.5 | 7.9 | ≈ 1500 | 2 | $\frac{3}{2}^+$ | 0.14 | A |
| 7.54 | 8.90 | 230 | 1 | $\frac{1}{2}^-$ | 0.02 | A |
| 8.18 | 9.49 | 30 | 1 | $\frac{3}{2}^-$ | 0.001 | A |
| 9.13 | 10.36 | 30 | 3 | $\frac{5}{2}^-$ | | A |
| 9.13 | 10.36 | 76 | 3 | $\frac{7}{2}^-$ | | A |
| | | | | | $\Gamma_p/\Gamma =$ | |
| 10.35 ± 50 | 11.49 | 430 ± 35 | 2 | $\frac{5}{2}^+$ | 0.70 ± 0.05 | A |
| 10.58 ± 30 | 11.70 | 115 ± 30 | 3 | $\frac{5}{2}^-$ | 0.60 ± 0.04 | A |
| 10.62 ± 40 | 11.74 | 250 ± 30 | 2 | $\frac{3}{2}^+$ | 0.30 ± 0.05 | A |
| 10.62 ± 50 | 11.74 | 530 ± 80 | 1 | $\frac{3}{2}^-$ | 0.55 ± 0.05 | A |
| 10.75 ± 40 | 11.86 | 380 ± 50 | 0 | $\frac{1}{2}^+$ | 0.35 ± 0.05 | A |
| 11.05 ± 50 | 12.13 | 250 ± 30 | 3 | $\frac{7}{2}^-$ | 0.30 ± 0.05 | (1973ME03) |
| 12.5 | 13.5 | ≈ 500 | | | | (1961NA02) |
| 13.13 ± 20 | 14.05 | 180 ± 35 | 2 | $\frac{3}{2}^+; T = \frac{1}{2}$ | 0.29 ± 0.07 | (1976ME18, 1969LE18) |

Table 13.22: ^{13}N levels from $^{12}\text{C}(\text{p}, \text{p})$, $^{12}\text{C}(\text{p}, \text{p}')$ and $^{12}\text{C}(\text{p}, \alpha)$ ^a (continued)

| E_p (MeV \pm keV) | $^{13}\text{N}^*$ (MeV) | Γ_{cm} (keV) | l_p | J^π | | Refs. |
|-----------------------------|----------------------------|--------------------------------|-------|--------------------------------------------------------|-----------------|---------------------------------------------|
| a 14.23075 \pm 0.2 | 15.06447 ^e | 0.932 \pm 0.028 ¹ | 1 | $\frac{3}{2}^-$; $T = \frac{3}{2}$ | | A, Table 13.7, (1980TH05) |
| 15.24 \pm 40 ^f | 15.99 | 135 \pm 90 | 4 | $\frac{7}{2}^+$; $T = \frac{1}{2}$ | 0.05 \pm 0.04 | (1976ME18, 1969LE18) |
| 15.2 | 16.0 | \approx 500 | | | | A |
| 16.8 | 17.4 | | | | f | (1976BE28) |
| 17.58 \pm 30 | 18.15 | 322 \pm 75 | 2 | $\frac{3}{2}^+$; $T = \frac{1}{2}$ | 0.08 \pm 0.02 | (1976ME18) |
| 17.60 \pm 20 | 18.17 | 225 \pm 50 | 1 | $\frac{1}{2}^-$; $T = \frac{1}{2}$ | 0.24 \pm 0.06 | (1976ME18, 1969LE18) |
| 17.857 \pm 5 | 18.406 | 66 \pm 8 | 2 | $\frac{3}{2}^+$; $T = \frac{3}{2}$ | g | A, (1976BE28, 1969LE18) |
| a 18.460 \pm 10 | 18.961 | 23 \pm 5 | | $\frac{3}{2}^-$ or $\frac{7}{2}^+$; $T = \frac{3}{2}$ | g | A, (1976BE28, 1969LE18) |
| a 19.40 ^j | 19.83 | 1000 | 3 | $\frac{5}{2}^-$; $T = \frac{1}{2}$ | | A, (1969LE18) |
| 19.46 | 19.88 | 750 | 4 | $\frac{7}{2}^+$; $T = \frac{1}{2}$ | | A, (1979GA13) |
| 19.8 | 20.2 | 1000 | | $\frac{5}{2}^-$ | g | (1976BE28, 1979GA13) |
| 20.6 \pm 300 | 20.9 | 1200 | | $\frac{1}{2}^+$ | f,g | (1973ME12, 1976BE28, 1979GA13) ⁱ |
| 21.1 | 21.4 | 750 | | $\frac{5}{2}^-$ | | (1979GA13) |
| 21.4 | 21.7 | | | $\frac{3}{2}^+$ | | A, (1979GA13) |
| 22.2 \pm 500 | 22.4 | \approx 1000 | | $\frac{1}{2}^+$ | | A, (1979GA13) |
| k 24.0 | 24.1 | \lesssim 500 | | | | A |
| 25.7 | 25.6 | | | $(\frac{3}{2})^-$ | | A |
| 27.02 | 26.84 | | | | | A |

Table 13.22: ^{13}N levels from $^{12}\text{C}(p, p)$, $^{12}\text{C}(p, p')$ and $^{12}\text{C}(p, \alpha)$ ^a (continued)

| E_p (MeV \pm keV) | $^{13}\text{N}^*$ (MeV) | Γ_{cm} (keV) | l_p | J^π | | Refs. |
|--------------------------|----------------------------|-------------------------------|-------|---------|---|------------|
| 32 | 31 | | | | j | (1976FE1C) |

A: See references listed in Table 13.27 in (1976AJ04).

^a See also Tables 13.25 in (1970AJ04) and 13.27 in (1976AJ04).

^b An R -matrix analysis by (1976ME22) leads to $E_x = 2.367, 3.501$ and 3.547 MeV, and $\Gamma_{\text{cm}} = 33, 55$ and 50 keV for these states. (1976ME22) also find that $^{13}\text{N}_{\text{g.s.}}$ has an appreciable effect on the low energy scattering.

^c See also (1976MA55).

^d A dispersion analysis leads to a spectroscopic factor of 0.53 ± 0.08 for $^{13}\text{N}_{\text{g.s.}}$. (1977ME05).

^e See also (1976IK01).

^f Resonance in yield of 12.7 MeV γ -rays (1976BE28).

^g Resonance in yield of 15.1 MeV γ -rays (1976BE28).

^h See however (1976BE28).

ⁱ See also (1976FE1C).

^j Resonance in yield of 4.4 MeV γ -rays (1976FE1C).

^k A $\frac{3}{2}^+$ state is indicated in this region by the work of (1976GA27).

^l $\Gamma_p = 263 \pm 15$ eV (1980TH05).

Yield curves for elastic protons, protons inelastically scattered to $^{12}\text{C}^*(4.4, 7.7, 9.6, 12.7, 15.1)$ and for γ -rays from $^{12}\text{C}^*(4.4, 12.7, 15.1)$ have been studied at many energies: see Table 13.22 for a display of the characteristics of the observed structure. Recent yield measurements include those of (1976ME22; $E_p = 0.3$ to 2.0 MeV), (1976BE28; 13.8 \rightarrow 24.4 MeV (12.7 MeV γ), 16.4 \rightarrow 24.4 MeV (15.1 MeV γ)), (1976FE1C; 16 \rightarrow 37 (4.4 MeV γ), 16 \rightarrow 31 (12.7 MeV γ), 17 \rightarrow 37 MeV (15.1 MeV γ)) and 22 to 27 MeV (1980HO07; (12.7 and 15.1 MeV γ)).

A phase shift analysis of the elastic scattering analyzing power for $E_{\bar{p}} = 11.5$ to 18.1 MeV shows four $T = \frac{1}{2}$ states with $E_x = 14.06, 16.00, 18.16$ and 18.18 MeV, with $J^\pi = \frac{3}{2}^+, \frac{7}{2}^+, \frac{3}{2}^+, \frac{1}{2}^-$: see Table 13.22 (1976ME18). At $E_{\bar{p}} = 19.15$ to 23.34 MeV, measurements of the elastic group and the protons to $^{12}\text{C}^*(4.4, 12.7)$ locate $\frac{1}{2}^+$ (E1), $\frac{5}{2}^-$ (E2) and $\frac{7}{2}^+$ (E3) resonances below 21 MeV, $\frac{3}{2}^+$ (E1) and $\frac{5}{2}^-$ resonances with $21 < E_x < 22$ MeV and $\frac{1}{2}^+$ and $\frac{3}{2}^+$ resonances above 22 MeV: see Table 13.22 (1979GA13). Elastic polarization measurements are also reported at $E_{\bar{p}} = 0.45$ to 0.60 MeV (1979KR18; study of Mott-Schwinger interaction), $E_p = 7.16$ to 7.43 MeV (1977ME06), $E_{\bar{p}} \approx 14.2$ MeV (1980WIZW), 36.2 MeV (1978BE49), 39.9 to 75.0 MeV (1980KA02), 52.3 to 68.2 MeV (1977EG1A), $E_p = 185$ MeV (1979IN01; $^{12}\text{C}^*(0, 4.4, 7.7, 9.6)$), $E_{\bar{p}} = 800$ MeV (1978HO05, 1978RA17) and $E_p = 1$ GeV (1977AL25, 1980AL09). Polarization measurements involving $^{12}\text{C}^*(4.4, 12.71, 15.11, 15.4, 16.1)$ are reported by (1978HO11) at $E_{\bar{p}} = 50$ and 65 MeV: a marked difference in the analyzing powers for the groups to the 1^+ states $^{12}\text{C}^*(12.71, 15.11)$ is observed. At $E_{\bar{p}} = 800$ MeV analyzing powers have been measured for $^{12}\text{C}^*(12.7, 15.1, 18.3, 19.4)$: those to $^{12}\text{C}^*(12.7, 18.3)$ are negative. The values of A_y appear to be characteristic of the isospin transfer (1980MO06). For other polarization measurements see (1979KA1N; $E_{\bar{p}} = 65$ MeV; p_0, p_2), (1979FU1H; $E_{\bar{p}} = 24.1, 26.2, 28.0$ MeV; p_1 ; measured spin flip probability), (1979BE44; $E_{\bar{p}} = 299, 380, 483$ and 561 MeV), (1979LI03; $^{12}\text{C}^*(4.4)$; $E_{\bar{p}} = 800$ MeV), (1977NA16, 1977NA29, 1978NA18: (p, 2p); $E_{\bar{p}} = 635$ MeV) and the older measurements [$E_p = 0.5$ MeV to 3.6 GeV] listed in Tables 13.26 in (1970AJ04) and 13.28 in (1976AJ04). See also (1978SE01).

For total cross section measurements see (1976AJ04) and (1979SC07; $E_p = 191$ to 550 MeV) and (1978JA16; $E_p = 0.87$ GeV/ c and 2.10 GeV/ c). See also (1979DE31). For reaction (b) see ^{11}B in (1980AJ01) and (1979KO21). For reaction (c) see ^8Be in (1979AJ01). For reaction (d) see (1979KO36).

Bremsstrahlung measurements are reported by (1978TR05) near the $E_p = 1.7$ MeV resonance and by (1976MA06, 1976MA55, 1977MU1C, 1979LE1G, 1979LE1H). See also (1979JA17, 1980LI01; theor.).

Inclusive cross sections have been measured by (1978FR12, 1979FR12, 1979KO21). For polarization measurements at $E_{\bar{p}} = 316$ and 516 MeV see (1978KA08). Studies of hadron multiple production are reported by (1977AV01, 1978AR1J, 1978AZ02). Spallation measurements have been carried out by (1976RO12, 1977FO04, 1978WE1D, 1979RA20). See also (1979VD02). The production of π^+ with $E_{\bar{p}} = 200$ MeV (reaction (e)) is reported by (1978AU07).

See also ^{12}C in (1980AJ01), (1975DE26, 1976CU08, 1976PL1C, 1976WE1E, 1976WE1F, 1976WUZZ, 1977BE1Y, 1977SA1B, 1978DE1Q, 1978GO05, 1979TR1D, 1980CO05, 1980WH1A), (1977AU1B, 1978DW1A, 1979RA1C; astophys.) and (1976AH09, 1976ES1B, 1976GO20, 1976JE1A,

1977BA29, 1976BR1L, 1977PH02, 1977WE1G, 1978AH04, 1978BA21, 1978BE66, 1978BH1B, 1978CA03, 1978CO1J, 1978FA04, 1978GR1D, 1978GU12, 1978RA20, 1978RE1D, 1978UC1A, 1978WU03, 1979AB13, 1979AM02, 1979DE1P, 1979ER02, 1979HE16, 1979LO12, 1979OS06, 1979PH05, 1979RA27, 1979RO1E, 1979TH1A, 1979WE1C, 1979YA1E, 1980AL12, 1980BR04, 1980CO06, 1980JA05, 1980RA1B; theor.).

| | | |
|----------------------------------------------------------|--------------------------|-------------------------|
| 15. (a) $^{12}\text{C}(\text{p}, \text{n})^{12}\text{N}$ | $Q_{\text{m}} = -18.120$ | $E_{\text{b}} = 1.9434$ |
| (b) $^{12}\text{C}(\text{p}, \text{pn})^{11}\text{C}$ | $Q_{\text{m}} = -18.721$ | |

The cross section for reaction (a) has been measured from threshold to $E_{\text{p}} = 50$ MeV. Resonant structure is observed corresponding to $E_{\text{x}} = 21, 24$ and , possibly ≈ 27 MeV (1968RI01). See also (1976RO10) and ^{12}N in (1980AJ01).

Cross sections for reaction (b) have been measured to 300 GeV: see (1976AJ04). At $E_{\text{p}} = 800$ MeV, $\sigma = 32.0 \pm 1.0$ mb (1976HO19). See also (1976BE1K).

| | | |
|----------------------------------------------------------|---------------------------|-------------------------|
| 16. (a) $^{12}\text{C}(\text{p}, \text{d})^{11}\text{C}$ | $Q_{\text{m}} = -16.497$ | $E_{\text{b}} = 1.9434$ |
| (b) $^{12}\text{C}(\text{p}, \text{t})^{10}\text{C}$ | $Q_{\text{m}} = -23.364$ | |
| (c) $^{12}\text{C}(\text{p}, ^3\text{He})^{10}\text{B}$ | $Q_{\text{m}} = -19.6940$ | |

For polarization measurements [reaction (a)] see (1970AJ04) and (1977HA1N; $E_{\text{p}} = 27.4$ MeV; d_0, d_1) and (1979HO1H; $E_{\text{p}} = 65$ MeV; d_0, d_1). See also ^{11}C in (1980AJ01).

Polarization measurements involving the t_0 and t_1 groups have been carried out at $E_{\text{p}} = 49.5$ MeV (1970NE17). See also $^{10}\text{B}, ^{10}\text{C}$ in (1979AJ01).

| | | |
|-------------------------------------------------|-------------------------|-------------------------|
| 17. $^{12}\text{C}(\text{p}, \alpha)^9\text{B}$ | $Q_{\text{m}} = -7.552$ | $E_{\text{b}} = 1.9434$ |
|-------------------------------------------------|-------------------------|-------------------------|

Yield curves for α_0 have been measured over the 14.2 MeV resonance, corresponding to the first $T = \frac{3}{2}$ state at $E_{\text{x}} = 15.06$ MeV, and from $E_{\text{p}} = 17$ to 20 MeV. The yield for the α_1 group has been determined for $E_{\text{p}} = 17$ to 21.5 MeV. Parameters of observed resonances are displayed in Table 13.22 (1969LE18). Excitation functions for α_0 have also been measured for $E_{\text{p}} = 18.5$ to 44 MeV at a number of angles: they exhibit structures which are typically 1 MeV broad (1971GU23). See also ^9B in (1979AJ01).

| | | |
|-------------------------------------------------------|--------------------------|-------------------------|
| 18. $^{12}\text{C}(\text{p}, ^6\text{Li})^7\text{Be}$ | $Q_{\text{m}} = -22.568$ | $E_{\text{b}} = 1.9434$ |
|-------------------------------------------------------|--------------------------|-------------------------|

Excitation functions have been measured for $E_p = 36$ to 43 MeV: no structure is apparent (1971HO25). See also (1979RA20) and ${}^6\text{Li}$, ${}^7\text{Be}$ in (1979AJ01).

19. (a) ${}^{12}\text{C}(\text{d}, \text{n}){}^{13}\text{N}$ $Q_m = -0.2812$
 (b) ${}^{12}\text{C}(\text{d}, \text{pn}){}^{12}\text{C}$ $Q_m = -2.2246$

Measurements of angular distributions of neutrons are tabulated in Table 13.27 of (1970AJ04) and in (1976AJ04). Recent measurements have been reported at $E_{\bar{\text{d}}} = 5.74$ to 9.70 MeV (1976TE03; n_0, n_1) and $E_{\text{d}} = 13.9$ and 15.25 MeV (1975AZ02; n_0, n_1, n_{2+3}).

Reaction (b) is dominated at $E_{\text{d}} = 5.0$ to 6.5 MeV and at 9.20 and 9.85 MeV by sequential decay via ${}^{13}\text{N}^*(3.51 + 3.55)$. At the lower energies ${}^{13}\text{N}^*(2.36)$ participates also: see (1976AJ04) and (1976GE20). See also ${}^{14}\text{N}$, (1977BA1L, 1979SI07), (1974LO1B), (1976LO1B; applications) and (1977TR09, 1978BA21; theor.).

20. (a) ${}^{12}\text{C}({}^3\text{He}, \text{d}){}^{13}\text{N}$ $Q_m = -3.5501$
 (b) ${}^{12}\text{C}({}^3\text{He}, \text{pd}){}^{12}\text{C}$ $Q_m = -5.4936$

Angular distributions have been studied to the first eight states of ${}^{13}\text{N}$ with $E({}^3\text{He})$ up to 40 MeV: see (1970AJ04, 1976AJ04). Recent measurements are reported at $E({}^3\text{He}) = 25.4$ MeV (1979SE07; $\text{d}_0, \text{d}_1, \text{d}_{2+3}$), $E({}^3\text{He}) = 33.3$ MeV (1976KA23; $\text{d}_0, \text{d}_{2+3}$) and $E({}^3\text{He}) = 43.6$ MeV (1980PE13; $\text{d}_0, \text{d}_1, \text{d}_{2+3}$ and d to ${}^{13}\text{N}^*(6.36, 6.89, 7.16, 7.38, 8.0, 9.0, 10.78)$) and 81.4 MeV (1976KO36; d to ${}^{13}\text{N}^*(0, 2.36, 3.51 + 3.55, 7.16 + 7.38, 7.9, 10.36, 11.1, 12.08)$). See also (1976RO1L). The spectroscopic factors derived by (1980PE13) are $S = 0.48, 0.14, 0.53, 0.007, 0.015, < 0.009, 0.024, 0.13$ and 0.064 for ${}^{13}\text{N}^*(0, 2.37, 3.55, 6.36, 6.89, 7.16, 7.38, 8.0, 10.78)$; $S < 0.005$ for ${}^{13}\text{N}^*(9.0)$ and evidence is presented for its $\frac{9}{2}^+$ character. For other values of S see (1976KA23, 1976KO36, 1979SE07). The energies and widths of the first three excited states are $E_x = 2368.2 \pm 2.8, 3507.8 \pm 7.6$ and 3549.2 ± 5.0 keV, with $\Gamma_{\text{cm}} = 36.1 \pm 2.8, 54.8 \pm 11.5$ and 46.5 ± 7.1 keV, respectively (1974BL06). Spectra obtained at $E({}^3\text{He}) = 36$ MeV at forward angles show the broad $\text{d}_{3/2}$ state at $E_x = 7.9$ MeV. Interfering with it is the 120 keV $\frac{3}{2}^+$ state at 6.89 MeV. The line shape of the $\frac{5}{2}^+$ state at 6.36 MeV is said to show a pronounced interference pattern (1979FU03). For reaction (b) see (1978MA42; $E({}^3\text{He}) = 70$ and 80 MeV). See also (1977BO30, 1977SA1L, 1978BA21; theor.) and ${}^{15}\text{O}$.

21. ${}^{12}\text{C}(\alpha, \text{t}){}^{13}\text{N}$ $Q_m = -17.8706$

Angular distributions of the t_0 groups have been measured at $E_\alpha = 56$ MeV (1969GA11) and 104 MeV (1972HA08).

$$22. {}^{12}\text{C}({}^7\text{Li}, {}^6\text{He}){}^{13}\text{N} \quad Q_m = -8.031$$

Angular distributions have been obtained at $E({}^7\text{Li}) = 36$ MeV for the ${}^6\text{He}$ ions to ${}^{13}\text{N}^*(0, 3.51 + 3.55)$ (1973SC26). The spectroscopic factor for ${}^{13}\text{N}_{\text{g.s.}}$ is 0.72 (1979ZE01).

$$23. {}^{12}\text{C}({}^{10}\text{B}, {}^9\text{Be}){}^{13}\text{N} \quad Q_m = -4.642$$

See (1976AJ04).

$$24. {}^{12}\text{C}({}^{11}\text{B}, {}^{10}\text{Be}){}^{13}\text{N} \quad Q_m = -9.285$$

See (1974AN36). See also (1976AJ04).

$$25. {}^{12}\text{C}({}^{12}\text{C}, {}^{11}\text{B}){}^{13}\text{N} \quad Q_m = -14.014$$

At $E({}^{12}\text{C}) = 93.8$ MeV angular distributions involving ${}^{13}\text{N}^*(0, 3.51 + 3.55)$ and various ${}^{11}\text{B}$ states have been measured by (1979FU04). See also (1979HEZX) and (1976AJ04).

$$26. {}^{12}\text{C}({}^{14}\text{N}, {}^{13}\text{C}){}^{13}\text{N} \quad Q_m = -5.607$$

Angular distributions involving ${}^{13}\text{N}_{\text{g.s.}}$ and various ${}^{13}\text{C}$ states [see reaction 43 in ${}^{13}\text{C}$] have been studied at $E({}^{14}\text{N}) = 28$ to 155 MeV: see (1976AJ04) and (1976BA16, 1976NA09). See also (1978MA1F) and (1977WE1H, 1978DZ1A, 1978NA15, 1978WE1F, 1979DO13; theor.).

$$27. {}^{12}\text{C}({}^{16}\text{O}, {}^{15}\text{N}){}^{13}\text{N} \quad Q_m = -10.184$$

At $E({}^{16}\text{O}) = 128$ MeV angular distributions are reported to ${}^{13}\text{N}^*(0, 2.36, 3.55)$ (1979PR07; also see for reduced widths).

$$28. {}^{12}\text{C}({}^{20}\text{Ne}, {}^{19}\text{F}){}^{13}\text{N} \quad Q_m = -10.901$$

See (1979OR01; $E({}^{20}\text{Ne}) = 150$ to 294 MeV).

$$29. \ ^{13}\text{C}(\pi^+, \pi^0)^{13}\text{N} \quad Q_m = 2.384$$

See (1979MO1N) and the “GENERAL” section here.

$$30. \text{ (a) } ^{13}\text{C}(\text{p}, \text{n})^{13}\text{N} \quad Q_m = -3.0030$$

$$\text{ (b) } ^{13}\text{C}(\text{p}, \text{pn})^{12}\text{C} \quad Q_m = -4.9464$$

Angular distributions of n_0, n_1, n_{2+3} have been measured for $E_p = 3.1$ to 50 MeV: see (1970AJ04, 1976AJ04). See also (1978WA1D). The 0° differential cross section for the n_0 group has been measured to be 4.5 ± 0.7 mb/sr at $E_p = 120$ MeV (1980GO07). In reaction (b) at $E_p = 7.9$ to 12 MeV one or both of the ^{13}N states at $E_x = 3.5$ MeV appear to be involved (1971OT02). See also (1980BA11; theor.).

$$31. \ ^{13}\text{C}(^3\text{He}, \text{t})^{13}\text{N} \quad Q_m = -2.2392$$

At $E(^3\text{He}) = 39.6$ MeV, angular distributions have been obtained for the tritons corresponding to the ground state of ^{13}N and to the excited states at $2.36, 3.53 \pm 0.03$ (unresolved), $6.36, 7.16, 7.38, 8.92 \pm 0.04, 11.85 \pm 0.04$ and 15.06 MeV. States at $E_x = 9.5, 10.78 \pm 0.04$ and 15.98 ± 0.05 MeV were also populated, the first of these weakly. The transitions to $^{13}\text{N}^*(7.38, 8.92, 11.85, 15.06)$ are $L = 2$ [$J^\pi = \frac{5}{2}^-, \frac{1}{2}^-, \frac{3}{2}^-, \frac{3}{2}^-$, respectively] (1969BA06). Angular distributions have also been reported at $E(^3\text{He}) = 14$ MeV (1970NU02; t_0, t_1). For a comparison of the angular distribution to $^{13}\text{N}^*(8.918)$ and $^{13}\text{C}^*(9.498)$ at $E(^3\text{He}) = 43.6$ MeV, see reaction 57 in ^{13}C (1980PE1E).

$$32. \ ^{13}\text{C}(^6\text{Li}, ^6\text{He})^{13}\text{N} \quad Q_m = -5.727$$

Angular distributions have been measured at $E(^6\text{Li}) = 31.8$ MeV (1970CH19; $^{13}\text{N}^*(0, 3.51 + 3.55)$). See also (1979ZE01) and (1976AJ04).

$$33. \ ^{13}\text{C}(^{14}\text{N}, ^{14}\text{C})^{13}\text{N} \quad Q_m = -2.377$$

See (1976AJ04) and (1978OS06; theor.).

$$34. \ ^{13}\text{O}(\beta^+)^{13}\text{N} \quad Q_m = 17.762$$

See ^{13}O .

$$35. \ ^{14}\text{N}(\gamma, n)^{13}\text{N} \quad Q_m = -10.554$$

See ^{14}N .

$$36. \ ^{14}\text{N}(n, 2n)^{13}\text{N} \quad Q_m = -10.554$$

See (1978RY02), (1977LE1J; applications) and ^{15}N .

$$37. \ ^{14}\text{N}(p, d)^{13}\text{N} \quad Q_m = -8.3290$$

Angular distributions have been measured for deuteron groups to $^{13}\text{N}^*(0, 2.36, 3.51 + 3.55, 7.38, 8.92, 11.88)$ at many energies up to $E_p = 155.6$ MeV: see (1976AJ04) and by (1977GU14; $E_p = 16.2$ and 17.7 MeV; p_0) and (1979AO02; $E_p = 21$ MeV; p_0). See also (1976WA15, 1978MA34; theor.).

$$38. \ ^{14}\text{N}(d, t)^{13}\text{N} \quad Q_m = -4.2963$$

Angular distributions of the tritons to $^{13}\text{N}^*(0, 3.51, 7.38, 8.92 + (9.00) + 9.48, 11.8)$ have been obtained at $E_d = 52$ MeV and analyzed by DWBA. The spectroscopic factors for the ^{13}N states [and the mirror states reached in the $^{14}\text{N}(d, ^3\text{He})^{13}\text{C}$ reaction] are in good agreement with theory and are additional evidence for the J^π assignments of $\frac{1}{2}^-$, $\frac{3}{2}^-$, $\frac{5}{2}^-$, $\frac{1}{2}^-$, $\frac{3}{2}^-$ and $\frac{3}{2}^-$ to these states (1968HI01). For comparisons of (d, t) and (d, ^3He) angular distributions see reaction 80 in ^{13}C . See also ^{16}O in (1982AJ01).

$$39. \text{ (a) } \ ^{14}\text{N}(^3\text{He}, \alpha)^{13}\text{N} \quad Q_m = 10.024$$

$$\text{ (b) } \ ^{14}\text{N}(^3\text{He}, p\alpha)^{12}\text{C} \quad Q_m = 8.0808$$

Alpha particle groups have been observed to the first seven excited states of ^{13}N , including two at $E_x = 7.166$ and 7.388 MeV (1962CL12; ± 8 keV). Angular distributions have been studied at many energies up to $E(^3\text{He}) = 45$ MeV: see (1976AJ04) and (1971GU22; 4 to 10.25 MeV; α_0, α_1). Reaction (b), studied at $E(^3\text{He}) = 8$ MeV, appears to involve some states of ^{13}N , possibly $^{13}\text{N}^*(7.93, 8.92, 11.87)$ (1969HO13). See also ^{17}F in (1982AJ01).

$$40. \ ^{14}\text{N}(^6\text{Li}, ^7\text{Li})^{13}\text{N} \quad Q_m = -3.303$$

An angular distribution has been measured at $E(^6\text{Li}) = 32$ MeV for the transition to $^{13}\text{N}_{\text{g.s.}}$ and $^7\text{Li}^*(0, 0.48)$. $^{13}\text{N}^*(2.36)$ was also populated (1971GR44). See also (1976AJ04) and reaction 83 in ^{13}C .

$$41. \ ^{14}\text{N}(^9\text{Be}, ^{10}\text{Be})^{13}\text{N} \quad Q_m = -3.741$$

See (1976AJ04).

$$42. \ ^{14}\text{N}(^{10}\text{B}, ^{11}\text{B})^{13}\text{N} \quad Q_m = 0.902$$

See (1976NA09).

$$43. \ ^{14}\text{N}(^{14}\text{N}, ^{15}\text{N})^{13}\text{N} \quad Q_m = 0.280$$

See (1976SW02). See also (1976AJ04).

$$44. \ ^{15}\text{N}(\text{p}, \text{t})^{13}\text{N} \quad Q_m = -12.9050$$

At $E_p = 43.7$ MeV, angular distributions have been obtained for the tritons corresponding to the ground state of ^{13}N and the excited states at 3.51 ($\frac{3}{2}^-$), 6.38 ± 0.03 ($\frac{5}{2}^+$), 7.38 ($\frac{5}{2}^-$), 8.93 ± 0.05 ($\frac{1}{2}^-$), 10.78 ± 0.06 ($\frac{1}{2}^-$), 11.88 ± 0.04 ($\frac{3}{2}^-$) and 15.06 ($\frac{3}{2}^-$; $T = \frac{3}{2}$) MeV [J^π values in parentheses, as determined by DWBA analyses using intermediate-coupling wave functions to obtain the two-nucleon structure factors] (1968FL03). Detailed comparisons have been made with the (p, ^3He) reaction to the mirror states in ^{13}C (1968FL02, 1968FL03, 1974MA12): see reaction 87 in ^{13}C . Angular distributions have also been measured for the triton groups to $^{13}\text{N}^*(0, 3.51 + 3.55, 7.38)$ at several energies in the range $E_p = 24.0$ to 43.5 MeV (1974PI05, 1975MI01). See also ^{16}O in (1982AJ01).

$$45. \ ^{16}\text{O}(\gamma, \text{t})^{13}\text{N} \quad Q_m = -25.033$$

See ^{16}O in (1982AJ01).

46. (a) $^{16}\text{O}(\text{p}, \alpha)^{13}\text{N}$ $Q_{\text{m}} = -5.2185$
 (b) $^{16}\text{O}(\text{p}, \text{p}\alpha)^{12}\text{C}$ $Q_{\text{m}} = -7.1620$

Angular distributions of the α_0 , α_1 and α_2 groups have been measured for E_{p} to 54.1 MeV: see (1970AJ04, 1976AJ04). In addition, (1972MA21) have measured the distribution of the α group to a state with $E_{\text{x}} = 12.13 \pm 0.06$ MeV, $\Gamma_{\text{cm}} \approx 300$ keV [$J^{\pi} = \frac{7}{2}^{-}$] at $E_{\text{p}} = 54.1$ MeV. See also (1976AJ04) and ^{17}F in (1982AJ01).

47. $^{16}\text{O}(^3\text{He}, ^6\text{Li})^{13}\text{N}$ $Q_{\text{m}} = -9.239$

Angular distributions have been studied at $E(^3\text{He}) = 30$ and 40.7 MeV for the ^6Li ions corresponding to $^{13}\text{N}_{\text{g.s.}}$ (1972OH01).

48. $^{16}\text{O}(\alpha, ^7\text{Li})^{13}\text{N}$ $Q_{\text{m}} = -22.566$

The angular distribution for the transition to $^{13}\text{N}_{\text{g.s.}} + ^7\text{Li}_{\text{g.s.}+0.48}$ has been measured at $E_{\alpha} = 42$ MeV (1972RU03).

49. $^{16}\text{O}(^{14}\text{N}, ^{17}\text{O})^{13}\text{N}$ $Q_{\text{m}} = -6.409$

This reaction has been studied at $E(^{14}\text{N}) = 79$ MeV (1976MO03) and 155 MeV (1975NA15, 1976NA09).

50. $^{18}\text{O}(\text{d}, ^7\text{Li})^{13}\text{N}$ $Q_{\text{m}} = -7.901$

See (1970AJ04).

51. $^{24}\text{Mg}(^{12}\text{C}, ^{23}\text{Na})^{13}\text{N}$ $Q_{\text{m}} = -9.747$

See (1978NO02).

¹³O
(Fig. 4)

GENERAL (See also (1976AJ04).):

Theoretical and review papers: (1975BE56, 1976AB04, 1977CE05, 1978GU10, 1979BE1H).

Mass of ¹³O: From the Q -value of the $^{16}\text{O}(^3\text{He}, ^6\text{He})^{13}\text{O}$ reaction [$Q_0 = -30.508 \pm 0.010$ MeV] (see reaction 4) the atomic mass excess of ^{13}O is determined to be 23.108 ± 0.010 MeV. ^{13}O is then bound with respect to $^{12}\text{N} + \text{p}$ and $^{11}\text{C} + 2\text{p}$ by 1.519 and 2.120 MeV, respectively.

1. $^{13}\text{O}(\beta^+)^{13}\text{N}$ $Q_m = 17.762$

The half-life of ^{13}O is 8.90 ± 0.20 msec: see (1970ES03) and (1976AJ04). ^{13}O decays to a number of states of ^{13}N some of which subsequently decay to $^{12}\text{C}^*(0, 4.4)$: see Table 13.24 (1970ES03). See also (1978RA2A) and (1977MA16, 1977RI08, 1979DE15; theor.).

2. $^{12}\text{C}(\text{p}, \pi^-)^{13}\text{O}$ $Q_m = -155.386$

At $E_p = 613$ MeV the ground state of ^{13}O and an excited state at $E_x = 2.82 \pm 0.24$ MeV are observed in addition to unresolved structures (1978CO15) [see Fig. 4 for analogue region in ^{13}B]. See also (1979HO1F).

3. $^{13}\text{C}(\pi^+, \pi^-)^{13}\text{O}$ $Q_m = -19.983$

See (1979GR1K).

4. $^{16}\text{O}(^3\text{He}, ^6\text{He})^{13}\text{O}$ $Q_m = -30.508$
 $Q_0 = -30.506 \pm 0.013$ (1971TR03);
 $Q_0 = -30.513 \pm 0.015$ (1970ME11); recalculated on basis
of ^9C mass excess of 28.912 MeV (see (1979AJ01)).

The ground state of ^{13}O has been populated at $E(^3\text{He}) = 62.6$ to 68.6 MeV (1970ME11, 1971TR03).

¹³F, ¹³Ne
(Not illustrated)

^{13}F and ^{13}Ne have not been observed. See (1975BE31; theor.).

Table 13.23: Energy levels of ^{13}O

| E_x (MeV) | $J^\pi; T$ | $\tau_{1/2}$ (msec) | Decay | Reactions |
|-------------------------|--------------------------------|---------------------|-----------|-----------------|
| g.s. 2.82 ± 0.24 | $(\frac{3}{2}^-); \frac{3}{2}$ | 8.90 ± 0.20 | β^+ | 1, 2, 3, 4 2 |

Table 13.24: Beta decay of ^{13}O ^a

| Decay to | | $E_p(\text{cm})$ (MeV) to | | Relative intensity | % of all β -decays | log ft |
|-------------------------|-----------------|--------------------------------|------------------------|------------------------------|-----------------------------|------------------------------|
| $^{13}\text{N}^*$ (MeV) | J^π | $^{12}\text{C}^*(\text{g.s.})$ | $^{12}\text{C}^*(4.4)$ | | | |
| g.s. | $\frac{1}{2}^-$ | | | | 88.1 ± 3.4 ^b | 4.10 ± 0.02 ^b |
| 3.51 | $\frac{3}{2}^-$ | observed | | 100 | 10.7 ± 3.1 | 4.52 ± 0.13 |
| 7.38 | $\frac{5}{2}^-$ | 5.48 ± 0.05 | not seen | 0.33 ± 0.10 | 0.40 ± 0.19 | 5.22 ± 0.23 |
| 8.92 | $\frac{1}{2}^-$ | observed | | 3.5 ± 0.3 | 0.54 ± 0.16 | 4.73 ± 0.14 |
| | | | 2.56 ± 0.05 | 1.5 ± 0.3 | | |
| 9.48 | $\frac{3}{2}^-$ | observed | | 0.8 ± 0.1 | 0.13 ± 0.04 | 5.18 ± 0.14 |
| | | | 3.12 ± 0.05 | 0.43 ± 0.15 | | |
| 10.36 | $\frac{5}{2}^-$ | observed | | 0.05 ± 0.03 ^c | 0.019 ± 0.012 | 5.8 ± 0.3 |
| | | | 3.97 ± 0.05 | 0.13 ± 0.07 | | |

^a (1970ES03). In addition there is some evidence for weak proton groups with $E_p = 3.44$ and 6.28 MeV (± 0.05 MeV). See also (1965MC09).

^b The ground state ft was taken to be 1.15 times that for ^{13}B (1970ES03). (1971WI07) find $(ft)^+/(ft)^- = 1.17 \pm 0.03$: see reaction 1 in ^{13}B .

^c Calculated value from the known ratio of the elastic and inelastic widths.

References

(Closed 01 August 1980)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1955MC75 J.N. McGruer, E.K. Warburton and R.S. Bender, Phys. Rev. 100 (1955) 235
- 1956CO72 B.C. Cook, A.S. Penfold and V.L. Telegdi, Phys. Rev. 104 (1956) 554
- 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., Phys. Rev. 104 (1956) 1064
- 1957CO57 B.C. Cook, Phys. Rev. 106 (1957) 300
- 1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
- 1960JA12 J. Janecke, Z. Naturforsch. A15 (1960) 593
- 1961NA02 Y. Nagahara, J. Phys. Soc. Jpn. 16 (1961) 133
- 1961NO05 E. Norbeck, Phys. Rev. 121 (1961) 824
- 1962CL12 D.D. Clayton, Phys. Rev. 128 (1962) 2254
- 1963DA12 E.A. Davis, T.W. Bonner, D.W. Worley, Jr. and R. Bass, Nucl. Phys. 48 (1963) 169
- 1963FI07 P.S. Fisher, D.F. Measday, F.A. Nikolaev, A. Kalmykov and A.B. Clegg, Nucl. Phys. 45 (1963) 113
- 1963HO19 H.D. Holmgren, L.M. Cameron and R.L. Johnston, Nucl. Phys. 48 (1963) 1
- 1963YO06 F.C. Young, J.C. Armstrong and J.B. Marion, Nucl. Phys. 44 (1963) 486; Erratum Nucl. Phys. 49 (1963) 695
- 1964BE24 A.M. Bernstein, R.A. Haberstroh, D.R. Hamilton, M. Posner and J.L. Snider, Phys. Rev. 136 (1964) B27
- 1964BR1A Breuer, Z. Phys. 178 (1964) 268
- 1964DE12 V.P. Denisov, A.V. Kulikov and L.A. Kulchitskii, Zh. Eksp. Teor. Fiz. 46 (1964) 1488; Sov. Phys. JETP 19 (1964) 1007
- 1964KU09 H.-M. Kuan, P.R. Almond, G.U. Din and T.W. Bonner, Nucl. Phys. 60 (1964) 509
- 1964MI04 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 50
- 1965AL17 P.R. Almond and J.R. Risser, Nucl. Phys. 72 (1965) 436

- 1965LI09 G.P. Lietz, S.F. Trevino, A.F. Behof and S.E. Darden, Nucl. Phys. 67 (1965) 193
- 1965MC09 R. McPherson, R.A. Esterlund, A.M. Poskanzer and P.L. Reeder, Phys. Rev. 140 (1965) B1513
- 1965PA05 J.R. Patterson, J.M. Poate and E.W. Titterton, Proc. Phys. Soc. 85 (1965) 1085
- 1965PA10 J.R. Patterson, J.M. Poate, E.W. Titterton and B.A. Robson, Proc. Phys. Soc. 86 (1965) 1297
- 1966BA01 F.C. Barker, J.R. Patterson, J.M. Poate and E.W. Titterton, Phys. Lett. 21 (1966) 318
- 1966BA13 G.C. Ball and J. Cerny, Phys. Lett. 21 (1966) 551
- 1966CE02 J. Cerny, R.H. Pehl, G. Butler, D.G. Fleming, C. Maples and C. Detraz, Phys. Lett. 20 (1966) 35
- 1966DE1C Detraz et al., Phys. Lett. 22 (1966) 638
- 1966GL01 R.N. Glover and A.D.W. Jones, Nucl. Phys. 84 (1966) 673
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966PA10 J.R. Patterson, J.M. Poate and E.W. Titterton, Proc. Phys. Soc. 88 (1966) 641
- 1966PO11 A.R. Poletti, J.W. Olness and E.K. Warburton, Phys. Rev. 151 (1966) 812
- 1966YO1A Young et al., Nucl. Instrum. Meth. Phys. Res. 44 (1966) 109
- 1967AL16 A.U. Aldzhauakhiri, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 6 (1967) 248; Sov. J. Nucl. Phys. 6 (1968) 180
- 1967DI01 G.U. Din, M.A. Nagarajan and R. Pollard, Nucl. Phys. A93 (1967) 190
- 1967KI03 K.G. Kibler, Phys. Rev. 155 (1967) 1110
- 1967PF02 L. Pfeiffer and L. Madansky, Phys. Rev. 163 (1967) 999
- 1967PR10 W.V. Prestwich, R.E. Cote and G.E. Thomas, Phys. Rev. 161 (1967) 1080
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968BO34 F. Boreli, B.B. Kinsey and P.N. Shrivastava, Phys. Rev. 174 (1968) 1147
- 1968CH05 L.F. Chase, Jr., R.E. McDonald, W.W. True and E.K. Warburton, Phys. Rev. 166 (1968) 997
- 1968DA05 C.N. Davids, Nucl. Phys. A110 (1968) 619
- 1968DI04 F.S. Dietrich, M. Suffert, A.V. Nero and S.S. Hanna, Phys. Rev. 168 (1968) 1169
- 1968FL02 D.G. Fleming, J. Cerny and N.K. Glendenning, Phys. Rev. 165 (1968) 1153
- 1968FL03 D.G. Fleming, J. Cerny, C.C. Maples and N.K. Glendenning, Phys. Rev. 166 (1968) 1012
- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161

- 1968HI01 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, P. Turek and G.J. Wagner, Nucl. Phys. A106 (1968) 161
- 1968MI05 P.F. Mizera and J.B. Gerhart, Phys. Rev. 170 (1968) 839
- 1968RI01 E.M. Rimmer and P.S. Fisher, Nucl. Phys. A108 (1968) 561
- 1968RI15 A.I.M. Ritchie, Nucl. Instrum. Meth. Phys. Res. 64 (1968) 181
- 1968RI16 F. Riess, P. Paul, J.B. Thomas and S.S. Hanna, Phys. Rev. 176 (1968) 1140
- 1968RO02 S.W. Robinson, C.P. Swann and V.K. Rasmussen, Phys. Lett. B26 (1968) 298
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, Nucl. Phys. A113 (1968) 395
- 1969AD02 E.G. Adelberger, A.B. McDonald and C.A. Barnes, Nucl. Phys. A124 (1969) 49
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969BA1F V.V. Balashov, N.M. Kabachnik and V.I. Markov, Nucl. Phys. A129 (1969) 369
- 1969CA1A R.R. Carlson and H.W. Wyborny, Phys. Rev. 178 (1969) 1529
- 1969FR03 E. Friedland and H. Verleger, Z. Phys. 222 (1969) 138
- 1969GA11 P. Gaillard, R. Bouche, L. Feuveys, M. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
- 1969HE22 S.H. Henson, D.B. Fossan, P. Paul and K.W. Jones, Phys. Rev. 188 (1969) 1967
- 1969HO13 K. Hofmann, W. Mausberg, M. Raabe, E. Rossle and M. Tauber, Z. Phys. 221 (1969) 417
- 1969JO21 K.W. Jones, W.R. Harris, M.T. McEllistrem and D.E. Alburger, Phys. Rev. 186 (1969) 978
- 1969LE18 M.J. LeVine and P.D. Parker, Phys. Rev. 186 (1969) 1021
- 1969RA20 V.K. Rasmussen and C.P. Swann, Phys. Rev. 183 (1969) 918
- 1969SN02 F.D. Snyder and M.A. Waggoner, Phys. Rev. 186 (1969) 999
- 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
- 1969VO01 W. von Oertzen, J.C. Jacmart, M. Liu, F. Pougheon, J.C. Roynette and M. Riou, Phys. Lett. B28 (1969) 482
- 1969WI22 G. Wittwer, H.-G. Clerc and G.A. Beer, Phys. Lett. B30 (1969) 634
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1970BA49 P.H. Barker, A. Huber, H. Knoth, U. Matter, A. Gobbi and P. Marmier, Nucl. Phys. A155 (1970) 401
- 1970CH19 V.I. Chuev, V.V. Davydov, V.I. Manko, B.G. Novatsky, S.B. Sakuta and D.N. Stepanov, Phys. Lett. B31 (1970) 624
- 1970DE14 G. Deconninck and J.-P. Meulders, Phys. Rev. C1 (1970) 1326

- 1970DI1A Dickens and Perey, Nucl. Sci. Eng. 40 (1970) 346
- 1970ES03 J.E. Esterl, J.C. Hardy, R.G. Sextro and J. Cerny, Phys. Lett. B33 (1970) 287
- 1970GA01 A. Gallmann, G. Frick, B. Heusch, F. Haas and E.K. Warburton, Nucl. Phys. A143 (1970) 326
- 1970HE24 J. Heisenberg, J.S. McCarthy and I. Sick, Nucl. Phys. A157 (1970) 435
- 1970KR09 N.N. Krasnov, P.P. Dmitriev, Z.P. Dmitrieva, I.O. Konstantinov and G.A. Molin, At. Energ. 28 (1970) 258; Sov. At. Energy 28 (1970) 335
- 1970ME11 R. Mendelson, G.J. Wozniak, A.D. Bacher, J.M. Loiseaux and J. Cerny, Phys. Rev. Lett. 25 (1970) 533
- 1970ME1C Meadows and Whalen, Nucl. Sci. Eng. 41 (1970) 351
- 1970NE17 J.M. Nelson, N.S. Chant and P.S. Fisher, Nucl. Phys. A156 (1970) 406
- 1970NU02 F. Nusslin and P. Braun-Munzinger, Z. Phys. 240 (1970) 217
- 1970SH06 N.K. Sherman, R.W. Gellie, K.H. Lokan and R.G. Johnson, Phys. Rev. Lett. 25 (1970) 114
- 1970TH01 M.N. Thompson, R.J.J. Stewart and J.E.M. Thompson, Phys. Lett. B31 (1970) 211
- 1970VO02 W. von Oertzen, M. Liu, C. Caverzasio, J.C. Jacmart, F. Pougheon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A143 (1970) 34
- 1970WE09 L.C. Welch, C.C. Chang, H.H. Forster, C.C. Kim, D.W. Devins and P.A. Deutchman, Nucl. Phys. A158 (1970) 644
- 1970WI04 G. Wittwer, H.-G. Clerc and G.A. Beer, Z. Phys. 234 (1970) 120
- 1971BA68 P.H. Barker, P.M. Cockburn, A. Huber, H. Knoth, U. Matter, H.-P. Seiler and P. Marmier, Ann. Phys. 66 (1971) 705
- 1971BE34 D. Bellman, Atomkernenergie 17 (1971) 145
- 1971BE51 J.C. Bergstrom, H. Crannell, F.J. Kline, J.T. O'Brien, J.W. Lightbody, Jr. and S.P. Fivozinsky, Phys. Rev. C4 (1971) 1514
- 1971CO14 W.R. Coker, P.D. Miller, J.L. Duggan, M.M. Duncan, R.L. Dangle and J. Lin, Nucl. Phys. A168 (1971) 307
- 1971GO24 V.Z. Goldberg, V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 1663; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 1514
- 1971GO40 N.B. Gove and M.J. Martin, Nucl. Data Tables A10 (1971) 205
- 1971GR44 K.O. Groeneveld, A. Richter, U. Strohmusch and B. Zeidman, Phys. Rev. Lett. 27 (1971) 1806
- 1971GU22 P. Guazzoni, S. Micheletti, M. Pignanelli, F. Gentilin and F. Pellegrini, Phys. Rev. C4 (1971) 1086

- 1971GU23 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Semenescu, Phys. Rev. C4 (1971) 1092
- 1971HO25 A.B. Holman, J.-L. Perrenoud, J.C. Young, M.B. Epstein and B.T. Wright, Nucl. Phys. A174 (1971) 161
- 1971HS03 T.H. Hsu, J.L. Honsaker, W.J. McDonald and G.C. Neilson, Nucl. Phys. A174 (1971) 365
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971LI11 M. Liu, W. von Oertzen, J.C. Jacmart, F. Pougheon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A165 (1971) 118
- 1971OT02 V.A. Otte, W. von Witsch, J. Sandler, D. Rendic and G.C. Phillips, Phys. Rev. C4 (1971) 322
- 1971RI19 S.M. Rizvi, P. Huber, F. Seiler and H.R. Striebel, Helv. Phys. Acta 44 (1971) 803
- 1971SC21 H. Schmidt-Bocking, G. Brommundt and K. Bethge, Z. Phys. 246 (1971) 431
- 1971SH09 Y.M. Shin, C.F. Wong and H.S. Caplan, Nucl. Phys. A166 (1971) 162
- 1971ST21 D.M. Stupin, R.A. Ristinen and P. Schwandt, Nucl. Phys. A173 (1971) 286
- 1971TR03 G.F. Trentelman, B.M. Preedom and E. Kashy, Phys. Rev. C3 (1971) 2205
- 1971VO01 W. von Oertzen, G. Bohlen, B. Gebauer, E.E. Habib and N. Marquardt, Phys. Lett. B34 (1971) 51
- 1971WI07 D.H. Wilkinson, D.E. Alburger, D.R. Goosman, K.W. Jones, E.K. Warburton, G.T. Garvey and R.L. Williams, Nucl. Phys. A166 (1971) 661
- 1971WY01 H.W. Wyborny and R.R. Carlson, Phys. Rev. C3 (1971) 2185
- 1971ZA04 N.I. Zaika, Y.V. Kibkalo, E.B. Levshin, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, P.L. Shmarin, A.M. Shchur and A.M. Yasnogorodskii, Yad. Fiz. 13 (1971) 928; Sov. J. Nucl. Phys. 13 (1971) 533
- 1972BE05 R.A.I. Bell, I.G. Graham and J.V. Thompson, Nucl. Phys. A179 (1972) 408
- 1972BE56 R.A.I. Bell, I.G. Graham, J.V. Thompson and E.W. Titterton, Nucl. Phys. A193 (1972) 385
- 1972BO68 H.G. Bohlen, M. Feil, A. Gamp, B. Kohlmeyer, N. Marquardt and W. von Oertzen, Phys. Lett. B41 (1972) 425
- 1972GE11 R.W. Gellie, K.H. Lokan, N.K. Sherman, R.G. Johnson and J.I. Lodge, Can. J. Phys. 50 (1972) 1689
- 1972HA08 G. Hauser, R. Lohken, G. Nowicki, H. Rebel, G. Schatz, G. Schweimer and J. Specht, Nucl. Phys. A182 (1972) 1
- 1972KE08 R.M. Keyser, R.A. Blue and H.R. Weller, Nucl. Phys. A186 (1972) 528

- 1972MA21 C. Maples and J. Cerny, Phys. Lett. B38 (1972) 504
- 1972OB01 A.W. Obst, T.B. Grandy and J.L. Weil, Phys. Rev. C5 (1972) 738
- 1972OH01 I.K. Oh, C.S. Zaidins, C.D. Zafiratos and S.I. Hayakawa, Nucl. Phys. A178 (1972) 497
- 1972PE11 C.A. Pearson, J.M. Covan, D. Zissermann, T.G. Miller, F.P. Gibson, R. Haglund, W. Morrison and G. Westley, Nucl. Phys. A191 (1972) 1
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430
- 1972SE09 F. Seiler, Nucl. Phys. A187 (1972) 379
- 1972TH14 G. Thornton, T. Joy and J.M. Calvert, Nucl. Phys. A198 (1972) 397
- 1972WY01 H.W. Wyborny, Nucl. Phys. A185 (1972) 669
- 1973AD02 E.G. Adelberger, A.B. McDonald, C.L. Cocke, C.N. Davids, A.P. Shukla, H.B. Mak and D. Ashery, Phys. Rev. C7 (1973) 889
- 1973DA17 S.E. Darden, S. Sen, H.R. Hiddleston, J.A. Aymar and W.A. Yoh, Nucl. Phys. A208 (1973) 77
- 1973DA24 S. Dahlgren, P. Grafstrom, B. Hoistad and A. Asberg, Nucl. Phys. A211 (1973) 243
- 1973DE14 D.C. De Martini, C.R. Soltesz and T.R. Donoghue, Phys. Rev. C7 (1973) 1824
- 1973GO03 J.D. Goss, A.A. Rollefson, G.L. Marolt and C.P. Browne, Phys. Rev. C7 (1973) 663
- 1973GO15 J.D. Goss, S.L. Blatt, D.R. Parsignault, C.D. Porterfield and F.L. Riffle, Phys. Rev. C7 (1973) 1837
- 1973HAVZ R.C. Haskell and L. Madansky, J. Phys. Soc. Jpn. Suppl. 34 (1973) 167
- 1973HE12 H.-D. Helb, P. Duck, G. Hartmann, G. Ischenko, F. Siller and H. Voit, Nucl. Phys. A206 (1973) 385
- 1973HU07 E. Huenges, H. Rosler and H. Vonach, Phys. Lett. B46 (1973) 361
- 1973ME03 H.O. Meyer and G.R. Plattner, Nucl. Phys. A199 (1973) 413
- 1973ME12 D.F. Measday, M. Hasinoff and D.L. Johnson, Can. J. Phys. 51 (1973) 1227
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973SA03 J. Sandler, G.S. Mutchler, W. von Witsch, V.A. Otte, G.C. Phillips and D. Rendic, Nucl. Phys. A202 (1973) 49
- 1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, Nucl. Phys. A212 (1973) 573
- 1973SM03 S.M. Smith, G. Tibell, A.A. Cowley, D.A. Goldberg, H.G. Pugh, W. Reichart and N.S. Wall, Nucl. Phys. A207 (1973) 273
- 1973VA25 L. Van der Zwan and K.W. Geiger, Nucl. Phys. A216 (1973) 188

- 1973WE12 H.R. Weller and R.A. Blue, Nucl. Phys. A211 (1973) 221
- 1974AL12 D.E. Alburger and D.R. Goosman, Phys. Rev. C10 (1974) 935
- 1974AN36 N. Anyas-Weiss, J.C. Cornell, P.S. Fisher, P.N. Hudson, A. Menchaca-Rocha, D.J. Millener, A.D. Panagiotou, D.K. Scott, D. Strottman, D.M. Brink et al., Phys. Rept. 12 (1974) 201
- 1974BA37 J.E.E. Baglin, E.J. Bentz and R.W. Carr, Phys. Rev. C10 (1974) 24
- 1974BE48 J.R. Beene, J. Asher, N. Ayres de Campos, R.D. Gill, M.A. Grace and W.L. Randolph, Nucl. Phys. A230 (1974) 141
- 1974BL06 S.L. Blatt, G.L. Marolt and J.D. Goss, Phys. Rev. C10 (1974) 1319
- 1974BO1E Bormann, Neuert and Scobel, IAEA, STI/DOC/10/156 (1974) 87
- 1974FI1D D. Fick, Nukleonika 19 (1974) 693
- 1974HO06 C.H. Holbrow, H.G. Bingham, R. Middleton and J.D. Garrett, Phys. Rev. C9 (1974) 902
- 1974KE06 K.W. Kemper, S. Cotanch, G.E. Moore, A.W. Obst, R.J. Puigh and R.L. White, Nucl. Phys. A222 (1974) 173
- 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325
- 1974LU06 E.J. Ludwig, C.E. Busch, T.B. Clegg, S.K. Datta and A.C. Watkins, Nucl. Phys. A230 (1974) 271
- 1974MA12 J.A. Macdonald, J. Cerny, J.C. Hardy, H.L. Harney, A.D. Bacher and G.R. Plattner, Phys. Rev. C9 (1974) 1694
- 1974MI05 C.A. Miller, J.W. Watson, D.I. Bonbright, F.J.S. Wilson and D.O. Wells, Phys. Rev. Lett. 32 (1974) 684
- 1974NO01 J.W. Noe, D.P. Balamuth and R.W. Zurmuhle, Phys. Rev. C9 (1974) 132
- 1974PI05 M. Pignanelli, S. Micheletti, I. Iori, P. Guazzoni, F.G. Resmini and J.L. Escudie, Phys. Rev. C10 (1974) 445
- 1974RO29 C. Rolfs and R.E. Azuma, Nucl. Phys. A227 (1974) 291
- 1974SA16 Z.A. Saleh, F. Machali, I.I. Bondouk and D.A. Darwish, Ann. Phys. (Leipzig) 31 (1974) 76
- 1974SP04 A.M.J. Spits and J. de Boer, Nucl. Phys. A224 (1974) 517
- 1974TH01 C. Thibault and R. Klapisch, Phys. Rev. C9 (1974) 793
- 1975AC1A Ackermann, Anders, Bormann and Scobel, Nucl. Cross Sect. and Tech., Pt. II, Washington, D.C., 1975 (1975) 819
- 1975AZ02 S.A. Azimov, U.R. Arifkhanov, M. Gulyamov, B.I. Islamov, T. Iskhakov, A.M. Mukhamedzhanov, E. Ergashov, Y.I. Faizullaev and R. Yarmukhamedov, Yad. Fiz. 22 (1975) 225; Sov. J. Nucl. Phys. 22 (1976) 115

- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BE56 W. Benenson, Nukleonika 20 (1975) 775
- 1975BL07 R.C. Block, Y. Fujita, K. Kobayashi and T. Oosaki, J. Nucl. Sci. Tech. 12 (1975) 1
- 1975CE04 F.E. Cecil, J.R. Shepard, R.E. Anderson, R.J. Peterson and P. Kaczkowski, Nucl. Phys. A255 (1975) 243
- 1975DE26 A.E. Denisov, R.P. Kolalis, V.S. Sadkovskii, G.A. Feofilov, O.V. Chubinskii, L.I. Vinogradov and Y.F. Andronov, Yad. Fiz. 21 (1975) 472; Sov. J. Nucl. Phys. 21 (1975) 247
- 1975GA1E Galimov, Doklady Akad. Nauk SSSR 225 (1975) 190
- 1975GE19 K.W. Geiger and L. van der Zwan, Nucl. Instrum. Meth. 131 (1975) 315
- 1975HI07 F. Hinterberger, P.V. Rossen, H.G. Ehrlich, B. Schuller, R. Jahn, J. Bisping and G. Welp, Nucl. Phys. A253 (1975) 125
- 1975HO1F Hollandsworth, Bucher, Youngblood and Niiler, Rept. BRL-1764 (1975)
- 1975HU1D Hufner, Phys. Rept. C21 (1975) 1
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, Phys. Rev. C11 (1975) 2049
- 1975KA1G Karol, Yester, Klobuchar and Caretto, Phys. Lett. B58 (1975) 489
- 1975LO1E Loumos, Lambert and Tomkin, Pub. Astron. Soc. Pac. 87 (1975) 859
- 1975MA21 R.E. Marrs, E.G. Adelberger, K.A. Snover and M.D. Cooper, Phys. Rev. Lett. 35 (1975) 202
- 1975MA41 G. Mairle and G.J. Wagner, Nucl. Phys. A253 (1975) 253
- 1975MI01 S. Micheletti, M. Pignanelli and P. Guazzoni, Phys. Rev. C11 (1975) 64
- 1975NA15 K.G. Nair, H. Voit, C.W. Towsley, M. Hamm, J.D. Bronson and K. Nagatani, Phys. Rev. C12 (1975) 1575
- 1975NI1C Nikolaev, Olkhovskii, Totskii and Chinarov, Ukr. Fiz. Zh. 20 (1975) 1882
- 1975PA09 B.H. Patrick, E.M. Bowey, E.J. Winhold, J.M. Reid and E.G. Muirhead, J. Phys. (London) G1 (1975) 874
- 1975PA1F Parks, Peek and Goldstein, Int. J. Appl. Rad. Isotopes 26 (1975) 683
- 1975PU01 R.J. Puigh, K.W. Kemper, G.E. Moore and R.L. White, Nucl. Phys. A237 (1975) 1
- 1975RA22 V.K. Rasmussen and F.R. Metzger, Phys. Rev. C12 (1975) 706
- 1975RA29 D.C. Radford and A.R. Poletti, Nucl. Phys. A254 (1975) 183
- 1975RA33 L. Ray, G. Westfall, S.A.A. Zaidi and W.R. Coker, Phys. Lett. B59 (1975) 427
- 1975RO30 V.C. Rogers, V.J. Orphan, C.G. Hoot and V.V. Verbinski, Nucl. Sci. Eng. 58 (1975) 298

- 1975SC35 W.F.W. Schneider, B. Kohlmeyer, F. Puhlhofer and R. Bock, Nucl. Phys. A251 (1975) 331
- 1975SC42 W.F.W. Schneider, B. Kohlmeyer and F. Puhlhofer, Z. Phys. A275 (1975) 249
- 1975SI22 M. Sieminski, M. Sosnowski and P. Zupranski, Nukleonika 20 (1975) 899
- 1975TR07 S. Tryti, T. Holtebekk and F. Ugletveit, Nucl. Phys. A251 (1975) 206
- 1975VA19 O.I. Vasileva, V.M. Lebedev, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 22 (1975) 446; Sov. J. Nucl. Phys. 22 (1976) 229
- 1975VO05 H. Voit, M. Hamm, K.G. Nair, C. Towsley, A.D. Bacher, K. Nagatani and F. Siller, Phys. Lett. B58 (1975) 152
- 1975WI26 K.H. Wilcox, R.B. Weisenmiller, G.J. Wozniak, N.A. Jelley, D. Ashery and J. Cerny, Phys. Lett. B59 (1975) 142
- 1975ZA08 N.I. Zaika, Y.V. Kibkalo, A.V. Mokhnach, O.F. Nemets, V.S. Semenov, F.P. Tokarev and P.L. Shmarin, Yad. Fiz. 21 (1975) 460; Sov. J. Nucl. Phys. 21 (1975) 241
- 1975ZI1A Zimmerman, Fowler and Caughlan, OAP-399 (1975)
- 1976AB04 A. Abul-Magd and J. Hufner, Z. Phys. A277 (1976) 379
- 1976AH09 I. Ahmad and Z.A. Khan, Nucl. Phys. A274 (1976) 519
- 1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1
- 1976AS1B Aslanides, Meson-Nucl. Phys., 1976 (1976) 204
- 1976AU1E Auerbach and Warszawski, Meson-Nucl. Phys., 1976 (1976) 322
- 1976BA16 J. Barrette, P. Braun-Munzinger, C.K. Gelbke, H.L. Harney, J. Kuzminski and I. Tseruya, Nucl. Phys. A261 (1976) 491
- 1976BE1C Beaudet, J. Roy. Astron. Soc. Can. 70 (1976) 247
- 1976BE1H Berman, Preprint UCRL-78482 (1976)
- 1976BE1K Bertini, Spallation Nucl. Reactions and Their Appl.; Eds., Shen and Merker (1976) 27
- 1976BE28 D. Berghofer, M.D. Hasinoff, R. Helmer, S.T. Lim, D.F. Measday and K. Ebisawa, Nucl. Phys. A263 (1976) 109
- 1976BI1A A. Billoire and M. Chemtob, Phys. Lett. B64 (1976) 389
- 1976BO1M Bouvier and Weibel, Arch. Sci. (Geneva) 29 (1976) 163
- 1976BO2C Bornec et al., Meson-Nucl. Phys., 1976 (1976) 260
- 1976BR1H Brown, Starrfield and Waller, Bull. Amer. Phys. Soc. 21 (1976) 587
- 1976BR1J Brook, Chisholm, Garrett and Duder, Polarization, Zurich, 1975 (1976) 473
- 1976BR1K Brown, Godfrey, Storey and Clark, Nature 262 (1976) 672
- 1976BR1L Brissaud, J. Phys. (London) G3 (1977) 1497

- 1976BR26 F. Brut and S. Jang, Phys. Rev. C14 (1976) 1638
- 1976BU16 M. Buenerd, C.K. Gelbke, B.G. Harvey, D.L. Hendrie, J. Mahoney, A. Menchaca-Rocha, C. Olmer and D.K. Scott, Phys. Rev. Lett. 37 (1976) 1191
- 1976BU1D Burov, Lukyanov and Titov, Proc. Dubna Conf., JINR D-9920, Vol. II (1976) 432
- 1976CA13 R. Casparis, B.T. Leemann, M. Preiswerk, H. Rudin, R. Wagner and P. Zupranski, Nucl. Phys. A263 (1976) 285
- 1976CO1N Cocu et al., CEA-R-4746 (1976)
- 1976CR1C Crawford, Fukuda and Syme, Lowell Conf., CONF-760715-P2 (1976) 1277
- 1976CU08 P. Cuzzocrea, G. Inghima, E. Perillo, E. Rosato, M. Sandoli and G. Spadaccini, Nuovo Cim. A31 (1976) 182
- 1976DE1F Dearborn, Eggleton and Schramm, Astrophys. J. 203 (1976) 455
- 1976DE1G Despain, OAP-459 (1976)
- 1976DE1H Dearborn and Eggleton, Quart. J. Royal Astron. Soc. 17 (1976) 448
- 1976DI11 M. Dillig and M.G. Huber, Lett. Nuovo Cim. 16 (1976) 299
- 1976DU04 J.L. Durell, P.J.A. Buttle, L.J.B. Goldfarb, W.R. Phillips, G.D. Jones, B.W. Hooton and M. Ivanovich, Nucl. Phys. A269 (1976) 443
- 1976DU08 R. Dumitrescu, A. Ciocanel and V. Corcalciuc, Rev. Roum. Phys. 21 (1976) 863
- 1976DY05 K. Dybdal, J.L. Eberhardt and N. Rud, Phys. Lett. B64 (1976) 414
- 1976ED1A Edge, Tompkins and Glenn, Nuovo Cim. A31 (1976) 641
- 1976EN02 H.D. Engelhardt, C.W. Lewis and H. Ullrich, Nucl. Phys. A258 (1976) 480
- 1976EP1A Epsteain, Arnett and Schramm, Astrophys. J. Suppl. 31 (1976) 111
- 1976ES1B Escusie, Fabroco, Pignanelli and Resmini, Polarization, Zuriich, 1975 (1976) 721
- 1976FE1B Feshbach, Meson-Nucl. Phys., 1976 (1976) 521
- 1976FE1C Feroni et al., Nuovo Cim. A35 (1976) 15
- 1976FI1E Fisher et al., Astrophys. J. 205 (1976) 938
- 1976FU06 G.H. Fuller, J. Phys. Chem. Ref. Data 5 (1976) 835
- 1976FU1C Fujita and Tsuji, Proc. Jpn. Acad. 52 (1976) 296
- 1976GA06 H. Garcilazo, Phys. Rev. C13 (1976) 1753
- 1976GA1F Gautier, Thompson, Fink and Larson, Astrophys. J. 205 (1976) 841
- 1976GA21 A. Gal and J.M. Eisenberg, Phys. Rev. C14 (1976) 1273
- 1976GA27 J. Gass and H.H. Muller, Nucl. Instrum. Meth. Phys. Res. 136 (1976) 559
- 1976GAYV D.I. Garber and R.R. Kinsey, BNL 325, 3rd Edition, Vol. 2 (1976)
- 1976GE1A Geiger and Van der Zwan, PXNR 2404 (1976)

- 1976GE20 H. Gemmeke, D. Baumgartner, A. Richter and L. Lassen, *Z. Phys.* A279 (1976) 425
- 1976GL11 D.W. Glasgow, F.O. Purser, H. Hogue, J.C. Clement, K. Stelzer, G. Mack, J.R. Boyce, D.H. Epperson, S.G. Buccino, P.W. Lisowski et al., *Nucl. Sci. Eng.* 61 (1976) 521
- 1976GO20 U. Gotz, F. Rosel, D. Trautmann and H. Jochim, *Z. Phys.* A278 (1976) 139
- 1976HA1J Hanna, Polarization, Zurich, 1975 (1976) 407
- 1976HA1N Haouat, Lachkar, Patin and Sigaud, *Acta Phys. Slovaca (Czech.)* 26 (1976) 51
- 1976HE17 E.F. Hefter and V.M. Semjonov, *Acta Phys. Pol.* B7 (1976) 427
- 1976HI05 M.D. High and B. Cujec, *Nucl. Phys.* A259 (1976) 513
- 1976HI1D Hinkle, Lambert and Snell, *Astrophys. J.* 210 (1976) 684
- 1976HO19 K.R. Hogstrom, *Phys. Rev.* C14 (1976) 753
- 1976IK01 P.G. Ikossi, W.J. Thompson, T.B. Clegg, W.W. Jacobs and E.J. Ludwig, *Phys. Rev. Lett.* 36 (1976) 1357
- 1976JA17 R. Jahn, D.P. Stahel, G.J. Wozniak, J. Cerny and H.P. Morsch, *Phys. Lett.* B65 (1976) 339
- 1976JA1D Jain, *Pramana* 5 (1976) 226
- 1976JE1A Jeukenne, Lejeune and Mahaux, Lowell Conf., Conf-760715-P1 (1976) 451
- 1976JO1B Joglekar, *Ann. Phys.* 101 (1976) 433
- 1976KA23 O. Karban, A.K. Basak, J.B.A. England, G.C. Morrison, J.M. Nelson, S. Roman and G.G. Shute, *Nucl. Phys.* A269 (1976) 312
- 1976KI1D King et al., Lowell Conf., CONF-760715-P2 (1976) 1344
- 1976KI1F Kirk, Lowell Conf., CONF-760715-P1 (1976) 389
- 1976KO22 R. Koch and H.H. Thies, *Nucl. Phys.* A272 (1976) 296
- 1976KO36 K. Koyama, *J. Phys. Soc. Jpn.* 41 (1976) 1445
- 1976LA1F Lamontagne et al., Polarization, Zurich, 1975 (1976) 707
- 1976LE12 J.R. Leigh, E.C. Pollacco, N. Shikazono, D.C. Weisser and C. Garrett, *J. Phys. (London)* G2 (1976) L39
- 1976LE1F Lefort, *Rept. Prog.* 39 (1976) 129
- 1976LE1K Lebedev et al., *Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682* (1976) 98
- 1976LE1M Lesiecki et al., Polarization, Zurich, 1975 (1976) 591
- 1976LI04 L.C. Liu and V. Franco, *Phys. Lett.* B61 (1976) 454
- 1976LI1H Liskien, Lowell Conf., CONF-760715-P2 (1976) 1110
- 1976LO1B Lorenzen, *Nucl. Instrum. Meth. Phys. Res.* 136 (1976) 289

- 1976MA06 C. Maroni, I. Massa and G. Vannini, Phys. Lett. B60 (1976) 344
- 1976MA1N Matsakis, Chui, Goldsmith and Townes, Astrophys. J. 206 (1976) L63
- 1976MA26 V.A. Matusevich, I.P. Chernov and V.V. Kozyr, Yad. Fiz. 23 (1976) 697; Sov. J. Nucl. Phys. 23 (1976) 367
- 1976MA55 C. Maroni, I. Massa and G. Vannini, Nucl. Phys. A273 (1976) 429
- 1976MA62 J.B. Martins, O.A.P. Tavares, M.L. Terranova and V. di Napoli, Lett. Nuovo Cim. 15 (1976) 364
- 1976ME18 H.O. Meyer, W.G. Weitkamp, J.S. Dunham, T.A. Trainor and M.P. Baker, Nucl. Phys. A269 (1976) 269
- 1976ME22 H.O. Meyer, G.R. Plattner and I. Sick, Z. Phys. A279 (1976) 41; Erratum Z. Phys. A282 (1977) 240
- 1976MI14 L.D. Miller and H.J. Weber, Phys. Lett. B64 (1976) 279
- 1976MO03 T. Motobayashi, I. Kohno, K. Katori, M. Yoshie, T. Ohi and H. Kamitsubo, Phys. Rev. Lett. 36 (1976) 390
- 1976NA09 K.G. Nair, K. Nagatani and H. Voit, Phys. Rev. Lett. 36 (1976) 1293
- 1976NI01 W. Niessner and K. Ilgen, Z. Phys. A277 (1976) 189
- 1976NI05 H. Nishimura and K. Kubodera, Prog. Theor. Phys. 56 (1976) 986
- 1976NO1C Norgaard and Fricke, Astron. Astrophys. 49 (1976) 337
- 1976NO1D Noble, Meson-Nucl. Phys., 1976 (1976) 221
- 1976NO1F Nordberg, Nilsson, Conde and Stromberg, Lowell Conf., CONF-760715-P2 (1976) 1318
- 1976OS06 E. Oset, Phys. Lett. B65 (1976) 46
- 1976OS07 A. Osman, Nuovo Cim. A35 (1976) 201
- 1976PA25 N.N. Pavlova and A.V. Yushkov, Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 820; Bull. Acad. Sci. USSR Phys. Ser. 40 (1976) 110
- 1976PL1C Plavko et al., Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 828
- 1976QU1A Querci and Querci, Astron. Astrophys. 49 (1976) 443
- 1976RE1B Reimers, Astron. Astrophys. 52 (1976) 457
- 1976RO10 J.L. Romero, F.P. Brady and J.A. Jungerman, Nucl. Instrum. Meth. 134 (1976) 537
- 1976RO12 C.T. Roche, R.G. Clark, G.J. Mathews and V.E. Viola, Jr., Phys. Rev. C14 (1976) 410
- 1976RO1L Roman, Polarization, Zurich, 1975 (1976) 255
- 1976RO1Q Roughton et al., Astrophys. 205 (1976) 302
- 1976SE1F Seyler, Polarization, Zurich, 1975 (1976) 397
- 1976SH13 R. Shyam and S. Mukherjee, Phys. Rev. C13 (1976) 2099

1976SL2A I. Slaus, Lowell Conf., CONF-760715-P1 (1976) 272
 1976SP1B Sparks, Starrfield and Truran, *Astrophys. J.* 208 (1976) 819
 1976ST12 R.G. Stokstad, Z.E. Switkowski, R.A. Dayras and R.M. Wieland, *Phys. Rev. Lett.* 37 (1976) 888
 1976ST22 A. Strazzeri, *Nuovo Cim.* A34 (1976) 229; Erratum *Nuovo Cim.* A41 (1976) 592
 1976SW02 Z.E. Switkowski, R.G. Stokstad and R.M. Wieland, *Nucl. Phys.* A274 (1976) 202
 1976TA07 M. Tanaka, S. Ochi, T. Minamisono, A. Mizobuchi and K. Sugimoto, *Nucl. Phys.* A263 (1976) 1
 1976TA1D Tanifuji, Mikoshiba and Terasawa, Lowell Conf., CONF-760715-P2 (1976) 1241
 1976TE03 R.K. Tenhaken and P.A. Quin, *Nucl. Phys.* A271 (1976) 173
 1976TH10 M. Thumm and H. Lesiecki, *Z. Phys.* A278 (1976) 77
 1976TH1B Thumm et al., Polarization, Zurich, 1975 (1976) 799
 1976TO1C Tornow, Lisowski, Byrd and Walter, Lowell Conf., CONF-760715-P2 (1976) 1358
 1976TO1D Tomkin, Luck and Lambert, *Astrophys. J.* 210 (1976) 694
 1976TSZZ T.L. Tsin, S.L. Blatt, T.R. Donoghue, L.J. Dries and R.G. Seyler, *Bull. Amer. Phys. Soc.* 21 (1976) 636, HG4
 1976VA1C Van Giai, *Meson-Nucl.*, 1976 (1976) 507
 1976VA1D Van Paradijs and Kester, *Astron. Astrophys.* 53 (1976) 1
 1976VA29 G.V. Valskii, *Yad. Fiz.* 24 (1976) 270; *Sov. J. Nucl. Phys.* 24 (1976) 140
 1976VI1B Vigroux, Audouze and Lequeux, *Astron. Astrophys.* 52 (1976) 1
 1976WA15 G.L. Wales and R.C. Johnson, *Nucl. Phys.* A274 (1976) 168
 1976WA1B Walter, Polarization, Zurich, 1975 (1976) 377
 1976WA1C Walter and Lisowski, Lowell Conf., CONF-760715-P2 (1976) 1061
 1976WA1G Watson, Anicich and Huntress, *Astrophys. J.* 205 (1976) L165
 1976WE01 H.R. Weller, R.A. Blue, S.E. Darden and W.A. Yoh, *Phys. Rev.* C13 (1976) 38
 1976WE1E Westkamp et al., Polarization, Zurich, 1975 (1976) 593
 1976WE1F Westkamp et al., Polarization, Zurich, 1975 (1976) 595
 1976WE21 G.D. Westfall and S.A.A. Zaidi, *Phys. Rev.* C14 (1976) 610
 1976WE28 U. Weiss, D. Fick, K.D. Hildenbrand, W. Weiss, G.R. Plattner and I. Sick, *Nucl. Phys.* A274 (1976) 253
 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, *Phys. Rev.* C14 (1976) 815
 1976WUZZ H.P. Wu, E.J. Ludwig, W.W. Jacobs, P.G. Ikossi, S.A. Tonsfeldt and W.J. Thompson, *Bull. Amer. Phys. Soc.* 21 (1976) 514, AF8

- 1977AJ02 F. Ajzenberg-Selove, Nucl. Phys. A281 (1977) 1
- 1977AL18 D.J. Albert, A. Nagl, J. George, R.F. Wagner and H. Uberall, Phys. Rev. C16 (1977) 503
- 1977AL1H Albert et al., Bull. Amer. Phys. Soc. 22 (1977) 612
- 1977AL1J Albert et al., Proc., Zurich-Sin. (1977) 246
- 1977AL25 G.D. Alkhazov, S.L. Belostotsky, O.A. Domchenkov, Y.V. Dotsenko, E.G. Kormin, N.P. Kuropatkin, V.N. Nikulin, S.S. Volkov, A.A. Vorobyov and M.A. Shuvaev, Phys. Lett. B70 (1977) 20
- 1977AM1C Amakawa, Mori and Yazaki, in Tokyo (1977) 485
- 1977AN1F Anttila, Bister, Fontell and Winterbon, Rad. Effects 33 (1977) 13
- 1977AN1G Antonov and Garistov, Bulg. J. Phys. 4 (1977) 6
- 1977AN21 I. Angeli and M. Csatlos, Nucl. Phys. A288 (1977) 480
- 1977AR06 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A283 (1977) 350
- 1977AS03 V.M. Asaturyan, E.O. Grigoryan, G.A. Vartapetyan, A.S. Danagulyan, N.A. Demekhina and A.G. Khudaverdyan, Yad. Fiz. 25 (1977) 1133; Sov. J. Nucl. Phys. 25 (1977) 601
- 1977AU01 N. Auerbach, Phys. Rev. Lett. 38 (1977) 804
- 1977AU1B Austin, MSUCL-254 (1977)
- 1977AU1E Audouze, CNO Isotopes in Astrophys.; Ed., J. Audouze (1977) 3
- 1977AU1F Audouze, Lequeux, Rocca-Volmerange and Vigroux, CNO Isotopes in Astrophys.; Ed., J. Audouze (1977) 155
- 1977AV01 V.V. Avdeychikov, V.I. Bogatin and O.V. Lozhkin, Yad. Fiz. 25 (1977) 3; Sov. J. Nucl. Phys. 25 (1977) 1
- 1977AZ01 G. Azuelos, J.E. Kitching and K. Ramavataram, Phys. Rev. C15 (1977) 1847
- 1977AZ02 G. Azuelos and J.E. Kitching, Nucl. Phys. A285 (1977) 19
- 1977BA1L Badovskii et al., Sov. At. Energy 43 (1977) 838
- 1977BA1P Batusov and Eramzhyan, Fiz. Elem. Chastits At. Yad. (USSR) 8 (1977) 229
- 1977BA1V Bahcall, AIP Conf. Proc. 37 (1977) 53
- 1977BA29 G. Baur, Nucl. Phys. A283 (1977) 521
- 1977BA2G Barnes, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 783
- 1977BA2H Bacher et al., in Tokyo (1977) 812

- 1977BA48 W. Bambynek, H. Behrens, M.H. Chen, B. Crasemann, M.L. Fitzpatrick, K.W.D. Ledingham, H. Genz, M. Mutterer and R.L. Intemann, *Rev. Mod. Phys.* 49 (1977) 77; Erratum *Rev. Mod. Phys.* 49 (1977) 961
- 1977BE1Y Berceanu et al., *Nucl. Instrum. Meth. Phys. Res.* 146 (1977) 319
- 1977BO30 I.I. Bondouk and A.S. Yousef, *Atomkernenergie* 30 (1977) 69
- 1977CA1H Casse, CAP-497 (1977)
- 1977CA1J Carbon, Butler, Kraft and Nocar, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 33
- 1977CA1K Caughlan, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 121
- 1977CE05 J. Cerny and J.C. Hardy, *Ann. Rev. Nucl. Sci.* 27 (1977) 333
- 1977CH1J Chruchwel, Walmsley and Winnewisser, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 99
- 1977CH22 R. Chechik, Z. Fraenkel, H. Stocker and Y. Eyal, *Nucl. Phys.* A287 (1977) 353
- 1977CI1A Ciric et al., *Fiz. Suppl.* 9 (1977) 39
- 1977CL1B Climenhaga, Harris, Holts and Smolinski, *Astrophys. J.* 215 (1977) 836
- 1977CL1C Clayton, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 15
- 1977CO1J Combes, Maillard and De. Bergh, *Astron. Astrophys.* 61 (1977) 531
- 1977DA18 J.R. Davis and G.U. Din, *Aust. J. Phys.* 30 (1977) 167
- 1977DA1B Davydov, Naumov and Shcherbachenko, INDC(CCP)-104/LN (1977)
- 1977DA1E Dahn et al., *Astrophys. J.* 216 (1977) 757
- 1977DE1N Dearborn, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 39
- 1977DI11 J.K. Dickens, G.L. Morgan, G.T. Chapman, T.A. Love, E. Newman and F.G. Perey, *Nucl. Sci. Eng.* 62 (1977) 515
- 1977DI1B M. Dillig and M.G. Huber, *Phys. Lett.* B69 (1977) 429
- 1977DI1C Dickman, Langer, McCutcheon and Shuter, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 95
- 1977DM1A Dmitrieva, Lebedev and Matusevich, *Izv. Akad. Nauk SSSR Ser. Fiz.* 41 (1977) 143
- 1977DO06 G.E. Dogotar, R.A. Eramzhan, H.R. Kissener and R.A. Sakaev, *Nucl. Phys.* A282 (1977) 474
- 1977EG1A Egawa et al., in Tokyo (1977) 489
- 1977FI1A Filleux, Tomerello and Burnett, 8th Lunar Sci. Conf., Houston, TX (1977)
- 1977FI1B Fitzgerald et al., *Bull. Amer. Phys. Soc.* 22 (1977) 541
- 1977FO04 P. Fontes, *Phys. Rev.* C15 (1977) 2159

- 1977FR20 T. Freye, H. Lorenz-Wirzba, B. Cleff, H.P. Trautvetter and C. Rolfs, *Z. Phys.* A281 (1977) 211
- 1977FU10 S. Furui, *Prog. Theor. Phys.* 57 (1977) 2001
- 1977FU1G Fujisawa et al., in Tokyo (1977) 510
- 1977FU1H Fujisawa et al., in Tokyo (1977) 511
- 1977FU1K R.C. Fuller, *Phys. Rev. C*16 (1977) 1865
- 1977FU1L Fujita and Tsuji, *Publ. Astron. Soc. Jpn.* 29 (1977) 711
- 1977GI06 W.R. Gibbs and A.T. Hess, *Phys. Lett.* B68 (1977) 205
- 1977GO1D Gonclaves and Santos, in Tokyo (1977) 597
- 1977GU07 H.P. Gubler, G.R. Plattner, I. Sick, A. Traber and W. Weiss, *Nucl. Phys.* A284 (1977) 114
- 1977GU14 M.G. Gulyamov, B.S. Mazitov, G.A. Radyuk, E.I. Rasulov and R. Yarmukhamedov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 41 (1977) 2214; *Bull. Acad. Sci. USSR Phys. Ser.* 41 (1977) 189
- 1977HA1L Hagen, Fisher and Ormes, *Astrophys. J.* 212 (1977) 262
- 1977HA1N Hasegawa et al., in Tokyo (1977) 539
- 1977HE12 J.A.J. Hermans, G.A.P. Engelbertink, L.P. Ekstrom, H.H. Eggenhuisen and M.A. Van Driel, *Nucl. Phys.* A284 (1977) 307
- 1977HE1D Hermans, Thesis, Utrecht Univ. (1977)
- 1977HE26 J.C.P. Heggie and Z.E. Switkowski, *Nucl. Instrum. Meth. Phys. Res.* 147 (1977) 425
- 1977HO1B Hoistad, *High Energy Phys. Nucl. Struct., Zurich, 1977* (1977) 215
- 1977HO32 J. Hohn, H.W. Barz and I. Rotter, *Nukleonika* 22 (1977) 1175
- 1977JO1C Jones et al., *Int. J. Appl. Rad. Isotopes* 28 (1977) 25
- 1977JO1D Joss, *Nature* 270 (1977) 310
- 1977KA26 D. Kalinsky, D. Melnik, U. Smilansky, N. Trautner, S. Weisrose and O. Dietzsch, *Nucl. Phys.* A293 (1977) 509
- 1977KO1P Koshel, *AIP Conf. Proc.* 36 (1977) 227
- 1977KU1G Kume and Ohtsubo, *Proc., Zurich-Sin* (1977) 18
- 1977LA1F Lambert and Sneden, *Astrophys. J.* 215 (1977) 597
- 1977LA1G Lambert and Ries, *Astrophys. J.* 217 (1977) 508
- 1977LA1H Lachkar, *Proc. Int. Specialists Symp. on Neutron Standards and Applications, 1977* (1977) 93
- 1977LE1H Lee, MacFarlane and Landau, *Bull. Amer. Phys. Soc.* 22 (1977) 81
- 1977LE1J Leach, Thomas and Vartsky, *Int. J. Appl. Rad. Isotopes* 28 (1977) 263

- 1977LI19 H. Liskien and A. Paulsen, *Atomkernenergie* 30 (1977) 59
- 1977LI1H Linke et al., *Astrophys. J.* 214 (1977) 50
- 1977LO19 K.P. Lohs and A. Gal, *Nucl. Phys.* A292 (1977) 375
- 1977MA06 M. Marangoni, P.L. Ottaviani and A.M. Saruis, *Nucl. Phys.* A277 (1977) 239
- 1977MA16 R.E. Marrs, E.G. Adelberger and K.A. Snover, *Phys. Rev.* C16 (1977) 61
- 1977MA1M Mausner et al., *Bull. Amer. Phys. Soc.* 22 (1977) 547
- 1977ME05 H.O. Meyer and G.R. Plattner, *Nucl. Phys.* A279 (1977) 53
- 1977ME06 H.O. Meyer, T.P. Cleary and W. Haeberli, *Nucl. Instrum. Meth.* 140 (1977) 85
- 1977ME1D Meeks et al., MSUCL-228 (1977)
- 1977ME1E Meder, *Bull. Amer. Phys. Soc.* 22 (1977) 577
- 1977MO1A T. Motobayashi, *Rept. Inst. Phys. Chem. Res.* 53 (1977) 81
- 1977MU1A Mukhopadhyay, *Phys. Rept.* C30 (1977) 1
- 1977MU1C Murnick and Kantner, in *Tokyo* (1977) 744
- 1977NA16 V.S. Nadezhdin, N.I. Petrov and V.I. Satarov, *Yad. Fiz.* 26 (1977) 230; *Sov. J. Nucl. Phys.* 26 (1977) 119
- 1977NA29 V.S. Nadezhdin, N.I. Petrov and V.I. Satarov, *Pisma Zh. Eksp. Teor. Fiz.* 26 (1977) 123; *JETP Lett.* 26 (1977) 115
- 1977NO07 J.M. Normand, *Nucl. Phys.* A291 (1977) 126
- 1977OS08 A. Osman, *Nuovo Cim.* A39 (1977) 138
- 1977PA1D Panagia, Bussoletti and Blanco, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 45
- 1977PE23 T. Pedersen, T. Engeland, I. Espe, J. Rekstad, C Ellegaard and S. Backe, *Nucl. Phys.* A293 (1977) 10
- 1977PH02 D.-L. Pham and R. de Swiniarski, *Nuovo Cim.* A41 (1977) 543
- 1977PR1E Prialnick, Shara and Shavin, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 63
- 1977PU02 J.E. Purcell and M.R. Meder, *Phys. Rev.* C16 (1977) 76
- 1977RE08 H. Reiss, H.V. Klapdor, G. Rosner, J.L.C. Ford, Jr. and S.T. Thornton, *Nucl. Phys.* A283 (1977) 149
- 1977RI08 W.A. Richter and P.R. de Kock, *Z. Phys.* A282 (1977) 207
- 1977SA19 A.M. Santana, H.T. Coelho and T.K. Das, *Phys. Rev.* C16 (1977) 1785
- 1977SA1B Sadkovskii, Feofilov, Denisov and Kolalis, *Yad. Fiz.* 26 (1977) 1200
- 1977SA1L Sakaguchi et al., in *Tokyo* (1977) 575
- 1977SA25 M. Sano, M. Tanaka and M. Wakai, *Prog. Theor. Phys.* 58 (1977) 1673

1977SC1G Schroder and Huizenga, *Ann. Rev. Nucl. Sci.* 27 (1977) 465
 1977SC1H Scalo, *Astrophys. J.* 215 (1977) 194
 1977SE09 F. Seiler, *Nucl. Phys.* A286 (1977) 1
 1977SE1D Seya et al., in *Tokyo* (1977) 149
 1977SH09 M.A.M. Shahabuddin, C.J. Webb and V.R.W. Edwards, *Nucl. Phys.* A284 (1977) 83
 1977SH1D Shimoda et al., in *Tokyo* (1977) 703
 1977SH1E Shackleton and Opdyke, *Nature* 270 (1977) 216
 1977SI1D Silberberg and Tsao, *Astrophys. J. Suppl.* 35 (1977) 129
 1977SM1D Smith and Whalen, *ANL/NDM-33* (1977)
 1977ST1H Starrfield, Trjran and Sparks, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 49
 1977ST1M Straatmann, *Int. J. Appl. Rad. Isotopes* 28 (1977) 13
 1977ST1N Stepancic, Ciric, Subotic and Stanojevic, *Fiz. Suppl.* 9 (1977) 41
 1977ST20 D.P. Stahel, G.J. Wozniak, M.S. Zisman, B.D. Jeltema and J. Cerny, *Phys. Rev.* C16 (1977) 1456
 1977SW05 Z.E. Switkowski, Shiu-Chin Wu, J.C. Overley and C.A. Barnes, *Nucl. Phys.* A289 (1977) 236
 1977TA08 M. Takai, T. Kambara, K. Tada, M. Nakamura and S. Kobayashi, *J. Phys. Soc. Jpn.* 43 (1977) 17
 1977TE01 W.D. Teeters and D. Kurath, *Nucl. Phys.* A275 (1977) 61
 1977TI1A Tinsley, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 175
 1977TR09 D. Trautmann and G. Baur, *Nukleonika* 22 (1977) 485
 1977TR1A Traber, Trautman and Alder, *Helv. Phys. Acta* 50 (1977) 195
 1977TR1D Truran, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 145
 1977TS1A Tsin, Seyler and Blatt, *Bull. Amer. Phys. Soc.* 22 (1977) 612
 1977UD1A Udagawa, *Proc. Int. Conf. Nucl. Struct.*, Tokyo (1977); *J. Phys. Soc. Jpn. Suppl.* 44 (1978) 667
 1977UH1A Uhlhorn, Lange and Gonsoir, in *Tokyo* (1977) 610
 1977WA02 J. Warszawski and N. Auerbach, *Nucl. Phys.* A276 (1977) 402
 1977WA1N Watson, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 105
 1977WA1P Wannier, *CNO Isotopes in Astrophys.*; Ed., J. Audouze (1977) 71
 1977WE1G Weber and Miller, *Proc.*, Zurich-Sin (1977) 195
 1977WE1H Werby, Nagarajan, Goldfinger and Tobocman, *Fiz. Suppl.* 9 (1977) 46
 1977WI1C Willett et al., in *Tokyo* (1977) 641

- 1977WI1D Wilson and Biegging, CNO Isotopes in Astrophys.; Ed., J. Audouze (1977) 89
- 1977WO04 J.G. Woodworth, K.G. McNeill, J.W. Jury, P.D. Georgopoulos and R.G. Johnson, Can. J. Phys. 55 (1977) 1704
- 1977YA1B Yakovlev, Fiz. Elem. Chastits At. Yad. (USSR) 8 (1977) 255
- 1977YO1D Yoro and Arima, in Tokyo (1977) 170
- 1977YO1E K. Yoro, Phys. Lett. B70 (1977) 147
- 1977YO1F Young, Golden and Kapetanakos, Rev. Sci. Instrum. 48 (1977) 432
- 1978AB08 A.Y. Abul-Magd and J. Hufner, Nucl. Phys, A308 (1978) 429
- 1978AD1A Adams and White, Bull. Amer. Phys. Soc. 23 (1978) 501
- 1978AH04 S.S. Ahmad, J. Phys. (London) G4 (1978) 1751
- 1978AJ02 F. Ajzenberg-Selove, E.R. Flynn and O. Hansen, Phys. Rev. C17 (1978) 1283
- 1978AL29 D.E. Alburger, S. Mordechai, H.T. Fortune and R. Middleton, Phys. Rev. C18 (1978) 2727
- 1978AL36 G.D. Alkhazov, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 2218; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 1
- 1978AN07 I. Angeli and M. Csatlos, Atomki Kozlem. 20 (1978) 1
- 1978AR1J Arefev et al., Yad. Fiz. 28 (1978) 1534
- 1978AR1N Arthur et al., Bull. Amer. Phys. Soc. 23 (1978) 62
- 1978AU07 E.G. Auld, A. Haynes, R.R. Johnson, G. Jones, T. Masterson, E.L. Mathie, D. Ottewell, P. Walden and B. Tatischeff, Phys. Rev. Lett. 41 (1978) 462
- 1978AV1A Avishai, Z. Phys. A285 (1978) 333
- 1978AZ02 L.S. Azhgirei, I.K. Vzorov, V.N. Zhmyrov, V.V. Ivanov, M.A. Ignatenko, A.S. Kuznetsov, M.G. Meshcheryakov, S.V. Razin and G.D. Stoletov, Yad. Fiz. 28 (1978) 1017; Sov. J. Nucl. Phys. 28 (1978) 522
- 1978BA1M Bading and Maxson, Bull. Amer. Phys. Soc. 23 (1978) 525
- 1978BA21 G. Baur, F. Roesel and D. Trautmann, J. Phys. (London) G4 (1978) 275
- 1978BA26 B.F. Bayman, A. Dudek-Ellis and P.J. Ellis, Nucl. Phys. A301 (1978) 141
- 1978BE49 G. Bendiscioli, E. Lodi Rizzini, A. Rotondi and A. Venaglioni, Nucl. Phys. A307 (1978) 22
- 1978BE66 Y.A. Bereznoi and N.A. Shlyakhov, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 743; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 47
- 1978BH1B Bhakar and Van Oers, Few Body Syst. Nucl. Forces, Graz, 1978, Springer Lect. Notes 87 (1978) 364

- 1978BI08 R. Billerey, C. Cerruti, A. Chevarier, N. Chevarier and A. Demeyer, *Z. Phys.* A285 (1978) 389
- 1978BL02 G.S. Blanpied, W.R. Coker, R.P. Liljestrang, G.W. Hoffmann, L. Ray, D. Madland, C.L. Morris, J.C. Pratt, J.E. Spencer, H.A. Thiessen et al., *Phys. Rev. C*18 (1978) 1436
- 1978BO1L Borders and Harris, *Nucl. Instrum. Meth. Phys. Res.* 149 (1978) 279
- 1978BO31 M. Bouten, E. Flerackers and M.C. Bouten, *Nucl. Phys.* A307 (1978) 413
- 1978BU1B Buffington, Orth and Mast, *Astrophys. J.* 226 (1978) 355
- 1978CA03 G. Cattapan, E. Maglione, G. Pisent and V. Vanzani, *Nucl. Phys.* A296 (1978) 263
- 1978CH02 H.C. Cheung, M.D. High and B. Cujec, *Nucl. Phys.* A296 (1978) 333
- 1978CH03 R. Chechik, Y. Eyal, H. Stocker and Z. Fraenkel, *Nucl. Phys.* A296 (1978) 307
- 1978CH16 R. Chechik, Z. Fraenkel, Y. Eyal and H. Stocker, *Nucl. Phys.* A304 (1978) 243
- 1978CH29 L.T. Chua, A. Gobbi, M.W. Sachs, D. Shapira, R. Wieland and D.A. Bromley, *Nuovo Cim.* A47 (1978) 430
- 1978CH30 L.T. Chua and D.A. Bromley, *Nuovo Cim.* A47 (1978) 443
- 1978CO07 J.P. Coffin, P. Engelstein, A. Gallmann, B. Heusch, P. Wagner and H.E. Wegner, *Phys. Rev. C*17 (1978) 1607
- 1978CO15 P. Couvert, G. Bruge, R. Beurtey, A. Boudard, A. Chaumeaux, M. Garcon, D. Garreta, P.C. Gugelot, G.A. Moss, S. Platchkov et al., *Phys. Rev. Lett.* 41 (1978) 530
- 1978CO1J J.R. Comfort and B.C. Karp, *Phys. Lett.* B80 (1978) 1
- 1978DA17 S.K. Datta, G.P.A. Berg and P.A. Quin, *Nucl. Phys.* A312 (1978) 1
- 1978DA1A Dalitz and Gal, *Ann. Phys.* 116 (1978) 167
- 1978DE15 P. Desgrolard, P.A.M. Guichon and J. Joseph, *Nuovo Cim.* A43 (1978) 475
- 1978DE1Q Delbrouck-Habaru, Robaye and Winand, *Nucl. Instrum. Meth. Phys. Res.* 152 (1978) 521
- 1978DE1R Deming, *Astrophys. J.* 223 (1978) L31
- 1978DO1B Dominy et al., *Astrophys. J.* 223 (1978) 949
- 1978DR07 W. Dreves, P. Zupranski, P. Egelhof, D. Kassen, E. Steffens, W. Weiss and D. Fick, *Phys. Lett.* B78 (1978) 36
- 1978DW1A Dwek, *Astrophys. J.* 225 (1978) L149
- 1978DY01 S.A. Dytman, J.F. Amann, P.D. Barnes, J.N. Craig, K.G.R. Doss, R.A. Eisenstein, J.D. Sherman, W.R. Wharton, G.R. Burleson, S.L. Verbeck et al., *Phys. Rev. C*18 (1978) 2316
- 1978DZ1A Dzhamalov, Dolinskii and Mukhamedzhanov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 2365

- 1978EM1A Emiliani, Hudson, Shinn and George, *Science* 202 (1978) 627
- 1978FA04 G. Faldt and I. Hulthage, *J. Phys. (London)* G4 (1978) 363
- 1978FI1F Firestone, Smith, Firestone and Tiedje, MSUCL-286 (1978)
- 1978FR12 S. Frankel, W. Frati, M. Gazzaly, G.W. Hoffmann, O. Van Dyck and R.M. Woloshyn, *Phys. Rev. Lett.* 41 (1978) 148
- 1978FU1G Fu and Perey, *At. Data Nucl. Data Tables* 22 (1978) 249
- 1978GE1C Gelbke et al., *Phys. Rept.* C42 (1978) 311
- 1978GO05 I.G. Golikov, M.N. Zhukov, I.I. Loshchakov and B.I. Ostroumov, *Yad. Fiz.* 27 (1978) 7; *Sov. J. Nucl. Phys.* 27 (1978) 3
- 1978GO1F J. Golden, R.A. Mahaffey, J.A. Pasour, F.C. Young and C.A. Kapetanacos, *Rev. Sci. Instrum.* 49 (1978) 1384
- 1978GR07 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.V. Ostashko, V.M. Pugach, V.A. Pilipchenko, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 147; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 126
- 1978GR1D Grigoryan and Shakhbazyan, *Yad. Fiz.* 28 (1978) 1372
- 1978GU10 K. Gul, *J. Phys. Soc. Jpn.* 44 (1978) 353
- 1978GU12 C. Gustafsson and E. Lambert, *Ann. Phys.* 111 (1978) 304
- 1978HA14 J.E. Halverson and W.H. Johnson, Jr., *Phys. Rev. C* 17 (1978) 1414; *Erratum Phys. Rev. C* 18 (1978) 2450
- 1978HA1P Haouat et al., *Nucl. Sci. Eng.* 65 (1978) 331
- 1978HA1Q Han, Chuang and He, *Phys. Energ. Fortis Phys. Nucl. (China)* 2 (1978) 249
- 1978HE1C Hendrie, *AIP Conf. Proc.* 47 (1978) 402
- 1978HE1D Heitzmann, *Atomkernenergie* 31 (1978) 262
- 1978HI06 F. Hinterberger, R. Schonhagen, P. von Rossen, B. Schuller, F.E. Blumenberg, P.D. Eversheim and R. Gorgen, *Nucl. Phys.* A308 (1978) 61
- 1978HI1D Hichwa and Nickles, *Bull. Amer. Phys. Soc.* 23 (1978) 1032
- 1978HO05 G.W. Hoffman, G.S. Blanpied, W.R. Coker, R.P. Liljestrang, N.M. Hintz, M.A. Oothoudt, T.S. Bauer, G. Igo, G. Pauletta, J. Soukup et al., *Phys. Rev. Lett.* 40 (1978) 1256
- 1978HO11 K. Hosono, M. Kondo, T. Saito, N. Matsuoka, S. Nagamachi, S. Kato, K. Ogino, Y. Kadota and T. Noro, *Phys. Rev. Lett.* 41 (1978) 621
- 1978HO1C Hodgson, *Riv. Nuovo Cim.* 1 (1978) 1
- 1978IB1A Iben and Truran, *Astrophys. J.* 220 (1978) 980

- 1978JA16 J. Jaros, A. Wagner, L. Anderson, O. Chamberlain, R.Z. Fuzesy, J. Gallup, W. Gorn, L. Schroeder, S. Shannon, G. Shapiro et al., Phys. Rev. C18 (1978) 2273
- 1978KA08 J. Kallne, A.W. Stetz and R.M. Woloshyn, Phys. Lett. B74 (1978) 170
- 1978KA1H Karadeniz, Rev. Fac. Sci., Univ. Istanbul C39 (1978) 108
- 1978KE1E Keeler, Mook and Tans, Nature 277 (1978) 121
- 1978KI08 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, Nucl. Phys. A302 (1978) 523
- 1978KI13 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, Nucl. Phys. A312 (1978) 394
- 1978KO01 D.G. Kovar, W. Henning, B. Zeidman, Y. Eisen, J.R. Erskine, H.T. Fortune, T.R. Ophel, P. Sperr and S.E. Vigdor, Phys. Rev. C17 (1978) 83
- 1978KR19 K.S. Krane, At. Data Nucl. Data Tables 22 (1978) 269
- 1978LA1H Lane, White, Knox and Auchampaugh, Bull. Amer. Phys. Soc. 23 (1978) 501
- 1978LE04 H.C. Lee, Nucl. Phys. A294 (1978) 473
- 1978LE10 P. Leleux, M. Bosman, P. Lipnik, P. Macq, J.P. Meulders, C. Pirart and G. Valenduc, Nucl. Instrum. Meth. 152 (1978) 495
- 1978LE15 R.D. Levine, S.G. Steadman, J.S. Karp and Y. Alhassid, Phys. Rev. Lett. 41 (1978) 1537
- 1978LE1N Legg and Phillips, Bull. Amer. Phys. Soc. 23 (1978) 616
- 1978LE1P Leventhal and Threlkeld, Science 202 (1978) 430
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, Table of Isotopes 7th Ed. (1978)
- 1978LI1D Li, Shen and Zhang, Phys. Energ. Fortis Phys. Nucl. (China) 3 (1979) 247
- 1978LU1C Luck, Astrophys. J. 219 (1978) 148
- 1978MA1F Malecki, Namyslowski, Reale and Minetti, Riv. Nuovo Cim. 1 (1978) 1
- 1978MA1P Markham, Austin and Styra, MSUCL-276 (1978)
- 1978MA34 B.S. Mazitov, G.A. Radyuk, E.N. Rasulov and R. Yarmukhamedov, Yad. Fiz. 28 (1978) 67; Sov. J. Nucl. Phys. 28 (1978) 33
- 1978MA42 N. Matsuoka, A. Shimizu, K. Hosono, T. Saito, M. Kondo, H. Sakaguchi, Y. Toba, A. Goto, F. Ohtani and N. Nakanishi, Nucl. Phys. A311 (1978) 173
- 1978MA44 J.F. Mateja, A.D. Frawley, A. Roy, J.R. Hurd and N.R. Fletcher, Phys. Rev. C18 (1978) 2622
- 1978MC04 A.B. McDonald and E.G. Adelberger, Phys. Rev. Lett. 40 (1978) 1692
- 1978ME12 E. Mezzetti, B. Minetti and A. Pasquarelli, Lett. Nuovo Cim. 22 (1978) 91

1978MI02 L.D. Miller and H.J. Weber, Phys. Rev. C17 (1978) 219
 1978MI04 M. Micklinghoff, Nucl. Phys. A295 (1978) 237
 1978MI1E M. Micklinghoff, Nucl. Phys. A295 (1978) 228
 1978NA15 P. Nagel, Phys. Rev. C18 (1978) 2617
 1978NA18 V.S. Nadezhdin, N.I. Petrov and V.I. Satarov, Yad. Fiz. 27 (1978) 1164; Sov. J. Nucl. Phys. 27 (1978) 617
 1978NE1B Needham et al., Bull. Amer. Phys. Soc. 23 (1978) 14
 1978NO02 R. Novotny, G. Hammer, D. Pelte, H. Emling and D. Schwalm, Nucl. Phys. A294 (1978) 255
 1978NO04 C. Nordborg, L. Nilsson, H. Conde and L.G. Stromberg, Nucl. Sci. Eng. 66 (1978) 75
 1978OS06 V.M. Osadchiev, Yad. Fiz. 27 (1978) 958; Sov. J. Nucl. Phys. 27 (1978) 507
 1978PA1B Park, Greiner and Scheid, Bull. Amer. Phys. Soc. 23 (1978) 538
 1978PE1E Peterson, Seglie and Ascutto, Bull. Amer. Phys. Soc. 23 (1978) 947
 1978PE1F Pettit, Bull. Amer. Phys. Soc. 23 (1978) 1034
 1978PO1A B. Povh, Ann. Rev. Nucl. Part. Sci. 28 (1978) 1
 1978QA01 S.M. Qaim and R. Wolfle, Nucl. Phys. A295 (1978) 150
 1978RA17 L. Ray, G.W. Hoffmann, G.S. Blanpied, W.R. Coker and R.P. Liljestrang, Phys. Rev. C18 (1978) 1756
 1978RA20 L. Ray and W.R. Coker, Phys. Lett. B79 (1978) 182
 1978RA2A S. Raman, C.A. Houser, T.A. Walkiewicz and I.S. Towner, At. Data Nucl. Data Tables 21 (1978) 567; Erratum At. Data Nucl. Data Tables 22 (1978) 369
 1978RE1D Redish, Alexander and Wallace, Bull. Amer. Phys. Soc. 23 (1978) 553
 1978RO1D Rolfs and Trautvetter, Ann. Rev. Nucl. Part. Sci. 28 (1978) 115
 1978RO1H Rost, Bull. Amer. Phys. Soc. 23 (1978) 964
 1978RY02 T.B. Ryves, P. Kolkowski and K.J. Zieba, J. Phys. (London) G4 (1978) 1783
 1978SC1E Scheid and Greiner, AIP Conf. Proc. 47 (1978) 374
 1978SE01 F. Seiler, F.N. Rad, H.E. Conzett and R. Roy, Nucl. Phys. A296 (1978) 205
 1978SH1B Shlomo, Rept. Prog. Phys. (GB) 41 (1978) 957
 1978SN1A Sneden, Lambert, Tomkin and Peterson, Astrophys. J. 222 (1978) 585
 1978SO1A Sokol, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 1829
 1978SO1B Sober et al., Bull. Amer. Phys. Soc. 24 (1978) 584
 1978ST1C Starrfield, Truran and Sparks, Astrophys. J. 226 (1978) 186
 1978ST1D Starrfield, Truran, Sparks and Arnould, Astrophys. J. 222 (1978) 600

1978SU1D Subramanian et al., Bull. Amer. Phys. Soc. 23 (1978) 12
 1978SU1E Subramanian et al., Bull. Amer. Phys. Soc. 23 (1978) 943
 1978TA09 T. Takemasa and H. Yoshida, Nucl. Phys. A304 (1978) 229
 1978TA1B Taras, AIP Conf. Proc. 47 (1978) 234
 1978TE02 G. Terlecki, W. Scheid, H.J. Fink and W. Greiner, Phys. Rev. C18 (1978) 265
 1978TI1A Tilbury et al., Bull. Amer. Phys. Soc. 23 (1978) 1033
 1978TI1B Tiedje et al., MSUCL-285 (1976)
 1978TR05 H.P. Trautvetter, M. Wiescher, K.-U. Kettner, C. Rolfs and J.W. Hammer, Nucl. Phys. A297 (1978) 489
 1978TR1C Truran, Cowan and Cameron, Astrophys. J. 222 (1978) L63
 1978UC1A Uchiyama, Prog. Theor. Phys. 59 (1978) 1025
 1978VA1B Vanysek and Rabe, Moom and the Planets 18 (1978) 441
 1978WA02 J. Warszawski, A. Gal and J.M. Eisenberg, Nucl. Phys. A294 (1978) 321
 1978WA1D Walter et al., Bull. Amer. Phys. Soc. 23 (1978) 926
 1978WA1E Wannier and Linke, Astrophys. J. 226 (1978) 817
 1978WE13 H.R. Weller, J. Szucs, P.G. Ikossi, J.A. Kuehner, D.T. Petty and R.G. Seyler, Phys. Rev. C18 (1978) 1120
 1978WE1D Wessels et al., Nucl. Instrum. Meth. 141 (1977) 157
 1978WE1F Werby and Bassichis, Bull. Amer. Phys. Soc. 23 (1978) 538
 1978WE1G H.J. Weber and J.M. Eisenberg, Nucl. Phys. A312 (1978) 201
 1978WI1F Weiland and Highfall, Bull. Amer. Phys. Soc. 23 (1978) 1033
 1978WO1C Wolf, Bull. Amer. Phys. Soc. 23 (1978) 1033
 1978WU03 J.R. Wu and C.C. Chang, Phys. Rev. C17 (1978) 1540
 1978ZA1D Zalm, Van Hienen and Glaudemans, Z. Phys. A287 (1978) 255
 1978ZE03 N.S. Zelenskaya, V.M. Lebedev and T.A. Yushchenko, Yad. Fiz. 28 (1978) 90; Sov. J. Nucl. Phys. 28 (1978) 44
 1979AB13 I. Abdul-Jalil and D.F. Jackson, J. Phys. (London) G5 (1979) 1699
 1979AD01 E.G. Adelberger, R.E. Marrs and K.A. Snover, Phys. Rev. C19 (1979) 1
 1979AJ01 F. Ajzenberg-Selove, Nucl. Phys. A320 (1979) 1
 1979AL1J Alster and Warszawski, Phys. Rept. 52 (1979) 87
 1979AL1P Allard, Nature 282 (1979) 56
 1979AL22 Y. Alhassid, R.D. Levine, J.S. Karp and S.G. Steadman, Phys. Rev. C20 (1979) 1789

- 1979AL26 G.D. Alkhazov, S.L. Belostotskii, A.A. Vorobev, O.A. Domchenkov, Y.V. Dotsenko, N.P. Kuropatkin, V.N. Nikulin and M.A. Shuvaev, *Pisma Zh. Eksp. Teor. Fiz.* 29 (1979) 88; *JETP Lett.* 29 (1979) 80
- 1979ALZW J. Alster, A. Doron, S. Gilad, M.A. Moinester, H. Baer, J.D. Bowman, M. Cooper, F. Cverna, R.H. Heffner, C.M. Hoffman et al., *Bull. Amer. Phys. Soc.* 24 (1979) 687, JM5
- 1979AM02 K. Amos and I. Morrison, *Phys. Rev.* C19 (1979) 2108
- 1979AN1J Aniol et al., *AIP Conf. Proc.* 54 (1979) 276
- 1979ANZM K. Aniol, D. Chiang, I. Halpern, D. Drake and G. Haouat, *Bull. Amer. Phys. Soc.* 24 (1979) 821, AD19
- 1979AO02 Y. Aoki, K. Nagano, S. Kunori and K. Yagi, *J. Phys. Soc. Jpn.* 47 (1979) 329
- 1979AR1G L.G. Arnold, *Phys. Rev. Lett.* 42 (1979) 1253
- 1979AU07 G.F. Auchampaugh, S. Plattard and N.W. Hill, *Nucl. Sci. Eng.* 69 (1979) 30
- 1979BA2D Baer et al., *Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C19*
- 1979BA2E Baer, *Bull. Amer. Phys. Soc.* 24 (1979) 823
- 1979BE1H Benenson and Kashy, *Rev. Mod. Phys.* 51 (1979) 527
- 1979BE1N Beer et al., *Conf. Proc. TRIUMF, Vancouver (1979) Paper 1B27*
- 1979BE1V Bell, Dickens and Gustafsson, *Astrophys. J.* 229 (1979) 604
- 1979BE1X Berman, *Suppl. Preprint UCRL-78482 (1979)*
- 1979BE44 D. Besset, Q.H. Do, B. Favier, L.G. Greeniaus, R. Hess, C. Lechanoine, D. Rapin, D.W. Werren and C. Weddigen, *Nucl. Instrum. Meth. Phys. Res.* 166 (1979) 379
- 1979BE50 A. Begum, R.B. Galloway and F.K. McNeil-Watson, *Nucl. Phys.* A332 (1979) 349
- 1979BJ1A Bjarle, Herrstrom, Jonnsson and Kristiansson, *Z. Phys.* A291 (1979) 383
- 1979BL1F Blatt et al., *Bull. Amer. Phys. Soc.* 24 (1979) 843
- 1979BO06 K. Bodek, M. Hugi, J. Lang, R. Muller, E. Ungricht, L. Jarczyk, B. Kamys and A. Strzalkowski, *Phys. Lett.* B82 (1979) 369
- 1979BO1K Bodek et al., *Helv. Phys. Acta* 52 (1979) 88
- 1979BO1U Booth, 1978 RPI Symp. Photopion Nucl. Phys. (1979) 129
- 1979BO22 V.I. Bogatin, O.V. Lozhkin and Y.P. Yakovlev, *Nucl. Phys.* A326 (1979) 508
- 1979BR03 H.S. Bradlow, W.D.M. Rae, P.S. Fisher, N.S. Godwin, G. Proudfoot and D. Sinclair, *Nucl. Phys.* A314 (1979) 171
- 1979CE1A L.S. Celenza and C.M. Shakin, *Phys. Rev.* C20 (1979) 385
- 1979CH07 Y.-d. Chan, H. Bohn, R. Vandenbosch, R. Sielemann, J.G. Cramer, K.G. Bernhardt, H.C. Bhang and D.T.C. Chiang, *Phys. Rev. Lett.* 42 (1979) 687; *Erratum Phys. Rev. Lett.* 43 (1979) 1625

- 1979CO1C Cooper, AIP Conf. Proc. 54 (1979) 222
- 1979CO1H Corfu et al., AIP Conf. Proc. 54 (1979) 528
- 1979CR1C Crannell et al., Bull. Amer. Phys. Soc. 24 (1979) 845
- 1979CRZZ H. Crannell, W.J. Briscoe, P.E. Burt, L.W. Fagg, J.T. O'Brien, D.I. Sober, S. Kowalski and C.F. Williamson Bull. Amer. Phys. Soc. 24 (1979) 683, JK7
- 1979DA1D Davis, Evans and Cleveland, AIP Conf. Proc. 52 (1979) 17
- 1979DE01 P. Desgrolard and P.A.M. Guichon, Phys. Rev. C19 (1979) 120
- 1979DE15 P. Desgrolard and P.A.M. Guichon, Z. Phys. A290 (1979) 373
- 1979DE1H Del Fiore et al., Int. J. Appl. Rad. Isotopes 30 (1979) 543
- 1979DE1N de Vries, Nukleonika 24 (1979) 75, 87
- 1979DE1P de Vries, BNL-51115 (1979) 513
- 1979DE31 R.M. DeVries and J.C. Peng, Phys. Rev. Lett. 43 (1979) 1373
- 1979DE34 D. Dehnhard, S.J. Tripp, M.A. Franey, G.S. Kyle, C.L. Morris, R.L. Boudrie, J. Piffaretti and H.A. Thiessen, Phys. Rev. Lett. 43 (1979) 1091
- 1979DO03 E.I. Dolinskii, S.P. Krekoten, A.M. Mukhamedzhanov and E.A. Romanovskii, Yad. Fiz. 29 (1979) 71; Sov. J. Nucl. Phys. 29 (1979) 34
- 1979DO13 L.R. Dodd, Aust. J. Phys. 32 (1979) 173
- 1979DO17 G.E. Dogotar, R.A. Eramzhyan, M. Gmitro, H.R. Kissener and E. Tinkova, J. Phys. (London) G5 (1979) L221
- 1979DU10 W. Dunnweber, P.D. Bond, C. Chasman and S. Kubono, Phys. Rev. Lett. 43 (1979) 1642; Erratum Phys. Rev. Lett. 44 (1980) 299
- 1979EN1D Engelmann and Loeuillet, IEEE Trans. on Nucl. Sci. 26 (1979) 1539
- 1979ER02 J. Ernst, Phys. Rev. C19 (1979) 896
- 1979FI1B Fisher et al., Conf. Proc. TRIUMF, Vancouver (1979) Paper 3E16
- 1979FR12 S. Frankel, W. Frati, M. Gazzaly, Y.D. Bayukov, V.I. Efremenko, G.A. Leksin, N.A. Nikiforov, V.I. Tchistilin, Y.M. Zaitsev and C.F. Perdrisat, Phys. Rev. C20 (1979) 2257
- 1979FU03 H. Fuchs, T.S. Bhatia, P. Doll, H. Hafner and C.J. Wagner, J. Phys. (London) G5 (1979) 797
- 1979FU04 C.B. Fulmer, R.M. Wieland, D.C. Hensley, S. Raman, G.R. Satchler, A.H. Snell, P.H. Stelson and R.G. Stokstad, Phys. Rev. C20 (1979) 670
- 1979FU1H Fujisawa et al., Tsukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 81
- 1979GA13 Y.R. Gaillard, P. Martin, G. Perrin, P. Desaintignon, M. Buenerd, J.M. Loiseaux, J.L. Escudie, A. Tarrats and H.V. Geramb, Nucl. Phys. A327 (1979) 349

- 1979GAZV E.N. Gazis, A.C. Xenoulis, P. Kakanis, X. Aslanoglou and A.D. Panagiotou, Bull. Amer. Phys. Soc. 24 (1979) 695, KH10
- 1979GE1D Genova and Schatzman, Astron. Astrophys. 78 (1979) 323
- 1979GI1D Gibbs, Gibson and Stephenson, Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C13
- 1979GR1J Greenwood and Chrien, Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (1979) 618
- 1979GR1K Grenne et al., Conf. Proc. TRIUMF, Vancouver (1979) Paper 1E23
- 1979GR20 P.J. Griffin and R.D. Koshel, Phys. Rev. C20 (1979) 2119
- 1979GU1D Guzik, Bull. Amer. Phys. Soc. 24 (1979) 692
- 1979GY1A Gyles et al., AIP Conf. Proc. 54 (1979) 522
- 1979GY1B Gyles et al., Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C16
- 1979HAYZ L.H. Harwood and G.M. Crawley, Bull. Amer. Phys. Soc. 24 (1979) 817, AC2
- 1979HE16 A.C. Heemskerck, L.P. Kok and H.T. Koster, Z. Phys. A293 (1979) 165
- 1979HE1D Hering et al., BNL-51115 (1979) 759
- 1979HE1F K.T. Hecht and W. Zahn, Nucl. Phys. A313 (1979) 77
- 1979HEZX B. Herman, T.M. Cormier and B.R. Fulton, Bull. Amer. Phys. Soc. 24 (1979) 664, HK1
- 1979HI1A Hirata, Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C10
- 1979HO17 J. Hohn, H.W. Barz and I. Rotter, Nucl. Phys. A330 (1979) 109
- 1979HO1F Hoistad et al., Conf. Proc. TRIUMF, Vancouver (1979) Paper 4B5
- 1979HO1H Hosono et al., Ysukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 117
- 1979IM02 B. Imanishi and W. von Oertzen, Phys. Lett. B87 (1979) 188
- 1979IN01 A. Ingemarsson, O. Jonsson and A. Hallgren, Nucl. Phys. A319 (1979) 377
- 1979IN1A Ingram, AIP Conf. Proc. 54 (1979) 455
- 1979JA04 L. Jarczyk, J. Okolowicz, A. Strzalkowski, K. Bodek, M. Hugi, J. Lang, R. Muller and E. Ungricht, Nucl. Phys. A316 (1979) 139
- 1979JA17 G.J. Jan, C.C. Perng and M.K. Liou, Phys. Lett. B85 (1979) 25
- 1979JA1F D.F. Jackson and D.J. Brenner, Prog. Part. Nucl. Phys. 5 (1981) 143
- 1979JA1G Jarczy et al., J. Phys. (London) G5 (1979) 565
- 1979JO08 R.R. Johnson, T. Masterson, B. Bassalleck, W. Gyles, T. Marks, K.L. Erdman, A.W. Thomas, D.R. Gill, E. Rost, J.J. Kraushaar et al., Phys. Rev. Lett. 43 (1979) 844
- 1979JO1C Jones, AIP Conf. Proc. 54 (1979) 116

- 1979JOZP D.L. Johnson, Bull. Amer. Phys. Soc. 24 (1979) 862, AB5
- 1979JU01 J.W. Jury, B.L. Berman, D.D. Faul, P. Meyer, K.G. McNeill and J.G. Woodworth, Phys. Rev. C19 (1979) 1684
- 1979KA07 S.B. Kaufman, M.W. Weisfield, E.P. Steinberg, B.D. Wilkins and D.J. Henderson, Phys. Rev. C19 (1979) 962
- 1979KA1N Kato, Tsukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 56
- 1979KI1J Kissenger and Eramzhyan, Conf. Proc. TRIUMF, Vancouver (1979) Paper 117
- 1979KO05 M.A. Kovash, S.L. Blatt, R.N. Boyd, T.R. Donoghue, H.J. Hausman and A.D. Bacher, Phys. Rev. Lett. 42 (1979) 700
- 1979KO20 D.G. Kovar, D.F. Geesaman, T.H. Braid, Y. Eisen, W. Henning, T.R. Ophel, M. Paul, K.E. Rehm, S.J. Sanders, P. Sperr et al., Phys. Rev. C20 (1979) 1305
- 1979KO21 V.I. Komarov, G.E. Kosarev, H. Muller, D. Netzband, V.D. Toneev, T. Stiehler, S. Tesch, K.K. Gudima and S.G. Mashnik, Nucl. Phys. A326 (1979) 297
- 1979KO26 L. Koester, K. Knopf and W. Waschkowski, Z. Phys. A292 (1979) 95
- 1979KO36 V.I. Komarov, G.E. Kosarev, H. Muller, D. Netzband, T. Stiehler and S. Tesch, J. Phys. (London) G5 (1979) 1717
- 1979KOZU P.E. Koehler, D.A. Resler, H.D. Knox, R.M. White and R.O. Lane, Bull. Amer. Phys. Soc. 24 (1979) 656, HF
- 1979KR18 H.H. Krause, W. Arnold, H. Berg, J. Ulbricht and G. Clausnitzer, Z. Phys. A293 (1979) 343
- 1979KU20 G.J. Kumbartzki, K. Hagemeyer, W. Knauer, G. Krosing, R. Kuhnen, V. Mertens, K.-H. Speidel, J. Gerber and W. Nagel, Hyperfine Interactions 7 (1979) 253
- 1979LA1H Lazareff, Audouze, Starrfield and Truran, Astrophys. J. 228 (1979) 875
- 1979LA22 R.H. Landau and A.W. Thomas, Phys. Lett. B88 (1979) 226
- 1979LAZP G.P. Lamaze, R.B. Schwartz and J.D. Kellie, Bull. Amer. Phys. Soc. 24 (1979) 862, AB4
- 1979LAZQ D.C. Larson, J.A. Harvey and N.W. Hill, Bull. Amer. Phys. Soc. 24 (1979) 862, AB1
- 1979LE1G Lesser, Trail and Liou, Bull. Amer. Phys. Soc. 24 (1979) 32
- 1979LE1H Lesser, Liou, Liu and Trail, Bull. Amer. Phys. Soc. 24 (1979) 32
- 1979LEZR J. Leroze, K. Min, D. Rowley, P. Stoler, E.J. Winhold, P.F. Yergin, A.M. Bernstein, K.I. Blomqvist, H. Caplan, G. Franklin et al., Bull. Amer. Phys. Soc. 24 (1979) 819, AD2
- 1979LI03 R.P. Liljestrang, G.S. Blanpied, W.R. Coker, C. Harvey, G.W. Hoffmann, L. Ray, C. Glashausser, G.S. Adams, T.S. Bauer, G. Igo et al., Phys. Rev. Lett. 42 (1979) 363
- 1979LI1G Lindner, Helmer and Brinkman, Int. J. Appl. Rad. Isotopes 30 (1979) 506

- 1979LI1H Liu, Jinag and Li, Phys. Energ. Fortis Phys. Nucl. (China) 3 (1979) 247
- 1979LO12 W.G. Love, Phys. Rev. C20 (1979) 1638
- 1979MA20 T.A.J. Maris, M.R. Teodoro and C.A.Z. Vasconcellos, Nucl. Phys. A322 (1979) 461
- 1979MA38 R.E. Marrs, R.E. Pollock and W.W. Jacobs, Phys. Rev. C20 (1979) 2308
- 1979MA39 R.E. Marrs and R.E. Pollock, Phys. Rev. C20 (1979) 2446
- 1979MAZW C.J. Martoff, J.A. Bistirlich, K.M. Crowe, J.P. Miller, S.S. Rosenblum, W.A. Zajc, H.W. Baer, G. Strassner, P. Truol and A.H. Wapstra, Bull. Amer. Phys. Soc. 24 (1979) 647, GK9
- 1979MC1B Mccutcheon et al., J. Roy. Astron. Soc. Can. 73 (1979) 296
- 1979ME1D Merrill and Ridgeway, Ann. Rev. Astron. Astrophys. 17 (1979) 9
- 1979MO14 T. Motobayashi, I. Kohno, T. Ooi and S. Nakajima, Nucl. Phys. A331 (1979) 193
- 1979MO1N Moinester et al., Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C20
- 1979MO1P Moniz and Toyama, Conf. Proc. TRIUMF, Vancouver (1979) Paper 1D9
- 1979MO1Q Moniz, Proc. Mainz, 1979, Springer Lect. Notes 108 (1979) 435
- 1979NE1B Neudachin, Romanovskii, Gubin and Makhmud, Sov. J. Nucl. Phys. 29 (1979) 170
- 1979OL1B Olsen and Richer, Astrophys. J. 227 (1979) 534
- 1979OR01 M.E. Ortiz, A. Dacal, A. Menchaca-Rocha, M. Buenerd, D.L. Hendrie, J. Mahoney, C. Olmer and D.K. Scott, Phys. Lett. B84 (1979) 63
- 1979OS01 R. Ost, M.R. Clover, R.M. DeVries, B.R. Fulton, H.E. Gove and N.J. Rust, Phys. Rev. C19 (1979) 740
- 1979OS06 P. Osland and R.J. Glauber, Nucl. Phys. A326 (1979) 255
- 1979OS10 A. Osman and G.U. Din, Ann. Phys. (Leipzig) 36 (1979) 56
- 1979OS11 A. Osman and G.U. Din, Ann. Phys. (Leipzig) 36 (1979) 64
- 1979PAZY J.Y. Park, W. Scheid and W. Greiner, Bull. Amer. Phys. Soc. 24 (1979) 665, HK3
- 1979PE1E Penzias, Rev. Mod. Phys. 51 (1979) 425
- 1979PH05 D.-L. Pham and R. de Swiniarski, Z. Phys. A291 (1979) 327
- 1979PI1C Pickar et al., Bull. Amer. Phys. Soc. 24 (1979) 819
- 1979PIZZ P.H. Pile, T.P. Sjoreen, R.E. Pollock, W.W. Jacobs, H.-O. Meyer, M.C. Green, F. Soga and R.D. Bent, Bull. Amer. Phys. Soc. 24 (1979) 614, DN3
- 1979PO16 E.C. Pollacco, D.M. Parkinson, J.R. Leigh, D.C. Weissner, N. Shikazono, G.S. Foote and D. Branford, Nucl. Phys. A330 (1979) 173
- 1979PR07 G. Proudfoot and W.D.M. Rae, Nucl. Phys. A329 (1979) 271
- 1979RA1C Ramaty, Kozlovsky and Lingenfelter, Astrophys. J. Suppl. 40 (1979) 487

1979RA20 J.R. Radin, E. Gradsztajn and A.R. Smith, Phys. Rev. C20 (1979) 787
 1979RA27 L. Ray, Phys. Rev. C20 (1979) 1857
 1979RE1A Redwine, AIP Conf. Proc. 54 (1979) 501
 1979RO12 N.A. Roughton, M.R. Fritts, R.J. Peterson, C.S. Zaidins and C.J. Hansen, At. Data Nucl. data Tables 23 (1979) 177
 1979RO1E Rotter, J. Phys. (London) G5 (1979) 75
 1979ROZV J.L. Romero, T.S. Subramanian, F.P. Brady, G.H. Herling and R.H. Bassel, Bull. Amer. Phys. Soc. 24 (1979) 656, HF1
 1979SA1M Sackman, CAP-551 (1979)
 1979SA21 A.N. Saharia and R.M. Woloshyn, Phys. Lett. B84 (1979) 401
 1979SA27 G.R. Satchler, Nucl. Phys. A329 (1979) 233
 1979SC07 P. Schwaller, M. Pepin, B. Favier, C. Richard-Serre, D.F. Measday and P.U. Renberg, Nucl. Phys. A316 (1979) 317
 1979SC25 E. Schwarz, J.-P. Egger, F. Goetz, P. Gretillat, C. Lunke, C. Perrin, B.M. Freedom and R.E. Mischke, Phys. Rev. Lett. 43 (1979) 1578
 1979SE06 B.D. Serot, Nucl. Phys. A322 (1979) 408
 1979SE07 R.R. Sercely, R.J. Peterson, P.A. Smith and E.R. Flynn, Nucl. Phys. A324 (1979) 53
 1979SI07 M. Sieminski, M. Sosnowski and P. Zupranski, Can. J. Phys. 57 (1979) 292
 1979SK04 D.M. Skopik, J.J. Murphy II, H.R. Weller, R.A. Blue, N.R. Roberson, S.A. Wender and D.R. Tilley, Phys. Rev. C20 (1979) 409
 1979SM08 A. Smith, R. Holt and J. Whalen, Nucl. Sci. Eng. 70 (1979) 281
 1979SM1B Smirnov et al., Nature 278 (1979) 630
 1979SN1A Snover, Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (1979) 319
 1979SOZY F. Soga, R.D. Bent, P.H. Pile, R.E. Pollak, M.C. Green and T.P. Sjoreen, Bull. Amer. Phys. Soc. 24 (1979) 614, DN7
 1979ST1D Stanley, Glover and Petrovich, Bull. Amer. Phys. Soc. 24 (1979) 816
 1979ST1J Stephenson, Gibbs, Gibson and Kaufmann, Bull. Amer. Phys. Soc. 24 (1979) 821
 1979SU1F Suzuki and Kubo, Tsukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 211
 1979SUZW T.S. Subramanian, J.L. Romero, F.P. Brady, D.H. Fitzgerald, R. Garrett, G. Needham, J.L. Ullmann, J.W. Watson and C. Zanelli, Bull. Amer. Phys. Soc. 24 (1979) 593, BM1
 1979SW1B Sweigart and Mengel, Astrophys. J. 229 (1979) 624
 1979TA1E Tanaka, Tsukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 220
 1979TH1A Thompson and Feagin, Bull. Amer. Phys. Soc. 24 (1979) 627

- 1979TR1B Truol, Proc. Mainz, 1979, Springer Lect. Notes 108 (1979) 351
- 1979TR1D Trail, Lesser, Bond and Liou, IEEE Trans. on Nucl. Sci. 26 (1979) 1174
- 1979TRZR S.J. Tripp, D. Dehnhard, M.A. Franey, G.S. Kyle, C.L. Morris, R.L. Boudrie, J. Piffaretti and H.A. Thiessen, Bull. Amer. Phys. Soc. 24 (1979) 819, AD1
- 1979TSZZ M. Tsangarides, J.G. Willis and P.H. Pile Bull. Amer. Phys. Soc. 24 (1979) 820, AD9
- 1979TU1A Tucker, Kutner and Massano, Astrophys. J. 227 (1979) L143
- 1979UN01 E. Ungricht, D. Balzer, M. Hugi, J. Lang, R. Muller, L. Jarczyk, B. Kamys and A. Strzalkowski, Nucl. Phys. A313 (1979) 376
- 1979URZZ P.P. Urone, C.I. Zanelli, F.P. Brady, C.L. Castaneda, M. Johnson, G.A. Needham, J.L. Romero, J. Ullmann and D.L. Johnson, Bull. Amer. Phys. Soc. 24 (1979) 657, HF5
- 1979VD02 A.I. Vdovin, I.G. Golikov, M.N. Zhukov, I.I. Loshchakov and V.I. Ostroumov, Izv. Akad. Nauk SSSR Ser. Fiz. 43 (1979) 148; Bull. Acad. Sci. USSR Phys. Ser. 43 (1979) 124
- 1979WE1C Weber and Eisenberg, AIP Conf. Proc. 54 (1979) 190
- 1979WE1E Weil, McDaniel and Burrows, Bull. Amer. Phys. Soc. 24 (1979) 889
- 1979WEZP J.L. Weil, F.D. McDaniel and T.W. Burrows, Bull. Amer. Phys. Soc. 24 (1979) 853, GC3
- 1979WI1G Wieland, Highfall and King, IEEE Trans. on Nucl. Sci. 26 (1979) 1713
- 1979WI1H Wiedenbeck et al., Bull. Amer. Phys. Soc. 24 (1979) 692
- 1979WO06 J.G. Woodworth, K.G. McNeill, J.W. Jury, P.D. Georgopoulos and R.G. Johnson, Nucl. Phys. A327 (1979) 53
- 1979YA1E Yabe, Mori and Kubo, Tsukuba Symp. on Polariz. Phenom., Nov. 1979; Ed., K. Yagi (1979) 70
- 1979ZE01 A.F. Zeller, K.W. Kemper, D.C. Weisser, T.R. Ophel, D.F. Hebbard and A. Johnston, Nucl. Phys. A323 (1979) 477
- 1979ZE1B Zelenskaya and Teplov, Izv. Akad. Nauk SSSR Ser. Fiz. 43 (1979) 2375
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, Nucl. Phys. A336 (1980) 1
- 1980AL09 G.D. Alkhazov, S.L. Belostotsky, O.A. Domchenkov, Y.V. Dotsenko, N.P. Kuropatkin, V.N. Nikulin, A.A. Vorobyov and M.A. Shuvaev, Phys. Lett. B90 (1980) 364
- 1980AL12 Y. Alexander, J.W. Van Orden, E.F. Redish and S.J. Wallace, Phys. Rev. Lett. 44 (1980) 1579
- 1980BA11 V.V. Balashov, H.W. Barz and H.U. Jager, J. Phys. (London) G6 (1980) L77
- 1980BA54 F.C. Baker and N. Ferdous, Aust. J. Phys. 33 (1980) 691
- 1980BL1B Blatt et al., Bull. Amer. Phys. Soc. 25 (1980) 603

- 1980BO03 J.D. Bowman, Nucl. Phys. A335 (1980) 375
- 1980BR04 I. Brissaud, Z. Phys. A294 (1980) 287
- 1980CH01 M.L. Chatterjee, L. Potvin and B. Cujec, Nucl. Phys. A333 (1980) 273
- 1980CH08 S. Chakravarti, Phys. Lett. B90 (1980) 350
- 1980CI03 S. Cierjacks, F. Hinterberger, G. Schmalz, D. Erbe, P.V. Rossen and B. Leugers, Nucl. Instrum. Meth. 169 (1980) 185
- 1980CO05 J.R. Comfort, S.M. Austin, P.T. Debevec, G.L. Moake, R.W. Finlay and W.G. Love, Phys. Rev. C21 (1980) 2147
- 1980CO06 J.R. Comfort and B.C. Karp, Phys. Rev. C21 (1980) 2162; Erratum Phys. Rev. C22 (1980) 1809
- 1980DE1L Devins et al., Bull. Amer. Phys. Soc. 25 (1980) 577
- 1980DE1M Dehnhard et al., Bull. Amer. Phys. Soc. 25 (1980) 593
- 1980DO1C Domogatsky and Nadyozhin, Astrophys. Space Sci. 70 (1980) 33, 77
- 1980ER01 R.A. Eramzhyan, M. Gmitro and H.R. Kissener, Nucl. Phys. A338 (1980) 436
- 1980FA07 E. Fabrici, S. Micheletti, M. Pignanelli, F.G. Resmini, R. De Leo, G. D'Erasmus and A. Pantaleo, Phys. Rev. C21 (1980) 844
- 1980FU04 H. Fuchs, J.A. Nolen, G.J. Wagner, H. Lenske and G. Baur, Nucl. Phys. A343 (1980) 133
- 1980GA12 H. Garcilazo, Phys. Rev. C21 (1980) 2094
- 1980GO07 C.D. Goodman, C.A. Goulding, M.B. Greenfield, J. Rapaport, D.E. Bainum, C.C. Foster, W.G. Love and F. Petrovich, Phys. Rev. Lett. 44 (1980) 1755
- 1980HE04 R.L. Helmer, M.D. Hasinoff, J.E. Bussioletti, K.A. Snover and T.A. Trainor, Nucl. Phys. A336 (1980) 219
- 1980HE1D Henkel, Walmsley and Wilson, Astron. Astrophys. 82 (1980) 41
- 1980HO07 R.H. Howell, F.S. Dietrich and F. Petrovich, Phys. Rev. C21 (1980) 1158
- 1980HO11 R.J. Holt, R.M. Laszewski, H.E. Jackson, J.E. Monahan and J.R. Specht, Phys. Rev. C21 (1980) 1699; Erratum Phys. Rev. C23 (1981) 1841
- 1980JA05 D.F. Jackson and I. Abdul-Jalil, J. Phys. (London) G6 (1980) 481
- 1980JO06 M.B. Johnson, Phys. Rev. C22 (1980) 192
- 1980KA02 S. Kato, K. Okada, M. Kondo, A. Shimizu, K. Hosono, T. Saito, N. Matsuoka, S. Nagamachi, K. Nisimura, N. Tamura et al., Nucl. Instrum. Meth. Phys. Res. 169 (1980) 589
- 1980LA1C R.H. Landau, Nucl. Phys. A335 (1980) 289
- 1980LA1D Larson, Hetrick and Harvey, Bull. Amer. Phys. Soc. 25 (1980) 543

- 1980LA1G Lambert, Dominy and Sivertsen, *Astrophys. J.* 235 (1980) 114
- 1980LE02 T.-S.H. Lee and D. Kurath, *Phys. Rev. C*21 (1980) 293
- 1980LI01 M.K. Liou, C.K. Liu, P.M.S. Lesser and C.C. Trail, *Phys. Rev. C*21 (1980) 518
- 1980LI1E Liu, *Bull. Amer. Phys. Soc.* 25 (1980) 606
- 1980MA1F Maleki, Rao and Levinger, *Bull. Amer. Phys. Soc.* 25 (1980) 606
- 1980MC1G McCutcheon, Dickman, Shuter and Roger, *Astrophys. J.* 237 (1980) 9
- 1980MI01 T. Mikumo, M. Sasagase, M. Sato, T. Ooi, Y. Higashi, Y. Nagashima and M. Yamanouchi, *Phys. Rev. C*21 (1980) 620
- 1980MO06 J.M. Moss, C. Glashauser, F.T. Baker, R. Boudrie, W.D. Cornelius, N. Hintz, G. Hoffmann, G. Kyle, W.G. Love, A. Scott et al., *Phys. Rev. Lett.* 44 (1980) 1189
- 1980MU1B N.C. Mukhopadhyay, *Nucl. Phys.* A335 (1980) 111
- 1980NA14 F. Naulin, C. Detraz, M. Bernas, D. Guillemaud, E. Kashy, M. Langevin, F. Pougheon, P. Roussel and M. Roy-Stephan, *J. Phys. Lett.* 41 (1980) L79
- 1980NA1B Nagl and Uberall, *Bull. Amer. Phys. Soc.* 25 (1980) 606
- 1980PE13 R.J. Peterson and J.J. Hamill, *Phys. Rev. C*22 (1980) 2282
- 1980PE1C R.J. Peterson, *Nucl. Phys.* A335 (1980) 365
- 1980PE1E Peterson et al., *Bull. Amer. Phys. Soc.* 25 (1980) 593
- 1980PE1F Penzias, *Science* 208 (1980) 663
- 1980RA1B L. Ray, *Nucl. Phys.* A335 (1980) 443
- 1980RI1A Rikus et al., UM-P-80/52 (1980)
- 1980SA04 A.N. Saharia and R.M. Woloshyn, *Phys. Rev. C*21 (1980) 1111
- 1980SC1D Scheid and Greiner, *Proc. Conf. in Nucl. Far from Stability* (1980) 295
- 1980SH01 O.D. Sharma and B.B. Srivastava, *J. Phys. (London)* G6 (1980) 209
- 1980SN01 K.A. Snover, P.G. Ikossi, E.G. Adelberger and K.T. Lesko, *Phys. Rev. Lett.* 44 (1980) 927
- 1980SO1B Sober et al., *Private Communications* (1980)
- 1980TH01 H.A. Thiessen, *Nucl. Phys.* A335 (1980) 329
- 1980TH05 W.J. Thompson, J.F. Wilkerson, T.B. Clegg, J.M. Feagin, E.J. Ludwig and E. Merzbacher, *Phys. Rev. Lett.* 45 (1980) 703
- 1980TR02 H.-J. Trost, A. Schwarz, U. Feindt, F.H. Heimlich, S. Heinzl, J. Hintze, F. Korber, R. Lekebusch, P. Lezoch, G. Mock et al., *Nucl. Phys.* A337 (1980) 377
- 1980TR1B Tripp et al., *Bull. Amer. Phys. Soc.* 25 (1980) 504
- 1980VO05 W. von Oertzen, B. Imanishi, H.G. Bohlen, W. Treu and H. Voit, *Phys. Lett.* B93 (1980) 21

- 1980WA24 E.K. Warburton, D.E. Alburger and D.J. Millener, Phys. Rev. C22 (1980) 2330
- 1980WH1A C.A., Jr. Whitten, Nucl. Phys. A335 (1980) 419
- 1980WI09 J.P. Wieleczko, S. Harar, M. Conjeaud and F. Saint-Laurent, Phys. Lett. B93 (1980)
35
- 1980WIZW Wilkerson, Bull. Amer. Phys. Soc. 25 (1980) 555, GE9
- 1980WO1A Woodworth, Dietrich and Jury, Bull. Amer. Phys. Soc. 25 (1980) 603
- 1982AJ01 F. Ajzenberg-Selove, Nucl. Phys. A375 (1982) 1

