

Energy Levels of Light Nuclei $A = 17$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 16-17$ was published in *Nuclear Physics A166* (1971), p. 1. This version of $A = 17$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed November 30, 1970)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 17$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{17}\text{B}\$](#) , [\$^{17}\text{C}\$](#) , [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#) , [\$^{17}\text{Ne}\$](#) , [\$^{17}\text{Na}\$](#)

B. Tables of Recommended Level Energies:

[Table 17.1](#): Energy levels of ^{17}N

[Table 17.5](#): Energy levels of ^{17}O

[Table 17.17](#): Energy levels of ^{17}F

[Table 17.22](#): Energy levels of ^{17}Ne

C. [References](#)

D. Figures: [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹⁷B
(Not illustrated)

¹⁷B has not been observed. (1966GA25) predict that it is unbound with respect to decay into ¹⁵B + 2n by 4.0 MeV. See also (1960ZE03).

¹⁷C
(Not illustrated)

¹⁷C has been observed in the 5.5 GeV proton bombardment of uranium: it is particle stable (1968PO04). (1966GA25) predict that it is bound, with respect to ¹⁶C + n, by 0.6 ± 0.4 MeV: $M - A$ is then 22.4 ± 0.4 MeV. See also (1960ZE03, 1969AR13).

¹⁷N
(Figs. 6 and 9)

GENERAL:

Theory and reviews: (1961BA1C, 1962GO31, 1966MA12, 1970HI15).

Experimental papers: (1962VO01, 1965CI01, 1966PO08, 1967AU1B, 1967CA1J, 1968AR1F, 1968DO20, 1969AR13).

1. (a) $^{17}\text{N}(\beta^-)^{17}\text{O}^* \rightarrow ^{16}\text{O} + \text{n}$ $Q_m = 4.536$
(b) $^{17}\text{N}(\beta^-)^{17}\text{O}$ $Q_m = 8.679$

The half-life of ¹⁷N is 4.14 ± 0.04 sec (1948KN24), 4.20 ± 0.08 sec (1961HI01), 4.16 ± 0.01 sec (1965DO13), 4.17 ± 0.02 sec (1970ME31): the mean of these values is 4.16 ± 0.01 sec. See also (1959AJ76).

The decay is primarily [$\approx 95\%$ (1964SI06)] to neutron unstable states of ¹⁷O, principally ¹⁷O*(4.55, 5.38, 5.94) [$J^\pi = \frac{3}{2}^-$, $\frac{3}{2}^-$ and $\frac{1}{2}^-$, respectively] (1961PE28, 1963GI04). The ratio of the intensities of these delayed neutrons from ¹⁷O*(5.38) and ¹⁷O*(4.55) is 1.6 (1961PE28). There are also weak branches to ¹⁷O*(0, 0.87, 3.06): see Table 17.2 (1964SI06). The character of the decay indicates $J^\pi = \frac{1}{2}^-$ for ¹⁷N(0). See also (1965MA16, 1970BE21, 1970HI15).

2. $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ $Q_m = 8.415$
 $Q_0 = 8.38 \pm 0.06$ (1959LI47).

Table 17.1: Energy levels of ^{17}N ^a

E_x in ^{17}N (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
0	$\frac{1}{2}^-; \frac{3}{2}$	$\tau_{1/2} = 4.16 \pm 0.01$ sec	β^-	1, 2, 3, 4, 5, 6, 9
1.3707 ± 0.8	$\leq \frac{3}{2}^+$	$\tau_m < 0.2$ psec	γ	2, 9
1.861 ± 10	$\frac{1}{2}^+$		γ	9
1.9080 ± 0.8	$\geq \frac{5}{2}^-$	$\tau_m > 0.5$ psec	γ	2, 9
2.5279 ± 3.6			γ	2, 9
3.1342 ± 4.6		$\tau_m < 0.3$ psec	γ	2, 9
3.220 ± 6		$\tau_m > 0.3$ psec	γ	2, 9
3.661 ± 6			γ	2, 9
3.684 ± 12				9
3.928 ± 12				9
4.023 ± 12			γ	2, 9
4.215 ± 25			γ	2, 9
4.470 ± 10			γ	2
5.176 ± 15			γ	2, 9
5.195 ± 15			γ	2, 9
5.523 ± 11	$(\frac{3}{2}^-)$		γ	2, 9
5.787 ± 15			γ	9
5.832 ± 11			γ	2, 9
6.08 ± 31				2
6.24 ± 24				2
6.43 ± 30				2
6.61 ± 24				2
6.99 ± 24				2
7.17 ± 40				2
7.37 ± 40				2
7.63 ± 40				2
7.78 ± 18				2
8.00 ± 24				2
8.14 ± 40				2
8.55 ± 40		broad		2
8.93 ± 40		broad		2
9.26 ± 40		broad		2
9.74 ± 40		broad		2

^a See also Table 17.4.

Table 17.2: Beta decay of ^{17}N ^a

Decay to $^{17}\text{O}^*$ (MeV)	J^π	Branch (%)	$\log ft$
0	$\frac{5}{2}^+$	1.6 ± 0.5	9.56 ± 0.13 ^c
0.87	$\frac{1}{2}^+$	2.6 ± 0.5	6.8 ± 0.1
3.06	$\frac{1}{2}^-$	0.46 ± 0.11	6.9 ± 0.1
3.84	$\frac{5}{2}^-$	< 0.1	> 7.4
> 3.84 ^b		≈ 95	

^a (1964SI06) and J.C. Hopkins, private communication.

^b See discussion in reaction 1 and (1961PE28).

^c $\log f_{1t}$: E.K. Warburton, private communication and (1971TO08).

Proton groups have been observed to many states in ^{17}N : see Table 17.3 (1959LI47, 1965HA05, 1966MC05). See also (1963MO1B). Angular distributions have been measured at $E(^7\text{Li}) = 5$ MeV to most of the states with $E_x < 6.5$ MeV (1966MC05).

Observed γ -transitions are displayed in Tables 17.3 and 17.4 (1965HA05, 1969TH01). The mean lifetimes of $^{17}\text{N}^*(1.37, 1.91, 3.13, 3.21)$ are, respectively, < 0.2 , > 0.5 , < 0.3 and > 0.3 psec, suggesting $J \leq \frac{3}{2}$ for $^{17}\text{N}^*(1.37)$ and $J \geq \frac{5}{2}$ for $^{17}\text{N}^*(1.91)$ (1969TH01). See also (1965CA05, 1967CA1D).

$$3. \ ^{14}\text{C}(\alpha, p)^{17}\text{N} \quad Q_m = -9.715$$

See (1961PE28).

$$4. \ ^{15}\text{N}(t, p)^{17}\text{N} \quad Q_m = -0.110$$

See (1963GI04, 1964SI06).

$$5. \ ^{17}\text{O}(n, p)^{17}\text{N} \quad Q_m = -7.896$$

See ^{18}O in (1972AJ02).

$$6. \ ^{18}\text{O}(\gamma, p)^{17}\text{N} \quad Q_m = -15.942$$

See ^{18}O in (1972AJ02).

Table 17.3: Levels of ^{17}N from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ and $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$

E_x (MeV \pm keV)					
(1959LI47) ^a	(1965HA05) ^a	(1966MC05) ^a	(1969TH01) ^b	(1960JA13) ^c	A
0				0	0 ^g
1.32 \pm 80			1.3707 \pm 0.8	1.374 \pm 18	1.381 \pm 12
				1.851 \pm 18	1.865 \pm 12 ^h
1.89 \pm 80			1.9080 \pm 0.8	1.906 \pm 18	1.921 \pm 12
2.50 \pm 80 ^d			2.5279 \pm 3.6	2.536 \pm 18	2.530 \pm 15
2.82 \pm 80					
			3.1342 \pm 4.6	3.132 \pm 18	3.138 \pm 12
3.27 \pm 90			3.2220 \pm 7.1	3.212 \pm 18	3.216 \pm 12
3.57 \pm 90			3.6656 \pm 6.5	3.652 \pm 25	3.650 \pm 12
					3.684 \pm 12
3.86 \pm 90					3.928 \pm 12
				4.010 \pm 25	4.023 \pm 12
4.18 \pm 90				(4.215 \pm 25) ^f	i
	4.47 \pm 10	4.47 \pm 40			i
					5.176 \pm 15
	5.21 \pm 20	5.23 \pm 40			5.195 \pm 15
	5.53 \pm 20	5.51 \pm 40			5.521 \pm 15 ^g
					5.787 \pm 15
	5.83 \pm 20	5.83 \pm 40			5.833 \pm 15
	6.07 \pm 50	6.09 \pm 40			
	6.25 \pm 30	6.23 \pm 40			
	6.45 \pm 40	6.41 \pm 40			
	6.60 \pm 30	6.62 \pm 40			
	6.99 \pm 30	6.99 \pm 40			
	(7.26 \pm 50)	7.17 \pm 40			
		7.37 \pm 40			
	(7.51 \pm 70)				
		7.63 \pm 40			
	7.79 \pm 20	7.73 \pm 40			
	8.00 \pm 30	8.00 \pm 40			

Table 17.3: Levels of ^{17}N from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ and $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ (continued)

E_x (MeV \pm keV)					
(1959LI47) ^a	(1965HA05) ^a	(1966MC05) ^a	(1969TH01) ^b	(1960JA13) ^c	A
	(8.25 \pm 30)	8.14 \pm 40			
		8.55 \pm 40 ^e			
		8.93 \pm 40			
		9.26 \pm 40			
		9.74 \pm 40			

A: A.D.W. Jones, private communication: $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$. See also (1968JO1G, 1969JO1L).

^a Measurement of proton groups from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$.

^b Measurement of γ -ray from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$.

^c Measurement of alpha groups from $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$.

^d The proton groups to this level and the ones below are not completely resolved.

^e This state and the ones below are broad.

^f This may represent a doublet.

^g $l = 1$.

^h $l = 0$.

ⁱ States observed but E_x not determined.

7. $^{18}\text{O}(\text{n}, \text{d})^{17}\text{N}$ $Q_m = -13.718$

See ^{19}O in (1972AJ02).

8. $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$ $Q_m = -10.449$

Not reported.

9. $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ $Q_m = 3.872$
 $Q_0 = 3.872 \pm 0.015$ (1960JA13).

Alpha-particle groups corresponding to ^{17}N states with $E_x < 5.9$ MeV are displayed in Table 17.3 (1960JA13, A.D.W. Jones, private communication). Angular distributions to most of these have been measured at $E_t = 12.9$ MeV (1968JO1G, 1969JO1L): the spectroscopic strength of the transition to $^{17}\text{N}^*(5.52)$ is equal to that of the ground state: it is probably the $p_{3/2}$ hole state.

Table 17.4: Radiative decays in ^{17}N ^a †

E_i (MeV)	E_f (MeV)	Branch (%)
1.37	0	100
1.86 + 1.91	0	75
	1.37	25
1.91	0	^b
2.53	0	13
	1.37	30
	1.91	57
3.13	0	< 14
	1.91	^b
3.22	0	50
3.66	1.86 + 1.91	^b
	3.13	^b
4.02	0	10
	1.86 + 1.91	40
	2.53	50
4.22	1.37	> 40
4.47	1.86 + 1.91	100
5.20	0	< 11
	1.86 + 1.91	^b
	3.22	^{b,c}
5.52	0	20
	1.37	30
	1.86 + 1.91	^b
5.83	0	< 12
	1.37	^b
	3.13 + 3.22	^{b,c}

† Recent work (reaction 9, A.D.W. Jones, private communication) shows a number of states in ^{17}N addition to the ones listed below: see Table 17.1.

^a (1965HA05, 1969TH01).

^b Transition seen but branching ratio not measured.

^c See, however, (1969TH01).

10. $^{19}\text{F}(\text{n}, ^3\text{He})^{17}\text{N}$

$$Q_m = -16.217$$

Not reported.

¹⁷O
(Figs. 7 and 9)

GENERAL: (See also (1959AJ76).)

Shell model: (1957WI1E, 1959BR1E, 1959FE1B, 1959KH1A, 1959SA11, 1960AK1A, 1960TA1C, 1961BA1E, 1961NE1B, 1962BH09, 1962TA1B, 1962TA1E, 1963CO12, 1963HA05, 1963KU1B, 1963PA03, 1964BR1H, 1964RI1A, 1965GI1B, 1965LE1E, 1965MA16, 1965ZA1B, 1966AR10, 1966BO1R, 1966BR04, 1966BR1R, 1966BR1Q, 1966DE18, 1966LA1E, 1966MA12, 1966QU1A, 1966RI1F, 1966SO05, 1966ZA1E, 1967BO1T, 1967EL03, 1967EN01, 1967FE01, 1967GO04, 1967LY02, 1967NI1D, 1967PA05, 1967PF01, 1968BI07, 1968DE13, 1968EL1C, 1968EL1A, 1968HE1H, 1968HO1H, 1968KA1E, 1968MA2B, 1968NI1A, 1968SU1E, 1968ZU02, 1969BO37, 1969CH1R, 1969EL1D, 1969EL1B, 1969GI1B, 1969GU1M, 1969KA09, 1969KU1G, 1969MA38, 1969PI1D, 1969SA1J, 1969UL03, 1969ZU1D, 1970BO2B, 1970EL1G, 1970GU1E, 1970HI15, 1970WA01).

Collective model: (1959YU1A, 1960RA1A, 1960SH1A, 1961NE1B, 1962AR1C, 1962DA1B, 1962MA1K, 1962MA23, 1962MA32, 1962MO1B, 1963CH02, 1964BA1L, 1964BR1H, 1965NE1C, 1966BR04, 1966BR1R, 1966BR1Q, 1967FE01, 1967GO04, 1967GR1D, 1968BI07, 1968EL1C, 1968MA2B, 1969BA2E, 1969MA38, 1970BA2B).

Electromagnetic transitions: (1959BA1C, 1959FA1C, 1960RA1A, 1962MA1K, 1962MA32, 1963CH02, 1964BA1L, 1965GR1H, 1965KA1D, 1966MA12, 1966QU1A, 1967KA21, 1969KH1C, 1969MA38, 1970EL08, 1970GO1H, 1970SI1C).

Special levels: (1961WI1D, 1962IN02, 1963CO12, 1965EJ1A, 1965GI1B, 1965LE1E, 1965NE1C, 1966SO05, 1967BE02, 1967GR1D, 1967PF01, 1967PI01, 1968MA2B, 1969BA2E, 1969GI1B, 1969HA1G, 1970BA2B, 1970EL08, 1970RY02, 1970WA01).

Other topics: (1959FE1B, 1959KH1A, 1959SA11, 1960GO1C, 1963WA1E, 1964LI1B, 1964NI1A, 1965HU1D, 1965ZA1B, 1966MI1F, 1966WA1K, 1967EL03, 1967GR1H, 1967KA1G, 1967LY02, 1967NI1D, 1968KA1E, 1968NI1A, 1968SU1F, 1968SU1C, 1968SU1E, 1968WO1C, 1969DE16, 1969FO1G, 1969HE1N, 1969IW1B, 1969JA1P, 1969KA09, 1969NO03, 1969NO1E, 1969PA1G, 1969PI1D, 1969SA1J, 1969SC14, 1969SC1Q, 1970BA1M, 1970HA1T, 1970MC17, 1970SU1B).

Ground state:

$$\mu = -1.89371 \pm 0.00009 \text{ nm (1967SH14);}$$

$$Q = 26.5 \pm 3.0 \text{ mb (1964LI14).}$$

See also (1959FA1C, 1960RA1A, 1962BE1D, 1963BE36, 1964ST1B, 1965IC1A, 1966MA1V, 1966MA19, 1966MI1F, 1967CO1D, 1967GO1A, 1967GR1D, 1967SH05, 1968BE1X, 1968LE1L, 1968PE16, 1968RO1E, 1969CH1R, 1969FU11, 1969SI1E, 1970EL08, 1970SI02, 1970SI1C).

Table 17.5: Energy levels of ^{17}O

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{1}{2}$			1, 2, 5, 6, 8, 9, 10, 12, 14, 19, 20, 22, 29, 30, 31, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48
0.87081 ± 0.22	$\frac{1}{2}^+$	$\tau_m = 258.6 \pm 2.6$ psec	γ	1, 2, 5, 6, 8, 9, 10, 14, 19, 20, 22, 29, 31, 32, 34, 41, 42, 43, 44, 45, 46, 47, 48
3.055 ± 2.5	$\frac{1}{2}^-$	$\tau_m = 120_{-60}^{+80}$ fsec	γ	5, 6, 8, 9, 12, 14, 22, 29, 31, 34, 41, 42, 45, 46
3.841 ± 3	$\frac{5}{2}^-$	$\tau_m \leq 25$ fsec	γ	5, 6, 8, 9, 12, 14, 19, 20, 29, 31, 37, 41, 42, 45, 46
4.554 ± 6	$\frac{3}{2}^-$	$\Gamma = 40 \pm 5$	n	5, 6, 8, 9, 14, 19, 20, 23, 29, 37, 41, 42, 46
5.083 ± 10	$\frac{3}{2}^+$	95 ± 5	n	8, 9, 14, 19, 23, 29, 41, 42
5.217 ± 5	$\frac{7}{2}^- \rightarrow \frac{11}{2}^-$	< 8		6, 8, 9, 14, 19, 20, 29, 37, 41, 46
5.377 ± 3	$\frac{3}{2}^-$	28 ± 7	n	6, 14, 19, 23, 29, 41, 42, 46
5.696 ± 3	$\frac{7}{2}^-$	3.4	n	6, 8, 9, 14, 19, 20, 23, 29, 37, 41, 46
5.731 ± 3	$\frac{1}{2}^+$	< 1	n	5, 6, 8, 9, 14, 23, 29, 46
5.867 ± 3	$\frac{3}{2}^+$	6.6	n	6, 8, 9, 14, 19, 23, 29, 46
5.935 ± 3	$\frac{1}{2}^-$	23 ± 10	n	5, 8, 9, 14, 19, 23, 29, 46
6.356 ± 8	$\frac{1}{2}^+$	135	n	5, 6, 14, 19, 23
6.859 ± 3		< 1	n	5, 6, 8, 9, 14, 19, 23, 29, 46

Table 17.5: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
6.970 \pm 3		< 1	n	8, 9, 14, 19, 23, 46
7.166 \pm 2.4	$\frac{5}{2}^-$	2.7 \pm 1	n, α	5, 7, 8, 9, 14, 19, 23, 28
7.29	$\frac{3}{2}^+$	500	n	23
7.378 \pm 2.4	$\frac{5}{2}^+$	0.5	n, α	5, 6, 7, 8, 9, 14, 20, 23, 28, 46
7.379 \pm 3	$\frac{5}{2}^-$	1.1	n	5, 6, 8, 9, 14, 19, 23, 28
(7.53)	$(\frac{3}{2})$	600	n	23
7.569 \pm 7	$\frac{7}{2}^-$	≤ 4	n, α	5, 6, 7, 8, 9, 14, 19, 28
7.67	$\frac{3}{2}^-$	400	n	19, 23
7.684 \pm 5	$\frac{7}{2}^-$	18	n, α	5, 7, 8, 9, 23, 28
(7.69)	$\frac{3}{2}^+$	3	n	8, 9, 23, 28
7.751 \pm 14	$(\frac{11}{2}^-)$			6, 8, 19, 20, 21, 37, 46
7.947 \pm 20	$\frac{1}{2}^-$	79 \pm 10	n, α	7, 23, 28
8.090 \pm 9	$\frac{3}{2}^+$	71 \pm 8	n, α	7, 19, 20, 23, 28
8.213 \pm 9	$\frac{3}{2}^-$	71 \pm 5	n, α	7, 19, 20, 23, 28, 29
8.347 \pm 5	$\frac{1}{2}^+$	9 \pm 3	n, α	7, 23, 28, 29
8.402 \pm 5	$\frac{5}{2}^+$	4 \pm 3	n, α	7, 23, 28
8.467 \pm 5	$\frac{7}{2}^+$	7 \pm 3	n, α	5, 7, 8, 9, 20, 23, 28
8.502 \pm 5	$\frac{5}{2}^-$	5 \pm 3	n, α	7, 8, 9, 19, 23, 28
(8.568 \pm 10)			n	23
8.703 \pm 8	$\frac{3}{2}^-$	50 \pm 3	n, α	6, 7, 8, 9, 19, 23, 28
8.87 \pm 20	$\frac{7}{2}^-$	6	n, α	7, 8, 9, 19, 20, 23
8.90 \pm 15	$\frac{3}{2}^+$	101 \pm 3	n, α	6, 7, 8, 9, 19, 20, 28
8.961 \pm 4	$\frac{7}{2}^-$	21 \pm 3	n, α	6, 7, 8, 9, 19, 20, 23, 28
9.14 \pm 30	$(\frac{9}{2}^-)$			6, 19, 20, 21
9.16 \pm 15	$\frac{1}{2}^-$	4 \pm 3	n, α	7, 28
9.18	$\frac{7}{2}^-$	3	α	7
9.19	$\frac{5}{2}^+$	5.5 \pm 1	n, α	7, 23, 28

Table 17.5: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
9.45	$\geq \frac{3}{2}$	140	n	23
9.50	$\frac{5}{2}^-$	15 ± 1	n, α	5, 7, 19, 28
9.705 ± 5	$\frac{7}{2}^+$	16 ± 1	n, α	6, 7, 19, 23
9.78	$\frac{3}{2}^+$	45	n, α	7, 19, 20, 23
9.88	$\frac{9}{2}^+$	12 ± 1	n, α	6, 7, 19, 20, 23, 28
9.95	$\frac{5}{2}^+$	107	n, α	7, 28
10.13	$\frac{5}{2}^+$	138	n, α	7
10.157 ± 7	$\frac{7}{2}^-$	42	n, α	7, 19, 23, 28
10.24	$\frac{7}{2}^+$	122	n, α	7, 28
10.32	$(\frac{5}{2}^+, \frac{7}{2}^-)$		n, α	7
10.422 ± 10		14 ± 3	n, α	7
10.49	$(\frac{5}{2}^+, \frac{7}{2}^-)$	75 ± 30	n, α	7
10.549 ± 6	$(\frac{7}{2}^-, \frac{9}{2}^+)$	47 ± 15	n, α	7, 19, 23, 24, 28
10.70	$(\frac{7}{2}^+)$	≤ 25	α	6, 7
10.769 ± 10	$(\frac{1}{2}^+, \frac{7}{2}^-)$	80 ± 20	n, α	7, 24, 28
10.909 ± 7	$\frac{5}{2}$	57 ± 15	n, α	7, 19, 23, 24, 28
11.026 ± 10	$T = \frac{1}{2}$	45 ± 10	n, α	7, 19, 24, 28
11.082 ± 6	$(\frac{1}{2})^-; \frac{3}{2}$	< 20		6, 19, 24, 41, 43
11.225 ± 10		100 ± 30	n, α	5, 7, 19, 28
11.51	$\geq \frac{3}{2}$	190	n, (α)	19, 23, 24, 28
11.615 ± 10		120 ± 30	n, α	7, 19, 24, 28
11.748 ± 10		40 ± 25	n, α	7, 28
11.813 ± 15		12 ± 3	n, α	6, 7, 28
11.97	$\geq \frac{3}{2}$	270	n	23
12.002 ± 15			n, α	6, 7, 24, 28
12.11 ± 20		150 ± 50	n, α	6, 7, 24
12.271 ± 15		100 ± 30	n, α	6, 7, 28
12.38 ± 20	$\geq \frac{1}{2}$	130	n, α	7, 23
12.417 ± 15			n, α	7
12.471 ± 5	$(\frac{3}{2})^-; \frac{3}{2}$		n	6, 24, 43

Table 17.5: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
12.592 \pm 15		75 \pm 30	n, α	7, 23
12.666 \pm 15		\approx 75	n, α	7, 23, 24
12.81 \pm 25			n, α	7
12.92 \pm 20		\gtrsim 150	n, α	7
12.950 \pm 8	$\frac{1}{2}^+; \frac{3}{2}$			6, 24, 43
12.994 \pm 8	$T = \frac{3}{2}$		n, α	6, 7, 43
13.073 \pm 15		16 \pm 4	n, α	7
13.481 \pm 15		\approx 120	n, α	7
13.606 \pm 15		250 \pm 100	n, α	7, 23
13.640 \pm 5	$(\frac{5}{2})^+; \frac{3}{2}$			6, 43
14.219 \pm 8	$T = \frac{3}{2}$			29, 43
14.282 \pm 12	$T = \frac{3}{2}$			6, 29, 43
14.62	$(\geq \frac{3}{2})$	340	n	6, 23
14.80	$(T = \frac{3}{2})$			6
14.99	$(\frac{5}{2}^+)$	\approx 150	n, d, α	17, 23
15.101 \pm 8	$T = \frac{3}{2}$		p, d, α	16, 17, 43
15.6		\approx 200	p, d, α	16, 17
20.5	$(\frac{1}{2}^+)$		n, ^3He	11, 23
21.1			n, ^3He	11
21.7	$(\frac{5}{2}^+)$		$^3\text{He}, \alpha$	11
22.1	$(\frac{1}{2}^-, \frac{3}{2}^-, \frac{7}{2}^-)$		n, $^3\text{He}, \alpha$	11
23.0	$(\frac{1}{2}^+)$		^3He	11

 1. $^7\text{Li}(^{14}\text{N}, \alpha)^{17}\text{O}$

$$Q_m = 16.154$$

The angular distribution of the α -particles corresponding to $^{17}\text{O}^*(0, 0.87)$ (unresolved) has been measured at $E(^{14}\text{N}) = 27.6$ MeV (1964WA1B).

 2. $^9\text{Be}(^{16}\text{O}, ^8\text{Be})^{17}\text{O}$

$$Q_m = 2.477$$

The lifetime of $^{17}\text{O}^*(0.87)$, τ_m , is 253 ± 6 psec (1969NI09): see Table 17.7. At $E(^{16}\text{O}) = 15$ and 18 MeV, the transitions to $^{17}\text{O}^*(0, 0.87)$ have been studied (1970BA1J). The excitation curve for the one-neutron transfer to $^{17}\text{O}^*(0.87)$ has been measured for $E(^{16}\text{O}) = 6$ to 22 MeV (1970BA55). See also (1968KN1A).

3. (a) $^{10}\text{B}(^7\text{Li}, \text{p})^{16}\text{N}$	$Q_m = 13.985$	$E_b = 27.767$
(b) $^{10}\text{B}(^7\text{Li}, \text{d})^{15}\text{N}$	$Q_m = 13.723$	
(c) $^{10}\text{B}(^7\text{Li}, \text{t})^{14}\text{N}$	$Q_m = 9.146$	
(d) $^{10}\text{B}(^7\text{Li}, \alpha)^{13}\text{C}$	$Q_m = 21.410$	

Cross sections to various of the final states have been measured at $E(^7\text{Li}) = 5.20$ MeV (1966MC05). See also (1963MO1B).

4. (a) $^{11}\text{B}(^6\text{Li}, \text{p})^{16}\text{N}$	$Q_m = 9.782$	$E_b = 23.564$
(b) $^{11}\text{B}(^6\text{Li}, \text{d})^{15}\text{N}$	$Q_m = 9.520$	
(c) $^{11}\text{B}(^6\text{Li}, \text{t})^{14}\text{N}$	$Q_m = 4.942$	
(d) $^{11}\text{B}(^6\text{Li}, \alpha)^{13}\text{C}$	$Q_m = 17.207$	

Cross sections to various of the final states have been measured at $E(^6\text{Li}) = 4.72$ MeV (1966MC05). See also (1963MO1B).

5. $^{12}\text{C}(^6\text{Li}, \text{p})^{17}\text{O}$	$Q_m = 7.607$
--	---------------

Angular distributions have been reported at $E(^6\text{Li}) = 3$ MeV (1963BA08; p_2, p_3, p_4), 3.4 to 4.0 MeV (1962BL13; $p_0 \rightarrow p_3$), 4.5 to 5.5 MeV (1966HE05; $p_0 \rightarrow p_4$), 5.6 to 6.6 MeV (1970JO09; $p_0 \rightarrow p_3$), 9.0 to 14.0 MeV (1970JO09; $p_0 \rightarrow p_4$) and 20 MeV (1968ME10; p to $^{17}\text{O}^*(8.47)$). Neither the direct reaction model nor the statistical compound nucleus model alone is adequate to describe the data (1970JO09). Proton groups to various states of ^{17}O with $E_x \leq 15.8$ MeV have been identified by (1966HE05, 1968ME10, 1970JO09). See also (1960SH05, 1967DZ01), (1967CA1D), (1968GA1J, 1969GI1B; theor.) and ^{18}F in (1972AJ02).

6. $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$	$Q_m = 2.579$
--	---------------

Angular distributions of the d_0 and d_1 groups have been measured at $E(^7\text{Li}) = 3.24$ to 3.64 MeV (1967MO23). At $E(^7\text{Li}) = 20$ and 24.4 MeV, deuteron groups are observed to many states of ^{17}O with $E_x < 15$ MeV, including states at $E_x = 8.50$ and 11.85 MeV and probable $T = \frac{3}{2}$ states at 11.08, 12.47, 12.99, 13.64, 14.28 and 14.80 MeV (1969BA2U, 1970COZA). See also (1970CA1N) and ^{19}F in (1972AJ02).

Table 17.6: Resonances in $^{13}\text{C}(\alpha, \alpha_0)^{13}\text{C}$, $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{16}\text{O}(n, \alpha)^{13}\text{C}$

E_α (MeV \pm keV)	E_n (MeV)	Γ_α/Γ	$\Gamma_{\text{c.m.}}$ (keV)	J^π	E_x^a (MeV)	Refs.
1.063 \pm 4			2.7 \pm 1	$\frac{5}{2}^+$	7.170	A
1.340 \pm 4			≤ 2	$\frac{5}{2}^+$	7.382	(A, 1969SC04)
1.585 \pm 7			≤ 4	$\frac{7}{2}^-$	7.569	(1956RU1A, 1957WA46, 1969SC04)
1.736 \pm 5			22	$\frac{5}{2}^+$	7.684	(1956RU1A, 1957WA46, 1969SC04)
2.080 \pm 20	4.0	0.03	79 \pm 10	$\frac{5}{2}^-$	7.947	(B, 1965BA32, 1967SE07)
2.250 \pm 20	4.15	0.05	71 \pm 8	$\frac{5}{2}^+$	8.077	(C, 1965BA32, 1967SE07)
2.420 \pm 20	4.25	0.11	71 \pm 5	$\frac{5}{2}^-$	8.207	(C, 1965BA32, 1967SE07)
2.603 \pm 5	4.46	0.44	9 \pm 3	$\frac{5}{2}^+$	8.347	(D, 1965BA32, 1967SE07)
2.675 \pm 5	4.53	0.08	4 \pm 3	$\frac{5}{2}^+$	8.402	(D, 1965BA32, 1967SE07)
2.760 \pm 5	4.59	0.97	7 \pm 3	$\frac{7}{2}^+$	8.467	(D, 1965BA32, 1967SE07)
2.805 \pm 5	4.62	0.26	5 \pm 3	$\frac{5}{2}^-$	8.502	(D, 1965BA32, 1967SE07)
3.08 \pm 15	4.85	0.06	50 \pm 3	$\frac{5}{2}^-$	8.71	(D, 1963DA12, 1965BA32, 1967SE07, 1971BA06)
(3.1)			broad	$\frac{5}{2}^-$	8.7	(1971BA06)
3.305		1.00	6	$\frac{7}{2}^-$	8.884	(1965BA32)
3.33 \pm 15	5.05	0.50	101 \pm 3	$\frac{5}{2}^+$	8.90	(D, 1963DA12, 1965BA32, 1967SE07, 1971BA06)
3.42 \pm 15	5.12	0.04	21 \pm 3	$\frac{7}{2}^-$	8.97	(D, 1963DA12, 1965BA32, 1967SE07, 1971BA06)
3.67 \pm 15	5.32	0.45	4 \pm 3	$\frac{5}{2}^-$	9.16	(1956BO61, 1963DA12, 1967SE07, 1968KE02)
3.69		1.00	3	$\frac{7}{2}^-$	9.18	(1968KE02)
3.72	5.37	0.20	5.5 \pm 1	$\frac{5}{2}^+$	9.20	(1956BO61, 1956SC1C, 1963DA12, 1967SE07, 1968KE02)
4.11	5.68	0.85	15 \pm 1	$\frac{5}{2}^-$	9.50	(1956BO61, 1956SC1C, 1963DA12, 1967SE07, 1968KE02)
(4.3)			broad	$\frac{5}{2}^-$	(9.6)	(1971BA06)
4.40		0.70	16 \pm 1	$\frac{7}{2}^+$	9.72	(1956BO61, 1956SC1C, 1967SE07, 1968KE02, 1971BA06)
	5.92					(1963DA12)
4.42		0.90	61	$\frac{5}{2}^+$	9.74	(1968KE02, 1971BA06)
4.58	6.08	0.18	12 \pm 1	$\frac{5}{2}^+$	9.86	(1963DA12, 1967SE07, 1968KE02, 1971BA06)
4.70	6.22	0.78	107	$\frac{5}{2}^+$	9.95	(1963DA12, 1968KE02, 1971BA06)
4.94		0.85	138	$\frac{5}{2}^+$	10.13	(1968KE02)

Table 17.6: Resonances in $^{13}\text{C}(\alpha, \alpha_0)^{13}\text{C}$, $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{16}\text{O}(n, \alpha)^{13}\text{C}$ (continued)

E_α (MeV \pm keV)	E_n (MeV)	Γ_α/Γ	$\Gamma_{\text{c.m.}}$ (keV)	J^π	E_x^a (MeV)	Refs.
4.995	6.41	0.15	46	$\frac{7}{2}^-$	10.176	(1963DA12, 1968KE02, 1970RO08, 1971BA06)
5.08	6.59	0.60	122	$\frac{7}{2}^+$	10.24	(1963DA12, 1967SE07, 1968KE02)
5.185				$\frac{5}{2}^+, \frac{7}{2}^-$	10.321	(1968KE02, 1970RO08)
5.317 ± 10			14 ± 3		10.422	(1963SP02, 1967SE07, 1968KE02)
5.40			75 ± 30	$\frac{5}{2}^+, \frac{7}{2}^-$	10.49	(1967SE07, 1968KE02)
5.496 ± 10	6.81		47 ± 15	$\frac{7}{2}^-, \frac{9}{2}^+$	10.559	(1963DA12, 1963SP02, 1968KE02, 1970RO08)
5.68		1.00	≤ 25	$(\frac{7}{2}^+)$	10.70	(1968KE02)
5.771 ± 10	7.04		80 ± 20	$\frac{1}{2}^+, \frac{7}{2}^-$	10.769	(1963DA12, 1963SP02, 1968KE02, 1970RO08)
5.945 ± 10	7.20		57 ± 15	$\frac{5}{2}^-$	10.902	(1963DA12, 1963SP02, 1968KE02, 1970RO08)
6.107 ± 10	7.32		45 ± 10		11.026	(1963DA12, 1963SP02, 1968KE02)
6.367 ± 10	7.59		100 ± 30		11.225	(1963DA12, 1963SP02)
	(7.79)				(11.47)	(1963DA12)
6.878 ± 10	7.90		120 ± 30		11.615	(1963DA12, 1963SP02)
7.051 ± 10	(8.06)		40 ± 25		11.748	(1963DA12, 1963SP02)
7.136 ± 15	8.22		12 ± 3		11.813	(1963DA12, 1963SP02)
7.384 ± 15	8.38				12.002	(1963DA12, 1963SP02)
7.52 ± 20			150 ± 50		12.11	(1963SP02)
7.736 ± 15	8.66		100 ± 30		12.271	(1963DA12, 1963SP02)
7.88 ± 20					12.38	(1963SP02)
7.927 ± 15					12.417	(1963SP02)
8.156 ± 15			75 ± 30		12.592	(1963SP02)
8.253 ± 15			≈ 5		12.666	(1963SP02)
8.44 ± 25					12.81	(1963SP02)
8.59 ± 20			$\gtrsim 150$		12.92	(1963SP02)
8.72 ± 20					13.02	(1963SP02)
8.785 ± 15			16 ± 4		13.073	(1963SP02)
9.319 ± 15			≈ 120		13.481	(1963SP02)

Table 17.6: Resonances in $^{13}\text{C}(\alpha, \alpha_0)^{13}\text{C}$, $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{16}\text{O}(n, \alpha)^{13}\text{C}$ (continued)

E_α (MeV \pm keV)	E_n (MeV)	Γ_α/Γ	$\Gamma_{\text{c.m.}}$ (keV)	J^π	E_x^a (MeV)	Refs.
9.483 \pm 15			250 \pm 100		13.606	(1963SP02)

A: (1953JO1A, 1956RU1A, 1957WA46).

B: (1955SE1A, 1956BO61, 1956RU1A, 1956SC1C, 1957WA46).

C: (1955SE1A, 1956BE98, 1956BO61, 1956RU1A, 1956SC1C, 1957WA46).

D: (1956BE98, 1956BO61, 1956SC1C, 1957WA46).

^a I am indebted to Dr. C.H. Johnson for a discussion of the nature of many of these states.

Table 17.7: Lifetime of $^{17}\text{O}^*(0.87)$ ^a

τ_m (psec)	Refs.
255 ± 13	(1960KA10)
263 ± 8	(1963LO03)
258.7 ± 4.2	(1964BE15)
233 ± 27	(1965AL14)
263 ± 7	(1965MC10)
232 ± 8	(1967BI05)
261 ± 7	(1969GO04)
253 ± 6	(1969NI09)
258.6 ± 2.6	mean ^b

^a See also (1953TH14, 1962GA15, 1967WA1C, 1968SC1E).

^b Mean does not include value of (1967BI05).

7. (a) $^{13}\text{C}(\alpha, n)^{16}\text{O}$

$$Q_m = 2.215$$

$$E_b = 6.357$$

(b) $^{13}\text{C}(\alpha, \alpha)^{13}\text{C}$

The yield of neutrons (reaction (a)) increases monotonically for $E_\alpha = 0.475$ to 0.700 MeV: $S(E) = [(5.48 \pm 1.77) + (12.05 \pm 3.91)E] \times 10^5 \text{ MeV} \cdot \text{b}$ (1968DA05). The astrophysical considerations are discussed by (1968DA1D). See also (1959AJ76).

Elastic scattering studies (reaction (b)) have been carried out for $E_\alpha = 2.0$ to 3.5 MeV (1965BA32) and 3.5 to 6.5 MeV (1968KE02). Recent measurements of yield curves for reaction (a) have been made at $E_\alpha = 2.0$ to 3.5 MeV (1965BA32), 2.0 to 5.6 MeV (1967SE07; σ_t), 3.5 to 6.5 MeV (1968KE02), 4.5 to 10.5 MeV (1970RO08; n_0), 5 to 10 MeV (1963SP02: 6.13 and 7 MeV γ) and 17.4 to 22.5 MeV (1963DE27; n_0). Angular distributions have been measured at many energies.

A number of resonances in the neutron yield and of anomalies in the elastic scattering have been observed: see Table 17.6 (1953JO1A, 1956BE98, 1956BO61, 1956RU1A, 1957WA46, 1963SP02, 1965BA32, 1967SE07, 1968KE02, 1970RO08). See also (1960ST02). Table 17.6 also shows the results from the inverse reaction $^{16}\text{O}(n, \alpha)^{13}\text{C}$. The study of the n_0 yield by (1970RO08) indicates more structure than is displayed in Table 17.6. In the range $E_n = 17.4$ to 22.4 MeV the n_0 cross section does not show any appreciable variation with energy. See also (1968LE1N) and (1963KE1A; theor.).

Table 17.8: ^{17}O levels from the study of $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$ and $^{13}\text{C}(^7\text{Li}, \text{t})^{17}\text{O}$ (1970BE31)

E_x^a (MeV)	J^π	$d\sigma/d\omega$ in ($\mu\text{b}/\text{sr}$) ^b	
		($^6\text{Li}, \text{d}$)	($^7\text{Li}, \text{t}$)
0 ^b	$\frac{5}{2}^+$	105	75
0.87 ^b	$\frac{1}{2}^+$	180	92
3.06 ^b	$\frac{1}{2}^-$	560	750
3.84 ^b	$\frac{5}{2}^-$	340	1400
4.55 ^b	$\frac{3}{2}^-$	285	1350
5.08 ^b	$\frac{3}{2}^+$	180	250
5.22 ^b	$\frac{7}{2}^- \rightarrow \frac{11}{2}^-$	245	230
5.38	$\frac{3}{2}^-$	c	c
5.70 ^b	$\frac{7}{2}^-$	230	530
5.73 ^b	$\frac{1}{2}^+$		
5.87	$\frac{3}{2}^+$		
5.94	$\frac{1}{2}^-$	90	150
6.24		c	c
6.36	$\frac{1}{2}^+$	c	c
6.86 ^b		92	125
6.97 ^b		200	320
7.17 ^b	$\frac{5}{2}^-$	350	1050
7.29	$\frac{3}{2}^+$	d	d
7.38 ^b	$\frac{5}{2}^+$	720	2000
7.57 ^b	$\frac{7}{2}^-$	98	310
7.68 ^b	$\frac{7}{2}^-$	620	1100
7.76 ± 0.02 ^{a,b}	$(\frac{11}{2}^-)$		
7.95	$\frac{1}{2}^-$	d	d
8.09	$\frac{3}{2}^+$	c	c
8.21	$\frac{3}{2}^-$	d	d
8.35	$\frac{1}{2}^+$	d	d
8.40	$\frac{5}{2}^+$	d	d
8.47 ^b	$\frac{7}{2}^+$	940	2400
8.50 ^b	$\frac{5}{2}^-$		

Table 17.8: ^{17}O levels from the study of $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$ and $^{13}\text{C}(^7\text{Li}, \text{t})^{17}\text{O}$ (1970BE31) (continued)

E_x^a (MeV)	J^π	$d\sigma/d\omega$ in ($\mu\text{b}/\text{sr}$) ^b	
		($^6\text{Li}, \text{d}$)	($^7\text{Li}, \text{t}$)
8.70	$\frac{3}{2}^-$	50	200
8.88	$\frac{7}{2}^-$	1400	2000
8.96	$\frac{7}{2}^-$		

^a These energies were not determined in this experiment, except for that of $^{17}\text{O}^*(7.76)$.

^b Angular distributions were obtained for these states. For these $d\sigma/d\omega$ were taken at the maximum of the distribution. For the other states $d\sigma/d\omega$ is the 30° value.

^c Group not seen: obscured by contaminant.

^d This level was not observed.

J^π assignments derived from polarization measurements at $E_\alpha = 3.36$ to 4.80 MeV are displayed in Table 17.6 (1971BA06). See also (1969SC04).

8. $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$ $Q_m = 4.885$

At $E(^6\text{Li}) = 18$ MeV, deuteron groups have been seen corresponding to many ^{17}O states with $E_x < 9$ MeV: see Table 17.8 (1970BE31). See also (1968OG1A, 1970OG1A).

9. $^{13}\text{C}(^7\text{Li}, \text{t})^{17}\text{O}$ $Q_m = 3.890$

This reaction has been studied at $E(^7\text{Li}) = 17$ MeV: see reaction 8 and Table 17.8 (1970BE31). See also (1968OG1A).

10. $^{13}\text{C}(^{16}\text{O}, ^{12}\text{C})^{17}\text{O}$ $Q_m = -0.804$

At $E(^{16}\text{O}) = 17$ and 20 MeV, the transitions to $^{17}\text{O}^*(0, 0.87)$ have been studied (1968KN1A, 1970BA1J). The excitation curve for the one-neutron transfer to $^{17}\text{O}^*(0.87)$ has been measured for $E(^{16}\text{O}) = 12$ to 25 MeV (1970BA55).

11. (a) $^{14}\text{C}(^3\text{He}, \text{n})^{16}\text{O}$ $Q_{\text{m}} = 14.616$ $E_{\text{b}} = 18.759$
 (b) $^{14}\text{C}(^3\text{He}, ^3\text{He})^{14}\text{C}$
 (c) $^{14}\text{C}(^3\text{He}, \alpha)^{13}\text{C}$ $Q_{\text{m}} = 12.402$

The n_0 yield (reaction (a)) for $E(^3\text{He}) = 1.6$ to 3.25 MeV indicates two resonances at $E(^3\text{He}) = 2.1$ and 2.8 MeV, corresponding to $^{17}\text{O}^*(20.5, 21.1)$ (1961JO24). The excitation function shows a resonance in the n_0 and n_{3+4} yields at $E(^3\text{He}) = 4.1$ MeV [$^{17}\text{O}^*(22.1)$], but not in the n_{1+2} yield: $J^\pi = \frac{1}{2}^-$ or $\frac{3}{2}^-$ is suggested (1970HO08).

Resonances are observed in the α -yield (reaction (c)) at $E(^3\text{He}) = 3.6$ and 4.1 MeV [$^{17}\text{O}^*(21.7, 22.1)$] and in the ^3He yield (reaction (b)) at 5.1 MeV [$^{17}\text{O}^*(23.0)$], with $J^\pi = (\frac{5}{2}^+)$, $(\frac{7}{2}^-)$ and $(\frac{1}{2}^+)$, respectively (1970KE1D). The variation of the ^3He optical parameters has been studied for $E(^3\text{He}) = 10$ to 18 MeV (1970DU07).

12. $^{14}\text{C}(\alpha, \text{n})^{17}\text{O}$ $Q_{\text{m}} = -1.819$

A study of $\text{n}-\gamma$ correlations leads to $J^\pi = \frac{1}{2}^-$ for $^{17}\text{O}^*(3.06)$ and probably to $J^\pi = \frac{5}{2}^-$ for $^{17}\text{O}^*(3.84)$. The upper limits to the decays $3.06 \rightarrow 0$ and $3.84 \rightarrow 0.87$ are, respectively, 2 and 5%. The lifetimes of $^{17}\text{O}^*(3.06, 3.84)$ are 120_{-60}^{+80} and < 25 fsec, respectively (1964AL11).

13. (a) $^{14}\text{N}(\text{t}, \text{p})^{16}\text{N}$ $Q_{\text{m}} = 4.840$ $E_{\text{b}} = 18.621$
 (b) $^{14}\text{N}(\text{t}, \text{d})^{15}\text{N}$ $Q_{\text{m}} = 4.577$
 (c) $^{14}\text{N}(\text{t}, \text{t})^{14}\text{N}$
 (d) $^{14}\text{N}(\text{t}, \alpha)^{13}\text{C}$ $Q_{\text{m}} = 12.264$

Excitation functions have been measured in the range $E_{\text{t}} = 1.0$ to 2.0 MeV for the p_0 , p_1 , p_2 , p_3 groups, the d_0 and t_0 groups and the α_0 and α_1 groups: the reactions appear to proceed primarily via a direct interaction mechanism (1964SC09). See also ^{16}N , and ^{13}C and ^{15}N in (1970AJ04).

14. (a) $^{14}\text{N}(\alpha, \text{p})^{17}\text{O}$ $Q_{\text{m}} = -1.193$
 (b) $^{14}\text{N}(\alpha, \text{p}\alpha)^{13}\text{C}$ $Q_{\text{m}} = -7.546$
 $Q_0 = -1.200 \pm 0.017$ (1967SP09).

Differential cross sections have been measured at $E_\alpha = 8.12$ to 11.52 MeV (1969SC21; $\text{p}\gamma_1$), 12.9 to 24.9 MeV (1970ZE01; p_0 , p_1), 13.0 , 13.4 , 14.25 , 14.5 and 18.1 MeV (1969RO07: states with $E_x < 7.6$ MeV), 17.3 to 24.9 MeV (1970ZE01; p_2), 22.3 MeV (1963YA1C; p_0 , p_1), 22.5

to 24.9 MeV (1970ZE01; p_3), and 26.8, 28.1 and 33.3 MeV (1961YA02; $p_0 \rightarrow p_3$). Using the $(2J + 1)$ rule, $J = \frac{7}{2}, \frac{9}{2}$ or $\frac{11}{2}$ for $^{17}\text{O}^*(5.22)$. The previously reported states at $E_x = 6.24$ and 7.29 MeV were not excited, but $^{17}\text{O}^*(6.97)$ is confirmed by the work of (1969RO07). The lifetime of $^{17}\text{O}^*(0.87)$, measured by this reaction and by reaction 29, $\tau_m = (4.30 \pm 0.21) \times 10^{-10}$ sec (1962GA15): see Table 17.7. See also (1959AJ76) and ^{18}F in (1972AJ02).

At $E_\alpha = 22.9$ MeV, the sequential decay (reaction (b)) appears to take place via a number of ^{17}O states with $8.46 \leq E_x \leq 13.57$ MeV. Those involved are believed to have $J \geq \frac{5}{2}$, $\Gamma_\alpha/\Gamma \geq 0.6$ (1969BA17). See also (1968KU1C).

15. $^{15}\text{N}(\text{d}, \text{n})^{16}\text{O}$ $Q_m = 9.901$ $E_b = 14.044$

The excitation function has been measured for $E_d = 0.5$ to 5.3 MeV. Above $E_d = 1.0$ MeV, pronounced peaks are observed, presumably to be ascribed to numerous overlapping resonances (1958WE31). See also (1970MU1H). Polarization measurements (n_0 group) are reported at $E_n = 1.6$ to 3.0 MeV (1966BR12), 2.65 MeV (1966BU09), 2.75 MeV (1969BU19), 3.09 to 3.83 MeV (1968ME15) and 4.35 and 5.50 MeV (1967BU16). See also (1966BR1E). See also ^{16}O .

16. $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$ $Q_m = 0.262$ $E_b = 14.044$

The excitation curve has been obtained for $E_d = 0.3$ to 2.7 MeV. There is some resonance structure at $E_d = 1.3$ and 1.9 MeV [$^{17}\text{O}^*(15.2, 15.7)$] (1957BO04). See also ^{16}N .

17. $^{15}\text{N}(\text{d}, \alpha)^{13}\text{C}$ $Q_m = 7.687$ $E_b = 14.044$

The α_0 yield curve for $E_d = 0.8$ to 1.8 MeV indicates two resonances at $E_d = 1.06$ and 1.25 MeV [$\Gamma \approx 100$ and 200 keV, respectively], attributed to an ^{17}O state at $E_x = 14.98$ MeV [$J^\pi = \frac{5}{2}^+$] and to one or more ^{17}O states at $E_x = 15.15$ MeV [$J^\pi = \frac{5}{2}^-$ or $\frac{7}{2}^-$] (1966TI03). In the range $E_d = 1.2$ to 2.5 MeV a broad maximum is observed in both the α_0 and α_1 yields at $E_d \approx 1.7$ MeV (1965MA59). See also (1959FI30).

18. $^{15}\text{N}(\text{t}, \text{n})^{17}\text{O}$ $Q_m = 7.787$

Not reported.

19. $^{15}\text{N}(^3\text{He}, \text{p})^{17}\text{O}$ $Q_m = 8.550$

Table 17.9: Decay properties of the lowest $T = \frac{3}{2}$ states in $A = 17$

	$^{17}\text{O}^*(11.08)$	$^{17}\text{F}^*(11.20)$
$\Gamma_{\text{c.m.}}$	$< 20 \text{ keV}^{\text{a}}$	$< 600 \text{ eV}^{\text{b}}$
Branching ratios (%):		
$^{16}\text{O}(\text{g.s.})$	$91 \pm 5^{\text{c}}$	$8.8 \pm 1.6^{\text{c}}$ $12 \pm 4^{\text{d}}$
$^{16}\text{O}^*(6.05)$	$5 \pm 2^{\text{c}}$	$23 \pm 5^{\text{c}}$ $26 \pm 8^{\text{d}}$
$^{16}\text{O}^*(6.92)$		$28 \pm 13^{\text{d}}$
$^{16}\text{O}^*(7.12)$		$34 \pm 14^{\text{d}}$
$\theta^2(\text{g.s.})/\theta^2(6.05)$	$3.4 \pm 1.4^{\text{c}}$	$0.16 \pm 0.05^{\text{c}}$

^a D.C. Hensley, quoted in (1970MC02).

^b (1967PA17).

^c (1970MC02).

^d (1969HAZE).

At $E(^3\text{He}) = 18 \text{ MeV}$, many ^{17}O states with $E_x < 11.6 \text{ MeV}$ have been observed. The levels at 5.70 MeV ($\frac{7}{2}^-$), 7.38 MeV ($\frac{5}{2}^-$) and 11.08 MeV ($\frac{1}{2}^-$; $T = \frac{3}{2}$) are particularly strongly populated (1969ME1K). See also (1965SE01, 1968BE1Y, 1968SE1C). At $E(^3\text{He}) = 8.4 \text{ MeV}$ a $T = \frac{1}{2}$ state at $E_x = 11.02 \text{ MeV}$ is excited (1970MC02): see also reaction 43 and Table 17.9. See also (1963PA01) and (1969BA1Z).

20. $^{15}\text{N}(\alpha, \text{d})^{17}\text{O}$

$$Q_{\text{m}} = -9.803$$

At $E_{\alpha} = 45.4 \text{ MeV}$, the deuteron spectrum is dominated by the groups corresponding to states with $E_x = 7.742 \pm 0.020$ and $9.137 \pm 0.030 \text{ MeV}$. These states are assigned $J^{\pi} = (\frac{11}{2}^-)$ and $(\frac{9}{2}^-)$, respectively and arise from a dominant $(\text{d}_{5/2})^2_5\text{p}_{1/2}^{-1}$ configuration. Angular distributions were measured as well for the deuterons corresponding to $^{17}\text{O}(0)$ and to states with $E_x = 0.87 \pm 0.05$, 5.208 ± 0.030 , 5.690 ± 0.030 , 7.367 ± 0.030 , 8.459 ± 0.030 , 8.890 ± 0.030 and $9.814 \pm 0.030 \text{ MeV}$. In addition the excitation of states with $E_x = 3.85 \pm 0.05$, 4.57 ± 0.05 and $8.147 \pm 0.030 \text{ MeV}$ is also reported (1969LU07). See also (1962HA40, 1966RI04) and (1963GL1C; theor.).

21. $^{15}\text{N}(^{11}\text{B}, ^9\text{Be})^{17}\text{O}$

$$Q_{\text{m}} = -1.775$$

Table 17.10: Recent angular distribution, total cross-section and polarization measurements in $^{16}\text{O}(n, n)^{16}\text{O}$ ^a

(a) *Angular distribution studies*

E_n (MeV)	Group	Refs.
0.3 – 1.7	n_0	(1961LA1A)
0.9	n_0	(1962LO09)
1.12 – 1.16, 1.64 – 1.70, 1.77 – 3.67	n_0	(1968JO1F)
1.51 – 2.25	n_0	(1962MA05)
1.96, 3.21 – 3.44	n_0	(1970FO03)
2.0 – 4.1	n_0	(1962HU07)
2.2 – 4.2	n_0	(1967JO12)
3.1 – 4.7	n_0	(1966LI03)
5.0 – 6.5	n_0	(1960SM02)
13.9	$n_0, n_{1+2+3+4}$	(1963BA46)
14	n_0	(1967BE75)
14.1	n_0, n_{1+2}, n_{3+4}	(1969ME15)
14.8	n_0	(1967LU1B)
15	n_0	(1959BE1B)

(b) *Cross-section measurements*

E_n (MeV)	Type	Refs.
0.003 – 10.0 eV	σ_t	(1960WA07)
1.44 eV	σ_{sc}	(1965RA15)
10 – 500 keV	σ_t	(1966MO09)
1.77 – 3.67 MeV	σ_t	(1968JO1F)
3 – 8	σ_t	(1969DA13)
3.3 – 5.1	σ_t	(1960TS02)
3.3 – 5.2	σ_t	(1968MA2D)
3.33 – 3.87	σ_t	(1967JO12)
3.4 – 16	σ_t	(1961FO07)
14 – 19	σ_{el}	(1970BO30)
14.1	σ_t, σ_{el}	(1969ME15)
18 – 27.6	σ_t	(1960PE25)

Table 17.10: Recent angular distribution, total cross-section and polarization measurements in $^{16}\text{O}(n, n)^{16}\text{O}$ ^a (continued)

(b) *Cross-section measurements* (continued)

E_n (MeV)	Type	Refs.
88 – 150	σ_t	(1966ME14)
13.9	$\sigma_{\text{el}}, \sigma_{\text{ne}}$	(1963BA46)
14.8	σ_{ne}	(1967CH42)
15	σ_{ne}	(1969NY1A)
6.4 – 9.8	$\sigma_n, n'\gamma_{6.1}$	(1959HA13)
6.72 – 11.0	$\sigma_n, n'\gamma_{6.1}, \sigma_n, n'\gamma_{6.9}$	(1970DI1C)
7.50 – 11.0	$\sigma_n, n'\gamma_{7.1}$	(1970DI1C)
7.7 – 9.4	$\sigma_n, n'\gamma_{6.9+7.1}$	(1959HA13)
9.0 – 14	$\sigma_n, n'\gamma_{8.9}$	(1970DI1C)
14.6	$\sigma_n, n'\gamma_{6.1}, \sigma_n, n'\gamma_{6.9+7.1}$	(1969BU08)
15	$\sigma_n, n'\gamma$	(1969NY1A)
17 – 37	$\sigma_n, 2n$	(1961BR1A)
11.08 – 19.04	σ_n, p	(1962DE13)
12.6 – 16.3	σ_n, p	(1961SE02, 1962SE1F)
14.7	σ_n, p	(1960DE19, 1962KA37)
14.8	σ_n, p	(1966MI14, 1966PR14)
3.77 – 4.67	$(\sigma_n, \alpha)_t$	(1966LI03)
5.0 – 8.8	σ_n, α_0	(1963DA12)
7.13 – 12.03	$\sigma_n, \alpha_0, \sigma_n, \alpha_1$	(1968DA1E)
7.6 – 8.7	σ_n, α_1	(1963DA12)
8.1 – 8.5	σ_n, α_{2+3}	(1963DA12)
7.1 – 12.0	$\sigma_n, \alpha_0, \sigma_n, \alpha_t$	(1968PA1X)
12 – 20	σ_n, α_0	(1963BO1E)
14.8 – 18.8	$\sigma_n, \alpha_0, \sigma_n, \alpha_{1+2+3}$	(1968SI06)
15	$\sigma_n, \alpha\gamma$	(1969NY1A)

(c) *Polarization measurements*

E_n (MeV)	Group	Refs.
0.2 – 2.2	n_0	(1962EL01)

Table 17.10: Recent angular distribution, total cross-section and polarization measurements in $^{16}\text{O}(n, n)^{16}\text{O}$ ^a (continued)

(c) *Polarization measurements* (continued)

E_n (MeV)	Group	Refs.
0.84	n ₀	(1962BE1C)
1.51 – 2.03	n ₀	(1962MA05)
2.84, 2.98	n ₀	(1963GL1D)
3.5	n ₀	(1962OT01)

^a See also (1959AJ76), (1964ST25, 1970GA1A).

At $E(^{11}\text{B}) = 115$ MeV, the states at $E_x = 7.7$ and 9.1 MeV [$J^\pi = \frac{11}{2}^-$ and $\frac{9}{2}^-$, respectively] are strongly populated. as in the $^{15}\text{N}(\alpha, d)^{17}\text{O}$ reaction (1967PO1E). See also (1966PO1E, 1967VO1A).

$$22. \ ^{16}\text{O}(n, \gamma)^{17}\text{O} \quad Q_m = 4.143$$

$$\sigma_{\text{capt.}} = 0.178 \pm 0.025 \text{ mb (1964JU05)}.$$

Capture γ -rays with $E_\gamma = 870 \pm 5, 1088 \pm 5, 2180 \pm 25$ and 3271 ± 15 keV, from the transitions to and decay of $^{17}\text{O}^*(0.87, 3.06)$, are reported by (1964JU05). See also (1969HO1X). Astrophysical considerations are discussed by (1968FOZY).

$$23. \ ^{16}\text{O}(n, n)^{16}\text{O} \quad E_b = 4.143$$

The scattering amplitude (bound) is $a = 5.80 \pm 0.05$ fm (1965DO14). The coherent scattering cross section is 4.23 ± 0.07 b (1964ST25). See also (1961WI1A, 1965RA15, 1969BA1P).

Recent cross section and angular distribution measurements are listed in Table 17.10. Cross section data are summarized in (1964ST25), while angular distribution data are displayed in (1970GA1A). See also (1968AL1G).

A large number of resonances have been observed for $E_n \leq 11.5$ MeV: these are displayed in Table 17.11 (1960TS02, 1961FO07, 1967JO12, 1968DA1F, 1968JO1F, 1969DA13, 1970BO30, 1970FO03) and (1959AJ76, 1964ST25). See also (1966LI03). Recent high-resolution cross section measurements and phase-shift analyses have led to a much better understanding of the ^{17}O level structure: see (1967JO12, 1970FO03) for a general review of the work on this reaction. See also (1960PE02, 1961GO1H, 1963KU1F), (1968AL1G) and (1959AU1A, 1960CO1A,

1960FO14, 1960MI1B, 1963JO1D, 1963KA27, 1963LU10, 1963PI03, 1965GA1F, 1965SL1B, 1966AG1A, 1966GA1L, 1967GR1J, 1967RE1C, 1967SC1E, 1967SC1H, 1967UN1A, 1968FO1D, 1969IW1A, 1969IW1C, 1969MA2B, 1970CO1U, 1970MU1C, 1970WA1J; theor.).

Polarization measurements have been carried out for $E_n = 0.2$ to 3.5 MeV: see Table 17.10 (1962BE1C, 1962EL01, 1962MA05, 1962OT01, 1963GL1D). See also the reviews by (1963HA1G, 1963PI03, 1966DA1B, 1966RO1B) and (1967BE1F; theor.)

$$24. \text{}^{16}\text{O}(n, n')\text{}^{16}\text{O}^* \qquad E_b = 4.143$$

Recent cross section measurements are listed in Table 17.10. The cross sections for production of 6.13 and (6.92 + 7.12) γ -rays in the range $E_n = 6.5$ to 10 MeV show a number of resonances: see Table 17.12 (1969HAZE). See also (1970SU01).

$$25. \text{}^{16}\text{O}(n, 2n)\text{}^{15}\text{O} \qquad Q_m = -15.668 \qquad E_b = 4.143$$

The cross section has been measured for $E_n = 17$ to 37 MeV (1961BR1A). See also (1964HE18).

$$26. \text{}^{16}\text{O}(n, p)\text{}^{16}\text{N} \qquad Q_m = -9.639 \qquad E_b = 4.143$$

Recent cross-section measurements are listed in Table 17.10. Resonances are reported at 11.8 and (15.2) MeV (1962DE13) and at 13.50 ± 0.01 , 14.10 ± 0.01 and 15.10 ± 0.05 MeV (1961SE02, 1962SE1F). See also (1958RO1A, 1969BR1F), (1959AJ76, 1960BU1C, 1963LE1D, 1963SH1C, 1966JE1B, 1968AL1G) and (1964DU1B; theor.).

$$27. \text{}^{16}\text{O}(n, d)\text{}^{15}\text{N} \qquad Q_m = -9.901 \qquad E_b = 4.143$$

See (1967VA12) and ^{15}N in (1970AJ04).

$$28. \text{}^{16}\text{O}(n, \alpha)\text{}^{13}\text{C} \qquad Q_m = -2.215 \qquad E_b = 4.143$$

The cross section has been measured from threshold to 20 MeV: see (1959AJ76) and Table 17.10. All of the ^{17}O levels observed in the inverse reaction $^{13}\text{C}(\alpha, n)^{16}\text{O}$ from $^{17}\text{O}^* = 7.95$ to 9.95 are also observed in this reaction: see Table 17.6 (1955SE1A, 1957WA46, 1963DA12). (1963BO1E) also report a broad maximum at $E_n \approx 16$ MeV. See also (1965FU1E), (1960BU1C, 1963CH1C, 1966JE1B) and (1964GA1A; theor.).

Table 17.11: Resonances in $^{16}\text{O}(n, n)^{16}\text{O}$ ^a

E_n (keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\frac{\gamma^2}{\hbar^2/\mu a^2}$ ^b	J^π	E_x (MeV)	Refs.
442	45	0.06	$\frac{3}{2}^+$	4.558	(1964ST25, 1970FO03)
1000	94	0.45	$\frac{3}{2}^+$	5.084	(1964ST25)
1312 ± 3.5	41	0.02	$\frac{3}{2}^-$	5.377	(1968DA1F, 1970FO03)
1651 ± 3	3.4	0.09	$\frac{7}{2}^-$	5.696	(1968JO1F, 1970FO03)
1689 ± 3	< 1			5.731	(1968JO1F, 1970FO03)
1833 ± 3	6.6	0.009	$\frac{3}{2}^+$	5.867	(1968JO1F, 1970FO03)
1906 ± 3	24.5	0.01	$\frac{1}{2}^-$	5.935	(1970FO03)
2353 ± 8	135	0.02	$\frac{1}{2}^+$	6.356	(1970FO03)
2888 ± 3	< 1			6.859	(1968JO1F, 1970FO03)
3006 ± 3	< 1			6.970	(1970FO03)
3212 ± 3	1.4	0.005	$\frac{5}{2}^-$	7.164	(1968JO1F, 1970FO03)
3350	500	0.23	$\frac{3}{2}^+$	7.294	(1967JO12)
3438 ± 3	0.5	0.0002	$\frac{5}{2}^+$	7.376	(1970FO03)
3441 ± 3	1.1	0.003	$\frac{5}{2}^-$	7.379	(1961FO07, 1970FO03)
(3600)	600		$(\frac{3}{2})$	(7.53)	(1961FO07)
3750	405		$\frac{3}{2}^-$	7.670	(1967JO12)
3769	14		$\frac{7}{2}^-$	7.688	(1967JO12)
3772	3		$\frac{3}{2}^+$	7.690	(1967JO12)
4000 ± 50			$\frac{1}{2}^-$	7.91	(1961FO07, 1967JO12) ^c
4200 ± 10	80		$\frac{3}{2}^+$	8.093	(1960TS02, 1961FO07, 1967JO12)
4330 ± 10	70		$\frac{3}{2}^-$	8.215	(1960TS02, 1961FO07, 1967JO12)
(4460)				(8.34)	(1960TS02, 1961FO07)
4540	≤ 10		$\geq \frac{5}{2}$	8.41	(1961FO07) ^c
4610	≤ 11		$\geq \frac{5}{2}$	8.48	(1961FO07) ^c
4650	≤ 13		$\geq \frac{5}{2}$	8.52	(1961FO07) ^c
4705 ± 10				8.568	(1960TS02)
4845 ± 10	55		$\frac{3}{2}$	8.700	(1960TS02, 1961FO07)
5010 ± 10	< 20			8.855	(1960TS02)
5122 ± 4	28		$\frac{7}{2}$	8.960	(1961FO07, 1969DA13)
5360	≤ 17		$\geq \frac{5}{2}$	9.18	(1961FO07) ^c

Table 17.11: Resonances in $^{16}\text{O}(n, n)^{16}\text{O}$ ^a (continued)

E_n (keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\frac{\gamma^2}{\hbar^2/\mu a^2}$ ^b	J^π	E_x (MeV)	Refs.
5640	140		$\geq \frac{3}{2}$	9.45	(1961FO07) ^c
5914 ± 5	28		$\geq \frac{3}{2}$	9.705	(1961FO07, 1969DA13)
6010	28		$\geq \frac{3}{2}$	9.80	(1961FO07) ^c
6100	25		$\geq \frac{1}{2}$	9.88	(1961FO07) ^c
6395 ± 7	38		$\geq \frac{3}{2}$	10.157	(1961FO07, 1969DA13)
6807 ± 7	40		$\geq \frac{3}{2}$	10.545	(1961FO07, 1969DA13)
7200 ± 8	70		$\geq \frac{3}{2}$	10.914	(1961FO07, 1969DA13)
7830	190		$\geq \frac{3}{2}$	11.51	(1961FO07) ^c
8320	270		$\geq \frac{3}{2}$	11.97	(1961FO07) ^c
8740	130		$\geq \frac{1}{2}$	12.36	(1961FO07) ^c
9050	95		$\geq \frac{1}{2}$	12.65	(1961FO07) ^c
10130	400		$(\geq \frac{1}{2})$	13.67	(1961FO07) ^c
11140	340		$(\geq \frac{3}{2})$	14.62	(1961FO07) ^c
11540	180		$(\geq \frac{1}{2})$	14.99	(1961FO07) ^c
17300			$(\frac{1}{2}^+)$	20.4	(1970BO30)

^a See also (1959AJ76).

^b See discussion in (1970FO03).

^c I am indebted to Dr. J.C. Davis for sending me revised energy values due to a recalibration of the analyzing magnet used by (1961FO07).

29. $^{16}\text{O}(d, p)^{17}\text{O}$

$$Q_m = 1.918$$

$$Q_0 = 1.920 \pm 0.003 \text{ (1967SP09).}$$

Angular distributions of the protons to many ^{17}O states have been studied for $E_d = 0.3$ to 26.3 MeV: see Table 17.13 (1959HA29, 1961HA19, 1961KE01, 1962MA25, 1963AL04, 1963AM1A, 1963YA03, 1964AM1A, 1964SC12, 1964TE1C, 1965DE1N, 1965GA1A, 1965LO02, 1965MO16, 1966AL09, 1966GA09, 1966SC09, 1967AL06, 1967OG1A, 1968DI06, 1968HO23, 1968NA06, 1968NG1B, 1969TH04, 1970DA14). See also (1969CO12, 1970CO1P) and ^{18}F in (1972AJ02).

(1970VI03) report a DWBA calculation to the unbound $\frac{3}{2}^+$ state at $E_x = 5.08$ MeV. The width of the corresponding neutron resonance [$E_n = 1.00$ MeV: see reaction 23] is determined to be 67 keV from the magnitude of the cross section.

Table 17.12: Resonances in $^{16}\text{O}(n, n'\gamma)^{16}\text{O}$
(1959HA13)

E_n (MeV)	E_x (MeV)
6.83	10.57
7.07	10.79
7.24	10.95
7.40	11.10
7.87	11.55
8.35	12.00
8.50	12.14
8.84	12.46
9.10	12.70
9.34	12.93

^{17}O levels derived from proton spectra measurements are displayed in Table 17.14 (1951BU1A, 1957BR82, 1965GA1A). The lifetime of $^{17}\text{O}^*(0.87)$ is 258.6 ± 2.6 psec: see Table 17.7 (1960KA10, 1963LO03, 1964BE15, 1965AL14, 1965MC10, 1967BI05, 1969GO04). $E_\gamma = 870.81 \pm 0.22$ keV (1966WI01), 870.5 ± 2.0 keV (1952TH24); the internal conversion coefficient is consistent with E2 (1952TH24). See also (1959LO59, 1960AL35, 1960GO20, 1961JA23, 1961LO1C, 1961PU1B, 1963DO1B, 1965HE1B, 1966BE1E, 1967LA1M, 1968GO1N), (1959AJ76) and (1960BE1B, 1960BI1B, 1960NA1A, 1961BU16, 1961WI1D, 1963GL1C, 1963LE1E, 1963SA1B, 1963SM05, 1963TA1A, 1965BU1G, 1965HU1E, 1965SI1D, 1965SM1A, 1967BA1R, 1967BA2E, 1967SC16, 1967SH1J, 1967SH1K, 1968EL1G, 1968PE1E, 1969HA1V, 1969IC02, 1969MA2C, 1969MC1G, 1969PE1K, 1969PE1L, 1970BU16, 1970OH06, 1970PE14, 1970PE1B; theor.).

$$30. \ ^{16}\text{O}(t, d)^{17}\text{O} \quad Q_m = -2.115$$

The angular distribution of the d_0 groups has been studied at $E_t = 5.5$ MeV (1961BA10). See also (1960MU07) and (1964EL1B; theor.).

$$31. \ ^{16}\text{O}(\alpha, ^3\text{He})^{17}\text{O} \quad Q_m = -16.435$$

Angular distributions of the ^3He particles corresponding to $^{17}\text{O}^*(0, 0.87, 3.06, 3.84)$ have been measured at $E_\alpha = 46$ MeV (1968PR1C).

Table 17.13: Recent $^{16}\text{O}(d, p)^{17}\text{O}$ angular distribution studies ^a

E_d (MeV)	Distribution of proton groups	Refs.
0.32 – 1.07	p ₁	(1968NG1B)
0.33 – 0.45	p ₁	(1965LO02)
0.75 – 1.50	p ₀ , p ₁	(1963AL04)
0.90 – 1.39	p ₀ , p ₁	(1963AM1A, 1964AM1A)
1.3, 4.0	p ₀ , p ₁	(1966GA09)
1.9 – 3.6	p ₀ , p ₁	(1968DI06)
4.0 – 6.0	p ₀ , p ₁ , p ₃	(1970DA14)
5.56	p ₀ , p ₁	(1965MO16)
6.00 – 8.55	p ₀ → p ₈	(1965DE1N, 1969TH04)
6.0 – 11.0	p ₀ , p ₁	(1968NA06)
7.73	p ₀ , p ₁	(1967OG1A)
10	p ₀ → p ₃	(1965GA1A)
10.2, 12.4, 14.8	p ₀ , p ₁	(1959HA29, 1961HA19)
11, 13	p ₀	(1966SC09)
11.8	p ₀ → p ₃	(1964SC12)
12	p ₀ , p ₁ , p ₅	(1966AL09, 1967AL06)
14.3	p ₀ → p ₈ , p ₁₀ , p ₁₃ , p ₁₅ → p ₁₈ , p ₂₀ , p ₂₂₊₂₃	(1968HO23)
14.95	p ₀ → p ₅ , p ₇ , p ₈	(1963YA03)
15	p ₀ → p ₅	(1961KE01)
26.3	p ₀ , p ₁	(1962MA25, 1964TE1C)

^a See also (1959AJ76).

Table 17.14: States of ^{17}O from $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$

E_x^a (MeV \pm keV)	$\Gamma_{\text{c.m.}}^a$ (keV)	E_x^b (MeV \pm keV)	E_x^c (MeV \pm keV)	S^d	$\theta_{\text{abs.}}^2^e$	J^π^f
0	< 8	0	0	0.81	0.045	$\frac{5}{2}^+$
0.871 ± 4^g	< 8	0.870 ± 20	0.883 ± 11	0.71	0.16	$\frac{1}{2}^+$
3.055 ± 4^g	< 8	3.060 ± 30	3.069 ± 10	0.032	0.0024	$\frac{1}{2}^-$
3.846 ± 5^g	< 8	3.850 ± 30	3.856 ± 11	0.028	0.0022	$\frac{7}{2}^-$ ⁱ
4.553 ± 6	40 ± 5	4.580 ± 20	4.567 ± 14		0.0071	$\frac{3}{2}^-$
5.083 ± 10	95 ± 5	5.070 ± 20			0.047	$\frac{3}{2}^+$
5.215 ± 5	< 8		5.229 ± 13			
5.378 ± 7	28 ± 7	5.310 ± 20	5.397 ± 14			$\frac{3}{2}^-$
5.695 ± 5^h	< 8				0.013	$\frac{7}{2}^-$
5.731 ± 5^h	< 8	5.760 ± 20	5.723 ± 14			
5.866 ± 5	< 8		5.875 ± 15			
5.940 ± 15	23 ± 10		5.957 ± 15			
		6.240 ± 20				
		6.890 ± 30	6.869 ± 14			
			(6.986 ± 15)			
			(7.371 ± 15)			
		7.510 ± 30				
		8.270 ± 40				
		(8.590 ± 40)				
		9.060 ± 40				

^a $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: (1957BR82).

^b $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$: (1951BU1A).

^c $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$: (1952WA1A).

^d $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: (1970DA14).

^e $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: (1961KE01, 1963YA03).

^f Assignments from (1955AJ61, 1956GR37, 1958RI1A, 1961KE01, 1963YA03, 1964SC12).

^g (1965GA1A) report $E_x = 873 \pm 5, 3056 \pm 4$ and 3838 ± 4 keV.

^h ΔE_x between $^{17}\text{O}^*(5.73, 5.70) = 34 \pm 2$ keV (1968BI09).

ⁱ $J^\pi = \frac{5}{2}^-$: see, for instance, reaction 46.

32. $^{16}\text{O}(^{11}\text{B}, ^{10}\text{B})^{17}\text{O}$ $Q_m = -7.313$

The excitation of $^{17}\text{O}^*(0, 0.87)$ is reported at $E(^{11}\text{B}) = 113.1$ MeV (1967PO13). See also (1969BR1D).

33. $^{16}\text{O}(^{14}\text{N}, ^{13}\text{N})^{17}\text{O}$ $Q_m = -6.410$

See (1965GA1B).

34. $^{17}\text{N}(\beta^-)^{17}\text{O}$ $Q_m = 8.679$

See ^{17}N .

35. $^{17}\text{O}(\gamma, \alpha)^{13}\text{C}$ $Q_m = -6.357$

See (1964GR08).

36. $^{17}\text{O}(e, e)^{17}\text{O}$

The ^{17}O charge radius, $r_{\text{rms}} = 2.662 \pm 0.026$ (using a distorted wave approximation), 2.700 ± 0.026 fm (using a Born approximation) (1970SI02). See also (1966DE1K, 1970TI1C).

37. $^{17}\text{O}(p, p)^{17}\text{O}$

The angular distribution of elastically scattered protons has been studied at $E_p = 11$ MeV (1967AL06). At $E_p = 29$ MeV, the odd parity states $^{17}\text{O}^*(3.84, 4.55, 5.22, 5.70, 7.75)$ are strongly populated (R. Mendelson, private communication). See also ^{18}F in (1972AJ02) and (1969IC02; theor.).

38. $^{17}\text{O}(^3\text{He}, ^3\text{He})^{17}\text{O}$

The elastic angular distribution has been measured at $E(^3\text{He}) = 17.3$ MeV (1968HA30). See also (1969HA1U, 1969MA1G).

39. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.760$

See ^{17}F .

40. $^{18}\text{O}(\gamma, n)^{17}\text{O}$ $Q_m = -8.046$

See (1963FU06).

41. $^{18}\text{O}(p, d)^{17}\text{O}$ $Q_m = -5.822$

Angular distributions have been measured at $E_p = 17.6$ MeV (1961LE1A, 1963LE03; d_0, d_1, d_2), 18.2 MeV (1967LU05; d_0, d_1, d_2, d_3) and at 27 and 31 MeV (R. Mendelson, private communication; $d_0 \rightarrow d_8$ and $^{17}\text{O}^*(11.08)$). $^{17}\text{O}^*(3.06, 5.38)$ contain most ($> 80\%$) of the $p_{1/2}$ and $p_{3/2}$ hole strength respectively (R. Mendelson). See also (1969IG1A, 1970BO1K, 1970HI15; theor.).

42. $^{18}\text{O}(d, t)^{17}\text{O}$ $Q_m = -1.789$

Angular distributions of the tritons corresponding to $^{17}\text{O}^*(0, 0.87, 3.84, 4.55, 5.08, 5.38)$ have been studied at $E_d = 15$ MeV (1961AR06). See also (1961VL02) and (1963OG1A, 1963RO12, 1968KA1E, 1970BO1K; theor.).

43. $^{18}\text{O}(^3\text{He}, \alpha)^{17}\text{O}$ $Q_m = 12.532$

Angular distributions of alpha particles are reported by (1965WA1D; α_0, α_1) at $E(^3\text{He}) = 2.68$ to 6.47 MeV and by (1969DE06: see Table 17.15) at $E(^3\text{He}) = 16$ MeV. The $T = \frac{3}{2}$ states reported by (1969DE06) are displayed in Table 17.15 [the isospin identification is based on the enhanced excitation and the narrow widths of these states]. The branching ratios for transitions to $^{16}\text{O}^*(0, 6.05)$ for $^{17}\text{O}^*(11.08)$ [the first $T = \frac{3}{2}$ state in ^{17}O] and for the analog state in ^{17}F are displayed in Table 17.9: the ratios of the reduced widths are quite different in the two mirror nuclei (1970MC02). See also (1969BA1Z).

44. $^{19}\text{F}(n, t)^{17}\text{O}$ $Q_m = -7.557$

Table 17.15: $T = \frac{3}{2}$ states of ^{17}O from $^{18}\text{O}(^3\text{He}, \alpha)^{17}\text{O}$ ^{a,b}

E_x (MeV \pm keV)	l_n	J^π	C^2S ^c
11.082 \pm 6	1	$(\frac{1}{2})^-$	0.49
12.471 \pm 5	1	$(\frac{3}{2})^-$	0.27
12.950 \pm 8	0	$\frac{1}{2}^+$	0.096
12.994 \pm 8			
13.640 \pm 5	2	$(\frac{5}{2})^+$	0.39
14.219 \pm 8			
14.282 \pm 12			
15.101 \pm 8			

^a See also Table 17.9.

^b (1969DE06).

^c Calculated assuming $C^2S = 4$ for $^{15}\text{O}^*(6.18)$.

Angular distributions of the t_0 and t_1 groups are reported at $E_n = 14.4$ MeV by (1968AN1F, 1968RE07). See also (1964VA1E).

$$45. \ ^{19}\text{F}(p, ^3\text{He})^{17}\text{O} \quad Q_m = -8.321$$

Angular distributions have been measured at $E_p = 30.5$ MeV (1967CO05, 1968CO1U: to $^{17}\text{O}^*(0, 0.87)$), 40 MeV (1966BR1X, 1966HO1F: to $^{17}\text{O}^*(0, 0.87, 3.06, 3.84)$) and 46 MeV (1969HU1H, 1970HU1J: to $^{17}\text{O}^*(0, 0.87, 3.06)$). The $E_p = 30.5$ MeV distributions have been compared by (1968CO1U) with those from the mirror (p, t) reaction.

$$46. \ ^{19}\text{F}(d, \alpha)^{17}\text{O} \quad Q_m = 10.033$$

$$Q_0 = 10.060 \pm 0.010 \text{ (1967SP09).}$$

Observed alpha groups are displayed in Table 17.14 (1951BU1A, 1952WA1A, 1968BI09). Angular distributions have been measured at many energies in the range $E_d = 0.3$ to 27.5 MeV: see Table 17.16 (1960HU10, 1961CI02, 1962FO02, 1962TA07, 1963HO1F, 1964JA08, 1964MA04, 1965CO07, 1965CO09, 1965ST14, 1966WE04, 1968BI09, 1968PR04, 1968TA02, 1969ME07, 1970BE68). The cross sections for formation of the low-lying states of ^{17}O obey the $2J + 1$ rule, when it is cautiously applied: see (1965CO07, 1965CO09, 1968TA1N). Application of this rule, together with the results of reaction 29, leads to $J^\pi = \frac{5}{2}^-$ for $^{17}\text{O}^*(3.84)$ (1965CO07, 1965CO09).

J for $^{17}\text{O}^*(5.22)$ is found to be $\frac{3}{2}$ (1968BI09). At $E_d = 1.2$ MeV, the γ -transition ($3.06 \rightarrow 0.87$) and ($3.84 \rightarrow 0$) are observed: the absence of ($3.06 \rightarrow 0$) is consistent with $J = \frac{1}{2}$ for $^{17}\text{O}^*(3.06)$ (1957MU1A). See also (1960KR02, 1960RI05, 1969LI22, 1970BI1D), (1969DA1L) and (1963MI1E; theor.).

Table 17.16: $^{19}\text{F}(d, \alpha)^{17}\text{O}$ angular distribution studies

E_d (MeV)	Distribution of alpha groups	Refs.
0.30 – 0.65	$\alpha_0 \rightarrow \alpha_3$	(1969ME07)
0.9 – 4.0	$\alpha_0 \rightarrow \alpha_4$	(1968TA02)
0.95 – 1.25	$\alpha_0 \rightarrow \alpha_3$	(1965ST14)
1.35 – 2.00	$\alpha_0 \rightarrow \alpha_3$	(1969BA2T)
1.5 – 2.5	$\alpha_0 \rightarrow \alpha_3$	(1963HO1F)
2.0, 2.2	$\alpha_0 \rightarrow \alpha_4, \alpha_6, \alpha_7, \alpha_{8+9}$	(1968BI09)
2.0 – 4.6	$\alpha_0 \rightarrow \alpha_2$	(1970BE68)
2.24, 2.33	$\alpha_0 \rightarrow \alpha_3$	(1964JA08)
5.5 – 11.5	$\alpha_0 \rightarrow \alpha_4$	(1966WE04)
9.2	$\alpha_0 \rightarrow \alpha_4$	(1965CO07, 1965CO09)
10	$\alpha_0 \rightarrow \alpha_3$	(1962FO02)
11.1, 11.4	α_0, α_1	(1960HU10)
13	α_0, α_1	(1961CI02)
14.7	$\alpha_0 \rightarrow \alpha_4$	(1962TA07)
20.9	$\alpha_0 \rightarrow \alpha_2$	(1968PR04)
27.5	α_0, α_1	(1964MA04)

47. $^{19}\text{F}(\alpha, ^6\text{Li})^{17}\text{O}$ $Q_m = -12.342$

At $E_\alpha = 42$ MeV, angular distributions are reported for the ^6Li ions corresponding to $^{17}\text{O}^*(0, 0.87)$ (1968MI05).

48. $^{20}\text{Ne}(n, \alpha)^{17}\text{O}$ $Q_m = -0.587$

At $E_n = 14.1$ MeV, angular distributions are reported for the α -particles, corresponding to $^{17}\text{O}^*(0, 0.87)$ (1966MC14). See also (1959BE66, 1966CE03, 1969KA1D).

¹⁷F
(Figs. 8 and 9)

GENERAL:

Shell model: (1957WI1E, 1960TA1C, 1962BH09, 1962TA1E, 1963KU1B, 1965LE1E, 1965MA16, 1966DE18, 1966MA12, 1966SO05, 1967EL03, 1968BI07, 1968EL1A, 1968HO1H, 1968MA2B, 1969EL1B, 1969KU1G, 1969MA38, 1970WA01).

Collective model: (1959FA1C, 1960RA1A, 1962AR1C, 1962MA1K, 1962MA23, 1962MA32, 1964BA1L, 1968BI07, 1968MA2B, 1969MA38).

Electromagnetic transitions: (1959BA1C, 1959FA1C, 1960RA1A, 1964BA1L, 1965GR1H, 1965KA1D, 1966MA12, 1967KA21, 1969KH1C, 1969MA38, 1970EL08, 1970GO1H, 1970SI1C).

Special levels: (1960EV1A, 1961WI1D, 1962BA1C, 1962IN02, 1965EJ1A, 1965LE1E, 1966SO05, 1967BE02, 1969HA1G, 1970EL08, 1970WA01).

Other topics: (1960GO1C, 1964LI1B, 1967NE1D, 1969HE1N, 1969NO1E, 1969SC14, 1969SC1Q, 1969TE1A, 1970BA1M).

Ground state:

$$\mu = 4.7224 \pm 0.0012 \text{ nm (1965SU02, 1966SU01)}.$$

See also (1964ST1B, 1966MA1V, 1967CO1D, 1967SH14, 1968LE1L, 1968PE16, 1968RO1E, 1969FU11).

1. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.760$

The decay is to the ground state of ^{17}O . The spectrum has the allowed shape down to 150 keV (1962VA27). The upper limit for the transition to $^{17}\text{O}^*(0.87)$ is $< 3.4 \times 10^{-4}$ per decay (1969GA05), 4×10^{-4} per decay (1964AR18) [$\log ft > 5.6$]. The half-life of ^{17}F is 66.0 ± 0.2 sec: see Table 17.18 and (1959KI99) $\log ft = 3.36$. See also (1970ST04). See also (1958VA31, 1959VA10, 1966SU01) and (1965GA1D, 1965HA31, 1965MI1B, 1966MI1F, 1967AM1H, 1969LE1D, 1969SU15; theor.).

2. $^{12}\text{C}(^{12}\text{C}, ^7\text{Li})^{17}\text{F}$ $Q_m = -16.859$

See (1962CH01).

Table 17.17: Energy levels of ^{17}F ^a

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{1}{2}$	$\tau_{1/2} = 66.0 \pm 0.2$ sec	β^+	1, 2, 4, 6, 13, 14, 15, 16, 17, 18, 19, 20, 21
0.49533 \pm 0.10	$\frac{1}{2}^+$	$\tau_m = 412 \pm 9$ psec	γ	4, 6, 13, 15, 16, 17, 18, 20
3.105 \pm 7	$\frac{1}{2}^-$	$\Gamma = 19 \pm 1$	p	6, 7, 13, 15, 20
3.86	$\frac{5}{2}^-$	$\tau_m = 6 \pm 1$ fsec	γ, p	6, 7, 13, 14, 15, 20
4.696 \pm 10	$\frac{3}{2}^-$	$\Gamma = 230$	p	7, 13, 20
5.103 \pm 10	$\frac{3}{2}^+$	1530	p	7
5.521 \pm 10	$\frac{3}{2}^-$	69	p	7, 14
5.672 \pm 10	$\frac{7}{2}^-$	40	p	7, 14
5.681 \pm 10	$\frac{1}{2}^+$	< 0.6	p	7
5.817 \pm 10	$\frac{3}{2}^+$	180	p	7
6.036 \pm 10	$\frac{1}{2}^-$	28	p	7
6.556 \pm 10	$\frac{1}{2}^+$	203	p	7
6.699 \pm 10	$\frac{3}{2}^-$	< 3	p	7
6.774 \pm 10	$\frac{3}{2}^+$	4.5	p	7
7.027 \pm 10	$\frac{3}{2}^-$	3.8	p	7
7.356 \pm 10	$\frac{3}{2}^+$	10 ± 2	p, α	7, 8
7.448 \pm 7		≤ 5	p	7
7.454 \pm 7		7 ± 2	p, α	7, 8
7.471 \pm 7		5 ± 2	p	7
7.478 \pm 10	$\frac{3}{2}^+$	795	p	7
7.546 \pm 10	$\frac{7}{2}^-$	28	p	7
7.75 \pm 20	$\frac{1}{2}^+$	179 ± 3	p, α	7, 8
7.95 \pm 15		10 ± 3	p	7
8.01 \pm 20		47 ± 20	p, α	7, 8
8.07 \pm 15	$\frac{5}{2}^+$	100 ± 20	p, α	7, 8
8.2	$\frac{3}{2}^-$	700 ± 240	p, α	7, 8
8.383 \pm 5	$\frac{5}{2}^-$	11 ± 5	p, α	7, 8
8.416 \pm 10	$\frac{7}{2}^+$	42 ± 10	p, α	7, 8
8.75 \pm 30	$\frac{5}{2}^+$	170 ± 30	p, α	7, 8

Table 17.17: Energy levels of ^{17}F ^a (continued)

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
8.95 \pm 20	$\frac{5}{2}^-$	120 \pm 20	p, α	7, 8
9.3 \pm 150	$\frac{3}{2}^+$	210 \pm 40	p, α	7, 8
9.62 \pm 20		310 \pm 70	p	7
9.88 \pm 20		130 \pm 20	p, α	7, 8
9.93 \pm 20		420 \pm 90	p	7
10.04 \pm 20	$\frac{7}{2}$	280 \pm 90	p	7
10.22 \pm 20		250 \pm 80	α	8
10.40 \pm 20		160 \pm 40	p	7
10.49 \pm 20	$\frac{7}{2}^-$	140 \pm 30	p, α	7, 8
(10.70 \pm 20)		140 \pm 30	p, α	7, 8
10.79 \pm 20		120 \pm 40	p, α	7, 8
10.95 \pm 20		190 \pm 50	p, α	7, 8
11.204 \pm 5	$\frac{1}{2}^-; \frac{3}{2}$	< 2	p, α	5, 7, 8
11.43 \pm 20		240 \pm 60	p, α	7, 8
11.57 \pm 40		160 \pm 30	p	7
(11.78 \pm 20)		190 \pm 90	p	7
(11.87 \pm 20)		40 \pm 20	α	8
12.00 \pm 20		120 \pm 40	p, α	7, 8
(12.19 \pm 20)		160 \pm 60	p, α	7, 8
12.25 \pm 20	$\frac{3}{2}^-$	190 \pm 50	p	7
12.35 \pm 20		260 \pm 50	p	7
12.556 \pm 7	$\frac{3}{2}^-; \frac{3}{2}$	≤ 3	p, α	5, 7, 8
(12.8)		≈ 3800	p	7
13.060 \pm 4	$(\frac{3}{2}, \frac{5}{2})^-; \frac{3}{2}$	≤ 4	p, α	5, 7, 8
13.082 \pm 5	$T = \frac{3}{2}$	≤ 5	p, α	7, 8
(13.15)		≈ 400	p	7
(13.7)		≈ 300	p	7
13.779 \pm 5	$T = \frac{3}{2}$	≈ 10	p, α	7, 8
14.310 \pm 5	$T = \frac{3}{2}$	≈ 14	p, α	7, 8
14.5		≈ 700	p	7

Table 17.17: Energy levels of ^{17}F ^a (continued)

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
14.8		≈ 600	p	7
(15.2)		≈ 500	p	7
15.6		≈ 500	p	7
(16.0)		≈ 400	p	7
(16.5)		≈ 500	p	7
(16.7)		≈ 400	p	7
(17.02)		≈ 400	p	7
17.2	$\frac{5}{2}^-$	≈ 450	p	7
17.4		≈ 400	p	7
(17.67)		≈ 400	p	7
(17.9)		≈ 450	p	7
(18.5)		≈ 400	p	7
19.8	$\frac{3}{2}^+$		p	7
20.7		≈ 400	p	7
21.6		≈ 400	p	7
22.4		≈ 400	p	7
25.7	$\frac{3}{2}^-$	broad	p	7
27.3	$\frac{5}{2}^-$		p	7

^a See also Table 17.20.

3. (a) $^{14}\text{N}(^3\text{He}, \text{n})^{16}\text{F}$ $Q_{\text{m}} = -0.969$ $E_{\text{b}} = 15.843$
 (b) $^{14}\text{N}(^3\text{He}, \text{p})^{16}\text{O}$ $Q_{\text{m}} = 15.243$
 (c) $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$ $Q_{\text{m}} = 1.799$
 (d) $^{14}\text{N}(^3\text{He}, \alpha)^{13}\text{N}$ $Q_{\text{m}} = 10.025$
 (e) $^{14}\text{N}(^3\text{He}, 2\alpha)^{11}\text{C}$ $Q_{\text{m}} = 2.297$

The p_0 yield (reaction (b)) does not show the presence of any resonances for $E(^3\text{He}) = 2.5$ to 5.5 MeV (1963GO09). See also (1962BI01, 1965BR42) and ^{16}O . The yields of α_0 , α_1 and $\alpha_2 + \alpha_3$ (reaction (d)) in the range $E(^3\text{He}) = 2.5$ to 8.5 MeV show broad uncorrelated fluctuations, except

Table 17.18: The half-life of ^{17}F

$\tau_{1/2}$ (sec)	Refs. ^a
66.0 ± 0.5	(1958AR15)
66 ± 1	(1949BR27)
66.3 ± 1	(1951LA15)
66.0 ± 0.5	(1954KO54)
66.0 ± 1.8	(1954WO20)
66.0 ± 0.2	(1960JA12)
66.0 ± 0.2	mean

^a See also (1952AJ38, 1954WA1A, 1962VA27).

for a structure at $E(^3\text{He}) = 4.5$ MeV (1970KN01). See also (1964BR1G, 1965BR42, 1966GO1E, 1967BE22, 1967HA20) and ^{13}N in (1970AJ04). For reaction (a) see (1966MA1R) and ^{16}F . For reaction (c) see (1967HA20) and ^{15}O in (1970AJ04). For reaction (e), see (1965BR42).

$$4. \ ^{14}\text{N}(\alpha, \text{n})^{17}\text{F} \quad Q_{\text{m}} = -4.735$$

See (1956DO1C, 1969SC21) and ^{18}F in (1972AJ02).

$$5. \ ^{15}\text{N}(^3\text{He}, \text{n})^{17}\text{F} \quad Q_{\text{m}} = 5.008$$

Neutron groups have been observed corresponding to ^{17}F states at $E_{\text{x}} = 11.195 \pm 0.007$, 12.540 ± 0.010 and 13.059 ± 0.009 MeV, with $\Gamma < 20$, < 25 and < 25 keV, respectively. Angular distributions at $E(^3\text{He}) = 10.36$ and 11.88 MeV lead to $J^{\pi} = \frac{1}{2}^{-}$ for $^{17}\text{F}^*(11.20)$ [$L = 0$], $\frac{3}{2}^{-}$ or $\frac{5}{2}^{-}$ for $^{17}\text{F}^*(12.54)$ and $(\frac{3}{2}^{-}, \frac{5}{2}^{-})$ for $^{17}\text{F}^*(13.06)$. These three states are probably the first three $T = \frac{3}{2}$ states in ^{17}F (1966AD07, 1969AD02). See also (1969BA1Z, 1969GA1P). The branching ratios for transitions to $^{16}\text{O}^*(0, 6.05, 6.13)$ for $^{17}\text{F}^*(11.20)$ and for the analog $T = \frac{3}{2}$ state in ^{17}O are displayed in Table 17.9: the ratios of the reduced widths are quite different in the two mirror nuclei (1970MC02).

$$6. \ ^{16}\text{O}(\text{p}, \gamma)^{17}\text{F} \quad Q_{\text{m}} = 0.601$$

Table 17.19: Recent $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$ yield curves and polarization studies ^a

(a) *Yield curves*

E_p (MeV)	Particles	Refs.
2.5 – 2.8	p_0	(1965GO08)
3.43 – 3.52	p_0	(1964SI04)
3.5 – 5.75	p_0	(1962HA37)
4.25 – 8.6	p_0	(1962SA11, 1962SA1C)
6.7 – 10.6	$\gamma_{6.13}, \gamma_{6.92}$	(1961ST17, 1962ST1A)
6.9 – 15.6	p_0	(1960KO09)
7.2 – 12.9	p_1, p_2, p_3, p_4	(1964DA02)
10.2 – 18	p_0	(1961MA15)
$\approx 11 - 15$	$p_0 \rightarrow p_5$	(1967VA1H, 1968TE1C)
$\approx 11.2, 12.7$	p_0, p_2, p_4	(1967PA17)
12 – 19	p_2, p_{3+4}	(1960KO09)
12.7 – 19.2	p_0	(1964DA07)
13.2 – 14.6	p_0, p_1, p_2, p_3, p_4	(1969SK1B)
13.5 – 19	p_0	(1964KE01)
14.6 – 19.2	p_2, p_{3+4}, p_5	(1964DA07)
16.0 – 30.6	p_0	(1969KA14)
18.5 – 30	$p_0, p_{1+2}, p_{3+4}, p_5$	(1970GU04)
20 – 30	p_0	(1968AP1A)
21 – 40	p_0, p_{1+2}, p_5	(1969BU1J)
23.4 – 46.1	p_{1+2}, p_5	(1970AU1C)
7.2 – 12.8	α_0	(1964DA02)
9.6 – 15	α	(1960FU06)
10.5 – 19	α_0	(1961MA15)
$\approx 11 - 15$	α_0	(1967VA1H, 1968TE1C)
12 – 18	$\sigma_{\text{act.}}$	(1961HI09)
13.20 – 14.08	α_0	(1969SK1B)
15.6 – 18.3	α_1	(1961MA15)
16.2 – 18.3	$\alpha_2 + \alpha_3$	(1961MA15)
21 – 40	α	(1969BU1J)

Table 17.19: Recent $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$ yield curves and polarization studies ^a (continued)

(b) *Polarization measurements*

E_p (MeV)	Particles	Refs.
1.51 – 2.99	p_0	(1967TR08)
2 – 12	p_0	(1965BL02)
3.76 – 4.65	p_0	(1962GO26, 1965BE1K)
4.27 – 4.84	p_0	(1964MA34, 1964MA43, 1965DR01)
4.5	p_0	(1965BO29)
7.9	p_0	(1961RO13)
8.5 – 11.9	p_0	(1962BL15)
8.7, 8.9	p_0	(1959AL08)
10	p_0	(1961RO05)
14.5	p_0	(1965RO22, 1966RO1R)
19.8 – 30.3	p_0	(1969KA14)
20.3	p_0, p_2, p_3, p_4, p_5	(1970BL03)
21.0 – 52.5	p_0	(1966BE1M, 1966BO1Q, 1966BO1T)
24.5 – 39.7	p_0	(1968EL12)

^a See also (1959AJ76).

Non-resonant capture has been studied for $E_p = 0.8$ to 2.1 MeV by (1954WA1A): three γ -rays are observed – γ_1 to the ground state, γ_2 to $^{17}\text{F}^*(0.50)$ and γ_3 from the $(0.50 \rightarrow 0)$ transition. The ratio of γ_2 to γ_1 is ≈ 10 over the above energy region. For $E_p = 2.56$ to 2.76 MeV, a study of the γ_1 and γ_2 yield shows an anomaly in the γ_2 yield [as predicted by (1961CH1C)]: it is deduced from it that $\Gamma_\gamma < 0.03$ eV for the capture transition $(3.10 \rightarrow 0.50)$, and that $\theta^2 = 0.38 \pm 0.08$ and 0.57 ± 0.10 , respectively, for $^{17}\text{F}^*(0, 0.50)$ (1965DO1G). The yield of 3.86 MeV γ -rays has been measured in the vicinity of the $E_p = 3.47$ MeV resonance, previously discussed by (1951LA15, 1951LA1B). The angular distribution of the γ -rays is characteristic of an almost pure dipole transition. The data lead to $J^\pi = \frac{5}{2}^-$ for $^{17}\text{F}^*(3.86)$, $\Gamma < 1.5$ keV, $\Gamma_\gamma = 0.11 \pm 0.02$ eV, $\tau_m = 6 \pm 1$ fsec (1963SE14). See also (1970AB1D).

For $E_p = 140$ to 170 keV, the cross section varies from 0.46 to 2.34×10^{-10} b (1958HE57). Astrophysical considerations are discussed by (1961CH1C, 1962GR1C, 1965DO1G).

The mean lifetime of $^{17}\text{F}^*(0.50)$ is 445 ± 22 psec (1960KA10). See also (1957LE28, 1958HO97). (1963TH1A) have calculated a single-particle lifetime of 320 ± 80 psec for this state.

See also (1959GR1A, 1965ZO1A, 1966ED1A, 1968RI1P) and (1959AJ76).

Table 17.20: Resonances in $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$

E_p (MeV \pm keV)	Γ_{lab} (keV)	Particles out	$^{17}\text{F}^*$ (MeV)	J^π, T	Refs.
2.663 ± 7^a	20 ± 1	p_0	3.105	$\frac{1}{2}^-$	(A, 1962SA1C, 1965BL02, 1965GO08)
3.47	1.63 ± 0.2	p_0	3.86	$\frac{5}{2}^-$	(A, 1962SA1C, 1964SI04, 1965BL02)
4.354 ± 10	240	p_0	4.696	$\frac{2}{2}^-$	(A, 1962HA37, 1962SA11, 1962SA1C, 1965BL02)
4.787 ± 10	1630	p_0	5.103	$\frac{2}{2}^+$	(A, 1962HA37, 1962SA11, 1962SA1C, 1965BL02)
5.231 ± 10	72.5	p_0	5.521	$\frac{2}{2}^-$	(A, 1962HA37, 1962SA11, 1962SA1C, 1965BL02)
5.392 ± 10	42.9	p_0	5.672	$\frac{7}{2}^-$	(A, 1962HA37, 1962SA11, 1962SA1C)
5.402 ± 10	< 0.6	p_0	5.681	$\frac{2}{2}^+$	(A, 1962SA11, 1962SA1C)
5.546 ± 10	191	p_0	5.817	$\frac{3}{2}^+$	(A, 1962HA37, 1962SA11, 1962SA1C, 1965BL02)
5.779 ± 10	30	p_0	6.036	$\frac{2}{2}^-$	(A, 1962SA11, 1962SA1C)
6.332 ± 10	216	p_0	6.556	$\frac{1}{2}^+$	(1962SA11, 1962SA1C)
6.484 ± 10	< 3	p_0	6.699	$\frac{2}{2}^-$	(A, 1962SA11, 1962SA1C)
6.564 ± 10	4.8	p_0	6.774	$\frac{3}{2}^+$	(A, 1962SA11, 1962SA1C, 1965BL02)
6.833 ± 10	4.0	$p_0, \gamma_{6.13}$	7.027	$\frac{2}{2}^-$	(1961ST17, 1962SA11, 1962SA1C, 1962ST1A, 1965BL02)
7.183 ± 10	10 ± 2	p_0, p_2, α_0	7.356	$\frac{2}{2}^+$	(1960KO09, 1962SA11, 1962SA1C, 1964DA02, 1965BL02)
7.280 ± 7	≤ 5	p_0	7.448		(1962SA11, 1962SA1C, 1964DA02)
7.287 ± 7	7 ± 2	p_0, p_1, p_2, α	7.454		(1962SA11, 1962SA1C, 1964DA02)
7.305 ± 7	5 ± 2	p_0, p_2	7.471		(1962SA11, 1962SA1C, 1964DA02)
7.313 ± 10	845	p_0	7.478	$\frac{2}{2}^+$	(1962SA11, 1962SA1C, 1964DA02)
7.385 ± 10	30	$p_0, p_2, \gamma_{6.13}$	7.546	$\frac{7}{2}^-$	(1961ST17, 1962SA11, 1962SA1C, 1962ST1A, 1964DA02)
7.60 ± 20	190 ± 3	p_0, p_1, α_0	7.75	$\frac{1}{2}^+$	(1962SA11, 1962SA1C, 1964DA02)
7.81 ± 15	10 ± 3	p_2	7.95		(1964DA02)
7.88 ± 20	50 ± 20	$p_0, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$	8.01		(1961ST17, 1962SA11, 1962SA1C, 1962ST1A, 1964DA02)
7.94 ± 15	110 ± 20	p_0, p_1, α_0	8.07	$\frac{5}{2}^+$	(1962SA11, 1962SA1C, 1964DA02)
8.1	750 ± 250	$(p_0), p_1, \alpha_0$	8.2	$\frac{2}{2}^-$	(1964DA02)
8.275 ± 5	12 ± 5	$p_0, p_1, p_2, p_3, \alpha_0$	8.383	$\frac{1}{2}^-$	(1962SA11, 1962SA1C, 1964DA02)
8.310 ± 10	45 ± 10	$p_0, p_1, p_2, p_3, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$	8.416	$\frac{2}{2}^+$	(1961ST17, 1962SA11, 1962SA1C, 1962ST1A, 1964DA02, 1965BL02)
8.66 ± 30^b	180 ± 30	p_2, p_3, p_4, α_0	8.75	$\frac{2}{2}^+$	(1963HA04, 1964DA02)

Table 17.20: Resonances in $^{16}\text{O}(\text{p}, \text{p})^{16}\text{O}$ and $^{16}\text{O}(\text{p}, \alpha)^{13}\text{N}$ (continued)

E_p (MeV \pm keV)	Γ_{lab} (keV)	Particles out	$^{17}\text{F}^*$ (MeV)	J^π, T	Refs.
8.88 ± 20	130 ± 20	$\text{p}_0 \rightarrow \text{p}_4, \alpha_0, \gamma_{6.13}, \gamma_{6.92}$	8.95	$\frac{5}{2}^-$	(1961ST17, 1962ST1A, 1963HA04, 1964DA02)
9.2 ± 150^b	220 ± 40	$\text{p}_0 \rightarrow \text{p}_4, \alpha_0, \gamma_{6.13}, \gamma_{6.92}$	9.3	$\frac{3}{2}^+$	(1961ST17, 1962ST1A, 1963HA04, 1964DA02)
9.59 ± 20	330 ± 75	$\text{p}_0, \text{p}_1, \text{p}_4$	9.62		(1964DA02)
9.87 ± 20	140 ± 20	$\text{p}_0, \text{p}_2, \alpha_0$	9.88		(1964DA02)
9.92 ± 20	450 ± 100	p_3	9.93		(1964DA02)
10.04 ± 20	300 ± 100	p_0, p_1	10.04	$\frac{7}{2}$	(1963HA04, 1964DA02)
10.23 ± 20	270 ± 85	α_0	10.22		(1964DA02)
10.42 ± 20	170 ± 40	p_1, p_3	10.40		(1964DA02)
10.52 ± 20	150 ± 30	$\text{p}_0, \text{p}_2, \alpha_0$	10.49	$\frac{7}{2}^-$	(1963HA04, 1964DA02, 1965TO03)
(10.74 \pm 20)	150 ± 30	p_1, α_0	(10.70)		(1964DA02)
10.83 ± 20	130 ± 40	$\text{p}_0, \text{p}_2, (\text{p}_3), (\alpha_0)$	10.79		(1963HA04, 1964DA02)
11.00 ± 20	200 ± 50	$(\text{p}_2), \text{p}_3, (\alpha_0)$	10.95		(1963HA04, 1964DA02)
11.276 ± 8^c	< 2	$\text{p}_0 \rightarrow \text{p}_5, \alpha_0$	11.204	$\frac{1}{2}^-; \frac{3}{2}$	(1967PA17, 1967VA1H, 1968TE1C) ^d
11.52 ± 20	260 ± 60	p_2, α_0	11.43		(1963HA04, 1964DA02)
11.67 ± 40^b	170 ± 30	p_0, p_3	11.57		(1963HA04, 1964DA02)
(11.89 \pm 20)	200 ± 100	p_2	(11.78)		(1964DA02)
(11.98 \pm 20)	40 ± 20	α_0	(11.87)		(1964DA02)
12.12 ± 20	130 ± 40	p_2, α_0	12.00		(1963HA04, 1964DA02)
(12.32 \pm 20)	170 ± 60	$(\text{p}_2), \alpha_0$	(12.19)		(1964DA02)
12.39 ± 20	200 ± 50	p_0, p_2	12.25	$\frac{3}{2}^-$	(1963HA04, 1964DA02, 1965TO03)
12.49 ± 20	280 ± 50	p_1, p_4	12.35		(1963HA04, 1964DA02)
12.714 ± 8	≤ 3	$\text{p}_0, \text{p}_1, \text{p}_2, \text{p}_4, \text{p}_5, \alpha_0$	12.556	$\frac{3}{2}^-; \frac{3}{2}$	(1963HA04, 1964DA02, 1967PA17, 1967VA1H, 1968TE1C)
($\approx 13.$)	≈ 4000	p_0	(12.8)		(1960KO09)
13.250 ± 4	≤ 4	$\text{p}_0 \rightarrow \text{p}_5, \alpha_0$	13.060	$T = \frac{3}{2}$	(1963HA04, 1964DA02, 1967VA1H, 1968TE1C, 1969SK1B) ^e
13.273 ± 5	≤ 5	$\text{p}_0 \rightarrow \text{p}_4, \alpha_0$	13.082	$T = \frac{3}{2}$	(1969SK1B)
(13.35)	≈ 450	p_0	(13.15)		(1964DA07)
(13.9)	≈ 350	p_0	(13.7)		(1964DA07, 1964KE01)

Table 17.20: Resonances in $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$ (continued)

E_p (MeV \pm keV)	Γ_{lab} (keV)	Particles out	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$	Refs.
14.015 \pm 5	\approx 11	$p_0 \rightarrow p_4, \alpha_0$	13.779	$T = \frac{3}{2}$	(1969SK1B)
14.579 \pm 7	\approx 15	$p_0 \rightarrow p_5, \alpha_0$	14.310	$T = \frac{3}{2}$	(1961HI09, 1967VA1H, 1968TE1C, 1969SK1B) ^e
14.8	\approx 700	p_0, p_{1+2}	14.5		(1960KO09, 1964DA07, 1964KE01)
15.1	\approx 600	p_{1+2}	14.8		(1964DA07)
(15.5)	\approx 550	$p_0, p_{1+2}, p_{3+4}, p_5$	(15.2)		(1964DA07)
15.9	\approx 550	p_0, p_{1+2}	15.6		(1964DA07, 1964KE01)
(16.4)	\approx 400	$p_0, p_{1+2}, p_{3+4}, p_5$	(16.0)		(1964DA07)
(16.9)	\approx 550	p_0	(16.5)		(1964DA07)
(17.1)	\approx 450	p_{1+2}	(16.7)		(1964DA07)
(17.45)	\approx 450	p_0, p_{3+4}	(17.02)		(1964DA07)
17.6	\approx 500	p_0, p_{3+4}	17.2	$\frac{5}{2}^-$	(1964DA07, 1969KA14)
17.9	\approx 450	p_0, p_{3+4}, p_5	17.4		(1964DA07, 1964KE01)
(18.15)	\approx 450	p_0, p_{1+2}	(17.67)		(1964DA07)
(18.4)	\approx 500	p_{3+4}, p_5	(17.9)		(1964DA07)
(19.0)	\approx 400	p_0, p_{1+2}, p_5	(18.5)		(1964DA07) ^f
20.4		p_0	19.8	$\frac{3}{2}$	(1969KA14)
21.4	\approx 400	p_0	20.7		(1968AP1A, 1969KA14)
22.3	\approx 400	p_0	21.6		(1968AP1A, 1969KA14)
23.2	\approx 400	p_0, p_5	22.4		(1968AP1A, 1969KA14)
26.7	broad	p_0	25.7	$\frac{5}{2}^-$	(1968AP1A, 1969KA14)
28.4		p_0	27.3	$\frac{5}{2}^-$	(1969KA14)

A: (1951LA15, 1951LA1B, 1953EP1A, 1954SE1A, 1954SE1B): see (1959AJ76).

^a A search for fine structure near this resonance was unsuccessful: (1968SE1D).

^b This may correspond to more than one state: see (1964DA02).

^c $\Gamma_{p_0} = 40_{-20}^{+10}$ eV (1967PA17).

^d And C.A. Barnes, private communication.

^e And P.D. Parker, private communication.

^f Structures in the yields of $p_0, p_{1+2}, p_{3+4}, p_5$ for $19 < E_p < 28$ MeV are also reported by (1970GU04).

7. (a) $^{16}\text{O}(p, p)^{16}\text{O}$		$E_b = 0.601$
(b) $^{16}\text{O}(p, p\alpha)^{12}\text{C}$	$Q_m = -7.161$	
(c) $^{16}\text{O}(p, pn)^{15}\text{O}$	$Q_m = -15.668$	

Yield curves for elastic protons, protons scattered to $^{16}\text{O}^*(6.05, 6.13, 6.92, 7.12, 8.87)$ and for γ -rays from $^{16}\text{O}^*(6.13, 6.92)$ have been studied at many energies up to $E_p = 46$ MeV: see Table 17.19 for a summary of the measurements and Table 17.20 for a display of the observed resonances (1960KO09, 1961MA15, 1961ST17, 1962HA37, 1962SA11, 1962SA1C, 1962ST1A, 1964DA02, 1964DA07, 1964KE01, 1964SI04, 1965GO08, 1967PA17, 1967VA1H, 1968AP1A, 1968SE1D, 1968TE1C, 1969BU1J, 1969KA14, 1969SK1B, 1970AU1C, 1970GU04). See also (1965TO03, 1967CA1Q, 1970AB1D). See also (1966DA1B), (1968CA30) and (1963CR10, 1963OP02, 1964VO1A, 1965HA28, 1966GA1L, 1969BO2A, 1969HA2C, 1969SO1A, 1969WA11, 1970SH14; theor.).

Total reaction cross sections have been measured at $E_p = 24.5$ to 45.9 MeV by (1967CA1R, 1969MC1A) and at 142 MeV by (1961TA06). At $E_p = 13.1$ MeV, cross sections for production of the p_{1+2} and p_{3+4} groups, as well as for the α_0 and α_1 groups (reaction (b)) have been measured by (1967CH41). At $E_p = 150$ MeV, cross sections for formation of $^{16}\text{O}^*(6.05, 6.13, 6.92)$ are reported by (1962RO25). See also (1962AL16). For a summary of angular distribution measurements, see Table 16.27 in ^{16}O .

Polarization measurements have been carried out for $E_p = 1.5$ to 52.5 MeV: see Table 17.19 (1959AL08, 1961RO05, 1962BL15, 1962GO26, 1964MA34, 1964MA43, 1965BE1K, 1965BL02, 1965BO29, 1965DR01, 1965RO22, 1966BE1M, 1966BO1Q, 1966BO1T, 1966RO1R, 1967TR08, 1968EL12, 1969KA14, 1970BL03). Below $E_p = 5$ MeV two very broad resonances give rise to polarizations which change slowly with energy except near the 2.66 and 3.47 MeV resonances (1965BL02). (1968EL12) report a drastic variation in the polarization angular distribution between $E_p = 21$ and 27 MeV, suggesting strong resonances in that region. See also (1967FA06), (1966DA1B, 1966RO1B) and (1959KE1A, 1960KA1E, 1963DU1B, 1963HO1D, 1965BA1M, 1965HA28, 1967LE13, 1968CH35, 1968RE03, 1969BO2A, 1969VA18; theor.).

For reaction (b) see also (1962AL16, 1962RO25); for reaction (c), see (1965VA23).

For discussions of spallation measurements, see (1967BE2D) and (1960GR1A, 1961TA10, 1962AL16, 1963AL1G, 1963VA1C, 1965ZO1A, 1967AU1A, 1967EP1A, 1967GR1K, 1967LI1B, 1968JA1M, 1968PA1J, 1968YI02, 1969AB1A, 1969DU10, 1969EP1B, 1970KO25). Astrophysical implications are considered by (1967BE2D) and (1966RE1D, 1967AU1A, 1967GR1K, 1967LI1B, 1968YI02). See also reaction 57 in ^{16}O .

8. $^{16}\text{O}(p, \alpha)^{13}\text{N}$	$Q_m = -5.218$	$E_b = 0.601$
--	----------------	---------------

Excitation functions of various α -groups and activation functions have been measured for $E_p = 7.2$ to 40 MeV: see Table 17.19 (1960FU06, 1961HI09, 1961MA15, 1964DA02, 1967VA1H, 1968TE1C, 1969BU1J, 1969SK1B). Observed resonances are displayed in Table 17.20 (1961HI09,

1964DA02, 1967VA1H, 1968TE1C, 1969SK1B). In addition to the sharp resonances reported by (1961HI09, 1967VA1H, 1968TE1C, 1969SK1B) a number of broad structures are observed above $E_p \approx 15$ MeV (1961HI09, 1961MA15, 1970GU04). Cross sections for formation of $^{13}\text{N}^*(0, 2.37, 3.51 + 3.55)$ have been measured at $E_p = 13.1$ MeV (1967CH41). See also ^{13}N in (1970AJ04).

$$9. \text{}^{16}\text{O}(\text{p}, \text{n})\text{}^{16}\text{F} \qquad Q_m = -16.212 \qquad E_b = 0.601$$

See ^{16}F .

$$10. \text{}^{16}\text{O}(\text{p}, \text{d})\text{}^{15}\text{O} \qquad Q_m = -13.443 \qquad E_b = 0.601$$

Excitation functions have been measured for $E_p = 21$ to 33 MeV (1969BU1J). At $E_p = 30.3$ MeV, polarization measurements have been made for the deuterons corresponding to $^{15}\text{O}^*(0, 6.18)$ (1967CH15). See also ^{15}O in (1970AJ04).

$$11. \text{}^{16}\text{O}(\text{p}, \text{t})\text{}^{14}\text{O} \qquad Q_m = -20.406 \qquad E_b = 0.601$$

Excitation functions have been measured for $E_p = 30$ to 33 MeV (1969BU1J). Polarization measurements for the tritons corresponding to $^{14}\text{O}(0)$ and the ^3He particles corresponding to the analog state, $^{14}\text{N}^*(2.31)$, have been carried out at $E_p = 49.5$ MeV: the analyzing powers of the two reactions are closely the same (1970NE1B). Asymmetry measurements are also reported at $E_p = 43.8$ MeV for the tritons corresponding to $^{14}\text{O}^*(0, 7.78, 9.74)$ and to $^{14}\text{O}^*(6.59)$. The latter disagree with DWBA predictions (1970HA23). See also ^{14}O in (1970AJ04).

$$12. \text{}^{16}\text{O}(\text{p}, \text{}^3\text{He})\text{}^{14}\text{N} \qquad Q_m = -15.243 \qquad E_b = 0.601$$

Polarization measurements have been reported at $E_p = 43.8$ MeV for the ^3He particles corresponding to $^{14}\text{N}^*(7.03)$ (1970HA23). See also reaction 11 (1970NE1B) and ^{14}N in (1970AJ04).

$$13. \text{}^{16}\text{O}(\text{d}, \text{n})\text{}^{17}\text{F} \qquad Q_m = -1.624$$

$$E_{\text{thresh.}} = 1.8292 \pm 0.0006; Q_0 = -1.6246 \pm 0.0005 \text{ (1960BO21);}$$

$$E_{\text{thresh.}} = 1.830 \pm 0.004 \text{ (1955MA85);}$$

$$E_{\text{thresh.}} = 1.832 \pm 0.003 \text{ (1961CH14).}$$

Table 17.21: $^{16}\text{O}(\text{d}, \text{n})^{17}\text{F}$ angular distribution studies ^a

E_{d} (MeV)	Angular distributions of groups	Refs.
1.93 – 2.39	n_0	(1961DI06)
2.20 – 3.30	n_0	(1968DI06)
2.70 – 3.30	n_1	(1968DI06)
3.00 – 5.50	n_0, n_1	(1970LO01)
4.75 – 6.0	n_0, n_1	(1970DA14)
5.02	n_0, n_1	(1961YA05)
7.73, 11, 12	n_0, n_1	(1969OL02)
8.0, 9.3	n_0, n_1	(1969TH04)

^a See also (1959AJ76).

Slow-neutron thresholds have been observed corresponding to the ground state and to the first excited state. For the latter, $Q_1 = -2.125 \pm 0.004$ MeV (1955MA85), -2.125 ± 0.003 MeV (1961CH14), leading to $E_x = 499 \pm 3$ keV (1955MA85), 498 ± 2 keV (1961CH14). See also (1965MA1K). (1966AL10) report $E_x = 495.33 \pm 0.10$ keV from the measurement of the corresponding E_γ . The lifetime for $^{17}\text{F}^*(0.50)$ is 406.8 ± 8.7 psec (1964BE15). See also (1967BI05, 1967WA1C, 1968SC1E).

Neutron groups have been observed corresponding to $^{17}\text{F}^*(0, 0.50)$, to states at 3.03 ± 0.06 and 4.74 ± 0.07 MeV ($\Gamma < 90$ keV) (1957GR1A), and to $^{17}\text{F}^*(3.86)$ (1969TH04). See also (1969OL02). Angular distributions have been measured for $E_{\text{d}} \leq 12$ MeV for the n_0 and n_1 groups: see Table 17.21 (1961DI06, 1961YA05, 1968DI06, 1969OL02, 1969TH04, 1970DA14, 1970LO01) and (1959AJ76). $l_p = 2$ and 0 , respectively for the transitions to $^{17}\text{F}^*(0, 0.5)$, in agreement with their assignments $J^\pi = \frac{5}{2}^+$ and $\frac{1}{2}^+$, respectively. Spectroscopic factors have been extracted and compared with those from $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: see (1969OL02, 1969TH04, 1970DA14) and (1970LO01). See also (1960MA21, 1962VA27, 1963KN04, 1963SA1C, 1965SU02, 1966AL12, 1966SU01, 1969OV01), (1963TA1A, 1963TR1A, 1969HA1V, 1970TE1B; theor.) and ^{18}F in (1972AJ02).

14. (a) $^{16}\text{O}(\text{}^3\text{He}, \text{d})^{17}\text{F}$ $Q_{\text{m}} = -4.893$
 (b) $^{16}\text{O}(\text{}^3\text{He}, \text{dp})^{16}\text{O}$ $Q_{\text{m}} = -5.494$
 $Q_0 = -4.90 \pm 0.09$ (1960WE04).

The angular distribution of the d_0 group has been measured at $E(^3\text{He}) = 9.16$ MeV: it is consistent with $l = 2$ (1959HI73). See also (1966AG1B, 1966EC1B). Reaction (b) appears to

proceed in part via excited states of ^{17}F at $E_x = 3.86$ and 5.6 MeV (1967HO14).

$$15. \ ^{16}\text{O}(\alpha, t)^{17}\text{F} \quad Q_m = -19.214$$

At $E_\alpha = 46$ MeV, angular distributions of triton groups to the first four states of ^{17}F have been measured (1968PR1C). See also (1963TR1A; theor.).

$$16. \ ^{16}\text{O}(^{11}\text{B}, ^{10}\text{Be})^{17}\text{F} \quad Q_m = -10.628$$

At $E(^{11}\text{B}) = 113$ MeV, the ^{10}Be spectrum shows the strong excitation of $^{17}\text{F}^*(0 + 0.5)$, in addition to some unidentified weaker groups (1967PO13). See also (1967VO1A, 1969BR1D).

$$17. \ ^{17}\text{O}(\text{p}, \text{n})^{17}\text{F} \quad Q_m = -3.542$$

Angular distributions of the n_0 and n_1 groups have been obtained for $E_p = 6.95$ to 13.50 MeV (n_0) and 6.95 to 12.45 MeV (n_1). There appears to be collective enhancement in the $L = 2$ transition to $^{17}\text{F}^*(0.5)$. A large spin-flip term in the effective two-body force is necessary to account for the strength of the ground state transition (1969AN06). See also (1966AN1A) and (1969SC1H).

$$18. \ ^{17}\text{O}(^3\text{He}, t)^{17}\text{F} \quad Q_m = -2.778$$

At $E(^3\text{He}) = 17.3$ MeV, angular distributions have been obtained for the tritons corresponding to $^{17}\text{F}^*(0, 0.50)$. The data have been analyzed using DWBA and a two-body interaction between the incident and target nucleons. An exact coupled-channel-equation calculation was also made for the ground state transition (1968HA30, 1969HA1U, 1969MA1G). See also (1966EC1B).

$$19. \ ^{19}\text{F}(\gamma, 2\text{n})^{17}\text{F} \quad Q_m = -19.581$$

See (1959OC07).

$$20. \ ^{19}\text{F}(\text{p}, \text{t})^{17}\text{F} \quad Q_m = -11.099$$

Table 17.22: Energy levels of ^{17}Ne

E_x in ^{17}Ne (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ (msec)	Decay	Reactions
0	$\frac{1}{2}^-; \frac{3}{2}$	108.0 ± 2.7	β^+	1, 2
1.35 ± 70				2
1.84 ± 70				2
2.77 ± 70				2
5.28 ± 70				2

Angular distributions of the triton groups corresponding to $^{17}\text{F}^*(0, 0.50)$ have been measured at $E_p = 16.4 - 18.0$ MeV (1963KA13), 17.5 MeV (1969HAZD), 22.8 MeV (1963HO24), 30.3 MeV (1967DI1C, 1968CO1U) and 46 MeV (1969HU1H, 1970HU1J). Angular distributions are also reported for the t_2 group (1963HO24, 1969HU1H) and for the t_3 and t_4 groups (1963HO24: partial distribution). See also (1962CO17, 1965RE1A, 1967AN1B, 1969AN1L).

$$21. \ ^{20}\text{Ne}(p, \alpha)^{17}\text{F} \quad Q_m = -4.129$$

At $E_p = 43$ MeV, the angular distribution of the α_0 group is reported by (1968FA1G). Some indication of a state at $E_x = 5.28$ MeV is reported by (1965PE1E).

^{17}Ne
(Fig. 9)

GENERAL:

Theory: (1964WI1B, 1965MA16, 1966MA12).

Reviews: (1960BA1F, 1960GO1B, 1961BA1C, 1962GO31, 1964GO1G, 1966GO1B, 1966GO1K, 1966MC1C, 1968CE01, 1969GA1P).

Mass of ^{17}Ne : $M - A$, determined from the Q -value of the $^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne}$ reaction, is 16.479 ± 0.050 MeV (1970ME11). See also (1967ES02). Then $^{17}\text{Ne} - ^{17}\text{F} = 14.53$ MeV and E_b for p, ^3He and α are, respectively, 1.50, 6.46 and 9.05 MeV. [E_b for an α -particle is calculated using the mass of (1970ME11) for ^{13}O .] The mass of ^{17}Ne predicted from mass formulae are $M - A = 16.508 \pm 0.023$ MeV (1970ME11), 16.517 ± 0.026 MeV (1968CE01), 16.63 MeV (1966KE16). See also (1964WI1B, 1965JA1C).




Table 17.23: Decay of ^{17}Ne [†]

Decay to $^{17}\text{F}^*$ (MeV)	J^π	Decay ^a (%)		$\log ft$ ^b (1969HAZE)	Decay to $^{16}\text{O}^*$ (MeV)	J^π	Decay (%) (1969HAZE)
		(1967ES02)	(1969HAZE)				
0	$\frac{5}{2}^+$	0.2 ^c		7.3 ^c			
0.50	$\frac{1}{2}^+$	0.6 ^c		6.8 ^c			
3.11	$\frac{1}{2}^-$	< 1.0	0.49	6.50 ± 0.07	0	0^+	100
4.32 ^e		2.0 ± 0.5					
4.70	$\frac{3}{2}^-$	19.2 ± 1.0	16.5	4.59 ± 0.02	0	0^+	100
5.52	$\frac{3}{2}^-$	56.2 ± 1.8	54.9	3.86 ± 0.01	0	0^+	100
6.04	$\frac{1}{2}^-$	12.4 ± 0.7	10.8	4.44 ± 0.02	0	0^+	100
6.43 ^e		0.9 ± 0.3	0.36	5.82 ± 0.10	0	0^+	100
6.70	$\frac{3}{2}^-$	0.4 ± 0.2					
7.03	$\frac{3}{2}^-$	≤ 0.5					
7.36 \rightarrow 7.48		≤ 0.7					
7.75	$\frac{1}{2}^+$		0.18	5.70 ± 0.20	0	0^+	
8.01	$\frac{5}{2}^+$ }	4.5 ± 0.4	6.9	4.00 ± 0.01	0	0^+	
8.07					6.05	0^+	
8.39	}	0.8 ± 0.1	6.6	3.85 ± 0.03	0	0^+	
8.42					6.05	0^+	
8.85			1.9	4.20 ± 0.03	0	0^+	
					6.05	0^+	
11.20 ^d	$\frac{1}{2}^-; T = \frac{3}{2}$	0.04 ± 0.01	0.61	3.35 ± 0.10	0	0^+	12 ± 4
					6.05	0^+	26 ± 8
					6.92	2^+	28 ± 13
					7.12	1^-	34 ± 14

[†] Note added in proof: For latest results, see (1971HA05).

^a See also (1964MC16, 1965HA20, 1965MA16).

^b Based on $\tau_{1/2} = 109 \pm 3$ msec.

^c From mirror $\log ft$ values (1967ES02).

^d No decay was observed to $^{16}\text{O}^*(6.13) [3^-]$ nor to $^{13}\text{N} (\text{O})$ [by α -emission]: upper limit is 10% of the observed proton decays (1969HAZE).

^e Proposed on the basis of the work of (1967ES02).

The half-life of ^{17}Ne is 102 ± 7 msec (1964MC16), 105 ± 5 msec (1967ES02), 107 ± 5 msec (1967FI10), 115 ± 5 msec (1969HAZE): the mean value is 108.0 ± 2.7 msec. See also (1964FL03, 1964FL05, 1965HA20). The decay is primarily to proton unstable states of ^{17}F at $E_x = 4.69, 5.52$ and 6.04 MeV, with $J^\pi = \frac{3}{2}^-, \frac{3}{2}^-$ and $\frac{1}{2}^-$, respectively: see Table 17.23 (1967ES02, 1969HAZE). The super-allowed decay to the analog state [$^{17}\text{F}^*(11.20)$] has $\log ft = 3.35 \pm 0.10$ (1969HAZE). The character of the decay leads to $J^\pi = \frac{1}{2}^-$ for $^{17}\text{Ne}(0)$. See also (1963BA63, 1963VL1B, 1964DA13, 1965HA31, 1965MA16, 1965MI1B, 1966HA22).

$$2. \ ^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne} \quad Q_m = -26.187$$

At $E(^3\text{He}) = 62$ MeV, ^6He groups are observed to five states of ^{17}Ne : see Table 17.22 (R. Mendelson, J.M. Loiseaux, G. Wozniak and J. Cerny, private communication).

^{17}Na
(Not illustrated)

^{17}Na has not been observed: its mass excess is predicted to be 35.61 MeV by (1966KE16). It is then unbound with respect to breakup into $^{16}\text{Ne} + p$ by 3.2 MeV and with respect to breakup into $^{14}\text{O} + 3p$ by 5.8 MeV.

References

(Closed 30 November 1970)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1948KN24 N. Knable, E.O. Lawrence, C.E. Leith, B.J. Moyer and R.L. Thornton, Phys. Rev. 74 (1948) 1217, F9
- 1949BR27 H. Brown and V. Perez-Mendez, Phys. Rev. 75 (1949) 1286, D5
- 1951BU1A Burrows, Powell and Rotblat, Proc. Roy. Soc. A209 (1951) 478
- 1951LA15 R.A. Laubenstein, M.J.W. Laubenstein, L.J. Koester and R.C. Mobley, Phys. Rev. 84 (1951) 12
- 1951LA1B Laubenstein and Laubenstein, Phys. Rev. 84 (1951) 18
- 1952AJ38 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 24 (1952) 321
- 1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
- 1952WA1A Watson and Buechner, Phys. Rev. 88 (1952) 1324
- 1953EP1A Eppling et al, Phys. Rev. 91 (1953) 438A
- 1953JO1A Jones and Wilkinson, Proc. Phys. Soc. A66 (1953) 1176
- 1953TH14 J. Thirion and V.L. Telegdi, Phys. Rev. 92 (1953) 1253
- 1954KO54 M.I. Korsunskii, Usp. Fiz. Nauk 52 (1954) 3
- 1954SE1A Sempert, Schneider and Martin, Helv. Phys. Acta 27 (1954) 313
- 1954SE1B Sempert, Thesis, Zurich ETH (1954)
- 1954WA1A Warren, Laurie, James and Erdman, Can. J. Phys. 32 (1954) 563
- 1954WO20 C. Wong, Phys. Rev. 95 (1954) 765
- 1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77
- 1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46
- 1955SE1A Seitz and Huber, Helv. Phys. Acta 28 (1955) 227
- 1956BE98 R.L. Becker and H.H. Barschall, Phys. Rev. 102 (1956) 1384
- 1956BO61 T.W. Bonner, A.A. Kraus, Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956DO1C Doyle and Robbins, Phys. Rev. 101 (1956) 1056
- 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. A69 (1956) 28

1956RU1A Rusbridge, Proc. Phys. Soc. A69 (1956) 830
 1956SC1C Schiffer, Kraus and Risser, Phys. Rev. 105 (1956) 1811
 1957BO04 N.A. Bostrom, E.L. Hudspeth and I.L. Morgan, Phys. Rev. 105 (1957) 1545
 1957BR82 C.P. Browne, Phys. Rev. 108 (1957) 1007
 1957GR1A Grismore and Parkinson, Rev. Sci. Instrum. 28 (1957) 245
 1957LE28 P. Lehmann, A. Leveque, T. Grjebine, J.-L. Picou and R. Barloutaud, Compt. Rend. 245 (1957) 2259
 1957MU1A Muller, Thesis, Univ. of Strasbourg (1957)
 1957WA46 R.B. Walton, J.D. Clement and F. Boreli, Phys. Rev. 107 (1957) 1065
 1957WI1E Winterberg, Z. Naturforsch. A12 (1957) 271
 1958AR15 S.E. Arnell, J. Dubois and O. Almen, Nucl. Phys. 6 (1958) 196
 1958HE57 R.E. Hester, R.E. Pixley and W.A.S. Lamb, Phys. Rev. 111 (1958) 1604
 1958HO97 R.E. Holland, F.J. Lynch and S.S. Hanna, Phys. Rev. 112 (1958) 903
 1958RI1A Richter and Ivash, Phys. Rev. 111 (1958) 265
 1958RO1A Roys and Shure, Nucl. Sci. Eng. 4 (1958) 536
 1958VA31 S.S. Vasilev and L.Y. Shavtvalov, Izv. Akad. Nauk SSSR Ser. Fiz. 22 (1958) 788; Columbia Tech. Transl. 22 (1959) 782
 1958WE31 J.L. Weil and K.W. Jones, Phys. Rev. 112 (1958) 1975
 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
 1959AL08 M.A. Al-Jeboori, M.S. Bokhari, B. Hird and A. Strzalkowski, Proc. Phys. Soc. A74 (1959) 705
 1959AU1A Auerbach, Reilly and Francis, Bull. Amer. Phys. Soc. 4 (1959) 272
 1959BA1C Barton, Brink and Delves, Nucl. Phys. 14 (1959) 256
 1959BE1B Berko, Groseclose, Stetson and Walker, Bull. Amer. Phys. Soc. 4 (1959) 257
 1959BE66 R.J. Bell, T.W. Bonner and F. Gabbard, Nucl. Phys. 14 (1959) 270
 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
 1959FA1C Fallieros and Ferrell, Phys. Rev. 116 (1959) 660
 1959FE1B Feingold, Phys. Rev. 114 (1959) 540
 1959FI30 G.E. Fischer and V.K. Fischer, Phys. Rev. 114 (1959) 533
 1959GR1A Griffiths, Congress Int. Phys. Nucl., Paris, 1958 (1959) 447
 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
 1959HA29 E.W. Hamburger, Thesis, Univ. of Pittsburgh (1959)
 1959HI73 S. Hinds and R. Middleton, Proc. Phys. Soc. A74 (1959) 775

- 1959KE1A Kerman, McManus and Thaler, *Ann. Phys.* 8 (1959) 551
- 1959KH1A Khanna and Green, *Bull. Amer. Phys. Soc.* 4 (1959) 387
- 1959KI99 O.C. Kistner and B.M. Rustad, *Phys. Rev.* 114 (1959) 1329
- 1959LI47 C.S. Littlejohn, *Phys. Rev.* 114 (1959) 250
- 1959LO59 G. Lopez, F. Alba, M. Mazari and M.E. Ortiz, *Rev. Mex. Fis.* 8 (1959) 17
- 1959OC07 J. O'Connell, P. Dyal and J. Goldemberg, *Phys. Rev.* 116 (1959) 173
- 1959SA11 J. Sawicki, *Nucl. Phys.* 13 (1959) 350
- 1959VA10 S.S. Vasilev and L.Y. Shavtvalov, *Zh. Eksp. Teor. Fiz.* 36 (1959) 317; *Sov. Phys. JETP* 9 (1959) 218
- 1959YU1A Yu, Teng, Chou and Lee, *Sci. Sin.* 8 (1959) 935
- 1960AK1A Akiyama, *Prog. Theor. Phys.* 23 (1960) 903
- 1960AL35 N.V. Alekseev, K.I. Zherebtsova, V.F. Litvin and Y.A. Nemilov, *Zh. Eksp. Teor. Fiz.* 39 (1960) 1508; *Sov. Phys. JETP* 12 (1961) 1049
- 1960BA1F Baz, *Sov. J. At. Energy* 6 (1960) 422
- 1960BE1B Belyaev, Zakhar'ev and Neudachin, *Atomn. Energ. (USSR)* 9 (1960) 298, *Sov. J. At. Energy* 9 (1961) 833
- 1960BI1B Biedenharn and Satchler, *Helv. Phys. Acta Suppl.* 6 (1960) 372
- 1960BO21 R.O. Bondelid, J.W. Butler and C.A. Kennedy, *Phys. Rev.* 120 (1960) 889
- 1960BU1C Bullock and Moore, *Phys. Rev.* 119 (1960) 721
- 1960CO1A Corman and Fowler, *Bull. Amer. Phys. Soc.* 5 (1960) 34
- 1960DE19 J.A. DeJuren and R.W. Stooksberry, *Phys. Rev.* 120 (1960) 901
- 1960EV1A Everling, *Z. Naturforsch.* A15 (1960) 84
- 1960FO14 J.L. Fowler, E.G. Corman and E.C. Campbell, *Proc. Int. Conf. on Nucl. Struct., Kingston, Canada* (1960) 474
- 1960FU06 M. Furukawa, Y. Ishizaki, Y. Nakano, T. Nozaki, Y. Saji and S. Tanaka, *J. Phys. Soc. Jpn.* 15 (1960) 2167
- 1960GO1B Goldansky, *Nucl. Phys.* 19 (1960) 482
- 1960GO1C Gove, *Proc. Int. Conf. on Nucl. Struct., Kingston, Canada* (1960) 438
- 1960GO20 S. Gorodetzky, G. Sutter, F. Scheibling, P. Chevallier and R. Armbruster, *J. Phys. Rad.* 21 (1960) 360
- 1960GR1A Gradstajn, *J. Phys. Rad.* 21 (1960) 761
- 1960HU10 C. Hu, *J. Phys. Soc. Jpn.* 15 (1960) 1741
- 1960JA12 J. Janecke, *Z. Naturforsch.* A15 (1960) 593

- 1960JA13 N. Jarmie and M.G. Silbert, Phys. Rev. 120 (1960) 914
- 1960KA10 J.V. Kane, R.E. Pixley, R.B. Schwartz and A. Schwarzschild, Phys. Rev. 120 (1960) 162
- 1960KA1E Karadzhev and Man'ko, Zh. Eksp. Teor. Fiz. 39 (1960) 416; Sov. Phys. JETP 12 (1961) 294
- 1960KO09 S. Kobayashi, J. Phys. Soc. Jpn. 15 (1960) 1164
- 1960KR02 T.H. Kruse, R.D. Bent and L.J. Lidofsky, Phys. Rev. 119 (1960) 289
- 1960MA21 B.E.F. Macefield and J.H. Towle, Proc. Phys. Soc. 76 (1960) 56
- 1960MI1B Mikhlin and Stavinskii, Atomn. Energ. (USSR) 8 (1960) 141; Sov. J. At. Energy 8 (1961) 127; Reactor Sci. Tech. 16 (1962) 119
- 1960MU07 J. Muto, F. de S. Barros and A.A. Jaffe, Proc. Phys. Soc. 75 (1960) 929
- 1960NA1A Nagarajan and Banerjee, Nucl. Phys. 17 (1960) 341
- 1960PE02 F.G.J. Perey and P. Lorrain, Bull. Amer. Phys. Soc. 5 (1960) 18, E6
- 1960PE25 J.M. Peterson, A. Bratenahl and J.P. Stoering, Phys. Rev. 120 (1960) 521
- 1960RA1A Raz, Phys. Rev. 120 (1960) 169
- 1960RI05 J. Rickards, Rev. Mex. Fis. 9 (1960) 35
- 1960SH05 S.M. Shafroth, J. Phys. Rad. 21 (1960) 353
- 1960SH1A Sheline and Wildermuth, Nucl. Phys. 21 (1960) 196
- 1960SM02 R.V. Smith, L.F. Chase, Jr., R.H. Abramson, J.B. Reagan and M. Walt, Bull. Amer. Phys. Soc. 5 (1960) 19, E8
- 1960ST02 R.L. Steele and J.R. Risser, Bull. Amer. Phys. Soc. 5 (1960) 108, M2
- 1960TA1C Talmi and Unna, Ann. Rev. Nucl. Sci. 10 (1960) 353
- 1960TS02 K. Tsukada and T. Fuse, J. Phys. Soc. Jpn. 15 (1960) 1994
- 1960WA07 R.B. Walton, N.F. Wikner, J.L. Wood and J.R. Beyster, Bull. Amer. Phys. Soc. 5 (1960) 288, WA5
- 1960WE04 H.E. Wegner and W.S. Hall, Phys. Rev. 119 (1960) 1654
- 1960ZE03 Ya.B. Zeldovich, Zh. Eksp. Teor. Fiz. 38 (1960) 1123; Sov. Phys. JETP 11 (1960) 812
- 1961AR06 J.C. Armstrong and K.S. Quisenberry, Phys. Rev. 122 (1961) 150
- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. 77 (1961) 853
- 1961BA1C Baz, Goldanskii and Zeldovich, Sov. Phys. Usp. 3 (1961) 729
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165

- 1961BR1A Brill', Vlasov, Kalinin and Sokolov, Dokl. Akad. Nauk SSSR 136 (1961) 55; Sov. Phys. Dokl. 6 (1961) 24
- 1961BU16 B. Buck and P.E. Hodgson, Phil. Mag. 6 (1961) 1371
- 1961CH14 R. Chiba, Phys. Rev. 123 (1961) 1316
- 1961CH1C Christy and Duck, Nucl. Phys. 24 (1961) 89
- 1961CI02 N. Cindro, M. Cerineo and A. Strzalkowski, Nucl. Phys. 24 (1961) 107
- 1961DI06 O. Dietzsch, Y. Hama, E.W. Hamburger and F.C. Zawislak, Nucl. Phys. 27 (1961) 103
- 1961FO07 D.B. Fossan, R.L. Walter, W.E. Wilson and H.H. Barschall, Phys. Rev. 123 (1961) 209
- 1961GO1H Gofman, Nemets and Tokarevskii, Sov. J. At. Energy 7 (1961) 946
- 1961HA19 E.W. Hamburger, Phys. Rev. 123 (1961) 619
- 1961HI01 S. Hinds, R. Middleton, A.E. Litherland and D.J. Pullen, Phys. Rev. Lett. 6 (1961) 113
- 1961HI09 H.A. Hill, E.L. Haase and D.B. Knudsen, Phys. Rev. 123 (1961) 1301
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, Rev. Mex. Fis. 10 (1961) 247
- 1961JO24 R.G. Johnson, L.F. Chase, Jr. and F.J. Vaughn, Proc. Rutherford Jub. Int. Conf. (1961) 591
- 1961KE01 E.L. Keller, Phys. Rev. 121 (1961) 820
- 1961LA1A Lane, Langsdorf, Monahan and Elwyn, Ann. Phys. 12 (1961) 135
- 1961LE1A Legg, Unpublished Thesis, Princeton Univ. (1961)
- 1961LO1C Lopez, Rev. Mex. Fis. 10 (1961) 283
- 1961MA15 D.R. Maxson, Phys. Rev. 123 (1961) 1304
- 1961NE1B Neudachin and Orlin, Zh. Eksp. Teor. Fiz. 41 (1961) 874
- 1961PE28 G.J. Perlow, W.J. Ramler, A.F. Stehney and J.L. Yntema, Phys. Rev. 122 (1961) 899
- 1961PU1B Pullen, Wilkinson and Whitehead, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 565
- 1961RO05 L. Rosen, J.E. Brolley, Jr. and L. Stewart, Phys. Rev. 121 (1961) 1423
- 1961RO13 L. Rosen, J.E. Brolley, Jr., M.L. Gursky and L. Stewart, Phys. Rev. 124 (1961) 199
- 1961SE02 K.W. Seemann and W.E. Moore, Bull. Amer. Phys. Soc. 6 (1961) 237, DA15
- 1961ST17 R.F. Sturgeon, Jr., J.A. Becker and J.D. Fox, Bull. Amer. Phys. Soc. 6 (1961) 506, E6
- 1961TA06 A.E. Taylor and E. Wood, Nucl. Phys. 25 (1961) 642
- 1961TA10 Tamers and Delibrias, Compt. Rend. 253 (1961) 1202

- 1961VL02 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.I. Chuev, *Izv. Akad. Nauk SSSR Ser. Fiz.* 25 (1961) 115; *Columbia Tech. Transl.* 25 (1961) 111
- 1961WI1A Wilkinson, Wollan and Koehler, *Ann. Rev. Nucl. Sci.* 11 (1961) 303
- 1961WI1D Wildermuth and Tang, *Bull. Amer. Phys. Soc.* 6 (1961) 78
- 1961YA02 H. Yamaguchi, *J. Phys. Soc. Jpn.* 16 (1961) 583
- 1961YA05 B. Yaramis, *Phys. Rev.* 124 (1961) 836
- 1962AL16 G. Albouy, J.P. Cohen, M. Gusakow, N. Poffe, H. Sergolle and L. Valentin, *Phys. Lett.* 2 (1962) 306
- 1962AR1C Arnurius, Buck and Satchler, *Electromag. Lifetime and Properties of Nucl. States*, N.A.S.-N.R.C. Pub. 974 (1962) 28
- 1962BA1C Barker and Treacy, *Nucl. Phys.* 38 (1962) 33
- 1962BE1C Beghian, Sugimoto, Wachter and Weber, *Bull. Amer. Phys. Soc.* 7 (1962) 883
- 1962BE1D Bessis, Lefebvre-Brion and Moser, *Phys. Rev.* 128 (1962) 213
- 1962BH09 K.H. Bhatt, *Nucl. Phys.* 39 (1962) 375
- 1962BI01 O.M. Bilaniuk and W.P. Alford, *Bull. Amer. Phys. Soc.* 7 (1962) 71, V1
- 1962BL13 J.M. Blair and R.K. Hobbie, *Phys. Rev.* 128 (1962) 2282
- 1962BL15 R.A. Blue, R.I. Brown, L.W. Morrow, W.G. Weitkamp and L.C. McIntyre, *Bull. Amer. Phys. Soc.* 7 (1962) 434, BA6
- 1962CH01 C. Chasman and D.A. Bromley, *Bull. Amer. Phys. Soc.* 7 (1962) 36, JA4
- 1962CO17 C.F. Coleman, P.E. Cavanagh, B.W. Ridley and J.F. Turner, *Conf. Low Energy Nucl. Phys. Harwell, AERE-R-4131*, 8, 3.3 (1962)
- 1962DA1B Davidson and Chi, *Electromag. Lifetimes and Properties of Nucl. States*; N.A.S.-N.R.C. Publ. 974 (1962) 237
- 1962DE13 J.A. DeJuren, R.W. Stooksberry and M. Wallis, *Phys. Rev.* 127 (1962) 1229
- 1962EL01 A.J. Elwyn and R.O. Lane, *Nucl. Phys.* 31 (1962) 78
- 1962FO02 J.M. Fowler, J.B. Reynolds, J.J. Wesolowski and R.J. Wilson, *Bull. Amer. Phys. Soc.* 7 (1962) 287, FB16
- 1962GA15 N.H. Gale, J.B. Garg and J.M. Calvert, *Nucl. Phys.* 38 (1962) 222
- 1962GO26 S. Gorodetzky, J. Ullman, G. Bergdolt and A. Gallmann, *Nucl. Phys.* 38 (1962) 177
- 1962GO31 V.I. Goldanskii, *Dokl. Akad. Nauk SSSR* 146 (1962) 1309; *Sov. Phys. Dokl.* 7 (1963) 922
- 1962GR1C Griffiths, Lal and Robertson, *Electromag. Lifetime and Properties of Nucl. States*, N.A.S.-N.R.C. Pub. 974 (1962) 205
- 1962HA37 R.W. Harris, G.C. Phillips and C.M. Jones, *Nucl. Phys.* 38 (1962) 259

- 1962HA40 B.G. Harvey, J. Cerny, R.H. Pehl and E. Rivet, Nucl. Phys. 39 (1962) 160
- 1962HU07 W. Hunzinger and P. Huber, Helv. Phys. Acta 35 (1962) 351
- 1962IN02 D.R. Inglis, Nucl. Phys. 30 (1962) 1
- 1962KA37 J. Kantele and D.G. Garoner, Nucl. Phys. 35 (1962) 353
- 1962LO09 G.N. Lovchikova, Atomn. Energ. (USSR) 13 (1962) 60
- 1962MA05 J.P. Martin and H.S. Zucker, Bull. Amer. Phys. Soc. 7 (1962) 72, V5
- 1962MA1K S. Matthies, V.G. Neudachin and Y.F. Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1962) 1060
- 1962MA23 S. Matthies, V.G. Neudachin and Yu.F. Smirnov, Zh. Eksp. Teor. Fiz. 42 (1962) 592; Sov. Phys. JETP 15 (1962) 411
- 1962MA25 S. Mayo and J.E. Testoni, Nucl. Phys. 36 (1962) 615
- 1962MA32 S. Matthies, V.G. Neudachin and Y.F. Smirnov, Nuclear Phys. 38 (1962) 63
- 1962MO1B Mottelson, Nucl. Spectroscopy; Ed., Racah (1962)
- 1962OT01 P.S. Otstavnov and V.I. Popov, Zh. Eksp. Teor. Fiz. 43 (1962) 385; Sov. Phys. JETP 16 (1963) 276
- 1962RO25 D.J. Rowe, A.B. Clegg, G.L. Salmon and P.S. Fisher, Proc. Phys. Soc. 80 (1962) 1205
- 1962SA11 S.R. Salisbury and H.T. Richards, Phys. Rev. 126 (1962) 2147
- 1962SA1C Salisbury, Hardie, Opplinger and Dangle, Phys. Rev. 126 (1962) 2143
- 1962SE1F Seeman and Moore, KAPL 2214 (1962)
- 1962ST1A Sturgeon, M. S. Thesis, F. S. Univ. (1962)
- 1962TA07 K. Takamatsu, J. Phys. Soc. Jpn. 17 (1962) 896; Erratum. J. Phys. Soc. Jpn. 18 (1963) 1234
- 1962TA1B Talmi and Unna, Nucl. Phys. 30 (1962) 280
- 1962TA1E Talmi, Nucl. Spectroscopy; Ed., Racah (1962)
- 1962VA27 S.S. Vasilev and L.Y. Shavtvalov, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1962) 1495; Columbia Tech. Transl. 26 (1963) 1521
- 1962VO01 V.V. Volkov, L. Pomorski, J. Tys and G.N. Flerov, Zh. Eksp. Teor. Fiz. 42 (1962) 635; Sov. Phys. JETP 15 (1962) 442
- 1963AL04 R. Almanza and M.E.O. de Lopez, Bull. Amer. Phys. Soc. 8 (1963) 125, U12
- 1963AL1G Albouy et al, J. Phys. Rad. 24 (1963) 67
- 1963AM1A Amsel, Thesis, Univ. of Paris (1963)
- 1963BA08 S. Bashkin, V.P. Hart and W.A. Seale, Phys. Rev. 129 (1963) 1750
- 1963BA46 R.W. Bauer, J.D. Anderson and L.J. Christensen, Nucl. Phys. 47 (1963) 241

- 1963BA63 R. Barton, R. McPherson, R.E. Bell, W.R. Frisken, W.T. Link and R.B. Moore, *Can. J. Phys.* 41 (1963) 2007
- 1963BE36 N. Bessis, H. Lefebvre-Brion and C.M. Moser, *Rev. Mod. Phys.* 35 (1963) 548
- 1963BO1E Bormann et al, *Z. Phys.* 174 (1963) 1
- 1963CH02 B.E. Chi and J.P. Davidson, *Phys. Rev.* 131 (1963) 366
- 1963CH1C Chatterjee, *Nucl. Phys.* 49 (1963) 686
- 1963CO12 B.L. Cohen, *Phys. Rev.* 130 (1963) 227
- 1963CR10 D.R. Croley, Jr. and L.D. Oppliger, *Bull. Amer. Phys. Soc.* 8 (1963) 538, S9
- 1963DA12 E.A. Davis, T.W. Bonner, D.W. Worley, Jr. and R. Bass, *Nucl. Phys.* 48 (1963) 169
- 1963DE27 G. Deconninck, M. De Vroey, J.P. Meulders and J. Simonet, *Nucl. Phys.* 49 (1963) 424
- 1963DO1B Douglas, Sala, Gomes and Polga, in Padua (1963) 558A
- 1963DU1B Duke, *Phys. Rev.* 129 (1963) 681
- 1963FU06 H. Fuchs, *Z. Phys.* 171 (1963) 416
- 1963GI04 J. Gilat, G.D. O'Kelley and E. Eichler, *Bull. Amer. Phys. Soc.* 8 (1963) 320, GA11
- 1963GL1C Glendenning, *Ann. Rev. Nucl. Sci.* 13 (1963) 191
- 1963GL1D Gleyvod, Durisch, Huber and Baumgartner, *Helv. Phys. Acta* 36 (1963) 287
- 1963GO09 S. Gorodetzky, G. Bassompierre, C.St. Pierre, A. Gallmann and P. Wagner, *Nucl. Phys.* 43 (1963) 92
- 1963HA04 G. Hardie, R.L. Dangle and L.D. Oppliger, *Phys. Rev.* 129 (1963) 353
- 1963HA05 M. Harvey, *Phys. Lett.* 3 (1963) 209
- 1963HA1G Haerberli, *Fast Neutron Phys.*; Eds., Marion and Fowler (1963) 1379
- 1963HO1D Hooton and Ashcroft, *Proc. Phys. Soc.* 81 (1963) 193
- 1963HO1F Hood, Scott and McEllistrem, *Bull. Amer. Phys. Soc.* 8 (1963) 548
- 1963HO24 H.D. Holmgren and C.B. Fulmer, *Phys. Rev.* 132 (1963) 2644
- 1963JO1D Joanou and Fenech, *J. Nucl. Energy* 17 (1963) 425
- 1963KA13 E. Kashy, J.C. Legg and N.R. Roberson, *Bull. Amer. Phys. Soc.* 8 (1963) 317, G6
- 1963KA27 V. Kaminsky, Y. Orlov, N. Marchenko and Y. Orevkov, *Nucl. Phys.* 48 (1963) 417
- 1963KE1A Kelley, *Bull. Amer. Phys. Soc.* 8 (1963) 605
- 1963KN04 G.F. Knoll, J.S. King and W.C. Parkinson, *Phys. Rev.* 131 (1963) 331
- 1963KU1B Kunz, *Can. J. Phys.* 41 (1963) 2187
- 1963KU1F Kukhtevich et al, *Sov. Prog. in Neutron Phys.*, Consult. Bureau (1963) 205
- 1963LE03 J.C. Legg, *Phys. Rev.* 129 (1963) 272

1963LE1D Levkovskii, Zh. Eksp. Teor. Fiz. 45 (1963) 305; Sov. Phys. JETP 18 (1964) 213
1963LE1E Levin, Private Communication (1963)
1963LO03 J. Lowe and C.L. McClelland, Phys. Rev. 132 (1963) 367
1963LU10 H.F. Lutz, J.B. Mason and M.D. Karvelis, Nucl. Phys. 47 (1963) 521
1963MI1E Mitra and Rook, in Padua (1963) 211A
1963MO1B Morrison, Gale, Hussain and Murray, 3rd Conf. on Reactions between Complex Nuclei (1963) 168
1963OG1A Ogloblin, Nucl. Phys. 47 (1963) 408
1963OP02 L.D. Oppliger and D.R. Croley, Jr., Bull. Amer. Phys. Soc. 8 (1963) 538, S10
1963PA01 G.M. Padawer and R.E. Benenson, Bull. Amer. Phys. Soc. 8 (1963) 25, F5
1963PA03 S.P. Pandya, Nucl. Phys. 43 (1963) 636
1963PI03 G. Pisent and A.M. Saruis, Nuovo Cim. 28 (1963) 600
1963RO12 J.R. Rook, Nucl. Phys. 41 (1963) 343
1963SA1B Satchler, in Padua (1963) 80
1963SA1C Sala et al, in Padua (1963) 563A
1963SE14 B.E. Segel, P.P. Singh, R.G. Allas and S.S. Hanna, Phys. Rev. Lett. 10 (1963) 345
1963SH1C Shure, Nucl. Sci. Eng. 16 (1963) 247
1963SM05 W.R. Smith and E.V. Ivash, Phys. Rev. 131 (1963) 304
1963SP02 R.H. Spear, J.D. Larson and J.D. Pearson, Nucl. Phys. 41 (1963) 353
1963TA1A Tanifuji, Nucl. Phys. 40 (1963) 357
1963TH1A Thomas and Tanner, Nucl. Phys. 44 (1963) 647
1963TR1A Truhlik, Nucl. Phys. 48 (1963) 329
1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163
1963VL1B Vlasov, Atomn. Energ. (USSR) 14 (1963) 45
1963WA1E Walecka and Willey, Nucl. Phys. 40 (1963) 282
1963YA03 K. Yagi, Y. Nakajima, K. Katori, Y. Awaya and M. Fujioka, Nucl. Phys. 41 (1963) 584
1963YA1C Yamazaki, Knodo and Yamabe, J. Phys. Soc. Jpn. 18 (1963) 620
1964AL11 T.K. Alexander, C. Broude and A.E. Litherland, Nucl. Phys. 53 (1964) 593
1964AM1A Amsel, Ann. Phys. 9 (1964) 297
1964AR18 S.E. Arnell and E. Wernborn, Ark. Fys. 25 (1963) 389
1964BA1L Barker, Nucl. Phys. 59 (1964) 513

- 1964BE15 J.A. Becker and D.H. Wilkinson, Phys. Rev. 134 (1964) B1200
- 1964BR1G Brill, Chuev and Ogloblin, in Paris (1964) 981
- 1964BR1H Brown, in Paris (1964) 129
- 1964DA02 R.L. Dangle, L.D. Oppliger and G. Hardie, Phys. Rev. 133 (1964) B647
- 1964DA07 W.W. Daehnick, Phys. Rev. 135 (1964) B1168
- 1964DA13 J.M. D'Auria and I.L. Preiss, Phys. Lett. 10 (1964) 300
- 1964DU1B Dunford and Francis, Bull. Amer. Phys. Soc. 9 (1964) 168
- 1964EL1B El-Nadi and Riad, Nucl. Phys. 50 (1964) 33
- 1964FL03 G.N. Flerov, V.A. Karnaukhov, G.M. Ter-Akopyan, L.A. Petrov and V.G. Subbotin, Nucl. Phys. 60 (1964) 129
- 1964FL05 G.N. Flerov, V.A. Karnaukhov, G.M. Ter-Akopyan, L.A. Petrov and V.G. Subbotin, Zh. Eksp. Teor. Fiz. 47 (1964) 419; Sov. Phys. JETP 20 (1965) 278
- 1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49
- 1964GO1G Goldsanskii, Dokl. Akad. Nauk SSSR 157 (1964) 321; Sov. Phys. Dokl. 9 (1965) 566
- 1964GR08 L.H. Greenberg, J.P. Roalsvig and R.N.H. Haslam, Can. J. Phys. 42 (1964) 731
- 1964HE18 I. Heertje, I. Delvenne, W. Nagel and A.H.W. Aten, Jr., Physica 30 (1964) 1762
- 1964JA08 M.F. Jahns, J.B. Nelson and E.M. Bernstein, Nucl. Phys. 59 (1964) 314
- 1964JU05 E.T. Journey and H.T. Motz, Bull. Amer. Phys. Soc. 9 (1964) 176, III5
- 1964KE01 R.A. Kenefick, W.S. Gray and J.J. Kraushaar, Bull. Amer. Phys. Soc. 9 (1964) 68, GC8
- 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 379
- 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 411
- 1964MA04 S. Mayo, J. Testoni and O.M. Bilaniuk, Phys. Rev. 133 (1964) B350
- 1964MA34 C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi and G. Zannoni, Phys. Lett. 9 (1964) 159
- 1964MA43 C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi and G. Zannoni, Nuovo Cim. 34 (1964) 860
- 1964MC16 R. McPherson, J.C. Hardy and R.E. Bell, Phys. Lett. 11 (1964) 65
- 1964NI1A Nigam, Phys. Rev. 133 (1964) B1381
- 1964RI1A Ripka, 3A (II)/C174, Paris (1964)
- 1964SC09 R.B. Schwartz, H.D. Holmgren, L.M. Cameron and A.R. Knudson, Phys. Rev. 134 (1964) B577

- 1964SC12 U. Schmidt-Rohr, R. Stock and P. Turek, Nucl. Phys. 53 (1964) 77
- 1964SI04 E.A. Silverstein, L.D. Oppliger and R.A. Blue, Bull. Amer. Phys. Soc. 9 (1964) 68, GC7
- 1964SI06 M.G. Silbert and J.C. Hopkins, Phys. Rev. 134 (1964) B16
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Edition, Suppl. 2 Vol. 1 (1964)
- 1964TE1C Testoni, Mayo and Hodgson, Nucl. Phys. 50 (1964) 479
- 1964VA1E Valkovic and Tomas, Congress Int. Phys. Nucl., Paris (1964)
- 1964VO1A Volkin, Bull. Amer. Phys., Soc. 9 (1964) 439
- 1964WA1B Wachter, Phys. Rev. 135 (1964) B1180
- 1964WI1B Wilkinson, Phys. Lett. 12 (1964) 348
- 1965AL14 T.K. Alexander and K.W. Allen, Can. J. Phys. 43 (1965) 1563
- 1965BA1M Barrett, Hill and Hodgson, Nucl. Phys. 62 (1965) 133
- 1965BA32 B.K. Barnes, T.A. Belote and J.R. Risser, Phys. Rev. 140 (1965) B616
- 1965BE1K Bergdolt, Ann. Phys. 10 (1965) 857
- 1965BL02 R.A. Blue and W. Haeberli, Phys. Rev. 137 (1965) B284
- 1965BO29 I. Boca, M. Cenja, E. Iliescu and N. Martalogu, Rev. Roum. Phys. 10 (1965) 415
- 1965BR42 O.D. Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965BU1G Butler, Hewitt and May, Phys. Rev. Lett. 15 (1965) 1032
- 1965CA05 R.R. Carlson and R.L. McGrath, Phys. Rev. Lett. 15 (1965) 173
- 1965CI01 S. Cierjacks, G. Markus, W. Michaelis and W. Ponitz, Phys. Rev. 137 (1965) B345
- 1965CO07 S.W. Cospers, B.T. Lucas and O.E. Johnson, Phys. Rev. 138 (1965) B51
- 1965CO09 S.W. Cospers and O.E. Johnson, Phys. Rev. 138 (1965) B610
- 1965DE1N de Forest, Bull. Amer. Phys. Soc. 10 (1965) 1125
- 1965DO13 I. Dostrovsky, R. Davis, Jr., A.M. Poskanzer and P.L. Reeder, Phys. Rev. 139 (1965) B1513
- 1965DO14 R.E. Donaldson, L. Passell, W. Bartolini and D. Groves, Phys. Rev. 138 (1965) B1116
- 1965DO1G Dominigo, Nucl. Phys. 61 (1965) 39
- 1965DR01 L. Drigo, C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi, G. Torielli and G. Zannoni, Nuovo Cim. 37 (1965) 1766
- 1965EJ1A Ejiri, INS Rept. 80 (1965)
- 1965FU1E Furst-Rauch and Munzer, Acta Phys. Aust. 20 (1965) 300

1965GA1A Garcia, Bolta, Lopez and Senent, An. Real Soc. Espan. Fiz. Quim 61 (1965) 341
1965GA1B Gaedke, Toth and Williams, Phys. Rev. 140 (1965) B296
1965GA1D Gaponov, Yad. Fiz. 2 (1965) 1002
1965GA1F Garside and MacDonald, Phys. Rev. 138 (1965) B582
1965GI1B Giraud, Nucl. Phys. 71 (1965) 373
1965GO08 V. Gomes, R.A. Douglas, T. Polga and O. Sala, Nucl. Phys. 68 (1965) 417
1965GR1H Grechukhin, Nucl. Phys. 62 (1965) 273
1965HA05 V.P. Hart, E. Norbeck and R.R. Carlson, Phys. Rev. 137 (1965) B17
1965HA20 J.C. Hardy and R.E. Bell, Can. J. Phys. 43 (1965) 1671
1965HA28 R.M. Haybron and H. McManus, Phys. Rev. 140 (1965) B638
1965HA31 J.C. Hardy and B. Margolis, Phys. Lett. 15 (1965) 276
1965HE1B Henkel, Bull. Amer. Phys. Soc. 10 (1965) 601
1965HU1D Hull and Shakin, Phys. Lett. 19 (1965) 506
1965HU1E Huby and Mines, Revs. Mod. Phys. 37 (1965) 406
1965IC1A Ichimura and Yazaki, Nucl. Phys. 63 (1965) 401
1965JA1C Janecke, Nucl. Phys. 61 (1965) 383
1965KA1D Kamerdzhiev, Yad. Fiz. 2 (1965) 415
1965LE1E Lee, Acta Phys. Sin. (China) 21 (1965) 1370
1965LO02 N. Longequeue, H. Beaumevieille, J.-P. Longequeue, M. Sandon and E. Ligeon, Compt. Rend. 260 (1965) 517
1965MA16 B. Margolis and N. De Takacsy, Phys. Lett. 15 (1965) 329
1965MA1K Marion, Nucl. Phys. 68 (1965) 463
1965MA59 N.A. Mansour, H.R. Saad, Z.A. Saleh, E.M. Sayed, I.I. Zaloubovsky and V.I. Gontchar, Nucl. Phys. 65 (1965) 433
1965MC10 R.E. McDonald, D.B. Fossan, L.F. Chase, Jr. and J.A. Becker, Phys. Rev. 140 (1965) B1198
1965MI1B Migdal and Khodel, Yad. Fiz. 8 (1965) 28
1965MO16 R. Moreh and T. Daniels, Nucl. Phys. 74 (1965) 403
1965NE1C Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 25
1965PE1E Pessoa and Dangle, Bull. Amer. Phys. Soc. 10 (1965) 463
1965RA15 L.A. Rayburn and E.O. Wollan, Nucl. Phys. 61 (1965) 381
1965RE1A Reynolds, Maxwell and Hintz, Bull. Amer. Phys. Soc. 10 (1965) 439
1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96

- 1965SE01 K.K. Seth, G. Walter, P.D. Miller and J.A. Biggerstaff, Bull. Amer. Phys. Soc. 10 (1965) 10, AC8
- 1965SI1D Sih-Tjuen, Acta Phys. Sin. (China) 21 (1965) 208
- 1965SL1B Slaggie, Bull. Amer. Phys. Soc. 10 (1965) 497
- 1965SM1A Smith, Nucl. Phys. 72 (1965) 593
- 1965ST14 D.M. Stanojevic, S.D. Cirilov and M.M. Ninkovic, Nucl. Phys. 73 (1965) 657
- 1965SU02 K. Sugimoto, A. Mizobuchi, K. Nakai and K. Matuda, Phys. Lett. 18 (1965) 38
- 1965TO03 W. Toutenhoofd, Aust. J. Phys. 18 (1965) 7
- 1965VA23 L. Valentin, Nucl. Phys. 62 (1965) 81
- 1965WA1D Waggoner and Jaffe, Nucl. Phys. 69 (1965) 305
- 1965ZA1B Zamick, Phys. Lett. 19 (1965) 580
- 1965ZO1A Zobel, Maienschein and Scroggs, ORNL 3506 (1965)
- 1966AD07 E.G. Adelberger and C.A. Barnes, Phys. Lett. 23 (1966) 474
- 1966AG1A Agee and Rosen, LA-3538-MS (1966)
- 1966AG1B Aguilar, de la Rubia, Sanchez and Martinez, An. Fis. y Quim. 62 (1966) 279
- 1966AL09 J.L. Alty, L.L. Green, R. Huby, G.D. Jones, J.R. Mines and J.F. Sharpey-Schafer, Phys. Lett. 20 (1966) 664
- 1966AL10 D.E. Alburger, A. Gallmann, J.B. Nelson, J.T. Sample and E.K. Warburton, Phys. Rev. 148 (1966) 1050
- 1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743
- 1966AN1A Anderson, F. S. U. Isobaris Spin Conf. (1966) 530
- 1966AR10 A. Armigliato, F. Brandolini, F. Pellegrini and E. Crescenti, Nuovo Cim. B45 (1966) 92
- 1966BE1E Beaumevieille et al, J. Phys. (Paris) 27 (1966) C1-150
- 1966BE1M Bercaw, Boschitz and Vincent, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 334
- 1966BO1Q Boschitz, Bull. Amer. Phys. Soc. 11 (1966) 355
- 1966BO1R Bohigas, J. Phys. (Paris) 27 (1966) C1-39
- 1966BO1T Boschitz, Chabre, Conzett and Slobodrian, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 331
- 1966BR04 G.E. Brown and A.M. Green, Nucl. Phys. 75 (1966) 401
- 1966BR12 R. Bruning, F.W. Busser, F. Niebergall and J. Christiansen, Phys. Lett. 21 (1966) 435
- 1966BR1E Bruning et al, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 146

1966BR1Q Brown, Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch (1969) 513
1966BR1R G.E. Brown and A.M. Green, Nucl. Phys. 85 (1966) 87
1966BR1X Brown et al, Bull. Amer. Phys. Soc. 11 (1966) 316
1966BU09 F.W. Busser, F. Niebergall, G. Sohngen and J. Christiansen, Nucl. Phys. 88 (1966) 593
1966CE03 M. Cevolani, S. Petralia and G. Di Caporiacco, Nucl. Phys. 79 (1966) 379
1966DA1B Darden, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 433
1966DE18 N. De Takacsy, Phys. Lett. 23 (1966) 260
1966DE1K De Forest and Walecka, Adv. Phys. 15 (1966) 1, 491
1966EC1B Eccles, Lutz and Rohn, Bull. Amer. Phys. Soc. 11 (1966) 735
1966ED1A J.A. Edgington and B. Rose, Nucl. Phys. 89 (1966) 523
1966GA09 A. Gallman, P. Fintz and P.E. Hodgson, Nucl. Phys. 82 (1966) 161
1966GA1L Garside and MacDonald, Antwerp 1965 Neutron Conf. (1966) 526
1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197
1966GO1B Goldanskii, Ann. Rev. Nucl. Sci. 16 (1966) 1
1966GO1E Gorodetzky et al, Nucl. Instrum. Meth. 42 (1966) 269
1966GO1K Goldanskii, Sov. Phys. Dokl. 11 (1966) 235
1966HA22 J.C. Hardy, R.I. Verrall and R.E. Bell, Nucl. Phys. 81 (1966) 113
1966HE05 D.W. Heikkinen, Phys. Rev. 141 (1966) 1007; Erratum Phys. Rev. 149 (1966) 990
1966HO1F Hoot, Bull. Amer. Phys. Soc. 11 (1966) 316
1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
1966LA1E Lawson, 2nd Symp. Struct. Low-Medium Mass Nuclei (1966) 6
1966LI03 D. Lister and A. Sayres, Phys. Rev. 143 (1966) 745
1966MA12 B. Margolis and N. De Takacsy, Can. J. Phys. 44 (1966) 1431
1966MA19 H.A. Mavromatis, L. Zamick and G.E. Brown, Nucl. Phys. 80 (1966) 545
1966MA1R J.H. Manley and W.E. Stein, Phys. Rev. 144 (1966) 956
1966MA1V H.A. Mavromatis and L. Zamick, Phys. Lett. 20 (1966) 171
1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
1966MC14 W.N. McDicken and W. Jack, Nucl. Phys. 88 (1966) 457
1966MC1C McPherson, F. S. U. Isobaric Spin Conf. (1966) 162

- 1966ME14 D.F. Measday and J.N. Palmieri, Nucl. Phys. 85 (1966) 129
- 1966MI14 B. Mitra and A.M. Ghose, Nucl. Phys. 83 (1966) 157
- 1966MI1F Migdal, Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch (1966) 171
- 1966MO09 F.P. Mooring, J.E. Monahan and C.M. Huddleston, Nucl. Phys. 82 (1966) 16
- 1966PO08 L. Pomorski, J. Tys and V.V. Volkov, Phys. Lett. 23 (1966) 369
- 1966PO1E Poth and Bromley, Bull. Amer. Phys. Soc. 11 (1966) 317
- 1966PR14 R. Prasad, D.C. Sarkar and C.S. Khurana, Nucl. Phys. 85 (1966) 476
- 1966QU1A C. Quesne, Nucl. Phys. 76 (1966) 268
- 1966RE1D Reeves, J. Phys. (Paris) 27 (1966) C1-28
- 1966RI04 E. Rivet, R.H. Pehl, J. Cerny and B.G. Harvey, Phys. Rev. 141 (1966) 1021
- 1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (1966) 237
- 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polarization Phenom. of Nucleons, Karlsruhe, 1965 (1966) 253
- 1966RO1R Rosen, Antwerp 1965 Neutron Conf. (1966) 379
- 1966SC09 J.P. Schiffer, L.L. Lee, Jr., A. Marinov and C. Mayer-Boricke, Phys. Rev. 147 (1966) 829
- 1966SO05 M. Soga, Nucl. Phys. 89 (1966) 697
- 1966SU01 K. Sugimoto, A. Mizobuchi, K. Nakai and K. Matuda, J. Phys. Soc. Jpn. 21 (1966) 213
- 1966TI03 Y.I. Titov, A.P. Klyucharev and V.D. Vypirailenko, Yad. Fiz. 4 (1966) 308; Sov. J. Nucl. Phys. 4 (1967) 221
- 1966WA1K G.E. Walker, Phys. Rev. 151 (1966) 745
- 1966WE04 J.J. Wesolowski, L.F. Hansen, J.G. Vidal and M.L. Stelts, Phys. Rev. 148 (1966) 1063
- 1966WI01 H.H. Williams, E.K. Warburton, K.W. Jones and J.W. Olness, Phys. Rev. 144 (1966) 801
- 1966ZA1E Zamick, Bull. Amer. Phys. Soc. 11 (1966) 56
- 1967AL06 J.L. Alty, L.L. Green, R. Huby, G.D. Jones, J.R. Mines and J.F. Sharpey-Schafer, Nucl. Phys. A97 (1967) 541
- 1967AM1H Amiet, Ebenhoh and Huguenin, Helv. Phys. Acta 40 (1967) 283
- 1967AN1B Anderson and Bevington, Bull. Amer. Phys. Soc. 12 (1967) 1144
- 1967AU1A Audouze, Epherre and Reeves, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 255
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967BA1R J.M. Bang, C.A. Pearson and L. Pocs, Nucl. Phys. A100 (1967) 24

1967BA2E J.M. Bang and C.A. Pearson, Nucl. Phys. A100 (1967) 1
1967BE02 W.P. Beres and W. MacDonald, Nucl. Phys. A91 (1967) 529
1967BE1F Beery, Harper, Stovall and Rosen, LA-3788 (1967)
1967BE22 V.V. Beloshitsky, Phys. Lett. B24 (1967) 559
1967BE2D Bernas, Gradsztajn, Reeves and Schatzman, Ann. Phys. 44 (1967) 426
1967BE75 P.L. Beach, R.W. Finlay, R.L. Cassola, and R.D. Koshel, Phys. Rev. 156 (1967) 1201
1967BI05 P.G. Bizzeti, A.M. Bizzeti-Sona, S. Kalbitzer and B. Povh, Z. Phys. 201 (1967) 295
1967BO1T Bohigas, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 940
1967BU16 W. Busse, B. Efken, D. Hilscher, J.A. Scheer and W.U. Schroder, Z. Phys. 206 (1967) 404
1967CA1D Carlson, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 475
1967CA1J Cabrespine, Gauvin, Lefort and Sauvage, Ark. Fys. 36 (1967) 463
1967CA1Q Cameron, Univ. of California, Los Angeles, Dept. Phys. Tech. Rept. P80 (1967)
1967CA1R Carlson et al, Bull. Amer. Phys. Soc. 12 (1967) 1190
1967CH15 N.S. Chant, P.S. Fisher and D.K. Scott, Nucl. Phys. A99 (1967) 669
1967CH41 R. Chapman and A.M. Macleod, Nucl. Phys. A94 (1967) 313
1967CH42 A. Chatterjee and A.M. Ghose, Phys. Rev. 161 (1967) 1181
1967CO05 R.K. Cole, R. Dittman, H.S. Sandhu, C.N. Waddell and J.K. Dickens, Nucl. Phys. A91 (1967) 665
1967CO1D Commins, Ann. Rev. Nucl. Sci. 17 (1967) 33
1967DI1C Dittman, Cole, Waddell and Sandhu, Bull. Amer. Phys. Soc. 12 (1967) 663
1967DZ01 T.G. Dzubay, Phys. Rev. 158 (1967) 977
1967EL03 J.P. Elliott, H.A. Mavromatis and E.A. Sanderson, Phys. Lett. B24 (1967) 358
1967EN01 T. Engeland and P.J. Ellis, Phys. Lett. B25 (1967) 57
1967EP1A Epherre and Gradsztajn, J. Phys. (France) 28 (1967) 745
1967ES02 R.A. Esterlund, R. McPherson, A.M. Poskanzer and P.L. Reeder, Phys. Rev. 156 (1967) 1094
1967FA06 J.A. Fannon, E.J. Burge, D.A. Smith and N.K. Ganguly, Nucl. Phys. A97 (1967) 263
1967FE01 P. Federman, Nucl. Phys. A95 (1967) 443
1967FI10 R.W. Fink, T.H. Braid and A.W. Friedman, Ark. Fys. 36 (1967) 471
1967GO04 A. Goswami and A.I. Sherwood, Nucl. Phys. A91 (1967) 64

- 1967GO1A G. Goldring, H.M. Loebenstein , I. Plesser and M.W. Sachs, Phys. Lett. B25 (1967) 538
- 1967GR1D L. Grunbaum and N.K. Ganguly, Nucl. Phys. A100 (1967) 645
- 1967GR1H Green and Berezdivin, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 600
- 1967GR1J A.E. Green, G. Darewych and R. Berezdivin, Phys. Rev. 157 (1967) 929
- 1967GR1K Gradsztajn, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 247
- 1967HA20 R.L. Hahn and E. Ricci, Nucl. Phys. A101 (1967) 353
- 1967HO14 R.A. Hoffswell, D. Jamnik, T.M. Noweir and A.I. Yavin, Phys. Rev. Lett. 19 (1967) 754
- 1967JO12 C.H. Johnson and J.L. Fowler, Phys. Rev. 162 (1967) 890
- 1967KA1G Kawai and Yazaki, Prog. Theor. Phys. 37 (1967) 638
- 1967KA21 S.P. Kamerdzhiev, Yad. Fiz. 5 (1967) 971; Sov. J. Nucl. Phys. 5 (1967) 693
- 1967LA1M B. Lawergren, Phys. Lett. B24 (1967) 395
- 1967LE13 H.K. Lee and H. McManus, Phys. Rev. 161 (1967) 1087
- 1967LI1B Lingenfelter and Ramaty, High Energy Nucl. Reactions in Astrophys.; Ed., B.S.P. Shen (1967) 99
- 1967LU05 H.F. Lutz, J.J. Wesolowski, S.F. Eccles and L.F. Hansen, Nucl. Phys. A101 (1967) 241
- 1967LU1B Lundberg, Schwarz and Zetterstrom, Ark. Fys. 34 (1967) 247
- 1967LY02 R.P. Lynch and T.T.S. Kuo, Nucl. Phys. A95 (1967) 561
- 1967MO23 J.F. Morgan and R.K. Hobbie, Phys. Rev. 163 (1967) 992
- 1967NE1D Nemirovskii, Sov. J. Nucl. Phys. 4 (1967) 334
- 1967NI1D W.K. Niblack and B.P. Nigam, Phys. Rev. 156 (1967) 1191
- 1967OG1A Ogloblin, Proc. Problem Symp. on Nucl. Phys., Tbilisi, 1967 (1967) 169
- 1967PA05 J.C. Parikh and N. Ullah, Nucl. Phys. A99 (1967) 529
- 1967PA17 J.R. Patterson, H. Winkler and C.S. Zaidins, Phys. Rev. 163 (1967) 1051
- 1967PF01 A. Pfitzner, Nucl. Phys. A100 (1967) 673
- 1967PI01 R.E. Pixley and W. Benenson, Nucl. Phys. A91 (1967) 177
- 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, Phys. Rev. 164 (1967) 1295
- 1967PO1E Poth and Bromley, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 94
- 1967RE1C Reynolds and Francis, Bull. Amer. Phys. Soc. 12 (1967) 516
- 1967SC16 H. Schulz, H.J. Wiebicke and R. Reif, Nucl. Phys. A101 (1967) 577
- 1967SC1E Schenter, Bull. Amer. Phys. Soc. 12 (1967) 11

- 1967SC1H Schenter, Bull. Amer. Phys. Soc. 12 (1967) 895
- 1967SE07 K.K. Sekharan, A.S. Divatia, M.K. Mehta, S.S. Kerekatte and K.B. Nambiar, Phys. Rev. 156 (1967) 1187
- 1967SH05 Y.Y. Sharon, Nucl. Phys. A99 (1967) 321
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SH1J Shanley and Aaron, Ann. Phys. 44 (1967) 363
- 1967SH1K Shukla, PUC 937 262 (1967)
- 1967SP09 A. Sperduto, Proc. 3rd Int. Conf. on At. Masses, Winnipeg, Canada, 1967 (1967) 657
- 1967TR08 W. Trachslin and L. Brown, Nucl. Phys. A101 (1967) 273
- 1967UN1A H.-J. Unger, Nucl. Phys. A104 (1967) 564
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
- 1967VA1H van Bree and Temmer, Bull. Amer. Phys. Soc. 12 (1967) 518
- 1967VO1A Volkov, Proc. Problem Symp. on Nucl. Phys., Tbilisi, 1967 (1967) 226
- 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators; Eds., Marion and van Patter (1967) 43
- 1968AL1G Alsmiller and Barish, Rept. NASA-CR-96371 (1968)
- 1968AN1F Antolkovic et al, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1658
- 1968AP1A Appel et al, Bull. Amer. Phys. Soc. 13 (1968) 680
- 1968AR1F Artyukh, Pomurski, Tys and Volkov, in Tokyo (1968) 214
- 1968BE1X G.F. Bertsch, Phys. Lett. B28 (1968) 302
- 1968BE1Y Bethge, Pullen and Middleton, Bull. Amer. Phys. Soc. 18 (1968) 1464
- 1968BI07 J. Birkholz and F. Beck, Phys. Lett. B28 (1968) 18
- 1968BI09 M. Bister, A. Fontell and P. Vikberg, Ann. Acad. Sci. Fennicae, Ser. A VI, No. 268 (1968) 2
- 1968CA30 J.M. Cameron, J.R. Richardson, W.T. van Oers and J.W. Verba, Phys. Rev. 167 (1968) 908
- 1968CE01 J. Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968CH35 J.S. Chalmers and A.M. Saperstein, Phys. Rev. 168 (1968) 1145; Addendum Phys. Rev. C2 (1970) 2034
- 1968CO1U Cole et al, in Tokyo (1968) 283
- 1968DA05 C.N. Davids, Nucl. Phys. A110 (1968) 619
- 1968DA1D Davids, Astrophys. J. 151 (1968) 775
- 1968DA1E Dandy, Wankling and Parnell, AWRE 060/68 (1968)

- 1968DA1F J.C. Davis and H.H. Barschall, Phys. Lett. B27 (1968) 636
- 1968DE13 N. de Takacsy, Can. J. Phys. 46 (1968) 2091
- 1968DI06 O. Dietzsch, R.A. Douglas, E.F. Pessoa, V.G. Porto, E.W. Hamburger, T. Polga, O. Sala, S.M. Perez and P.E. Hodgson, Nucl. Phys. A114 (1968) 330
- 1968DO20 I. Dostrovsky, H. Gauvin and M. Lefort, Phys. Rev. 169 (1968) 836
- 1968EL12 H.B. Eldridge, S.N. Bunker, J.M. Cameron, J.R. Richardson and W.T.H. van Oers, Phys. Rev. 167 (1968) 915
- 1968EL1A Elliott, 3rd Symp. Struct. Low-Medium Mass Nuclei (1968) 48
- 1968EL1C Elliott and Wilsdon, Proc. Roy. Soc. 302 (1968) 509
- 1968EL1G M. El-Nady and M.A. Sharaf, Phys. Lett. B28 (1968) 165
- 1968FA1G Falk, Kidney, Tandon and Kulisic, Bull. Amer. Phys. Soc. 13 (1968) 1464
- 1968FO1D Fowler, Bull. Amer. Phys. Soc. 13 (1968) 1462
- 1968FOZY D.T. Fowler, Neutron Cross Sections Tech., NBS Special Pub. 299, Vol. I (1968) 1
- 1968GA1J E. Gadioli, I. Iori and L. Zetta, Phys. Rev. 174 (1968) 1140
- 1968GO1N Gofman et al, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 690
- 1968HA30 L.F. Hansen, M.L. Stelts, J.G. Vidal, J.J. Wesolowski and V.A. Madsen, Phys. Rev. 174 (1968) 1155
- 1968HE1H K.T. Hecht, P.J. Ellis and T. Engeland, Phys. Lett. B27 (1968) 479
- 1968HO1H Horie and Hsieh, in Tokyo (1968) 51
- 1968HO23 K. Hosono, J. Phys. Soc. Jpn. 25 (1968) 36
- 1968JA1M Jacquot et al, Compt. Rend. B266 (1968) 1286
- 1968JO1F Johnson et al, Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 851
- 1968JO1G Jones, Chase and Nightingale, Bull. Amer. Phys. Soc. 13 (1968) 651
- 1968KA1E Kawai and Yazaki, in Tokyo (1968) 238
- 1968KE02 G.W. Kerr, J.M. Morris and J.R. Risser, Nucl. Phys. A110 (1968) 637
- 1968KN1A Knoth et al, Helv. Phys. Acta 41 (1968) 1278
- 1968KU1C Kuhlmann, BMWF FBK 68 06 (1968)
- 1968LE1L R. Leonardi and M. Rosa-Clot, Phys. Rev. Lett. 21 (1968) 377
- 1968LE1N Lehman, Int. J. Appl. Rad. Isotopes 19 (1968) 841
- 1968MA2B Manakos, Z. Phys. 214 (1968) 57
- 1968MA2D Manero, An. Real Soc. Espan. Fis. Quim. A64 (1968) 63
- 1968ME10 K. Meier-Ewert, K. Bethge and K.-O. Pfeiffer, Nucl. Phys. A110 (1968) 142
- 1968ME15 M.M. Meier, R.S. Thomason and R.L. Walter, Nucl. Phys. A115 (1968) 540

- 1968MI05 P.F. Mizera and J.B. Gerhart, Phys. Rev. 170 (1968) 839
- 1968NA06 I.M. Naqib and L.L. Green, Nucl. Phys. A112 (1968) 76
- 1968NG1B Nguyen Van Sen et al, J. Phys. 29 (France) (1968) 15
- 1968NI1A W.K. Niblack and B.P. Nigam, Phys. Rev. 167 (1968) 996
- 1968OG1A Ogloblin, Nucl. Struct. Dubna Symp. 1968, IAEA (1968) 204
- 1968PA1J Palevsky, Proc. of Symp. on Use of Nimrod, 1968, RHEL-R166 (1968) 19
- 1968PA1X Parnell, Dandy and Wankling, Phys. Med. Bio. 13 (1968) 665
- 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8 (1969) 270
- 1968PE1E D.C. Peaslee, Phys. Rev. 171 (1968) 1208
- 1968PO04 A.M. Poskanzer, G.W. Butler, E.K. Hyde, J. Cerny, D.A. Landis and F.S. Goulding, Phys. Lett. B27 (1968) 414
- 1968PR04 J.R. Priest and J.S. Vincent, Phys. Rev. 167 (1968) 933
- 1968PR1C Proctor, Benenson, Paddock and Locard, Bull. Amer. Phys. Soc. 13 (1968) 1464
- 1968RE03 R. Reif, J. Slotta and J. Hohn, Phys. Lett. B26 (1968) 484
- 1968RE07 D. Rendic, B. Antolkovic, G. Paic, M. Turk and P. Tomas, Nucl. Phys. A117 (1968) 113
- 1968RI1P Riess et al, in Tokyo (1968) 141
- 1968RO1E Rohl, Z. Phys. 215 (1968) 56
- 1968SC1E Schwarzschild and Warburton, Ann. Rev. Nucl. Sci. 18 (1968) 265
- 1968SE1C Seth, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, 1967 (1968) 179
- 1968SE1D Sellin, Beres, Bilpuch and Newson, Bull. Amer. Phys. Soc. 13 (1968) 651
- 1968SI06 I. Sick, E. Baumgartner, P. Huber and T. Stammbach, Helv. Phys. Acta 41 (1968) 573
- 1968SU1C K. Sugawara, Nucl. Phys. A110 (1968) 305
- 1968SU1E Sukhoruchkin, Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 923
- 1968SU1F Suekane, in Tokyo (1968) 57
- 1968TA02 Y. Takeuchi, Y. Hiratate, K. Miura, T. Tohei and S. Morita, Nucl. Phys. A109 (1968) 105
- 1968TA1N Takeuchi and Sanada, TUENS-4 (1968)
- 1968TE1C Temmer, Teitelman, van Bree and Ogata, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 299
- 1968WO1C S.S.M. Wong, Nucl. Phys. A120 (1968) 625

- 1968YI02 F. Yiou, M. Baril, J. Dufaure de Citres, P. Fontes, E. Gradsztajn and R. Bernas, Phys. Rev. 166 (1968) 968
- 1968ZU02 A.P. Zuker, B. Buck and J.B. McGrory, Phys. Rev. Lett. 21 (1968) 39
- 1969AB1A Abdinov and Barashenkov, Joint Inst. Nucl. Res., Lab. Theor. Phys., USSR, Rept. P2 4788 (1969)
- 1969AD02 E.G. Adelberger, A.B. McDonald and C.A. Barnes, Nucl. Phys. A124 (1969) 49
- 1969AN06 J.D. Anderson, S.D. Bloom, C. Wong, W.F. Hornyak and V.A. Madsen, Phys. Rev. 177 (1969) 1416
- 1969AN1L Anderson, Thesis, Stanford Univ. (1969)
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969BA17 K. Bahr, T. Becker, R. Jahr and W.R. Kuhlmann, Nucl. Phys. A129 (1969) 388
- 1969BA1P Bacon et al, Acta Cryst. A25 (1969) 391
- 1969BA1Z Barnes, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 179
- 1969BA2E A.I. Baz and V.I. Manko, Phys. Lett. B28 (1969) 541
- 1969BA2T Baldeweg, Beckert and Schneidereit, Zent. Kernf. Ross. Dresden, Rept. No. ZFK 184 (1969)
- 1969BA2U Ballini, Delaunay, Fouan and Tellez, Addendum to Contrib., Montreal (1969) 6
- 1969BO2A Boschitz, Rept. NASA-TN-D-5067 (1969)
- 1969BO37 J. Bobker, Phys. Rev. 185 (1969) 1294
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (1969) 242
- 1969BR1F Bruninx and Crombeen, Int. J. Appl. Rad. Isotopes 20 (1969) 255
- 1969BU08 E.M. Burymov, Yad. Fiz. 9 (1969) 933; Sov. J. Nucl. Phys. 9 (1969) 546
- 1969BU19 F.W. Busser, H. Dubenkropp, F. Niebergall and K. Sinram, Nucl. Phys. A129 (1969) 666
- 1969BU1J Bunker et al, Bull. Amer. Phys. Soc. 14 (1969) 529
- 1969CH1R M. Chemtob, Nucl. Phys. A123 (1969) 449
- 1969CO12 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A134 (1969) 561
- 1969DA13 J.C. Davis and F.T. Noda, Nucl. Phys. A134 (1969) 361
- 1969DA1L Daehnick, Bull. Amer. Phys. Soc. 14 (1969) 102; Private Communication
- 1969DE06 C. Detraz and H.H. Duhm, Phys. Lett. B29 (1969) 29
- 1969DE16 J. Dey, J.P. Elliott, A.D. Jackson, H.A. Mavromatis, E.A. Sanderson and B. Singh, Nucl. Phys. A134 (1969) 385

- 1969DU10 V.E. Dudkin, V.N. Kuzmin, L.N. Smirenniyi, N.S. Shimanskaya and R.M. Yakovlev, *Yad. Fiz.* 9 (1969) 925; *Sov. J. Nucl. Phys.* 9 (1969) 541
- 1969EL1B Elliott, *Proc. Int. Conf., Montreal* (1969) 277
- 1969EL1D Ellis and Engeland, *Contrib., Montreal* (1969) 344
- 1969EP1B M. Epherre, E. Gradsztajn, R. Klapisch and H. Reeves, *Nucl. Phys.* A139 (1969) 545
- 1969FO1G Fowler, *Contrib., Montreal* (1969) 334
- 1969FU11 G.H. Fuller and V.W. Cohen, *Nucl. Data Tables* A5 (1969) 433
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, *Phys. Rev.* 179 (1969) 921
- 1969GA1P Garvey, *Nucl. Isospin, Proc. 1969 Asilomar Conf.* (1969) 703
- 1969GI1B Gillet, *Proc. Int. Conf., Montreal* (1969) 483
- 1969GO04 S. Gorodetzky, J.C. Merdinger, N. Schulz and A. Knipper, *Nucl. Phys.* A129 (1969) 129
- 1969GU1M V.N. Guman, L.A. Sliv and Yu.I. Kharitonov, *Phys. Lett.* B28 (1969) 575
- 1969HA1G Hanna, *Proc. Int. Conf., Montreal* (1969) 443
- 1969HA1U Hansen and Stelts, *Nucl. Isospin, Proc. 1969 Asilomar Conf.* (1969) 269
- 1969HA1V K.A. Hamza and S. Edwards, *Phys. Rev.* 181 (1969) 1494
- 1969HA2C J. Habanec, *Phys. Lett.* B30 (1969) 402
- 1969HAZD R.C. Haight, *Thesis, Princeton Univ.* (1969)
- 1969HAZE J.C. Hardy, J.E. Esterl, R.G. Sextro and J. Cerny, *Nucl. Isospin, Proc. 1969 Conf.* (1969) 725
- 1969HE1N E.M. Henley and C.E. Lacy, *Phys. Rev.* 184 (1969) 1228
- 1969HO1X Hoot et al, *Bull. Amer. Phys. Soc.* 14 (1969) 494
- 1969HU1H Huber, Cole, Waddell and Spitzer, *Bull. Amer. Phys. Soc.* 14 (1969) 1168
- 1969IC02 M. Ichimura, M. Kawai, T. Ohmura and B. Imanishi, *Phys. Lett.* B30 (1969) 143
- 1969IG1A Igarashi, Kawai and Yazaki, *Prog. Theor. Phys.* 42 (1969) 245
- 1969IW1A Iwao, *Prog. Theor. Phys.* 42 (1969) 1304
- 1969IW1B Iwao, *DPKU-027-68* (1969)
- 1969IW1C Iwao, *DPKU-026-68* (1969)
- 1969JA1P R.L. Jaffe and W.J. Gerace, *Nucl. Phys.* A125 (1969) 1
- 1969JO1L Jones, Chase and Nightingale, *Bull. Amer. Phys. Soc.* 14 (1969) 1221
- 1969KA09 S. Kahana, H.C. Lee and C.K. Scott, *Phys. Rev.* 180 (1969) 956
- 1969KA14 O. Karban, P.D. Greaves, V. Hnizdo, J. Lowe, N. Berovic, H. Wojciechowski and G.W. Greenlees, *Nucl. Phys.* A132 (1969) 548

- 1969KA1D Karge, *Wiss. Z. Friedrich. Schiller Univ. Jena. Math. Naturwiss. Reihe (Germany)* 18 (1969) 51
- 1969KH1C Khanna and Harvey, *Bull. Amer. Phys. Soc.* 14 (1969) 604
- 1969KU1G Kulkarni and Pandya, *Nuovo Cim.* B60 (1969) 199
- 1969LE1D Leonardi and Rosa-Clot, *Lett. Nuovo Cim.* 1 (1969) 829
- 1969LI22 Y.-C. Liu, W.-S. Hou and C. Chang, *Chin. J. Phys. (Taiwan)* 7 (1969) 63
- 1969LU07 C.C. Lu, M.S. Zisman and B.G. Harvey, *Phys. Rev.* 186 (1969) 1086
- 1969MA1G Madsen, *Nucl. Isospin, Proc. 1969 Asilomar Conf.* (1969) 149
- 1969MA2B MacKellar, Reading and Kerman, *Bull. Amer. Phys. Soc.* 14 (1969) 572
- 1969MA2C MacFarlane, *Proc. Int. Conf., Montreal* (1969) 385
- 1969MA38 H.A. Mavromatis and B. Singh, *Nucl. Phys.* A139 (1969) 451
- 1969MC1A McGill, *UCLA P-83* (1969)
- 1969MC1G B.H. McKellar, *Phys. Rev.* 181 (1969) 1502
- 1969ME07 L. Mesko, B. Schlenk, G. Somogyi and A. Valek, *Nucl. Phys.* A130 (1969) 449
- 1969ME15 D. Meier, M. Brullmann, H. Jung and P. Marmier, *Helv. Phys. Acta* 42 (1969) 813
- 1969ME1K Mermaz, Lemaire, Scheurer and Seth, *Addendum to Contrib., Montreal* (1969) 10
- 1969NI09 R.J. Nickles, *Nucl. Phys.* A134 (1969) 308
- 1969NO03 J.A. Nolen, Jr. and J.P. Schiffer, *Phys. Lett.* B29 (1969) 396
- 1969NO1E Nolen and Schiffer, *Ann. Rev. Nucl. Sci.* 19 (1969) 471
- 1969NY1A Nyberg, Jonsson and Bergqvist, *NP 6902, Research Inst. Nat. Defence, Stockholm* (1969)
- 1969OL02 C.J. Oliver, P.D. Forsyth, J.L. Hutton, G. Kaye and J.R. Mines, *Nucl. Phys.* A127 (1969) 567
- 1969OV01 J.C. Overley, P.D. Parker and D.A. Bromley, *Nucl. Instrum. Meth.* 68 (1969) 61
- 1969PA1G Parikh, *Contrib., Montreal* (1969) 212
- 1969PE1K C.A. Pearson, J.M. Bang and L. Pocs, *Phys. Rev.* 179 (1969) 1082
- 1969PE1L Pearson, Bang and Pocs, *Ann. Phys.* 52 (1969) 83
- 1969PI1D Pinkston, Iano and Satchler, *Bull. Amer. Phys. Soc.* 14 (1969) 573
- 1969RO07 F.A. Rose, *Nucl. Phys.* A124 (1969) 305
- 1969SA1J Saunier and Pearson, *Bull. Amer. Phys. Soc.* 14 (1969) 36
- 1969SC04 H. Scholermann, *Z. Phys.* 220 (1969) 211
- 1969SC14 J.P. Schiffer, J.A. Nolen, Jr. and N. Williams, *Phys. Lett.* B29 (1969) 399
- 1969SC1H Schmittroth, *Thesis, Oregon State Univ.* (1969)

1969SC1Q Schiffer, Nucl. Isospin, Proc. 1969 Asilomar Conf. (1969) 733
 1969SC21 W.A. Schier and J.D. Reber, Phys. Rev. 181 (1969) 1371
 1969SI1E S.G. Siegel and L. Zamick, Phys. Rev. 184 (1969) 1230
 1969SK1B Skwiersky, Chesterfield and Parker, Contrib., Montreal (1969) 317
 1969SO1A Soroka and Pucherov, Ukr. Fiz. Zh. 14 (1969) 688
 1969SU15 K. Sugimoto, Phys. Rev. 182 (1969) 1051
 1969TE1A Temmer, Isospin in Nucl. Phys.; Ed., D.H. Wilkinson (1969) 693
 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
 1969TH04 S.T. Thornton, Nucl. Phys. A137 (1969) 531
 1969UL03 N. Ullah and S.S.M. Wong, Phys. Rev. 188 (1969) 1645
 1969VA18 W.T.H. van Oers and J.M. Cameron, Phys. Rev. 184 (1969) 1061
 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
 1969ZU1D Zuker and Buck, Contrib., Montreal (1969) 193
 1970AB1D Abraham, Gray and Legg, Bull. Amer. Phys. Soc. 15 (1970) 1685
 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
 1970AU1C Austin et al, Private Communication (1970)
 1970BA1J Barker et al, Nucl. Reactions induced by Heavy Ions, Heidelberg, 1969 (1970) 152
 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119
 1970BA2B Baz and Manko, Sov. J. Nucl. Phys. 10 (1970) 46
 1970BA55 P.H. Barker, P.M. Cockburn, H.P. Seiler and P. Marmier, Phys. Rev. Lett. 25 (1970) 1350
 1970BE21 G. Bertsch and A. Molinari, Nucl. Phys. A148 (1970) 87
 1970BE31 K. Bethge, D.J. Pullen and R. Middleton, Phys. Rev. C2 (1970) 395
 1970BE68 M.G. Betigeri, C.M. Lamba, N. Sarma, D.K. Sood and N.S. Thampi, Energ. Nucl. (Milan) 17 (1970) 256
 1970BI1D Bigler et al, Bull. Amer. Phys. Soc. 15 (1970) 1685
 1970BL03 A.G. Blair, C. Glashauser, R. de Swiniarski, J. Goudergues, R. Lombard, B. Mayer, J. Thirion and P. Vaganov, Phys. Rev. C1 (1970) 444
 1970BO1K Borbei and Dolinski, Sov. J. Nucl. Phys. 10 (1970) 173
 1970BO2B J. Bobker, Phys. Rev. C2 (1970) 322
 1970BO30 F. Boreli, Fizika 2 (1970) 97
 1970BU16 V.E. Bunakov, K.A. Gridnev and L.V. Krasnov, Phys. Lett. B32 (1970) 587
 1970CA1N Carlson and Johnson, Bull. Amer. Phys. Soc. 15 (1970) 629

- 1970CO1P Corrigan, Prior and Darden, Bull. Amer. Phys. Soc. 15 (1970) 483
- 1970CO1U M. Coz, A.D. MacKellar and L.Z. Arnold, Ann. Phys. 58 (1970) 504
- 1970COZA E. Cotton, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 289
- 1970DA14 N.E. Davison, W.K. Dawson, G. Roy and W.J. McDonald, Can. J. Phys. 48 (1970) 2235
- 1970DI1C Dickens and Perey, Nucl. Sci. Eng. 40 (1970) 283
- 1970DU07 J.L. Duggan, J.Y. Park, S.D. Danielopoulos, P.D. Miller, J. Lin, M.M. Duncan and R.L. Dangle, Nucl. Phys. A151 (1970) 107
- 1970EL08 P.J. Ellis and S. Siegel, Nucl. Phys. A152 (1970) 547
- 1970EL1G Ellis and Engeland, Nucl. Phys. A144 (1970) 161
- 1970FO03 J.L. Fowler and C.H. Johnson, Phys. Rev. C2 (1970) 124
- 1970GA1A Garber et al, BNL 400, 3rd Edition, Vol. 1 (1970)
- 1970GO1H P. Goode and S. Siegel, Phys. Lett. B31 (1970) 418
- 1970GU04 P. Guazzoni, S. Micheletti and M. Pignanelli, Lett. Nuovo Cim. 3 (1970) 639
- 1970GU1E Guman, Sliv and Kharitonov, Sov. J. Nucl. Phys. 10 (1970) 302
- 1970HA1T Harvey and Khanna, Nucl. Phys. A152 (1970) 588
- 1970HA23 J.C. Hardy, A.D. Bacher, G.R. Plattner, J.A. Macdonald and R.G. Sextro, Phys. Rev. Lett. 25 (1970) 298
- 1970HI15 M. Hirata, Prog. Theor. Phys. 43 (1970) 1526
- 1970HO08 J.L. Honsaker, T.H. Hsu, W.J. McDonald and G.C. Neilson, Nucl. Phys. A144 (1970) 473
- 1970HU1J Huber, Cole, Waddell and Spitzer, Bull. Amer. Phys. Soc. 15 (1970) 767
- 1970JO09 D.J. Johnson and M.A. Waggoner, Phys. Rev. C2 (1970) 41
- 1970KE1D Keyser, Blue, Ramirez and Weller, Bull. Amer. Phys. Soc. 15 (1970) 1685
- 1970KN01 A.R. Knudson and F.C. Young, Nucl. Phys. A149 (1970) 323
- 1970KO25 V.I. Komarov, G.E. Kosarev and O.V. Savchenko, Yad. Fiz. 11 (1970) 711; Sov. J. Nucl. Phys. 11 (1970) 399
- 1970LO01 G. Lodin and L. Nilsson, Z. Phys. 233 (1970) 181
- 1970MC02 A.B. McDonald, E.G. Adelberger, H.B. Mak, D. Ashery, A.P. Shukla, C.L. Cocke and C.N. Davids, Phys. Lett. B31 (1970) 119
- 1970MC17 B.H.J. McKellar, Phys. Lett. B32 (1970) 246
- 1970ME11 R. Mendelson, G.J. Wozniak, A.D. Bacher, J.M. Loiseaux and J. Cerny, Phys. Rev. Lett. 25 (1970) 533
- 1970ME31 H.O. Menlove, R.H. Augustson, C.N. Henry, Nucl. Sci. Eng. 40 (1970) 136

1970MU1C Munro, Bull. Amer. Phys. Soc. 15 (1970) 193
1970MU1H Murphy and Ritter, Bull. Amer. Phys. Soc. 15 (1970) 483
1970NE1B J.M. Nelson, N.S. Chant and P.S. Fisher, Phys. Lett. B31 (1970) 445
1970OG1A Ogloblin, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (1970) 231
1970OH06 T. Ohmura, B. Imanishi, M. Ichimura and M. Kawai, Prog. Theor. Phys. 43 (1970) 347
1970PE14 C.A. Pearson, D. Zissermann and J.M. Covan, Nucl. Phys. A152 (1970) 449
1970PE1B Pearson, Rickel and Zissermann, Nucl. Phys. A148 (1970) 273
1970RO08 A.D. Robb, W.A. Schier and E. Sheldon, Nucl. Phys. A147 (1970) 423
1970RY02 V.F. Rybachenko and A.A. Sadovoi, Yad. Fiz. 11 (1970) 1192; Sov. J. Nucl. Phys. 11 (1970) 663
1970SH14 R. Shanta and R.K. Satpathy, Phys. Rev. C2 (1970) 1279
1970SI02 R.P. Singhal, J.R. Moreira and H.S. Caplan, Phys. Rev. Lett. 24 (1970) 73
1970SI1C Siegel and Zamick, Nucl. Phys. A145 (1970) 89
1970ST04 H.J. Strubbe and D.K. Callebaut, Nucl. Phys. A143 (1970) 537
1970SU01 B.I. Sukhanov and N.P. Tkach, Yad. Fiz. 11 (1970) 33; Sov. J. Nucl. Phys. 11 (1970) 17
1970SU1B Sukhoruchkin, Sov. J. Nucl. Phys. 10 (1970) 144
1970TE1B Tepel, Z. Phys. 234 (1970) 107
1970TI1C Tishchenko and Kresnin, Sov. J. Nucl. Phys. 10 (1970) 442
1970VI03 C.M. Vincent and H.T. Fortune, Phys. Rev. C2 (1970) 782
1970WA01 G.E. Walker and D. Schlobohm, Nucl. Phys. A140 (1970) 49
1970WA1J W.L. Wang and C.M. Shakin, Phys. Lett. B32 (1970) 421
1970ZE01 N.S. Zelenskaya, N.V. Karabanov, A.V. Spassky, K.G. Tarov, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 11 (1970) 722; Sov. J. Nucl. Phys. 11 (1970) 405
1971BA06 W.L. Baker, C.E. Busch, J.A. Keane and T.R. Donoghue, Phys. Rev. C3 (1971) 494
1971HA05 J.C. Hardy, J.E. Esterl, R.G. Sextro and J. Cerny, Phys. Rev. C3 (1971) 700
1971TO08 I.S. Towner, E.K. Warburton and G.T. Garvey, Ann. Phys. 66 (1971) 674
1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1

